

# UC Riverside

## UC Riverside Previously Published Works

### Title

Tris(1,3-dichloro-2-propyl)phosphate Induces Genome-Wide Hypomethylation within Early Zebrafish Embryos

### Permalink

<https://escholarship.org/uc/item/4q52835v>

### Journal

Environmental Science and Technology, 50(18)

### ISSN

0013-936X

### Authors

Volz, David C  
Leet, Jessica K  
Chen, Albert  
et al.

### Publication Date

2016-09-20

### DOI

10.1021/acs.est.6b03656

Peer reviewed

## Tris(1,3-dichloro-2-propyl)phosphate Induces Genome-Wide Hypomethylation within Early Zebrafish Embryos

David C. Volz,<sup>\*,†</sup> Jessica K. Leet,<sup>‡,#</sup> Albert Chen,<sup>§</sup> Heather M. Stapleton,<sup>§</sup> Neerja Katiyar,<sup>||</sup> Rakesh Kaundal,<sup>||</sup> Yang Yu,<sup>⊥</sup> and Yinsheng Wang<sup>⊥</sup>

<sup>†</sup>Department of Environmental Sciences, University of California, Riverside, Riverside, California 92521, United States

<sup>‡</sup>Department of Environmental Health Sciences, University of South Carolina, Columbia, South Carolina 29208, United States


<sup>§</sup>Division of Environmental Sciences and Policy, Duke University, Durham, North Carolina 27708, United States

<sup>||</sup>Bioinformatics Facility, Institute for Integrative Genome Biology, University of California, Riverside, Riverside, California 92521, United States

<sup>⊥</sup>Department of Chemistry, University of California, Riverside, Riverside, California 92521, United States

### Supporting Information

**ABSTRACT:** Tris(1,3-dichloro-2-propyl)phosphate (TDCIPP) is a high-production volume organophosphate-based plasticizer and flame retardant widely used within the United States. Using zebrafish as a model, the objectives of this study were to determine whether (1) TDCIPP inhibits DNA methyltransferase (DNMT) within embryonic nuclear extracts; (2) uptake of TDCIPP from 0.75 h postfertilization (hpf, 2-cell) to 2 hpf (64-cell) or 6 hpf (shield stage) leads to impacts on the early embryonic DNA methylome; and (3) TDCIPP-induced impacts on cytosine methylation are localized to CpG islands within intergenic regions. Within this study, 5-azacytidine (5-azaC, a DNMT inhibitor) was used as a positive control. Although 5-azaC significantly inhibited zebrafish DNMT, TDCIPP did not affect DNMT activity *in vitro* at concentrations as high as 500  $\mu$ M. However, rapid embryonic uptake of 5-azaC and TDCIPP from 0.75 to 2 hpf resulted in chemical- and chromosome-specific alterations in cytosine methylation at 2 hpf. Moreover, TDCIPP exposure predominantly resulted in hypomethylation of positions outside of CpG islands and within intragenic (exon) regions of the zebrafish genome. Overall, these findings provide the foundation for monitoring DNA methylation dynamics within zebrafish as well as identifying potential associations among TDCIPP exposure, adverse health outcomes, and DNA methylation status within human populations.



## INTRODUCTION

Early embryonic development is a dynamic, complex process that is dependent on the timing and extent of epigenetic reprogramming of the zygotic genome. Immediately following fertilization, early embryogenesis progresses through two phases of the maternal-to-zygotic transition (MZT): (1) rapid degradation of maternally loaded transcripts and (2) minor and major waves of zygotic genome activation. The MZT is highly conserved across a wide range of invertebrate and vertebrate species, including sea urchins, nematodes, fruit flies, zebrafish, frogs, mice, and humans.<sup>1,2</sup> In humans, the MZT occurs *in utero* at the 8-cell stage (3 d postfertilization) prior to implantation within the uterine wall<sup>2</sup> whereas, in zebrafish, the MZT commences *ex utero* near the end of cleavage (2 h postfertilization, hpf) and terminates at approximately midblastula (2.75–3 hpf).<sup>3</sup> After rapid erasure of maternal and paternal methylation marks following fertilization, zygotic genome activation, tissue-specific gene expression, and normal somatic development of mammalian and zebrafish embryos are dependent on steady, *de novo* genome-wide cytosine methylation by DNA methyltransferases (DNMT) during the MZT.<sup>4,5</sup> Therefore, alterations

in *de novo* DNA methylation marks, via impacts on DNMT activity or methyl donor concentrations, during early embryonic development have the potential to change the trajectory of normal development and, if inherited, persist across multiple generations.<sup>6</sup>

Over the last 10–15 years, the role of environmental factors, including nutrition, chemical exposures, and physical variables such as temperature, in mediating epigenetic modifications during early development has been intensely investigated, albeit the number of environment-related epigenetic studies pales in comparison to studies considering nonenvironmental factors.<sup>7</sup> To date, the most well-studied environmental chemical known to alter DNA methylation and, as a result, offspring phenotypes is bisphenol A (BPA),<sup>8–13</sup> a high-production volume chemical used in the production of polycarbonate plastics and epoxy resins. In a seminal study published by Dolinoy et al. in 2007,<sup>13</sup>

Received: July 21, 2016

Revised: August 29, 2016

Accepted: August 30, 2016

Published: August 30, 2016

maternal exposure of yellow agouti (Avy) mice to BPA resulted in CpG hypomethylation at the Avy locus and a corresponding shift in coat color phenotypes within offspring, an effect that was rescued by maternal supplementation with methyl donors (folic acid or genistein). Importantly, this was one of the first animal studies to demonstrate that maternal exposure to an environmental chemical during *in utero* development has the potential to reprogram the offspring epigenome, leading to immediate phenotypic consequences (e.g., change in coat color) and potential long-term effects on adult disease susceptibility. These findings have since provided the basis for investigating the potential association of BPA exposure and epigenetic changes within human populations,<sup>14–16</sup> raising the possibility that epigenetic marks within certain loci may be used as diagnostic indicators of prior environmental exposures.

Tris(1,3-dichloro-2-propyl)phosphate (TDCIPP) is a chlorinated phosphate ester used as a high-production volume plasticizer and flame retardant within polyurethane (rigid and flexible), plastics, resins, and acrylic latexes.<sup>17</sup> As TDCIPP persists within indoor and outdoor environmental media,<sup>18</sup> environmental exposure to TDCIPP may pose a health risk to humans and ecological species, particularly during sensitive windows of early development (e.g., during epigenetic reprogramming) that have the potential to coincide with higher internal doses of TDCIPP.<sup>19</sup> Using zebrafish as a model, we previously demonstrated that the cleavage period (0.75–2.25 h postfertilization, hpf) during embryogenesis is susceptible to TDCIPP-induced delays in methylation of the zygotic genome.<sup>20</sup> Within various life-stages of zebrafish, TDCIPP exposure also results in a wide range of adverse effects, including disruption of thyroid hormone regulation,<sup>21–23</sup> gene expression,<sup>24,25</sup> behavior,<sup>26–29</sup> and reproduction.<sup>30–32</sup> In addition, exposure of early zebrafish embryos to 3  $\mu\text{M}$  TDCIPP, an identical nominal aqueous concentration used within our 2012 study, arrests epiboly and induces severe malformations later in development.<sup>24</sup> However, to our knowledge, no studies have investigated the potential impacts of TDCIPP on DNMT activity and cytosine methylation within zebrafish embryos (or any other animal model). Therefore, the objectives of this study were to determine whether (1) TDCIPP inhibits zebrafish DNMT within embryonic nuclear extracts; (2) uptake of TDCIPP from 0.75 hpf (2-cell) to 2 hpf (64-cell) or 6 hpf (shield stage) leads to impacts on the DNA methylome during cleavage; and (3) TDCIPP-induced impacts on cytosine methylation are localized to CpG islands within intergenic regions. For DNMT activity assays and all embryonic exposures, 5-azacytidine (5-azaC), an inhibitor of DNMT activity, was used as a positive control.<sup>4</sup>

## MATERIALS AND METHODS

**Animals.** Adult wild-type (strain 5D) zebrafish were maintained and bred on a 14 h light:dark cycle within a five-shelf stand-alone system (Aquatic Habitats, Inc., Apopka, FL, USA) containing photoperiod light-cycle enclosures and recirculating conditioned reverse osmosis (RO) water as previously described.<sup>33</sup> For all experiments described below, newly fertilized eggs were staged according to previously described methods.<sup>34</sup> All fish were handled and treated in accordance with approved Institutional Animal Care and Use Committee protocols at the University of South Carolina, Columbia.

**Chemicals.** TDCIPP (99% purity) and 5-azaC (>98% purity) were purchased from ChemService (West Chester, PA)

and Tocris (Bristol, United Kingdom), respectively. Stock and working solutions of each chemical were prepared and stored as previously described.<sup>33</sup>

**DNMT Activity Assays.** Newly fertilized eggs were collected immediately after spawning and placed in groups of approximately 100 per Petri dish within a light- and temperature-controlled incubator. Embryos (100 per time point) were collected at 2, 10, and 24 hpf and stored at  $-20\text{ }^{\circ}\text{C}$ . Nuclear proteins were extracted from whole embryo pools using an EpiQuik Nuclear Extraction Kit (Epigentek Group, Farmingdale, NY). Nuclear extract was kept on ice and immediately quantified using a BCA Protein Assay (Pierce Biotechnology, Rockford, IL) following the manufacturer's instructions. Optical density of the colorimetric reaction was quantified using a VICTOR X3Multilabel Plate Reader (PerkinElmer, Waltham, MA), and total protein was quantified using a standard curve generated from bovine serum albumin.


DNMT inhibition was quantified using an EpiQuik DNMT Activity/Inhibition Assay Ultra Fluorometric Kit (Epigentek Group, Farmingdale, NY). DNMT inhibition within nuclear extracts (6.5  $\mu\text{g}$  of protein per reaction) derived from 2-, 10-, and 24-hpf embryos was quantified in the presence of vehicle (0.1% dimethyl sulfoxide, DMSO), 250  $\mu\text{M}$  5-azaC, or TDCIPP (63, 125, 250, or 500  $\mu\text{M}$ ). All reactions were conducted in triplicate. Fluorescence was measured by a VICTOR X3Multilabel Plate Reader (PerkinElmer, Waltham, MA) and data were corrected for background and reported as relative fluorescent units.

### Embryonic Exposures for Genomic DNA Extractions.


Newly fertilized eggs were collected immediately after spawning and placed in groups of approximately 100 per Petri dish within a light- and temperature-controlled incubator until 0.75 hpf (2-cell stage). Prior to each experiment, 50 mL glass beakers were thoroughly rinsed with DMSO and RO water. Viable 5D embryos were exposed to 10 mL of vehicle (0.1% DMSO), 250  $\mu\text{M}$  5-azaC, or 2  $\mu\text{M}$  TDCIPP in triplicate glass beakers (70 embryos per replicate beaker) under a 14 h:10 h light:dark cycle and static conditions at  $28\text{ }^{\circ}\text{C}$  from 0.75 hpf to 2 or 6 hpf. To minimize the potential for false negative findings and confounding effects of systemic toxicity at 2 and 6 hpf, 250  $\mu\text{M}$  5-azaC and 2  $\mu\text{M}$  TDCIPP were selected based on the (1) absence of effects on cell cycle, overall embryo size, or cell morphology from 0.75 to 2 hpf; (2) absence of developmental delays from 0.75 to 6 hpf; but (3) presence of severe malformations (trunk curvature, tail malformations, craniofacial malformations, decreased body length, pericardial edema, and yolk sac edema) at 24 hpf (data not shown). Embryos (25/replicate) were collected at either 2 or 6 hpf, transferred from beakers to 2 mL cryovials, snap-frozen in liquid nitrogen, and stored at  $-80\text{ }^{\circ}\text{C}$ . These experiments resulted in three independent replicate samples for each time point and treatment group.

### Quantification of Internal Embryonic Doses of TDCIPP and BDCIPP.

Embryos (50 per replicate) were treated with vehicle (0.1% DMSO) or 2  $\mu\text{M}$  TDCIPP within six replicate beakers per treatment starting at 0.75 hpf. At 2 and 6 hpf, embryos were transferred from three replicate beakers to three 2 mL cryovials, snap-frozen in liquid nitrogen, and stored at  $-80\text{ }^{\circ}\text{C}$  until analysis. Frozen embryos were then processed and analyzed for TDCIPP and BDCIPP concentrations using previously described protocols.<sup>20</sup> Method detection limits (MDLs) were defined as three times the standard deviation of lab blanks (if present) or three times the noise. MDLs for



**Figure 1.** TDCIPP does not inhibit zebrafish DNMT *in vitro*. Mean relative fluorescence units (RFU)  $\pm$  standard deviation.  $N =$  three replicates/group.




**Figure 2.** TDCIPP uptake occurs by 2 hpf following initiation of exposure at 0.75 hpf. Mean internal dose (per 50 embryos)  $\pm$  standard deviation. BDCIPP was less than the method detection limit (MDL) across all groups.  $N =$  three replicate pools/group. Asterisk (\*) denotes significant treatment effect ( $p < 0.05$ ) relative to vehicle control (0.1% DMSO).

TDCIPP and BDCIPP were 0.1 and 1.0 ng, respectively. Recoveries averaged  $103 \pm 8\%$  and  $118 \pm 17\%$  for the internal standards, D15-TDCIPP and D10-BDCIPP, respectively.


**Quantification of 5-Methyl-2'-deoxycytidine (5-mdC) and 2'-Deoxyguanosine (dG).** Genomic DNA (1  $\mu\text{g}$  per replicate sample) was digested and processed similar to previously described methods.<sup>35</sup> Enzymes were removed by extraction with chloroform, and the resulting aqueous layer was subjected directly to LC-MS/MS and LC-MS/MS/MS analysis for quantification of 5-mdC and dG, respectively, as previously described.<sup>35</sup> The amounts of 5-mdC and dG (in moles) in nucleoside mixtures were calculated from area ratios of peaks found in selected-ion chromatograms for the analytes over their corresponding isotope-labeled standards (SI Figures S1 and S2), amounts of the labeled standards added (in moles), and calibration curves (SI Figure S3). 5-mdC levels (% of dG) were calculated by comparing the moles of 5-mdC relative to the moles of dG.

**DNA Methylome Profiling Using Whole-Genome Bisulfite Sequencing (WGBS).** Genomic DNA was extracted



**Figure 3.** 5-azaC and TDCIPP do not significantly impact total 5-mdC levels at 2 or 6 hpf (A) nor global cytosine methylation at 2 hpf (B) following initiation of exposure at 0.75 hpf. Mean 5-mdC (A) or cytosine methylation (B)  $\pm$  standard deviation.  $N =$  three replicates per group within Panel A.  $N =$  two replicates per control group and three replicates per treatment group within Panel B.

from pooled 2-hpf embryos (25 per replicate) using a Wizard Genomic DNA Purification Kit (Promega, Madison, WI). After elution of DNA in 25  $\mu\text{L}$  of nuclease-free 1 $\times$  Tris-EDTA buffer, the concentration and quality of all DNA samples were quantified using a Qubit 2.0 fluorometer (Invitrogen, Carlsbad, CA) and Agilent 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA), respectively; as the DNA concentration within one of the vehicle (0.1% DMSO) samples was  $<0.5$  ng/ $\mu\text{L}$ , only two vehicle control samples (vs three samples for 250  $\mu\text{M}$  5-azaC and 2  $\mu\text{M}$  TDCIPP) were prepared for sequencing. DNA samples (8 total) were then processed using an EZ DNA Methylation-Lightning Kit (Zymo Research Corp., Irvine, CA) for bisulfite conversion, and nonbarcoded sequencing libraries were then prepared using a TruSeq DNA Methylation Kit (Illumina, San Diego, CA). The concentration and quality of sequencing libraries were quantified using a Qubit 2.0 fluorometer (Invitrogen, Carlsbad, CA) and Agilent 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA), respectively, and eight libraries were then paired-end (2 $\times$ 50) sequenced on eight lanes (one library per lane) of a HiSeq 2500 Sequencing System (Illumina, San Diego, CA). This sequencing strategy generated 160–185 M short reads per replicate, providing sufficient



**Figure 4.** 5-azaC (A) and TDCIPP (B) exposure results in chromosome-specific alterations in cytosine methylation at 2 hpf following initiation of exposure at 0.75 hpf. All signal-to-noise ratios (SNRs) are relative to vehicle controls and reflect variation among two replicates per control group and three replicates per treatment group. SNRs > 0 and SNRs < 0 represent hypermethylation and hypomethylation, respectively, relative to vehicle controls.

coverage (~6–7× per sample) for differential methylation analysis by WGBS.<sup>36</sup>

All bisulfite-converted, 2×50 sequence reads (Reads 1 and 2) were first checked for quality control using FastQC. Approximately 97% of Read 1 sequences resulted in a quality score  $\geq$ Q30, whereas only 76% of Read 2 sequences resulted in a quality score  $\geq$ Q30. Therefore, Read 1 sequences were aligned against the current zebrafish genome assembly (GRCz10) using Bismark v0.14.3<sup>37</sup> and Bowtie 2 v2.2.5<sup>38</sup> to determine the methylation status of all cytosines within each sample. Bismark's output included BAM files (.bam) containing alignments, methylation call strings, and a report summarizing alignment statistics and percent methylated cytosines within a CpG, CHG, and CHH context, where H = A, T, or C. Bismark-generated BAM files were then sorted and converted into SAM files (.sam) using Samtools.<sup>39</sup> SAM files were then imported into methylKit<sup>40</sup> using the read.bismark function; the read.context option was set to "CpG", whereas all other options were set to default values for the read.bismark function. This

approach allowed us to determine percent cytosine methylation at base-pair resolution within a CpG context on each of 25 chromosomes. Using percent cytosine methylation data, signal-to-noise ratios ( $SNR = (\mu_1 - \mu_0) / (\sigma_1 + \sigma_0)$ , where  $\mu_1$  = treatment mean;  $\mu_0$  = vehicle control mean;  $\sigma_1$  = treatment standard deviation;  $\sigma_0$  = vehicle control standard deviation) were then calculated at all available positions on each of 25 chromosomes to account for the magnitude and variation of cytosine methylation relative to vehicle controls. SNRs > 0 and SNRs < 0 represent hypermethylation and hypomethylation, respectively, relative to vehicle controls.

#### Secondary Analysis of TDCIPP-Specific WGBS Data.

TDCIPP-specific data were sorted by SNR, and SNRs between -5 and +5 were removed from the data set. SNR thresholds of >5 or <5 were selected in order to (1) filter out highly variable vs consistent, TDCIPP-related impacts on cytosine methylation across independent treatment replicates and (2) prioritize regions of interest that were reliably impacted by TDCIPP exposure. Across all 25 chromosomes, 189 out of 25 270 positions were

identified and mapped to the current zebrafish genome assembly (GRCz10) hosted by the National Center for Biotechnology Information (NCBI) (<http://www.ncbi.nlm.nih.gov/genome?term=danio%20rerio>). After mapping each position to the zebrafish genome, we then (1) determined whether the position mapped to an intragenic (exon or intron) or intergenic region; (2) identified the NCBI Gene ID if mapped to an intragenic region; (3) identified the NCBI Gene ID corresponding to genes immediately upstream and downstream of the mapped position; and 4) determined whether the position mapped to a NCBI-predicted CpG island. Genomic sequences for all identified CpG islands were also compared against NCBI's nucleotide collection (nr/nt) database using Basic Local Alignment Search Tool (BLASTn) to identify potential human, mouse, or rat orthologs and, if mapped to an intragenic region, corresponding NCBI Gene IDs.

**Statistical Analyses.** Statistical procedures for DNMT data and internal embryonic doses were performed using SPSS Statistics 22.0 (Chicago, IL). A general linear model (GLM) analysis of variance (ANOVA) ( $\alpha = 0.05$ ) was used for all data, as these data did not meet the equal variance assumption for non-GLM ANOVAs. Pair-wise Tukey-based multiple comparisons of least-squares means were performed to identify significant differences among treatment groups.


**Accession Number.** Raw Illumina (fastq.gz) sequencing files (eight files totaling 70.98 GB) are available via NCBI's BioProject database under BioProject ID PRJNA330715 (<http://www.ncbi.nlm.nih.gov/bioproject/330715>).

## RESULTS AND DISCUSSION


Since our previous study suggested that normal DNA methylation at the end of cleavage (2 hpf) was decreased in TDCIPP-treated embryos,<sup>20</sup> we tested the hypothesis that TDCIPP inhibits zebrafish DNMT activity *in vitro*. We relied on an assay containing universal DNA substrate in order to identify potential impacts of TDCIPP on multiple zebrafish DNMTs present during early embryonic development.<sup>41–44</sup> Although 5-azaC significantly inhibited zebrafish DNMT within 2-, 10-, and 24-hpf nuclear extracts, TDCIPP did not affect DNMT activity up to 500  $\mu$ M (Figure 1), suggesting that TDCIPP-induced impacts on DNA methylation during cleavage are likely not due to direct DNMT inhibition. Rather, these findings point to the possibility that TDCIPP may be impacting the concentration of available methyl donors (e.g., S-adenosyl-L-methionine) during early embryonic development.

To help guide interpretation of our findings, we quantified mean internal doses of TDCIPP and BDCIPP within 2- and 6-hpf embryos following initiation of exposure to vehicle (0.1% DMSO) or 2  $\mu$ M TDCIPP at 0.75 hpf. Although mean internal doses (per 50 embryos) of TDCIPP were <8.5 ng within vehicle controls, which is consistent with typical background concentrations, and BDCIPP was not detected across all treatment groups, mean internal doses (per 50 embryos) of TDCIPP at 2 and 6 hpf were 80.86 and 129.97 ng, respectively (Figure 2). Therefore, these data demonstrate that TDCIPP uptake is rapid, TDCIPP is not metabolized to BDCIPP during the first 6 h of embryonic development, and TDCIPP (but not BDCIPP) is likely responsible for any impacts on DNA methylation.

To address the potential for 5-azaC- and TDCIPP-induced impacts on DNA methylation, we relied on (1) LC-MS/MS and LC-MS/MS/MS to quantify 5-mdC and dG, respectively, within genomic DNA extracted from vehicle-, 5-azaC-, and



**Figure 5.** TDCIPP exposure from 0.75 to 2 hpf results in genome-wide hypomethylation within intragenic regions of the zebrafish genome. Intragenic regions include exons and introns, whereas intergenic regions represent sequences between genes.



**Figure 6.** TDCIPP exposure from 0.75 to 2 hpf predominantly impacts positions mapped outside of CpG islands (A) and, for the remaining positions mapped within CpG islands, TDCIPP-induced effects were localized to intragenic regions of the zebrafish genome (B).

TDCIPP-exposed 2- and 6-hpf embryos and (2) WGBS on Illumina's HiSeq 2500 platform to identify the magnitude and extent, if any, of 5-azaC- and TDCIPP-induced effects on the entire DNA methylome at 2 hpf (the end of cleavage) at single

NC\_007115.6| Danio rerio strain Tuebingen chromosome 4, GRCz10 (Length=76625712)  
 NR\_046235.1| Homo sapiens RNA, 45S pre-ribosomal 5 (RNA45S5), ribosomal RNA (Length=13357)

NC_007115.6	31043007	CGGTGCTGCAGTATCGTTGCTTTTAGGCGGGATTCTGACTTAGAGGCGTTCAGTCATAAT	31043066
NR_046235.1	12664	CGGCGCTGCCGATCGTTCGCTGGGCGGGATTCTGACTTAGAGGCGTTCAGTCATAAT	12605
NC_007115.6	31043067	CCCACAGATGGTGGCTTCGCGCCAGTGGCTCCTCGGCCAAGCGCA-CGAGCCAAATGTCT	31043125
NR_046235.1	12604	CCCACAGATGGTAGCTTCGCCCCATTGGCTCCTCAGCCAAGCACATACA-CCAAATGTCT	12546
NC_007115.6	31043126	GAACC CGCGTTCCTCTCGTACTGAGCGGGATTACTATTGCAACAA CGTTCA-CATCAG	31043184
NR_046235.1	12545	GAACCTG CGGTTCCTCTCGTACTGAGCAGGATTACCATGGCAACAC---ACATCATCAG	12489
NC_007115.6	31043185	TAGGGTAAAATAACCTGTCTCA CGACGGTCTAAACCCAGCTCA CGTTCCTGTAGCGG	31043244
NR_046235.1	12488	TAGGGTAAAATAACCTGTCTCA CGACGGTCTAAACCCAGCTCA CGTTCCTATTAGTGG	12429
NC_007115.6	31043245	GTGAACAATCCAA CGCTTGGTGTATTCTGTTTACAAATGATAGGAAGAGC CGACATCGAA	31043304
NR_046235.1	12428	GTGAACAATCCAA CGCTTGGTGAATTCTGCTTACAAATGATAGGAAGAGC CGACATCGAA	12369
NC_007115.6	31043305	GTATCAAAA-----G--G--ATGAA CGCTTGGC CGCCACAAGCCAGTTATCCCTGTGGT	31043354
NR_046235.1	12368	GGATCAAAAAG CGACGT CGCTATGAA CGCTTGGC CGCCACAAGCCAGTTATCCCTGTGGT	12309
NC_007115.6	31043355	AACTTTCTGACACCTCTGCTTAAACCCCAAAGC CGGAAGGAT CGTGAGGCCCGCT	31043414
NR_046235.1	12308	AACTTTCTGACACCTCTGCTTAAACCCCAAAGGTCAGAAGGAT CGTGAGGCCCGCT	12249
NC_007115.6	31043415	TT CGCGGTC CGTACTCATACTGAAAATCAAGATCAAGCAAGCTTTTGCCTTCTGCTCTA	31043474
NR_046235.1	12248	TTCA CGGTCTGTATT CGTACTGAAAATCAAGATCAAG CGAGCTTTTGCCTTCTGCTCCA	12189
NC_007115.6	31043475	CGGGAGGTTTGTGTCCTCCCTGAGCT CGCCTTAGGACACCTG CGTTACCGTTTGACAGGT	31043534
NR_046235.1	12188	CGGGAGGTTTGTGTCCTCCCTGAGCT CGCCTTAGGACACCTG CGTTACCGTTTGACAGGT	12129
NC_007115.6	31043535	GTAC CGCCCCAGTCAAACCTCCACCTGCCACTGTCCC CGGGCAGGT CGCGCGC CGGGG	31043594
NR_046235.1	12128	GTAC CGCCCCAGTCAAACCTCCACCTGCCACTGTCCC CGGAGCGGT CGCGC-C-CG GC	12071
NC_007115.6	31043595	AGGCTGGCCC CGGGGCGCTTGACACAGAGGTGGGAAAGCCCTGACCCAGAGC-CACCC	31043653
NR_046235.1	12070	CGGCGCGCGCGCGGGCGCTTGGCGCCAGAAGCGAG-A--GCC--CTCGGGGT CGCCC	12017
NC_007115.6	31043654	AACCTCCCTGCCTAAC CGGGTGAAG---AAA CGATAAGAGTAGTGGTATTTCAC CGGCG	31043709
NR_046235.1	12016	--CC-CC--GCCTCAC CGGGTCAAGTGAAGAAA CGATCAGAGTAGTGGTATTTCAC CGGCG	11962
NC_007115.6	31043710	GCCAGC 31043715	
NR_046235.1	11961	GCC CGC 11956	

**Figure 7.** BLASTn identifies a CpG island localized to human 45S pre-ribosomal 5 as an ortholog (87% identity) to a zebrafish-specific, TDCIPP-susceptible 726-bp fragment of a 1373-bp intergenic CpG island on chr4. CpG sites are highlighted yellow.

base-pair resolution. Although 5-azaC and TDCIPP did not significantly impact 5-mdC levels at 2 or 6 hpf (Figure 3A) nor global cytosine methylation at 2 hpf (Figure 3B), 5-azaC and TDCIPP exposure resulted in chemical- and chromosome-specific alterations in cytosine methylation within a CpG context at 2 hpf (Figure 4), suggesting that aggregated analysis of global cytosine methylation lacks sufficient sensitivity to detect spatially resolved, position-specific effects. Interestingly, exposure to 5-azaC from 0.75 to 2 hpf resulted in a stronger impact than TDCIPP on cytosine methylation at 2 hpf, leading to strong hypermethylation (SNR > 20) within four chromosomes and strong hypomethylation (SNR < 20) within 19 chromosomes (Figure 4A). In contrast, although the overall magnitude of TDCIPP-induced impacts on cytosine methylation was less than 5-azaC, TDCIPP only resulted in strong hypomethylation (SNR < 20) across 13 chromosomes (in particular, chr1, chr22, and chr25) (Figure 4B).

On the basis of secondary analysis of our TDCIPP-specific WGBS data, we identified 39 out of 189 positions (20%) with

a SNR > 5 and 150 out of 189 positions (80%) with a SNR < 5 (Supplemental File 3), indicating that, based on a SNR threshold that filters out positions with highly variable responses, hypomethylation (relative to vehicle controls) represented the large majority of reproducible, TDCIPP-induced impacts on the zebrafish DNA methylome at 2 hpf. Moreover, with the exception of chr24, TDCIPP-induced alterations in cytosine methylation occurred on all other 24 chromosomes (in particular, chr1, chr 2, chr 3, chr 5, chr 7, and chr 20) (Supplemental File 3), suggesting that TDCIPP-induced epigenetic modifications are genome-wide and not localized to certain loci. Interestingly, contrary to our working hypothesis, 154 out of 189 positions (81.5%) mapped to intragenic regions of the zebrafish genome (151 positions mapped to exons, whereas only 3 positions mapped to introns), whereas only 35 out of 189 positions (18.5%) mapped to intergenic regions (which includes promoter regions) (Figure 5).

In addition to revealing the location of TDCIPP-induced impacts on the zebrafish genome, we also determined whether

these effects occurred within or outside of CpG islands. Surprisingly, 155 out of 189 positions (82%) mapped outside of CpG islands (Figure 6A). For the remaining positions (34 out of 189) that mapped within CpG islands, 79% of CpG islands identified (27 out of 34) were localized to intragenic regions of the zebrafish genome (with the majority being mapped to exons) (Figure 6B). For all 34 CpG islands, we then used BLASTn to identify potential orthologs within the human, mouse, and rat genome. Although no orthologs were identified within the rat or mouse genome, one ortholog was identified within the human genome (Supplemental File 3), where a zebrafish-specific 726-bp fragment of a 1373-bp intergenic CpG island on chr4 was 87% identical to an exon-localized CpG island within human 45S preribosomal 5 (RNA45SS, NCBI GeneID: 100861532) on chr22 (Figure 7).

This study yielded two important yet unexpected conclusions: (1) TDCIPP exposure during cleavage predominantly results in non-CpG island hypomethylation localized to intragenic regions of the zebrafish genome; and (2) WGBS-based identification of contaminant-susceptible CpG islands within the zebrafish methylome has the potential to identify human orthologs for epigenetic-focused environmental epidemiology studies. To date, the majority of the scientific literature has focused on the role of CpG island methylation within promoter (intergenic) regions in regulating gene transcription.<sup>7</sup> Moreover, most investigations have been limited to studying environmentally induced changes in DNA methylation within small, targeted regions of the genome rather than nontargeted, unbiased profiling of the entire methylome.<sup>45</sup> By using high-resolution WGBS to scan the entire zebrafish DNA methylome, we revealed that, contrary to our working hypothesis, non-CpG island hypomethylation within exons accounted for the majority of TDCIPP-induced effects, suggesting that strategies for enrichment of CpG islands prior to sequencing, such as methylated DNA affinity capture coupled with sequencing (MethylCap-seq) or reduced representation bisulfite sequencing (RRBS), may, although more cost-effective, lead to a high proportion of false-negative results. Moreover, the use of methods to eliminate non-CpG islands prior to sequencing may limit the probability of identifying potential candidate loci for future investigations in zebrafish, other animal models, or human populations.

In conclusion, we have revealed that TDCIPP-induced impacts to the zebrafish DNA methylome at 2 hpf are complex and primarily localized to regions outside of classical CpG islands. As our data are based on a single TDCIPP concentration and developmental stage, future research should focus on monitoring DNA methylation dynamics within WGBS-identified, TDCIPP-susceptible loci across multiple TDCIPP concentrations and developmental stages using a cost-effective, targeted sequencing-based strategy such as bisulfite amplicon sequencing.<sup>46</sup> In addition, as we have identified a CpG island within the human genome that may be susceptible to TDCIPP (or other flame retardants and plasticizers), future environmental epidemiological studies should consider measuring DNA methylation within this region to determine whether there are potential associations between chemical exposure, adverse health outcomes, and DNA methylation status.

## ■ ASSOCIATED CONTENT

### 📄 Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.est.6b03656.

Spreadsheets containing processed and analyzed WGBS data (XLSX)

Supporting data for 5-mdC and dG analysis (Figures S1–S3), and position-specific signal-to-noise ratios for cytosine methylation on each of 25 chromosomes (Figures S4–S28) (PDF)

Spreadsheets containing secondary analysis of TDCIPP-specific cytosine methylation data (XLSX)

## ■ AUTHOR INFORMATION

### Corresponding Author

\*D. C. Volz. Phone: (951) 827-4450; Fax: (951) 827-3993; E-mail: david.volz@ucr.edu.

### Present Address

#J. K. Leet. Columbia Environmental Research Center, U.S. Geological Survey, Columbia, Missouri 65201, United States.

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

Funding was provided by the National Institutes of Health (R21ES022797 and R21ES025392). We gratefully thank Dr. Robert Tanguay (Oregon State University) for providing founder fish to establish our 5D zebrafish colony, Dr. R. Sean Norman (University of South Carolina) for use of the VICTOR X3Multilabel Plate Reader, and John Weger and Clay Clark (Institute for Integrative Genome Biology, University of California, Riverside) for Illumina sequencing services.

## ■ REFERENCES

- (1) Tadros, W.; Lipshitz, H. D. The maternal-to-zygotic transition: a play in two acts. *Development* **2009**, *136* (18), 3033–42.
- (2) Niakan, K. K.; Han, J.; Pedersen, R. A.; Simon, C.; Pera, R. A. Human pre-implantation embryo development. *Development* **2012**, *139* (5), 829–41.
- (3) Mhanni, A. A.; McGowan, R. A. Global changes in genomic methylation levels during early development of the zebrafish embryo. *Dev. Genes Evol.* **2004**, *214* (8), 412–7.
- (4) Martin, C. C.; Laforest, L.; Akimenko, M. A.; Ekker, M. A role for DNA methylation in gastrulation and somite patterning. *Dev. Biol.* **1999**, *206* (2), 189–205.
- (5) Ding, Y. B.; Long, C. L.; Liu, X. Q.; Chen, X. M.; Guo, L. R.; Xia, Y. Y.; He, J. L.; Wang, Y. X. 5-aza-2'-deoxycytidine leads to reduced embryo implantation and reduced expression of DNA methyltransferases and essential endometrial genes. *PLoS One* **2012**, *7* (9), e45364.
- (6) Daxinger, L.; Whitelaw, E. Understanding transgenerational epigenetic inheritance via the gametes in mammals. *Nat. Rev. Genet.* **2012**, *13* (3), 153–62.
- (7) Burrell, H. H.; Baccarelli, A. A. Environmental epigenetics: from novelty to scientific discipline. *J. Appl. Toxicol.* **2014**, *34* (2), 113–6.
- (8) Kundakovic, M.; Gudsnuik, K.; Franks, B.; Madrid, J.; Miller, R. L.; Perera, F. P.; Champagne, F. A. Sex-specific epigenetic disruption and behavioral changes following low-dose in utero bisphenol A exposure. *Proc. Natl. Acad. Sci. U. S. A.* **2013**, *110* (24), 9956–61.
- (9) Anderson, O. S.; Nahar, M. S.; Faulk, C.; Jones, T. R.; Liao, C.; Kannan, K.; Weinhouse, C.; Rozek, L. S.; Dolinoy, D. C. Epigenetic responses following maternal dietary exposure to physiologically relevant levels of bisphenol A. *Environ. Mol. Mutagen* **2012**, *53* (5), 334–42.
- (10) Zhou, R.; Chen, F.; Chang, F.; Bai, Y.; Chen, L. Persistent overexpression of DNA methyltransferase 1 attenuating GABAergic inhibition in basolateral amygdala accounts for anxiety in rat offspring exposed perinatally to low-dose bisphenol A. *J. Psychiatr. Res.* **2013**, *47* (10), 1535–44.



- (11) Strakovsky, R. S.; Wang, H.; Engeseth, N. J.; Flaws, J. A.; Helfferich, W. G.; Pan, Y. X.; Lezmi, S. Developmental bisphenol A (BPA) exposure leads to sex-specific modification of hepatic gene expression and epigenome at birth that may exacerbate high-fat diet-induced hepatic steatosis. *Toxicol. Appl. Pharmacol.* **2015**, *284* (2), 101–12.
- (12) Mao, Z.; Xia, W.; Chang, H.; Huo, W.; Li, Y.; Xu, S. Paternal BPA exposure in early life alters Igf2 epigenetic status in sperm and induces pancreatic impairment in rat offspring. *Toxicol. Lett.* **2015**, *238* (3), 30–8.
- (13) Dolinoy, D. C.; Huang, D.; Jirtle, R. L. Maternal nutrient supplementation counteracts bisphenol A-induced DNA hypomethylation in early development. *Proc. Natl. Acad. Sci. U. S. A.* **2007**, *104* (32), 13056–61.
- (14) Nahar, M. S.; Liao, C.; Kannan, K.; Harris, C.; Dolinoy, D. C. In utero bisphenol A concentration, metabolism, and global DNA methylation across matched placenta, kidney, and liver in the human fetus. *Chemosphere* **2015**, *124*, 54–60.
- (15) Nahar, M. S.; Kim, J. H.; Sartor, M. A.; Dolinoy, D. C. Bisphenol A-associated alterations in the expression and epigenetic regulation of genes encoding xenobiotic metabolizing enzymes in human fetal liver. *Environ. Mol. Mutagen* **2014**, *55* (3), 184–95.
- (16) Kim, J. H.; Rozek, L. S.; Soliman, A. S.; Sartor, M. A.; Hablas, A.; Seifeldin, I. A.; Colacino, J. A.; Weinhouse, C.; Nahar, M. S.; Dolinoy, D. C. Bisphenol A-associated epigenomic changes in prepubescent girls: a cross-sectional study in Gharbiah, Egypt. *Environ. Health* **2013**, *12*, 33.
- (17) WHO, Environmental Health Criteria 209-Flame Retardants: Tris(chloropropyl)phosphate and tris(2-chloroethyl)phosphate. In Geneva, Switzerland, 1998; p 129pp.
- (18) van der Veen, I.; de Boer, J. Phosphorus flame retardants: properties, production, environmental occurrence, toxicity and analysis. *Chemosphere* **2012**, *88* (10), 1119–53.
- (19) Hoffman, K.; Butt, C. M.; Chen, A.; Limkakeng, A. T., Jr.; Stapleton, H. M. High Exposure to Organophosphate Flame Retardants in Infants: Associations with Baby Products. *Environ. Sci. Technol.* **2015**, *49* (24), 14554–9.
- (20) McGee, S. P.; Cooper, E. M.; Stapleton, H. M.; Volz, D. C. Early zebrafish embryogenesis is susceptible to developmental TDCPP exposure. *Environ. Health Perspect* **2012**, *120* (11), 1585–91.
- (21) Xu, T.; Wang, Q.; Shi, Q.; Fang, Q.; Guo, Y.; Zhou, B. Bioconcentration, metabolism and alterations of thyroid hormones of Tris(1,3-dichloro-2-propyl) phosphate (TDCPP) in Zebrafish. *Environ. Toxicol. Pharmacol.* **2015**, *40* (2), 581–6.
- (22) Wang, Q.; Lai, N. L.; Wang, X.; Guo, Y.; Lam, P. K.; Lam, J. C.; Zhou, B. Bioconcentration and transfer of the organophorous flame retardant 1,3-dichloro-2-propyl phosphate causes thyroid endocrine disruption and developmental neurotoxicity in zebrafish larvae. *Environ. Sci. Technol.* **2015**, *49* (8), 5123–32.
- (23) Wang, Q.; Liang, K.; Liu, J.; Yang, L.; Guo, Y.; Liu, C.; Zhou, B. Exposure of zebrafish embryos/larvae to TDCPP alters concentrations of thyroid hormones and transcriptions of genes involved in the hypothalamic-pituitary-thyroid axis. *Aquat. Toxicol.* **2013**, *126*, 207–13.
- (24) Fu, J.; Han, J.; Zhou, B.; Gong, Z.; Santos, E. M.; Huo, X.; Zheng, W.; Liu, H.; Yu, H.; Liu, C. Toxicogenomic responses of zebrafish embryos/larvae to tris(1,3-dichloro-2-propyl) phosphate (TDCPP) reveal possible molecular mechanisms of developmental toxicity. *Environ. Sci. Technol.* **2013**, *47* (18), 10574–10582.
- (25) Liu, C.; Wang, Q.; Liang, K.; Liu, J.; Zhou, B.; Zhang, X.; Liu, H.; Giesy, J. P.; Yu, H. Effects of tris(1,3-dichloro-2-propyl) phosphate and triphenyl phosphate on receptor-associated mRNA expression in zebrafish embryos/larvae. *Aquat. Toxicol.* **2013**, *128–129*, 147–57.
- (26) Jarema, K. A.; Hunter, D. L.; Shaffer, R. M.; Behl, M.; Padilla, S. Acute and developmental behavioral effects of flame retardants and related chemicals in zebrafish. *Neurotoxicol. Teratol.* **2015**, *52* (Pt B), 194–209.
- (27) Noyes, P. D.; Haggard, D. E.; Gonnerman, G. D.; Tanguay, R. L. Advanced morphological - behavioral test platform reveals neurodevelopmental defects in embryonic zebrafish exposed to comprehensive suite of halogenated and organophosphate flame retardants. *Toxicol. Sci.* **2015**, *145* (1), 177–95.
- (28) Wang, Q.; Lam, J. C.; Man, Y. C.; Lai, N. L.; Kwok, K. Y.; Guo, Y.; Lam, P. K.; Zhou, B. Bioconcentration, metabolism and neurotoxicity of the organophorous flame retardant 1,3-dichloro-2-propyl phosphate (TDCPP) to zebrafish. *Aquat. Toxicol.* **2015**, *158*, 108–15.
- (29) Dishaw, L. V.; Hunter, D. L.; Padnos, B.; Padilla, S.; Stapleton, H. M. Developmental exposure to organophosphate flame retardants elicits overt toxicity and alters behavior in early life stage zebrafish (*Danio rerio*). *Toxicol. Sci.* **2014**, *142* (2), 445–54.
- (30) Wang, Q.; Lam, J. C.; Han, J.; Wang, X.; Guo, Y.; Lam, P. K.; Zhou, B. Developmental exposure to the organophosphorus flame retardant tris(1,3-dichloro-2-propyl) phosphate: estrogenic activity, endocrine disruption and reproductive effects on zebrafish. *Aquat. Toxicol.* **2015**, *160*, 163–71.
- (31) Liu, X.; Ji, K.; Jo, A.; Moon, H. B.; Choi, K. Effects of TDCPP or TPP on gene transcriptions and hormones of HPG axis, and their consequences on reproduction in adult zebrafish (*Danio rerio*). *Aquat. Toxicol.* **2013**, *134–135*, 104–11.
- (32) Zhu, Y.; Ma, X.; Su, G.; Yu, L.; Letcher, R. J.; Hou, J.; Yu, H.; Giesy, J. P.; Liu, C. Environmentally Relevant Concentrations of the Flame Retardant Tris(1,3-dichloro-2-propyl) Phosphate Inhibit Growth of Female Zebrafish and Decrease Fecundity. *Environ. Sci. Technol.* **2015**, *49* (24), 14579–87.
- (33) Isales, G. M.; Hipszer, R. A.; Rafferty, T. D.; Chen, A.; Stapleton, H. M.; Volz, D. C. Triphenyl phosphate-induced developmental toxicity in zebrafish: potential role of the retinoic acid receptor. *Aquat. Toxicol.* **2015**, *161*, 221–30.
- (34) Kimmel, C. B.; Ballard, W. W.; Kimmel, S. R.; Ullmann, B.; Schilling, T. F. Stages of embryonic development of the zebrafish. *Dev. Dyn.* **1995**, *203* (3), 253–310.
- (35) Yu, Y.; Guerrero, C. R.; Liu, S.; Amato, N. J.; Sharma, Y.; Gupta, S.; Wang, Y. Comprehensive Assessment of Oxidatively Induced Modifications of DNA in a Rat Model of Human Wilson's Disease. *Mol. Cell. Proteomics* **2016**, *15* (3), 810–7.
- (36) Ziller, M. J.; Hansen, K. D.; Meissner, A.; Aryee, M. J. Coverage recommendations for methylation analysis by whole-genome bisulfite sequencing. *Nat. Methods* **2015**, *12* (3), 230–232 1 p following 232.
- (37) Krueger, F.; Andrews, S. R. Bismark: a flexible aligner and methylation caller for Bisulfite-Seq applications. *Bioinformatics* **2011**, *27* (11), 1571–2.
- (38) Langmead, B.; Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **2012**, *9* (4), 357–9.
- (39) Li, H.; Handsaker, B.; Wysoker, A.; Fennell, T.; Ruan, J.; Homer, N.; Marth, G.; Abecasis, G.; Durbin, R. Genome Project Data Processing, S., The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **2009**, *25* (16), 2078–9.
- (40) Akalin, A.; Kormaksson, M.; Li, S.; Garrett-Bakelman, F. E.; Figueroa, M. E.; Melnick, A.; Mason, C. E. methylKit: a comprehensive R package for the analysis of genome-wide DNA methylation profiles. *Genome biology* **2012**, *13* (10), R87.
- (41) Aluru, N.; Kuo, E.; Helfrich, L. W.; Karchner, S. I.; Linney, E. A.; Pais, J. E.; Franks, D. G. Developmental exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin alters DNA methyltransferase (dnmt) expression in zebrafish (*Danio rerio*). *Toxicol. Appl. Pharmacol.* **2015**, *284* (2), 142–51.
- (42) Fang, X.; Corrales, J.; Thornton, C.; Scheffler, B. E.; Willett, K. L. Global and gene specific DNA methylation changes during zebrafish development. *Comp. Biochem. Physiol., Part B: Biochem. Mol. Biol.* **2013**, *166* (1), 99–108.
- (43) Takayama, K.; Shimoda, N.; Takanaga, S.; Hozumi, S.; Kikuchi, Y. Expression patterns of dnmt3aa, dnmt3ab, and dnmt4 during development and fin regeneration in zebrafish. *Gene Expression Patterns* **2014**, *14* (2), 105–10.
- (44) Mhanni, A. A.; McGowan, R. A. Variations in DNA (cytosine-5)-methyltransferase-1 expression during oogenesis and early development of the zebrafish. *Dev. Genes Evol.* **2002**, *212* (11), 530–3.

(45) Rozek, L. S.; Dolinoy, D. C.; Sartor, M. A.; Omenn, G. S. Epigenetics: relevance and implications for public health. *Annu. Rev. Public Health* **2014**, *35*, 105–22.

(46) Masser, D. R.; Stanford, D. R.; Freeman, W. M. Targeted DNA methylation analysis by next-generation sequencing. *J. Visualized Exp.* **2015**, (96), e5248810.3791/52488.