UC San Diego UC San Diego Previously Published Works

Title

Altered lateralization of the cingulum in deployment-related traumatic brain injury: An ENIGMA military-relevant brain injury study

Permalink

https://escholarship.org/uc/item/4g68m68c

Journal Human Brain Mapping, 44(5)

ISSN

1065-9471

Authors

Dennis, Emily L Newsome, Mary R Lindsey, Hannah M <u>et al.</u>

Publication Date

2023-04-01

DOI

10.1002/hbm.26179

Copyright Information

This work is made available under the terms of a Creative Commons Attribution-NonCommercial License, available at <u>https://creativecommons.org/licenses/by-nc/4.0/</u>

Peer reviewed

RESEARCH ARTICLE

WILEY

Altered lateralization of the cingulum in deployment-related traumatic brain injury: An ENIGMA military-relevant brain injury study

Emily L. Dennis ^{1,2} Mary R. Newsome ^{3,4} Hannah M. Lindsey ^{1,2}
Maheen Adamson ^{5,6,7} 💿 Tara A. Austin ⁸ Seth G. Disner ^{9,10}
Blessen C. Eapen ^{11,12} Carrie Esopenko ¹³ Carol E. Franz ^{14,15}
Elbert Geuze ^{16,17} [] Courtney Haswell ¹⁸ Sidney R. Hinds II ¹⁹
Cooper B. Hodges ²⁰ Andrei Irimia ^{21,22} Kimbra Kenney ^{19,23}
Inga K. Koerte ^{24,25} William S. Kremen ^{14,15,26} Harvey S. Levin ^{3,4†}
Rajendra A. Morey ^{18,27,28} John Ollinger ²³ Jared A. Rowland ^{28,29,30}
Randall S. Scheibel ^{3,4} Martha E. Shenton ^{24,31} Danielle R. Sullivan ^{32,33}
Leah D. Talbert ³⁴ Sophia I. Thomopoulos ³⁵ Maya Troyanskaya ^{3,4}
William C. Walker ^{20,36} Xin Wang ³⁷ Ashley L. Ware ^{1,38} 🛛
John Kent Werner ¹⁹ Wright Williams ³ Paul M. Thompson ^{35,39}
David F. Tate ^{1,2} Elisabeth A. Wilde ^{1,2,4}

Correspondence

Emily L. Dennis, Department of Neurology, TBI and Concussion Center, University of Utah School of Medicine, Salt Lake City, UT, USA. Email: emily.dennis@hsc.utah.edu

Funding information

National Institutes of Health, Grant/Award Numbers: R61NS120249, K99NS096116; VA RR&D, Grant/Award Number: IK2RX002922; South Central VA Healthcare Network Veterans Health Administration: VA RR&D SPIRE, Grant/Award Number: I21RX001608: DOD PRARP: Dutch Ministry of Defence. Grant/Award Numbers: R01AG058822, R01NS100973; DoD, Grant/Award Number: W81XWH-18-1-0413; James J. and Sue Femino Foundation: Hanson-Thorell Research Scholarship; the USC CURVE & SURE programs: the Leonard Davis School of Gerontology, Grant/Award Number: R01NS100952; European Research Council (ERC), Grant/Award Number: 804326; VA

Abstract

Traumatic brain injury (TBI) in military populations can cause disruptions in brain structure and function, along with cognitive and psychological dysfunction. Diffusion magnetic resonance imaging (dMRI) can detect alterations in white matter (WM) microstructure, but few studies have examined brain asymmetry. Examining asymmetry in large samples may increase sensitivity to detect heterogeneous areas of WM alteration in mild TBI. Through the Enhancing Neuroimaging Genetics Through Meta-Analysis Military-Relevant Brain Injury working group, we conducted a mega-analysis of neuroimaging and clinical data from 16 cohorts of Active Duty Service Members and Veterans (n = 2598). dMRI data were processed together along with harmonized demographic, injury, psychiatric, and cognitive measures. Fractional anisotropy in the cingulum showed greater asymmetry in individuals with deployment-related TBI, driven by greater left lateralization in TBI. Results remained significant after accounting for potentially confounding variables including posttraumatic stress disorder, depression, and handedness, and were driven primarily by

[†] Deceased

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes. © 2022 The Authors. *Human Brain Mapping* published by Wiley Periodicals LLC.

For affiliations refer to page 1896

Rehabilitation Research and Development Service (VA RR&D), Grant/Award Numbers: I01RX003443, I01RX003442; VA Health Services Research and Development Service: Research Career Scientist, Grant/Award Number: IK6HX002608: Merit Review. Grant/Award Number: I01 CX001820; United States (U.S.) Department of Veterans Affairs; Clinical Sciences R&D (CSRD) Service: VA CSR&D Career Development; VA Career Development, Grant/Award Number: 5IK2CX001508; Defense and Veterans Brain Injury Centers; U.S. Army Medical Research and Materiel Command (USAMRMC), Grant/Award Numbers: W81XWH-13-2-0025 13129004 Chronic Effects of Neurotrauma Consortium (CENC), Grant/Award Number: PT108802-SC104835 U54EB020403 K99 MH119314; NIMH, Grant/Award Number: NARSAD 27786; Department of Defense; Psychological Health/ Traumatic Brain Injury Research Program LongTerm Impact of Military Relevant Brain Injury Consortium (LIMBIC), Grant/Award Number: W81XWH18PH; TBIRPLIMBIC; U.S. Department of Veterans Affairs, Grant/Award Numbers: 101 CX002097, 101 CX002096, 101 HX003155 I01 RX003444 I01 RX003443 I01 RX003442, I01 CX001135, I01 CX001246, I01 RX001774, I01 RX001135, I01 RX002076, I01 RX001880, I01 RX002172, I01 RX002173, I01 RX002171, I01 RX002174, I01 RX002170, W81XWH1920067, W81XWH1320095

1 | INTRODUCTION

The Iraq and Afghanistan conflicts have brought traumatic brain injury (TBI) to the forefront of military medicine. Service Members (SM) experienced ~450,000 TBIs between 2000 and 2021 (Military Health System, n.d), the majority of which were mild TBIs (mTBIs). While many warfighters appear to recover from these "invisible injuries" within weeks (Helmick et al., 2015), some continue to report somatic, cognitive, and psychological symptoms years later (Mac Donald et al., 2017). Deployment-related TBI is associated with worse outcomes than TBI sustained outside deployment (Martindale et al., 2018, 2021; Stein et al., 2015, 2016; Yurgil et al., 2014). Understanding brain structural changes and their relation to neurobehavioral sequelae following blast-related injuries, however, requires further elucidation as research on impact mTBI may not apply as the injury mechanisms differ.

Neuroimaging has revealed persistent alterations in brain structure and function after extracranial injuries have healed (Wilde et al., 2015). Diffusion magnetic resonance imaging (dMRI) is particularly sensitive to axonal neuropathology of TBI, but evidence is mixed regarding alterations in white matter (WM) organization after mTBI. Several studies have reported no TBI-related differences in dMRI metrics (Bolzenius et al., 2018; Davenport et al., 2015; Levin et al., 2010), and those that have reported significant results generally show lower fractional anisotropy (FA) in spatially distributed regions (Davenport et al., 2012; Hayes et al., 2015; Jorge et al., 2012). With the "pothole"

individuals whose worst TBI occurred before age 40. Alterations in the cingulum were also associated with slower processing speed and poorer set shifting. The results indicate an enhancement of the natural left laterality of the cingulum, possibly due to vulnerability of the nondominant hemisphere or compensatory mechanisms in the dominant hemisphere. The cingulum is one of the last WM tracts to mature, reaching peak FA around 42 years old. This effect was primarily detected in individuals whose worst injury occurred before age 40, suggesting that the protracted development of the cingulum may lead to increased vulnerability to insults, such as TBI.

KEYWORDS

DTI, military, traumatic brain injury

approach, group differences irrespective of location can be shown (Davenport et al., 2012; Jorge et al., 2012; Taber et al., 2015). Larger sample sizes may also help address heterogeneity, and big data approaches could facilitate sophisticated modeling to identify patient clusters with different patterns of WM disruption. One prior analysis on a subset of the cohorts included here used nonnegative matrix factorization (NMF), a data-driven approach, to reduce WM measurements to a set of components accounting for a large amount of the variance. This revealed an age-dependent effect of TBI on FA (a common proxy for WM organization), such that the age-related decreases in FA were steeper than expected after TBI (Bouchard et al., 2022). In addition to the pothole approach applied in previous papers and our prior investigation using the NMF approach, examining tract asymmetry may be another method to identify group differences in tract organization that is more robust to spatial heterogeneity.

Several WM tracts have well-established asymmetry, such as left lateralization of the cingulum and arcuate fasciculus, while asymmetry-related findings for other tracts have been mixed (Gong et al., 2005; Park et al., 2004; Song et al., 2015; Takao et al., 2013; Takao, Abe, et al., 2011; Takao, Hayashi, & Ohtomo, 2011; Yin et al., 2013). Tract asymmetry may jointly contribute to the functional lateralization of the brain along with gray matter asymmetries (Ocklenburg et al., 2016). Arcuate fasciculus asymmetry is closely tied to the lateralization of language (Ocklenburg et al., 2013), while greater cingulum asymmetry has been linked to attention and response/set shifting (Gläscher et al., 2012; Luna et al., 2021). We 1890 WILEY-

examined asymmetry of lateralized tracts to be more sensitive to patient heterogeneity and reduce the number of statistical tests. Asymmetry standardizes measures to the individual, making each his/her own reference, thus increasing sensitivity to subtle, heterogeneous effects (Toga & Thompson, 2003).

The Enhancing Neuroimaging Genetics Through Meta-Analysis (ENIGMA) Military-Relevant Brain Injury working group is part of the ENIGMA Brain Injury working group (https://enigma.ini.usc.edu/ ongoing/enigma-tbi/), a global collaboration among neuroimaging researchers focusing on various TBI patient populations (Dennis et al., 2020; Tate et al., 2021; Wilde et al., 2021). It was built on the framework created by the ENIGMA consortium, which seeks to achieve greater statistical power through harmonized image processing and meta/mega-analysis (Thompson et al., 2020). The ENIGMA diffusion tensor imaging (DTI) workflow (Jahanshad et al., 2013) has been used to identify altered WM organization across a range of disorders (Kelly et al., 2017; Kochunov et al., 2020), including moderate/severe TBI in pediatric patients (Dennis et al., 2021), and posttraumatic stress disorder (PTSD; Dennis et al., 2019). We used the ENIGMA DTI workflow to test the hypothesis that some structures would display greater asymmetry in participants with a history of TBI.

2 | MATERIALS AND METHODS

2.1 | Study samples

Our analysis included 16 cohorts of Veteran and Active Duty SMs, totaling 1775 participants with TBI history and 823 comparison

IABLE 1 Demographic information across cono	ABLE 1	Demographic	information	across	cohorts
---	--------	-------------	-------------	--------	---------

participants without TBI history. Of the 1775 participants, 1080 had at least one deployment-related injury, 480 had history of nonmilitary TBI only, and 215 did not have information regarding injury context (military vs. nonmilitary). While the majority of cohorts focused on Iraq/Afghanistan Veterans and SMs, two focused on Vietnam Veterans, a significantly older population. Across cohorts, the age range was 18–85 ($M = 41.7 \pm 12.7$) years. Table 1 provides demographic and clinical details on the cohorts. Inclusion and exclusion criteria for each study are in Table S1. All participants provided institutional review board-approved written informed consent. The group with a history of TBI was significantly younger than the group without TBI history (M = 40.6 vs. 43.8 years, $p = 3.6 \times 10^{-8}$) and had a greater proportion of males (91% vs. 85%, $p = 2.8 \times 10^{-5}$). The gender-

2.2 | Harmonizing injury data

potential exposure to TBI.

Details on injuries were collected using a range of tools (see Tables 1 and S2). From these disparate scales, we extracted common variables, as detailed in Note S1.

related differences match existing epidemiological trends in military

TBI, and the age difference is likely related to military rank and thus

2.3 | Image acquisition and processing

Acquisition parameters are provided in Table S3. Preprocessing, including eddy current correction, echo-planar imaging-induced

Note: Number	s are shown t	for total sample,	TBI (any) ar	id contro	l groups, and	l male/fema	le. The age	range, ave	erage age in	years with	SD, sample	TBI
severity, and t	he TBI scale	collected are also	o listed.									

Abbreviation: TBI, traumatic brain injury.

TABLE I Demog	graphic inform	nation ac	ross conort	S					
Site	Total no.	тві	Control	м	F	Age range	Average age (SD)	Severity	TBI scale
ADNIDoD	109	67	42	109	0	60-85	69.6 (4.6)	Mod/severe	Custom
BETTER/UMCU	84	12	72	84	0	21-57	36.2 (9.5)	Mild	Custom
LIMBIC-CENC	1028	837	191	891	137	22-70	40.1 (9.8)	Mild	PCE
Chronic Effects	80	64	16	72	8	23-54	34.7 (7.4)	Mild	Polytrauma
Duke1	105	70	35	86	19	24-46	39.6 (8.8)	Mild	MN-BEST/QCuBE
Duke2	84	50	34	58	26	23-67	40.1 (11.3)	Mild	MN-BEST/QCuBE
HDFT	28	23	5	23	5	24-58	36.9 (8.7)	Mild-moderate	Custom
Houston Blast	42	32	10	39	3	21-54	32.4 (8.0)	Mild	Polytrauma
INTRuST	85	54	31	69	16	18-69	39.4 (12.8)	Mild	Custom
iSCORE	101	30	71	88	13	19-51	35.8 (7.7)	Mild	Custom
NICoE	258	210	48	246	12	19-59	39.4 (6.5)	Mild	OSU
ROVER WISER	53	23	30	34	19	21-58	31.4 (6.4)	Mild	Polytrauma
SPIRE/GBET	53	39	14	47	6	24-50	35.3 (6.9)	Mild-severe	Polytrauma
VA Minneapolis1	124	92	32	120	4	23-62	34.3 (8.5)	Mild	MN-BEST
VA Minneapolis2	130	99	31	121	9	22-59	32.8 (8.4)	Mild	MN-BEST
VETSA	234	73	161	234	0	56-66	61.8 (2.6)	Mild	Custom
Total	2598	1775	823	2321	277	19-85	41.7 (12.7)		

distortion correction, and tensor fitting, was completed at the University of Utah for sites sharing raw data, or locally for others. Recommended protocols and quality control procedures are available on the ENIGMA-DTI and Neuroimaging Informatics Tools and Resources Clearinghouse webpages. These procedures were recommended, but coordination of preprocessing schemes accommodated site- and acquisition-specific pipelines. Once tensors were estimated, they were mapped and projected onto the ENIGMA-DTI template, and averaged within regions of interest (ROIs; http:// enigma.ini.usc.edu/protocols/dti-protocols/) in a TBSS-based approach (Smith et al., 2006). Further details and ROI abbreviations are in Note S2. In the seven sites for whom raw data was shared, we extracted motion parameters from the eddy current correction step to determine if motion played a role in our case-control findings. We compared rotation and translation averaged across the X, Y, and Z axes, finding no significant between-group differences (all ps >.05). We calculated FA lateralization index and asymmetry for each lateralized ROI:

$$Lateralization \ index = \left(FA_{Left} - FA_{Right}\right) / \left(0.5 * \left(FA_{Left} + FA_{Right}\right)\right)$$

where asymmetry was the absolute value of the lateralization index. Significant effects with asymmetry were followed post hoc with examinations of the lateralization index. Asymmetry was the primary measure as it would detect alterations irrespective of direction.

2.4 | Statistics

Mega-analysis was performed on individual-level ROI data. Linear mixed effects models were implemented with Ime in R 3.1.3. Nested random effects controlled for cohort and site, as some studies included multiple data collection sites. Age and gender were included as covariates in all analyses. The average correlation in asymmetry between all ROIs was r = .13. For multiple comparisons correction, we used a modified Bonferroni threshold, following recent ENIGMA analyses (Dennis et al., 2019) to calculate the effective number of independent tests based on the observed correlation structure between alternate responses. The equation of Li and Ji (Li & Ji, 2005) yielded an effective number of tests of $V_{eff} = 16$, yielding a significance threshold of p < .05/16 = .003125. Results that did not pass correction for multiple comparisons (.05 > p > .003125) are reported in the Supplement for completeness, but not interpreted. Across analyses, Cohen's d statistics are reported for group comparisons and unstandardized betas (b) are reported for linear regressions.

We calculated corrected *p*-values using the following equation:

$$P_{adj} = 1 - (1 - p)^{V_{eff}}$$

where p is the unadjusted p-value. Corrected p-values are shown for primary analyses. We used FDR to correct p-values for post hoc analyses.

2.5 | Quadratic age term

We first conducted analyses to determine whether a quadratic age term, age², should be included in statistical models along with age and gender, as age has a nonlinear effect on FA (Kochunov et al., 2012). The effect of this term upon the regression was not significant, so age² was not included in subsequent analyses.

2.6 | Primary group comparisons

Our primary analysis compared participants with a history of deployment-related TBI to those with no history of TBI, excluding individuals reporting a history of only nonmilitary TBI and participants whose records did not specify the source of TBI. Deployment-related TBI included both combat and noncombat injuries. As a specificity analysis, we examined group differences between participants with a history of nonmilitary TBI to a non-TBI group as well as individuals with a history of blast-related TBI to the non-TBI group.

2.7 | Supplementary analyses

Supplementary analyses on participant subgroups, interactions, injury variables, and symptom inventories are summarized in Note S3.

2.8 | Cognitive function

Across the 16 cohorts included in this study, seven collected the trail making test (TMT), including 1613 participants, 676 of whom had a history of deployment-related TBI. TMT Part A measures visual search and motor speed and Part B measures set shifting (Sánchez-Cubillo et al., 2009). Participants with Part A or Part B completion times greater than 3 SD above the study-wide mean were winsorized to 3 SD above the mean.

3 | RESULTS

3.1 | Primary group comparisons

We found significantly greater FA asymmetry in deployment-related TBI compared with no history of TBI in the cingulum (d = 0.18, corrected p = .0024), along with greater asymmetry in the superior longitudinal fasciculus (SLF), although this did not survive multiple comparisons correction (d = 0.10, corrected p = .35). Post hoc analysis of the cingulum revealed greater left lateralization in the TBI group (d = 0.15, FDR-corrected p = .0042; Figure 1; Benjamini & Hochberg, 1995). We further examined group differences in FA of the left and right cingula and found no significant differences. The higher asymmetry was also present in the blast-related TBI vs. non-TBI comparison, although it did not survive multiple comparisons correction



FIGURE 1 Group differences in cingulum asymmetry. Violin plots are shown for the nontraumatic brain injury (TBI) group (red), and deployment-related TBI group (blue) with group *t*-test *p*-values (corrected). The asymmetry analyses were corrected for multiple comparisons across all regions of interest tested using the Li and Ji adjusted Bonferroni correction. The laterality and cingulum fractional anisotropy (FA) analyses were false discovery rate-corrected across the three post hoc tests. The values on the y-axis are the normalized residuals for cingulum asymmetry, laterality index, left FA, and right FA, accounting for age, gender, and the nested random effects of cohort and site. As the cingulum is generally left-lateralized, actual laterality index values are generally positive, but that is not reflected here due to the adjustments and normalization

(d = 0.11, corrected p = .26). As an exploratory analysis in the LIMBIC-CENC cohort, we examined primary blast-related TBI and found no significant group difference in cingulum asymmetry. Primary blast-related TBI is TBI due to overpressure with no simultaneous blunt injury. Blast-related TBI is more commonly secondary or tertiary (with blunt impacts from projectiles or with force propelling the individual into a solid surface). Post hoc, the differences in the deployment-related TBI group remained significant when covarying for PTSD, depression, and both. Comparing individuals with a history of only nonmilitary TBI to the non-TBI group, the effect in the cingulum was not significant. Last, to rule out differences in scan parameters across sites, we examined the effect in the cingulum separately in sites using isotropic versus anisotropic voxels, and in sites using high angular resolution versus low angular resolution (low resolution being 32 directions or fewer). Results were consistent across subanalyses. Results are summarized in Tables 2 and S4 and S5.

3.2 | Cognitive function

In the non-TBI group, there were no detectable significant associations between TMT and cingulum FA. We examined the effect in the cingulum post hoc to test for functional consequences of the shift in asymmetry. As a post hoc test, we corrected for multiple comparisons across eight tests using FDR (TMT-A and TMT-B with cingulum asymmetry, laterality, and left and right cingulum FA). There were no significant associations with asymmetry in the deployment-related TBI group. There were slight trend associations in the deployment-related TBI group between the lateralization index of the cingulum and TMT-A and TMT-B time ($b = -5.5 \times 10^{-4}$, p = .16; $b = -1.8 \times 10^{-4}$, p = .16, respectively; Figure S1). Within the deployment-related TBI group, FA of the left cingulum was negatively associated with TMT-A and TMT-B ($b = -5.0 \times 10^{-4}$, p = .042; $b = -1.7 \times 10^{-4}$, p = .042, respectively, Figure 2).

4 | DISCUSSION

We examined WM asymmetry in 2598 military SM and Veterans, reporting a primary finding of greater asymmetry in the cingulum among individuals with a history of deployment-related TBI compared to those with no lifetime history of TBI. This was driven by greater left lateralization in the TBI group. The cingulum is a key structure in the limbic system, connecting medial aspects of the frontal and parietal lobes (not including the hippocampal cingulum, as the dorsal and hippocampal were distinct in this study; Budisavljevic et al., 2016) and supporting a number of important executive functions, including working memory, inhibition, and processing speed (Bettcher et al., 2016). The cingulum is also one of the last tracts to mature, with an average age-at-peak FA of 42 years old (Lebel et al., 2012). This

ROIs	Deployment-related 1	rbi (N = 1872)		Blast-related TBI (N =	1509)		Primary blast TBI (N =	= 337)		HX of TBI with LOC		
	d [95% CI]	Uncorrected <i>p</i>	Corrected <i>p</i>	d [95% CI]	Uncorrected <i>p</i>	Corrected <i>p</i>	d [95% CI]	Uncorrected <i>p</i>	Corrected <i>p</i>	d [95% CI]	Uncorrected <i>p</i>	Corrected <i>p</i>
ACR	$-0.01 \left[-0.11, 0.08\right]$.75	Ţ	-0.02 [-0.11, 0.07]	0.63	1	-0.06 [-0.16, 0.03]	.17	.95	0.06 [-0.04, 0.15]	.24	66.
ALIC	0.04 [-0.05, 0.13]	.35	1	0.03 [-0.06, 0.12]	0.52	1	0.05 [-0.04, 0.14]	.32	1	0.05 [-0.04, 0.14]	.29	1
000 000	0.18 [0.08, 0.27]	.00015	.0024	0.11 [0.02, 0.20]	0.019	.26	-0.06 [-0.15, 0.04]	.23	.98	0.06 [-0.04, 0.15]	.25	66.
CGH	0.01 [-0.08, 0.10]	.81	1	0.01 [-0.08, 0.10]	0.82	1	$-0.04 \left[-0.13, 0.05\right]$.39	1	0.02 [-0.07, 0.11]	.66	1
CST	-0.08 [-0.17, 0.01]	.086	.76	-0.05 [-0.14, 0.04]	0.24	66.	-0.07 [-0.16, 0.02]	.12	.87	-4.1 e-3 [-0.09, 0.09]	.93	1
CR	0.06 [-0.03, 0.15]	Ż	1	0.04 [-0.05, 0.13]	0.41	1	0.04 [-0.05, 0.14]	.34	1	0.12 [0.03, 0.21]	.009	.13
EC	$-0.04 \left[-0.13, 0.05\right]$.42	1	$-0.02 \left[-0.11, 0.07\right]$	0.73	1	0.09 [0.00, 0.18]	.056	9.	-0.04 [-0.13, 0.05]	.39	1
FX/ ST	0.03 [-0.06, 0.12]	.46	Ł	0.02 [-0.07, 0.11]	0.62	ħ	-0.07 [-0.16, 0.02]	.14	.91	0.03 [-0.06, 0.12]	.54	L
<u>ں</u>	0.05 [-0.04, 0.14]	ų	1	0.04 [-0.05, 0.13]	0.39	1	0.11 [0.02, 0.20]	.017	.24	0.02 [-0.07, 0.11]	.68	1
PCR	0.04 [-0.05, 0.13]	.39	1	0.04 [-0.05, 0.13]	0.42	1	0.02 [-0.07, 0.12]	9.	1	0.06 [-0.03, 0.16]	.17	.95
PLIC	1.0 e-3 [-0.09, 0.09]	.98	1	-0.01 [-0.10, 0.08]	0.78	1	5.1 e-3 [-0.09, 0.10]	.91	1	-0.03 [-0.12, 0.06]	.46	1
PTR	0.07 [-0.02, 0.16]	.12	.87	0.06 [-0.03, 0.15]	0.19	.97	0.05 [-0.04, 0.14]	.27	66:	0.07 [-0.02, 0.16]	.15	.93
RLIC	3.0 e-3 [-0.09, 0.09]	.95	1	0.02 [-0.07, 0.11]	0.71	1	0.08 [-0.01, 0.17]	.088	.77	0.04 [-0.05, 0.13]	.42	1
SCR	0.06 [-0.03, 0.15]	.22	.98	0.09 [0.00, 0.18]	0.055	9 .	0.05 [-0.04, 0.14]	.31	1	-1.5 e-3 [-0.09, 0.09]	.97	1
SFO	-0.03 [-0.12, 0.06]	.55	1	0.02 [-0.07, 0.11]	0.69	1	0.02 [-0.07, 0.11]	.73	1	0.04 [-0.06, 0.13]	.44	1
SLF	0.10 [0.01, 0.19]	.027	.35	0.06 [-0.03, 0.15]	0.18	.96	0.03 [-0.06, 0.12]	.57	1	0.14 [0.05, 0.23]	.0018	.028
SS	0.07 [-0.02, 0.16]	.13	.89	0.05 [-0.04, 0.14]	0.26	.99	-0.06 [-0.15, 0.04]	.24	.99	0.06 [-0.03, 0.15]	.18	.96
TAP	-0.08 [-0.18, 0.01]	.068	.68	-0.02 [-0.11, 0.07]	0.63	1	0.03 [-0.06, 0.12]	.53	1	-0.02 [-0.11, 0.07]	.66	1
UNC	0.02 [-0.07, 0.11]	.67	1	0.02 [-0.07, 0.11]	0.67	£1	0.03 [-0.06, 0.13]	.45	÷	0.04 [-0.05, 0.13]	.35	t.
<i>Note</i> : Re: uncorrec	sults from group analyse ted <i>p</i> -values are shown,	s are shown cor bolded statistic	mparing deploy s are those tha	ment-related TBI to no it survive correction for	TBI, blast-relat multiple compa	ed TBI to no T Irisons, italicize	BI, and primary blast T ed statistics are those t	Bl to no TBl, alor hat do not surviv	ng with total s e multiple com	ample size for each comp parisons correction (.05	arison. Cohen's > <i>p</i> > .003125).	d, 95% Cl, and Primary blast

Group differences in tract asymmetry. **TABLE 2** TBI is blast overpressure with no concurrent blunt injury.

Abbreviations: ACR, Anterior corona radiata; ALIC, anterior limb of internal capsule; CGC, cingulum; CGH, hippocampal cingulum; CR, corona radiata; CST, corticospinal tract; EC, external/extreme capsule; FX/ST, fornix-stria terminalis; IC, internal capsule; LOC, loss of consciousness; PCR, posterior corona radiata; PLIC, posterior limb of internal capsule; PTR, posterior thalamic radiation; RLIC, retrolenticular limb of internal capsule; ROI, Region of interest; SCR, superior corona radiata; SFO, superior fronto-occipital fasciculus; SLF, superior longitudinal fasciculus; SS, sagittal stratum; TAP, tapetum; TBI, traumatic brain injury; UNC, uncinate.



FIGURE 2 Group differences in tract asymmetry. Results from group analyses are shown comparing deployment-related traumatic brain injury (TBI) to no TBI, blast-related TBI to no TBI, and primary blast TBI to no TBI, along with total sample size for each comparison. Cohen's *d*, 95% CI, and uncorrected *p*-values are shown, bolded statistics are those that survive correction for multiple comparisons, italicized statistics are those that do not survive multiple comparisons correction (.05 > p > .003125). Primary blast TBI is blast overpressure with no concurrent blunt injury. ACR, anterior corona radiata; ALIC, anterior limb of internal capsule; CGC, cingulum; CGH, hippocampal cingulum; CR, corona radiata; CST, corticospinal tract; EC, external/extreme capsule; FX/ST, fornix-stria terminalis; IC, internal capsule; PCR, posterior corona radiata; PLIC, posterior limb of internal capsule; PCR, posterior thalamic radiation; RLIC, retrolenticular limb of internal capsule; ROI, Region of interest; SCR, superior corona radiata; SFO, superior fronto-occipital fasciculus; SLF, superior longitudinal fasciculus; SS, sagittal stratum; TAP, tapetum; UNC, uncinate.

prolonged maturation may render the cingulum more vulnerable to environmental impacts, and prior structural equation modeling did not show significant genetic contributions to cingulum asymmetry, supporting the hypothesis that environment plays an outsized role (Jahanshad et al., 2010). This age-at-peak happens to coincide with the average age of our sample (41.7 years). Indeed, when we ran the analysis after separating participants whose worst injury occurred before age 40 versus those whose injury occurred after age 45, only the comparison in the younger group remained significant. This analysis excluded Vietnam Veterans as the remoteness of their injuries may implicate different injury-related and age-related processes. Heterogeneity across cohorts and the general reliance on self-reports of injuries, including their timing and severity, mean that this must be interpreted with caution, but this does present a hypothesis that can be addressed more directly in individual studies. As the sample size differed greatly between these analyses (855 younger vs. 106 older TBI participants), we cannot rule out decrease in statistical power as the reason for the difference; however, the result does lend support to the interpretation that the cingulum is particularly vulnerable to injury because of its long maturation.

The length, orientation, and proximity of the cingulum to the falx and ventricles may contribute to strain and shear forces on the

tract during impact (Bigler, 2007; Ware et al., 2019; Zhou et al., 2022). Disentangling the effects of blast-related versus impact-related TBI is difficult, especially in our analysis of heterogeneous legacy datasets. The differences between blast-related TBI and no TBI were not significant after multiple comparisons correction and the effect size was smaller, but the sample size was smaller, and blast-related injuries in military settings often also involve a co-occurring impact. An exploratory analysis in the small subset of the sample with data on primary blast TBI (blast TBI due to overpressure with no concurrent blunt injury; 146 vs. 191) did not yield significant results for the cingulum, lending support to the interpretation that the biomechanical forces of TBI may affect the cingulum more than the pressure forces of blast-related TBI. Targeted analysis in animal studies and well-characterized cohort studies are necessary to confirm this finding.

Left lateralization of cingulum FA is a well-documented phenomenon in healthy (primarily right-handed) individuals (Gong et al., 2005; Takao et al., 2013; Takao, Hayashi, & Ohtomo, 2011) and is linked with cingulum-related cognitive functions (Caeyenberghs et al., 2016; Luna et al., 2021; Metzler-Baddeley et al., 2012). Development, aging, and sex do not appear to contribute to laterality of the whole cingulum (Takao, Abe, et al., 2011; Trivedi et al., 2009), although aging effects have been reported for the subcallosal cingulum (Sibilia et al., 2017). In tracts that are more asymmetric, the nondominant hemisphere may be more susceptible to damage, as neural resources may be diverted to maintaining integrity of the dominant hemisphere. Lower FA in the right cingulum may be due to less myelination, or to less organized or less densely packed fibers. A recent, large study of young adults using multishell dMRI showed that the neuronal density of the left cingulum was greater than that of the right, while the orientation dispersion index was higher on the right, indicating less coherence in fiber direction (Tsuchida et al., 2021). Lower fiber density could mean greater elasticity and thus more stretch during impact; finite element modeling consistently indicates high axonal strain in the cingulum (Chatelin et al., 2011; Zhang & Gennarelli, 2011; Zhao et al., 2019). Less asymmetry may partially explain the lack of significant group differences in females, as some studies have found lower tract asymmetry in women than men (Jahanshad et al., 2010; Thiebaut de Schotten et al., 2011), but the much smaller female sample size (n = 199) limited statistical power and thus limits our ability to draw conclusions from the female-only analyses.

Alterations in cingulum microstructure have been reported previously in mTBI (Kinnunen et al., 2011; Kraus et al., 2007; Niogi et al., 2008; Rutgers et al., 2008; Wu et al., 2010), including militaryspecific mTBI (Mac Donald et al., 2011; Yeh et al., 2014; Yurgelun-Todd et al., 2011), as well as across psychiatric disorders (Abdul-Rahman et al., 2011; Kochunov et al., 2020; Lochner et al., 2012). In healthy individuals, greater cingulum asymmetry has been associated with better attentional control and executive function (Luna et al., 2021). Other studies have reported that alterations in WM asymmetry are associated with cognitive deficits (Gómez-Gastiasoro et al., 2019). Two prior studies examining WM asymmetry in mTBI reported increased asymmetry in the corticospinal tract (CST; Maruta et al., 2020; Vakhtin et al., 2020). Unfortunately, this midline tract was not separated into left and right components in the ENIGMA atlas, so we did not have the opportunity to replicate that finding. Several studies in participants with TBI have reported associations between WM microstructural organization in the right cingulum and cognitive function, including reaction time and set shifting (Bonnelle et al., 2011; Kinnunen et al., 2011; Sugiyama et al., 2009). Prior studies have shown that deployment-related TBI is associated with poorer TMT performance, even when compared to nondeployment TBI (Martindale et al., 2021). We report that lower FA of the left cingulum is linked with slower processing speed and set shifting in the TBI group, echoing the findings of Gläscher et al., 2012 and Luna et al., 2021, but this association was not present in the non-TBI comparison group. Altered asymmetry of the cingulum in our sample was driven by an exaggerated laterality index, but no significant changes in either hemisphere. This suggests that asymmetry differences may be due to minor changes in both hemispheres, or that the anatomic location of the changes differ across participants.

Our study has several limitations. First, as a retrospective analysis of data collected across different studies, there are many sources of

heterogeneity in our data that we cannot fully characterize or account for. Differences in TBI history assessment limited the analyses to the most common denominators, which were often more general variables than individual sites collected. Our results present several interesting hypotheses that future studies can interrogate in greater detail, such as whether the cingulum is vulnerable because of its extended maturation, whether females are less vulnerable to alterations in cingulum asymmetry because they have less natural cingulum asymmetry, or whether the right cingulum is more vulnerable due to less densely packed fibers. Second, most of the TBI history for our cohorts was self-reported, which has inherent issues with reliability. We report all analyses of injury variables (such as duration of loss of consciousness) for completeness, but do not interpret these results. Third, TBSS is limited as a tensor-based approach compared to a tractography approach, so we cannot fully attribute results to particular fiber bundles. Fourth, we were limited in the clinical endpoints we could examine to those that were most general (e.g., categorical variable for current depression), most commonly collected (e.g., TMT), or already harmonized (e.g., PCL versions; Kennedy, 2022). Work is ongoing in the ENIGMA Clinical Endpoints working group (a subgroup of the Brain Injury working group) to develop additional harmonized measures across multiple domains. Thus, future studies will be able to examine how TBI-related alterations in brain structure and function correspond to changes in memory, attention, and inhibition, among others. Fifth, as a cross-sectional study, we cannot examine whether asymmetry differences predate injury, or how the differences in asymmetry evolve over the lifespan. Sixth, there were significant differences between our groups in age and gender. These match epidemiological trends, and age and gender were included as covariates in all analyses. Lastly, while one cohort came from the Netherlands, the rest of the cohorts were from the United States, which may limit the generalizability of our results to other military contexts.

5 | CONCLUSIONS

Our effect sizes are small, but in line with other ENIGMA analyses (Dennis et al., 2019; Kochunov et al., 2020). Given the many sources of heterogeneity in our sample, and the existing, often inconclusive, literature on dMRI alterations in military-relevant TBI (predominantly mTBI), small effect sizes are expected. The necessary sample size to find an effect of this size with 80% power and a significance threshold of 5% is 972, far larger than most published work on dMRI in military brain injury, highlighting the importance of collaborative projects. Our work points to subtle alterations in the balance of the brain after TBI, and suggests a phenomenon of heightened vulnerability of the nondominant hemisphere, perhaps as neural resources are diverted to support the dominant hemisphere. These alterations have functional consequences, as they were associated with slower processing speed and set shifting. Future targeted studies in individual, wellcharacterized cohorts or animal models will further elucidate specific underlying mechanisms and functional implications for alterations in tract asymmetry.

FUNDING INFORMATION

This work was supported by the NIH R61NS120249 K99NS096116; U.S. Army Medical Research and Materiel Command (USAMRMC) award 13129004: VA RR&D IK2RX002922: South Central VA Healthcare Network Veterans Health Administration: VA RR&D SPIRE Award I21RX001608: DOD PRARP: Dutch Ministry of Defence: R01AG058822 R01NS100973. DoD contract W81XWH-18-1-0413, a grant from the James J. and Sue Femino Foundation, a Hanson-Thorell Research Scholarship, the USC CURVE & SURE programs, and the Leonard Davis School of Gerontology: R01NS100952; European Research Council (ERC) (Starting Grant 804326); VA Rehabilitation Research and Development Service (VA RR&D) I01RX003443 I01RX003442: VA Health Services Research and Development Service Research Career Scientist Award IK6HX002608: Merit Review Award Number I01 CX001820 from the United States (U.S.) Department of Veterans Affairs Clinical Sciences R&D (CSRD) Service: VA CSR&D Career Development Award VA Career Development Award 5IK2CX001508: VA CSR&D Career Development Award 1IK2CX001772-01: Defense and Veterans Brain Injury Centers, the U.S. Army Medical Research and Materiel Command (USAMRMC): W81XWH-13-2-0025 and the Chronic Effects of Neurotrauma Consortium (CENC): PT108802-SC104835 U54EB020403 K99 MH119314; (NIMH); NARSAD 27786. This work was supported by the Assistant Secretary of Defense for Health Affairs endorsed by the Department of Defense, through the Psychological Health/Traumatic Brain Injury Research Program LongTerm Impact of Military Relevant Brain Injury Consortium (LIMBIC) Award W81XWH18PH/TBIRPLIMBIC under Awards W81XWH1920067 W81XWH1320095, No and by the U.S. Department of Veterans Affairs Awards No. I01 CX002097, I01 CX002096, I01 HX003155, I01 RX003444, I01 RX003443, I01 RX003442, I01 CX001135, I01 CX001246, I01 RX001774, I01 RX001135, I01 RX002076, I01 RX001880, I01 RX002172, I01 RX002173, I01 RX002171, I01 RX002174 I01 RX002170.

CONFLICT OF INTEREST

The views expressed in this article are those of the author(s) and do not reflect the official policy of the Department of Army/Navy/Air Force, Department of Defense, or US Government. Inga K. Koerte receives funding for a collaborative project unrelated to this article and serves as a paid scientific advisor for Abbott. She receives royalties for book chapters. Her spouse is an employee at Siemens AG. Paul M. Thompson received a research grant from Biogen, Inc., for research unrelated to this article.

AFFILIATIONS

¹Department of Neurology, University of Utah School of Medicine, Salt Lake City, Utah, USA

²George E. Wahlen Veterans Affairs Salt Lake City Healthcare System, Salt Lake City, Utah, USA ³Michael E. DeBakey Veterans Affairs Medical Center, Houston, Texas, USA

⁴H. Ben Taub Department of Physical Medicine and Rehabilitation, Baylor College of Medicine, Houston, Texas, USA

⁵Rehabilitation Department, VA Palo Alto Health Care System, Palo Alto, California, USA

⁶Neurosurgery, Stanford School of Medicine, Stanford, California, USA

⁷Operational Military Exposure Network (WOMEN), VA Palo Alto Healthcare System, California, Palo Alto, 94304, USA

⁸The VA Center of Excellence for Research on Returning War Veterans, Waco, Texas, USA

⁹Minneapolis VA Health Care System, Minneapolis, Minnesotta, USA ¹⁰Department of Psychiatry and Behavioral Sciences, University of Minnesota Medical School, Minneapolis, Minnesotta, USA

¹¹Department of Physical Medicine and Rehabilitation, VA Greater Los Angeles Health Care System, Los Angeles, California, USA
¹²Department of Medicine, David Geffen School of Medicine at

UCLA, Los Angeles, California, USA

¹³Department of Rehabilitation and Human Performance, Icahn School of Medicine at Mount Sinai, New York, New York, USA ¹⁴Department of Psychiatry, University of California, San Diego, La Jolla, California, USA

¹⁵Center for Behavior Genetics of Aging, University of California, San Diego, La Jolla, California, USA

¹⁶University Medical Center Utrecht, Utrecht, The Netherlands
¹⁷Brain Research and Innovation Centre, Ministry of Defence, Utrecht, The Netherlands

¹⁸Department of Psychiatry and Behavioral Sciences, Duke University, Durham, North Carolina, USA

¹⁹Department of Neurology, Uniformed Services University, Bethesda, Maryland, USA

²⁰Department of Physical Medicine and Rehabilitation, Virginia Commonwealth University, Richmond, Virginia, USA

²¹Ethel Percy Andrus Gerontology Center, Leonard Davis School of Gerontology, University of Southern California, Los Angeles, California, USA

²²Department of Biomedical Engineering, Viterbi School of Engineering, University of Southern California, Los Angeles, California, USA

²³National Intrepid Center of Excellence, Walter Reed National Military Medical Center, Bethesda, Maryland, USA

²⁴Psychiatry Neuroimaging Laboratory, Brigham and Women's Hospital, Boston, Massachusetts, USA

²⁵Department of Child and Adolescent Psychiatry, Psychosomatics and Psychotherapy, Ludwig-Maximilians-Universität, Munich, Germany

²⁶Center of Excellence for Stress and Mental Health, VA San Diego Healthcare System, La Jolla, California, USA

²⁷Duke-UNC Brain Imaging and Analysis Center, Duke University, Durham, North Carolina, USA

²⁸VA Mid-Atlantic Mental Illness Research Education and Clinical Center (MA-MIRECC), Durham, North Carolina, USA ²⁹W.G. (Bill) Hefner VA Medical Center, Salisbury, North

Carolina, USA

³⁰Department of Neurobiology & Anatomy, Wake Forest School of Medicine, Winston-Salem, North Carolina, USA

³¹VA Boston Healthcare System, Boston, Massachusetts, USA

³²National Center for PTSD, VA Boston Healthcare System, Boston, Massachusetts, USA

³³Department of Psychiatry, Boston University School of Medicine, Boston, Massachusetts, USA

³⁴Department of Psychology, Brigham Young University, Provo, Utah, USA

³⁵Imaging Genetics Center, Stevens Neuroimaging & Informatics Institute, Keck School of Medicine of USC, Marina del Rey, California, USA

³⁶Hunter Holmes McGuire Veterans Affairs Medical Center, Richmond, Virginia, USA

³⁷Department of Psychiatry, University of Toledo, Toledo, Ohio, USA
 ³⁸Department of Psychology, Georgia State University, Atlanta,

Georgia, USA

³⁹Departments of Neurology, Pediatrics, Psychiatry, Radiology, Engineering, and Ophthalmology, USC, Los Angeles, California, USA

DATA AVAILABILITY STATEMENT

Researchers interested in accessing the data described here must first join the ENIGMA Military-Relevant Brain Injury group and agree to abide by the Memorandum of Understanding governing data sharing and authorship. All projects are opt-in, so the specific cohorts available will differ.

ORCID

Emily L. Dennis ^b https://orcid.org/0000-0001-7112-4009 Maheen Adamson ^b https://orcid.org/0000-0002-1534-4063 Elbert Geuze ^b https://orcid.org/0000-0003-3479-2379 Andrei Irimia ^b https://orcid.org/0000-0002-9254-9388 Sophia I. Thomopoulos ^b https://orcid.org/0000-0002-0046-4070 Ashley L. Ware ^b https://orcid.org/0000-0002-8390-4915

REFERENCES

- Abdul-Rahman, M. F., Qiu, A., & Sim, K. (2011). Regionally specific white matter disruptions of fornix and cingulum in schizophrenia. *PLoS One*, *6*, e18652.
- Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. J R Stat Soc Series B Stat Methodol, 57, 289–300.
- Bettcher, B. M., Mungas, D., Patel, N., Elofson, J., Dutt, S., Wynn, M., Watson, C. L., Stephens, M., Walsh, C. M., & Kramer, J. H. (2016). Neuroanatomical substrates of executive functions: Beyond prefrontal structures. *Neuropsychologia*, 85, 100–109.
- Bigler, E. D. (2007). Anterior and middle cranial fossa in traumatic brain injury: Relevant neuroanatomy and neuropathology in the study of neuropsychological outcome. *Neuropsychology*, 21, 515–531.
- Bolzenius, J. D., Velez, C. S., Lewis, J. D., Bigler, E. D., Wade, B. S. C., Cooper, D. B., Kennedy, J. E., Reid, M. W., Ritter, J. L., York, G. E., & Tate, D. F. (2018). Diffusion imaging findings in US Service members with mild traumatic brain injury and posttraumatic stress disorder. *The Journal of Head Trauma Rehabilitation*, 33, 393–402.

- Bonnelle, V., Leech, R., Kinnunen, K. M., Ham, T. E., Beckmann, C. F., De Boissezon, X., Greenwood, R. J., & Sharp, D. J. (2011). Default mode network connectivity predicts sustained attention deficits after traumatic brain injury. *Journal of Neuroscience*, 31, 13442– 13451.
- Bouchard, H. C., Sun, D., Dennis, E. L., Newsome, M. R., Disner, S. G., Elman, J., Silva, A., Velez, C., Irimia, A., Davenport, N. D., Sponheim, S. R., Franz, C. E., Kremen, W. S., Coleman, M. J., Williams, M. W., Geuze, E., Koerte, I. K., Shenton, M. E., Adamson, M. M., ... Morey, R. A. (2022). Age-dependent white matter disruptions after military traumatic brain injury: Multivariate analysis results from ENIGMA brain injury. *Human Brain Mapping*, *43*, 2653– 2667. https://doi.org/10.1002/hbm.25811
- Budisavljevic, S., Kawadler, J. M., Dell'Acqua, F., Rijsdijk, F. V., Kane, F., Picchioni, M., McGuire, P., Toulopoulou, T., Georgiades, A., Kalidindi, S., Kravariti, E., Murray, R. M., Murphy, D. G., Craig, M. C., & Catani, M. (2016). Heritability of the limbic networks. *Social Cognitive* and Affective Neuroscience, 11, 746–757.

Caeyenberghs, K., Metzler-Baddeley, C., Foley, S., & Jones, D. K. (2016). Dynamics of the human structural connectome underlying working memory training. *The Journal of Neuroscience*, 36, 4056–4066.

- Chatelin, S., Deck, C., Renard, F., Kremer, S., Heinrich, C., Armspach, J.-P., & Willinger, R. (2011). Computation of axonal elongation in head trauma finite element simulation. *Journal of the Mechanical Behavior of Biomedical Materials*, 4, 1905–1919.
- Davenport, N. D., Lim, K. O., Armstrong, M. T., & Sponheim, S. R. (2012). Diffuse and spatially variable white matter disruptions are associated with blast-related mild traumatic brain injury. *NeuroImage*, 59, 2017– 2024.
- Davenport, N. D., Lim, K. O., & Sponheim, S. R. (2015). White matter abnormalities associated with military PTSD in the context of blast TBI. Human Brain Mapping, 36, 1053–1064.
- Dennis, E. L., Baron, D., Bartnik-Olson, B., Caeyenberghs, K., Esopenko, C., Hillary, F. G., Kenney, K., Koerte, I. K., Lin, A. P., Mayer, A. R., Mondello, S., Olsen, A., Thompson, P. M., Tate, D. F., & Wilde, E. A. (2020). ENIGMA brain injury: Framework, challenges, and opportunities. *Human Brain Mapping*, 43, 149–166. https://doi.org/10.1002/ hbm.25046
- Dennis, E. L., Caeyenberghs, K., Hoskinson, K. R., Merkley, T. L., Suskauer, S. J., Asarnow, R. F., Babikian, T., Bartnik-Olson, B., Bickart, K., Bigler, E. D., Ewing-Cobbs, L., Figaji, A., Giza, C. C., Goodrich-Hunsaker, N. J., Hodges, C. B., Hovenden Aa, E. S., Irimia, A., Königs, M., Levin, H. S., ... Wilde, E. A. (2021). White matter disruption in pediatric traumatic brain injury: Results from ENIGMA pediatric moderate to severe traumatic brain injury. *Neurology*, *97*, e298–e309. https://doi.org/10.1212/WNL.000000000012222
- Dennis, E. L., Disner, S. G., Fani, N., Salminen, L. E., Logue, M., Clarke, E. K., Haswell, C. C., Averill, C. L., Baugh, L. A., Bomyea, J., Bruce, S. E., Cha, J., Choi, K., Davenport, N. D., Densmore, M., du Plessis, S., Forster, G. L., Frijling, J. L., Gonenc, A., ... Morey, R. A. (2019). Altered white matter microstructural organization in posttraumatic stress disorder across 3047 adults: Results from the PGC-ENIGMA PTSD consortium. *Molecular Psychiatry*, *26*, 4315–4330. https://doi.org/10. 1038/s41380-019-0631-x
- Gläscher, J., Adolphs, R., Damasio, H., Bechara, A., Rudrauf, D., Calamia, M., Paul, L. K., & Tranel, D. (2012). Lesion mapping of cognitive control and value-based decision making in the prefrontal cortex. *Proceedings of the National Academy of Sciences of the United States of America*, 109, 14681–14686.
- Gómez-Gastiasoro, A., Zubiaurre-Elorza, L., Peña, J., Ibarretxe-Bilbao, N., Rilo, O., Schretlen, D. J., & Ojeda, N. (2019). Altered frontal white matter asymmetry and its implications for cognition in schizophrenia: A tractography study. *Neuroimage Clinical*, 22, 101781.
- Gong, G., Jiang, T., Zhu, C., Zang, Y., Wang, F., Xie, S., Xiao, J., & Guo, X. (2005). Asymmetry analysis of cingulum based on scale-invariant

1898 WILEY-

parameterization by diffusion tensor imaging. Human Brain Mapping, 24, 92-98.

- Hayes, J. P., Miller, D. R., Lafleche, G., Salat, D. H., & Verfaellie, M. (2015). The nature of white matter abnormalities in blast-related mild traumatic brain injury. *Neuroimage Clinical*, *8*, 148–156.
- Helmick, K. M., Spells, C. A., Malik, S. Z., Davies, C. A., Marion, D. W., & Hinds, S. R. (2015). Traumatic brain injury in the US military: Epidemiology and key clinical and research programs. *Brain Imaging and Behavior*, 9, 358–366.
- Jahanshad, N., Kochunov, P. V., Sprooten, E., Mandl, R. C., Nichols, T. E., Almasy, L., Blangero, J., Brouwer, R. M., Curran, J. E., de Zubicaray, G. I., Duggirala, R., Fox, P. T., Hong, L. E., Landman, B. A., Martin, N. G., McMahon, K. L., Medland, S. E., Mitchell, B. D., Olvera, R. L., ... Glahn, D. C. (2013). Multi-site genetic analysis of diffusion images and voxelwise heritability analysis: A pilot project of the ENIGMA-DTI working group. *Neuro-Image*, 81, 455–469.
- Jahanshad, N., Lee, A., Barysheva, M., & McMahon, K. (2010). Genetic influences on brain asymmetry: A DTI study of 374 twins and siblings. *NeuroImage*, 52(2), 455–469.
- Jorge, R. E., Acion, L., White, T., Tordesillas-Gutierrez, D., Pierson, R., Crespo-Facorro, B., & Magnotta, V. A. (2012). White matter abnormalities in veterans with mild traumatic brain injury. *The American Journal* of Psychiatry, 169, 1284–1291.
- Kelly, S., van Velzen, L., Veltman, D., Thompson, P., Jahanshad, N., Schmaal, L., & ENIGMA-MDD-DTI Working Group. (2017). White matter microstructural differences in major depression: Meta-analytic findings from enigma-MDD DTI. *Biol Psychiatry*, 81, S381.
- Kennedy, E., Dennis, E. L., Lindsey, H. M., deRoon-Cassini, T., Du Plessis, S., Fani, N., Kaufman, M. L., Koen, N., Larson, C. L., Laskowitz, S., Lebois, L. A. M., Morey, R. A., Newsome, M. R., Palermo, C., Pastorek, N. J., Powers, A., Scheibel, R., Seedat, S., Seligowski, A., ... Tate, D. F. (2022). Harmonizing PTSD severity scales across instruments and sites. *Neuropsychology*. https://doi.org/10.1037/ neu0000823
- Kinnunen, K. M., Greenwood, R., Powell, J. H., Leech, R., Hawkins, P. C., Bonnelle, V., Patel, M. C., Counsell, S. J., & Sharp, D. J. (2011). White matter damage and cognitive impairment after traumatic brain injury. *Brain*, 134, 449–463.
- Kochunov, P., Hong, L. E., Dennis, E. L., Morey, R. A., Tate, D. F., Wilde, E. A., Logue, M., Kelly, S., Donohoe, G., Favre, P., Houenou, J., Ching, C. R. K., Holleran, L., Andreassen, O. A., Velzen, L. S., Schmaal, L., Villalón-Reina, J. E., Bearden, C. E., Piras, F., ... Jahanshad, N. (2020). ENIGMA-DTI: Translating reproducible white matter deficits into personalized vulnerability metrics in crossdiagnostic psychiatric research. *Human Brain Mapping*, 7, 1.
- Kochunov, P., Williamson, D. E., Lancaster, J., Fox, P., Cornell, J., Blangero, J., & Glahn, D. C. (2012). Fractional anisotropy of water diffusion in cerebral white matter across the lifespan. *Neurobiology of Aging*, 33, 9–20.
- Kraus, M. F., Susmaras, T., Caughlin, B. P., Walker, C. J., Sweeney, J. A., & Little, D. M. (2007). White matter integrity and cognition in chronic traumatic brain injury: A diffusion tensor imaging study. *Brain*, 130, 2508–2519.
- Lebel, C., Gee, M., Camicioli, R., Wieler, M., Martin, W., & Beaulieu, C. (2012). Diffusion tensor imaging of white matter tract evolution over the lifespan. *NeuroImage*, 60, 340–352.
- Levin, H. S., Wilde, E., Troyanskaya, M., Petersen, N. J., Scheibel, R., Newsome, M., Radaideh, M., Wu, T., Yallampalli, R., Chu, Z., & Li, X. (2010). Diffusion tensor imaging of mild to moderate blast-related traumatic brain injury and its sequelae. *Journal of Neurotrauma*, 27, 683–694.
- Li, J., & Ji, L. (2005). Adjusting multiple testing in multilocus analyses using the eigenvalues of a correlation matrix. *Heredity*, *95*, 221–227.

- Lochner, C., Fouché, J.-P., du Plessis, S., Spottiswoode, B., Seedat, S., Fineberg, N., Chamberlain, S. R., & Stein, D. J. (2012). Evidence for fractional anisotropy and mean diffusivity white matter abnormalities in the internal capsule and cingulum in patients with obsessive-compulsive disorder. *Journal of Psychiatry & Neuroscience*, 37, 193–199.
- Luna, F. G., Lupiáñez, J., & Martín-Arévalo, E. (2021). Microstructural white matter connectivity underlying the attentional networks system. *Beha*vioural Brain Research, 401, 113079.
- Mac Donald, C. L., Johnson, A. M., Cooper, D., Nelson, E. C., Werner, N. J., Shimony, J. S., Snyder, A. Z., Raichle, M. E., Witherow, J. R., Fang, R., Flaherty, S. F., & Brody, D. L. (2011). Detection of blast-related traumatic brain injury in U.S. military personnel. *The New England Journal* of *Medicine*, 364, 2091–2100.
- Mac Donald, C. L., Johnson, A. M., Wierzechowski, L., Kassner, E., Stewart, T., Nelson, E. C., Werner, N. J., Adam, O. R., Rivet, D. J., Flaherty, S. F., Oh, J. S., Zonies, D., Fang, R., & Brody, D. L. (2017). Outcome trends after US military concussive traumatic brain injury. *Journal of Neurotrauma*, 34, 2206–2219.
- Martindale, S. L., Epstein, E. L., Taber, K. H., VA Mid-Atlantic MIRECC Workgroup, & Rowland, J. A. (2018). Behavioral and health outcomes associated with deployment and nondeployment Acquisition of Traumatic Brain Injury in Iraq and Afghanistan veterans. Archives of Physical Medicine and Rehabilitation, 99, 2485–2495.
- Martindale, S. L., Ord, A. S., Lad, S. S., Miskey, H. M., Taber, K. H., & Rowland, J. A. (2021). Differential effects of deployment and nondeployment mild TBI on neuropsychological outcomes. *Rehabilitation Psychology*, 66, 128–138.
- Maruta, J., Mallott, J. M., Sulioti, G., Ghajar, J., Palacios, E. M., & Mukherjee, P. (2020). Concussion disrupts Normal brain White matter microstructural symmetry. *Frontiers in Neurology*, 11, 548220.
- Metzler-Baddeley, C., Jones, D. K., Steventon, J., Westacott, L., Aggleton, J. P., & O'Sullivan, M. J. (2012). Cingulum microstructure predicts cognitive control in older age and mild cognitive impairment. *The Journal of Neuroscience*, 32, 17612–17619.
- Military Health System. (n.d.). DOD TBI Worldwide Numbers. https:// health.mil/Military-Health-Topics/Centers-of-Excellence/Traumatic-Brain-Injury-Center-of-Excellence/DOD-TBI-Worldwide-Numbers
- Niogi, S. N., Mukherjee, P., Ghajar, J., Johnson, C., Kolster, R. A., Sarkar, R., Lee, H., Meeker, M., Zimmerman, R. D., Manley, G. T., & McCandliss, B. D. (2008). Extent of microstructural white matter injury in postconcussive syndrome correlates with impaired cognitive reaction time: A 3T diffusion tensor imaging study of mild traumatic brain injury. *AJNR. American Journal of Neuroradiology*, *29*, 967–973.
- Ocklenburg, S., Friedrich, P., Güntürkün, O., & Genç, E. (2016). Intrahemispheric white matter asymmetries: The missing link between brain structure and functional lateralization? *Reviews in the Neurosciences*, 27, 465–480.
- Ocklenburg, S., Hugdahl, K., & Westerhausen, R. (2013). Structural white matter asymmetries in relation to functional asymmetries during speech perception and production. *NeuroImage*, 83, 1088–1097.
- Park, H.-J., Westin, C.-F., Kubicki, M., Maier, S. E., Niznikiewicz, M., Baer, A., Frumin, M., Kikinis, R., Jolesz, F. A., McCarley, R. W., & Shenton, M. E. (2004). White matter hemisphere asymmetries in healthy subjects and in schizophrenia: A diffusion tensor MRI study. *NeuroImage*, 23, 213–223.
- Rutgers, D. R., Toulgoat, F., Cazejust, J., Fillard, P., Lasjaunias, P., & Ducreux, D. (2008). White matter abnormalities in mild traumatic brain injury: A diffusion tensor imaging study. AJNR. American Journal of Neuroradiology, 29, 514–519.
- Sánchez-Cubillo, I., Periáñez, J. A., Adrover-Roig, D., Rodríguez-Sánchez, J. M., Ríos-Lago, M., Tirapu, J., & Barceló, F. (2009). Construct validity of the trail making test: Role of task-switching, working memory, inhibition/interference control, and visuomotor abilities. *Journal of the International Neuropsychological Society*, 15, 438–450.

- Sibilia, F., Kehoe, E. G., Farrell, D., Kerskens, C., O'Neill, D., McNulty, J. P., Mullins, P., & Bokde, A. L. W. (2017). Aging-related microstructural alterations along the length of the cingulum bundle. *Brain Connectivity.*, 7, 366–372. https://doi.org/10.1089/brain. 2017.0493
- Smith, S. M., Jenkinson, M., Johansen-Berg, H., Rueckert, D., Nichols, T. E., Mackay, C. E., Watkins, K. E., Ciccarelli, O., Cader, M. Z., Matthews, P. M., & Behrens, T. E. (2006). Tract-based spatial statistics: Voxelwise analysis of multi-subject diffusion data. *NeuroImage*, 31, 1487–1505.
- Song, J. W., Mitchell, P. D., Kolasinski, J., Ellen Grant, P., Galaburda, A. M., & Takahashi, E. (2015). Asymmetry of White matter pathways in developing human brains. *Cerebral Cortex*, 25, 2883–2893.
- Stein, M. B., Kessler, R. C., Heeringa, S. G., Jain, S., Campbell-Sills, L., Colpe, L. J., Fullerton, C. S., Nock, M. K., Sampson, N. A., Schoenbaum, M., Sun, X., Thomas, M. L., Ursano, R. J., & Army STARRS collaborators. (2015). Prospective longitudinal evaluation of the effect of deployment-acquired traumatic brain injury on posttraumatic stress and related disorders: Results from the Army study to assess risk and resilience in servicemembers (Army STARRS). *The American Journal of Psychiatry*, 172, 1101–1111.
- Stein, M. B., Ursano, R. J., Campbell-Sills, L., Colpe, L. J., Fullerton, C. S., Heeringa, S. G., Nock, M. K., Sampson, N. A., Schoenbaum, M., Sun, X., Jain, S., & Kessler, R. C. (2016). Prognostic indicators of persistent post-concussive symptoms after deployment-related mild traumatic brain injury: A prospective longitudinal study in U.S. Army Soldiers. *Journal of Neurotrauma*, *33*, 2125–2132.
- Sugiyama, K., Kondo, T., Oouchida, Y., Suzukamo, Y., Higano, S., Endo, M., Watanabe, H., Shindo, K., & Izumi, S.-I. (2009). Clinical utility of diffusion tensor imaging for evaluating patients with diffuse axonal injury and cognitive disorders in the chronic stage. *Journal of Neurotrauma*, 26, 1879–1890.
- Taber, K. H., Hurley, R. A., Haswell, C. C., Rowland, J. A., Hurt, S. D., Lamar, C. D., & Morey, R. A. (2015). White matter compromise in veterans exposed to primary blast forces. *The Journal of Head Trauma Rehabilitation*, 30, E15–E25.
- Takao, H., Abe, O., Yamasue, H., Aoki, S., Sasaki, H., Kasai, K., Yoshioka, N., & Ohtomo, K. (2011). Gray and white matter asymmetries in healthy individuals aged 21-29 years: A voxel-based morphometry and diffusion tensor imaging study. *Human Brain Mapping*, 32, 1762–1773.
- Takao, H., Hayashi, N., & Ohtomo, K. (2011). White matter asymmetry in healthy individuals: A diffusion tensor imaging study using tract-based spatial statistics. *Neuroscience*, 193, 291–299.
- Takao, H., Hayashi, N., & Ohtomo, K. (2013). White matter microstructure asymmetry: Effects of volume asymmetry on fractional anisotropy asymmetry. *Neuroscience*, 231, 1–12.
- Tate, D. F., Dennis, E. L., Adams, J. T., Adamson, M., Belanger, H. G., Bigler, E. D., Bouchard, H. C., Clark, A. L., Delano-Wood, L. M., Disner, S. G., Eapen, B. C., Franz, C. E., Geuze, E., Goodrich-Hunsaker, N. J., Han, K., Hayes, J. P., Hinds, S. R., 2nd, Hodges, C. B., Hovenden, E. S., ... Wilde, E. A. (2021). Coordinating global multi-site studies of military-relevant traumatic brain injury: Opportunities, challenges, and harmonization guidelines. *Brain Imaging Behavior*, 15(2), 585–613.
- Thiebaut de Schotten, M., Ffytche, D. H., Bizzi, A., Dell'Acqua, F., Allin, M., Walshe, M., Murray, R., Williams, S. C., Murphy, D. G. M., & Catani, M. (2011). Atlasing location, asymmetry and inter-subject variability of white matter tracts in the human brain with MR diffusion tractography. *NeuroImage*, 54, 49–59.
- Thompson, P. M., Jahanshad, N., Ching, C. R. K., Salminen, L. E., Thomopoulos, S. I., Bright, J., Baune, B. T., Bertolín, S., Bralten, J., Bruin, W. B., Bülow, R., Chen, J., Chye, Y., Dannlowski, U., de

Kovel, C. G. F., Donohoe, G., Eyler, L. T., Faraone, S. V., Favre, P., ... Consortium, E. N. I. G. M. A. (2020). ENIGMA and global neuroscience: A decade of large-scale studies of the brain in health and disease across more than 40 countries. *Translational Psychiatry*, *10*, 100.

- Toga, A. W., & Thompson, P. M. (2003). Mapping brain asymmetry. Nature Reviews. Neuroscience, 4, 37–48.
- Trivedi, R., Agarwal, S., Rathore, R. K. S., Saksena, S., Tripathi, R. P., Malik, G. K., Pandey, C. M., & Gupta, R. K. (2009). Understanding development and lateralization of major cerebral fiber bundles in pediatric population through quantitative diffusion tensor tractography. *Pediatric Research*, 66, 636–641.
- Tsuchida, A., Laurent, A., Crivello, F., Petit, L., Pepe, A., Beguedou, N., Debette, S., Tzourio, C., & Mazoyer, B. (2021). Changes in regional white matter volumetry and microstructure during the postadolescence period: A cross-sectional study of a cohort of 1,713 university students. *Frontiers in Systems*, 15, 692152.
- Vakhtin, A. A., Zhang, Y., Wintermark, M., Massaband, P., Robinson, M. T., Ashford, J. W., & Furst, A. J. (2020). White matter asymmetry: A reflection of pathology in traumatic brain injury. *Journal of Neurotrauma*, 37, 373–381.
- Ware, A. L., Biekman, B., Hachey, R., MacLeod, M., Bird, W., Pathak, S., Clarke, E., Borrasso, A., Puccio, A. M., Glavin, K., Pomiecko, K., Moretti, P., Beers, S. R., Levin, H. S., Schneider, W., Okonkwo, D. O., & Wilde, E. A. (2019). A preliminary high-definition fiber tracking study of the executive control network in blast-induced traumatic brain injury. *Journal of Neurotrauma*, 36, 686–701.
- Wilde, E. A., Bouix, S., Tate, D. F., Lin, A. P., Newsome, M. R., Taylor, B. A., Stone, J. R., Montier, J., Gandy, S. E., Biekman, B., Shenton, M. E., & York, G. (2015). Advanced neuroimaging applied to veterans and service personnel with traumatic brain injury: State of the art and potential benefits. *Brain Imaging and Behavior*, *9*, 367–402.
- Wilde, E. A., Dennis, E. L., & Tate, D. F. (2021). The ENIGMA brain injury working group: Approach, challenges, and potential benefits. *Brain Imaging Behavior*, 15(2), 465–474.
- Wu, T. C., Wilde, E. A., Bigler, E. D., Yallampalli, R., McCauley, S. R., Troyanskaya, M., Chu, Z., Li, X., Hanten, G., Hunter, J. V., & Levin, H. S. (2010). Evaluating the relationship between memory functioning and cingulum bundles in acute mild traumatic brain injury using diffusion tensor imaging. *Journal of Neurotrauma*, *27*, 303–307.
- Yeh, P.-H., Wang, B., Oakes, T. R., French, L. M., Pan, H., Graner, J., Liu, W., & Riedy, G. (2014). Postconcussional disorder and PTSD symptoms of military-related traumatic brain injury associated with compromised neurocircuitry. *Human Brain Mapping*, 35, 2652–2673.
- Yin, X., Han, Y., Ge, H., Xu, W., Huang, R., Zhang, D., Xu, J., Fan, L., Pang, Z., & Liu, S. (2013). Inferior frontal white matter asymmetry correlates with executive control of attention. *Human Brain Mapping*, 34, 796–813.
- Yurgelun-Todd, D. A., Bueler, C. E., McGlade, E. C., Churchwell, J. C., Brenner, L. A., & Lopez-Larson, M. P. (2011). Neuroimaging correlates of traumatic brain injury and suicidal behavior. *The Journal of Head Trauma Rehabilitation*, 26, 276–289.
- Yurgil, K. A., Barkauskas, D. A., Vasterling, J. J., Nievergelt, C. M., Larson, G. E., Schork, N. J., Litz, B. T., Nash, W. P., Baker, D. G., & Marine Resiliency Study Team. (2014). Association between traumatic brain injury and risk of posttraumatic stress disorder in active-duty marines. JAMA Psychiatry, 71, 149–157.
- Zhang, L., & Gennarelli, T. (2011). Mathematical modeling of diffuse brain injury: Correlations of foci and severity of brain strain with clinical symptoms and pathology IRCOBI impact Biomechanics, 315–324. http://www.ircobi.org/wordpress/downloads/irc0111/2011/Session1/ 13.pdf
- Zhao, W., Bartsch, A., Benzel, E., Miele, V., & Stemper, B. D. (2019). Regional brain injury vulnerability in football from two finite element

models of the human head. IRCOBI Florence. http://www.ircobi.org/ wordpress/downloads/irc19/pdf-files/89.pdf

Zhou, Z., Li, X., Domel, A. G., Dennis, E. L., Georgiadis, M., Liu, Y., Raymond, S. J., Grant, G., Kleiven, S., Camarillo, D., & Zeineh, M. (2022). The presence of the temporal horn exacerbates the vulnerability of hippocampus during head impacts. *Frontiers in Bioengineering and Biotechnology*, 10, 754344.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article. How to cite this article: Dennis, E. L., Newsome, M. R., Lindsey, H. M., Adamson, M., Austin, T. A., Disner, S. G., Eapen, B. C., Esopenko, C., Franz, C. E., Geuze, E., Haswell, C., Hinds, S. R. II, Hodges, C. B., Irimia, A., Kenney, K., Koerte, I. K., Kremen, W. S., Levin, H. S., Morey, R. A., ... Wilde, E. A. (2023). Altered lateralization of the cingulum in deployment-related traumatic brain injury: An ENIGMA military-relevant brain injury study. *Human Brain Mapping*, 44(5), 1888–1900. <u>https://doi.org/10.1002/hbm.26179</u>