UC Riverside

International Organization of Citrus Virologists Conference Proceedings (1957-2010)

Title

Molecular Characterization of Citrus tristeza virus Isolates from Epirus (Greece)

Permalink

<https://escholarship.org/uc/item/4gg24059>

Journal

International Organization of Citrus Virologists Conference Proceedings (1957-2010),
17(17)

ISSN

2313-5123

Authors

Vovlas, C.
Barbarossa, L.

Publication Date

2010

DOI

10.5070/C54gg24059

Peer reviewed

Molecular Characterization of *Citrus tristeza virus* Isolates from Epirus (Greece)

L. Barbarossa¹ and C. Vovlas²

¹Istituto di Virologia Vegetale del CNR, Sezione di Bari, Via Amendola 165/A, 70126 Bari, Italy

²Dipartimento di Protezione delle Piante e Microbiologia Applicata, Università degli Studi, Via Amendola 165/A, 70126 Bari, Italy

ABSTRACT. Since June 2000, *Citrus tristeza virus* (CTV) has been reported in Greece including Crete, in some cases introduced accidentally in budwood from Spain. Here we present a preliminary molecular characterization of some CTV isolates based on amplification of the coat protein gene and use of RFLP. Samples were collected from symptomless Washington navel and Navelina trees about 20 yr old in commercial groves in Epirus (western Greece), grafted on sour orange. Analysis of the CP gene sequences showed 98-99% identity to the T30 isolate from Florida and the Spanish isolate T385, both mild isolates. RFLP analysis confirmed the characterization of the Greek isolates as mild. A phylogenetic tree drawn using Clustal X and TREEVIEW, and including a worldwide variety of CTV isolates, again confirmed close identity with T30 and T385, bringing additional evidence that the Greek isolates came from Spain, followed by local spread of the virus, largely through human activity.

In 2005 a survey for virus and virus-like diseases conducted in the main citrus-growing areas of Epirus (western Greece) revealed the presence of *Citrus tristeza virus* (CTV) (3). In June 2000, the virus was detected for the first time in Argolis (north-east Peloponnese) and in Chania (Crete), due to accidental introduction of CTV-infected budwood from Spain (4). However little is known about the characteristics of Greek isolates, in spite of the recognized importance of isolate identities in defining disease management and control strategies in order to avoid potential epidemic outbreaks. The present study is a preliminary molecular characterization of some CTV isolates detected in Arta valley, based on amplification of the the coat protein (CP) gene by RT-PCR jointly with the use of RFLP (5).

Isolates were collected from Washington navel and Navelina orange trees, about 20-yr-old, inspected in commercial groves. Plants were symptomless although grafted on sour orange rootstock. Samples, collected from randomly selected trees, were tested by dot-blot hybridization assay to detect CTV-infected plants (2). cDNA synthesis and cloning of the CP gene were done as previously described (3). Clones were

automatically sequenced (MWG Biotech, Germany) and the complete CP sequence of isolates CTV-G17, CTV-G9, CTV-G73 and CTV-G74 was deposited in the EMBL database under accession numbers AM406802, AM406803, AM746968 and AM746969, respectively. They were compared to existing sequences using the Blast (NCBI) program (1). Analysis of the CP gene nucleotide sequences, compared to published sequences, showed 98-99% nt identity to the T30 isolate from Florida (accession no. AF260651) and the Spanish isolate T385 (accession no. Y18420). Restriction analysis of the CP gene, based on *HinfI* and *RsaI* enzyme sites, revealed considerable resemblance to mild CTV isolates. When classified by the *HinfI* RFLP pattern (5), the Greek isolates fell in group 2 (nt positions 73, 111, 410, 501), which also contained mild isolates T30 and T385. When classified by the *RsaI* products, group 2 with three restriction sites (nt positions 145, 312, 576) contained the Greek isolates and also mild isolates T30 and T385. According to Mendoza et al. (6), the CP gene of CTV strains designated as mild strains does not contain the *HaeIII* and *KpnI* restriction sites; the absence of these sites in the CP gene sequence of CTV-G17, CTV-G9, CTV-

G73 and CTV-G74 confirmed their characterization as mild isolates.

Using Clustal X (9), a multiple alignment was generated with the CP gene sequences of the Greek CTV isolates and several worldwide CTV isolates, showing different biological characteristics (Table 1). The phylogenetic tree, displayed using TREEVIEW version 1.6.6 (7), showed that the Greek isolates were closely related to known mild isolates and very similar to T385 and T30 strains (Fig. 1). Interestingly, their deduced CP amino acid

sequence contained, at positions 122 and 124, isoleucine (I) and tyrosine (Y) respectively; these are present at the same positions in the CP amino acid sequence of most mild CTV isolates, including T385 and T30 (8). These results are additional evidence that introduction of CTV in Greece has occurred due to infected propagative material imported from Spain (4), and suggest further local spread of the virus, and the strong influence of human activity in its dissemination.

TABLE 1
CITRUS TRISTEZA VIRUS (CTV) COAT PROTEIN GENE SEQUENCES SELECTED
FROM

GEN BANK AND USED IN SEQUENCE AND PHYLOGENETIC ANALYSIS		
Isolates	Country	Accession number
B35	California, USA	L12175
CTV-G9	Greece	AM406803
CTV-G17	Greece	AM406802
CTV-G73	Greece	AM746968
CTV-G74	Greece	AM746969
CTV-0032	Italy	AJ518842
TN-2	Iran	AY707460
T6-31	Montenegro	AY764154
T30	Florida, USA	AF260651
T385	Spain	Y18420
2-93	Portugal	AF184116
CTV-B	India	AF501867
Cheju Island	South Korea	AF249279
Cy95-14	Cyprus	EF491668
Jordan	Jordan	AY550252
NuagA	Japan	AB046398
Qaha	Egypt	AY340974
SY568	California, USA	AF001623
T36	Florida, USA	U16304
VT	Israel	U56902
13C	Portugal	AF184113
19-121	Portugal	AF184114
28C	Portugal	AF184118
446-6	Croatia	AY791842

Fig. 1 Dendrogram showing the genetic relationship among CP gene sequences of Greek and other *Citrus tristeza virus* (CTV) isolates (see Table 1). Scale bar indicates changes per nucleotide.

LITERATURE CITED

1. Altschul, S. F., F. Stephen, W. Gish, W. Miller, E. W. Myers, and D. J. Lipman
1990. Basic logic alignment search tool. *J. Mol. Biol.* 215: 403-410.
2. Barbarossa, L., and V. Savino
2006. Sensitive and specific digoxigenin-labeled RNA probes for routine detection of *Citrus tristeza virus* by dot-blot hybridisation. *J. Phytopathol.* 154: 329-335.
3. Barbarossa, L., G. Loconsole, and C. Vovlas
2007. Virus and virus-like diseases of citrus in Epirus. *J. Plant Pathol.* 89: 273-276.
4. Dimou, D., J. Drosopoulou, E. Moschos, C. Spanou, and P. Dermatas
2002. First report of *Citrus tristeza virus* (CTV) in Greece. In: *Proc. 15th Conf. IOCV*, 78-82. IOCV, Riverside, CA.
5. Gillings, M., P. Broadbent, J. Indsto, and R. Lee
1993. Characterisation of isolates and strains of *Citrus tristeza closterovirus* using analysis of the coat protein gene amplified by the polymerase chain reaction. *J. Virol. Methods* 44: 305-317.
6. Mendoza, A., C. Salazar, O. Alvarado, M. A. Cruz, M. A. Rodríguez, and H. A. Barrera Saldaña

2005. Molecular differentiation of mild and severe *Citrus tristeza virus* isolates in Mexico. In: *Proc. 16th Conf. IOCV*, 61-67. IOCV, Riverside, CA.
7. Page, R. D. M.
1996. TREEVIEW: An application to display phylogenetic trees on personal computers. *Comput. Appl. Biosci.* 12: 357-358.
8. Roy, A., P. Ramachandran, and R. H. Brlansky
2003. Grouping and comparison of Indian citrus tristeza virus isolates based on coat protein gene sequences and restriction analysis patterns. *Arch. Virol.* 148: 707-722.
9. Thompson, J. D., T. J. Gibson, F. Plewniak, F. Jeanmougin, and D. G. Higgins
1997. The ClustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* 24: 4876-4882.