

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

PROPERTIES OF THE  $w$  MESON FROM  $K^- + p \rightarrow A + w$

### Permalink

<https://escholarship.org/uc/item/4gr580xv>

### Authors

Murray, Joseph J.  
Ferro-Luzzi, Massimiliano  
Huwe, Darrell O.  
[et al.](#)

### Publication Date

1963-03-14

University of California  
Ernest O. Lawrence  
Radiation Laboratory

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545*

PROPERTIES OF THE  $\omega$  MESON  
FROM  $K^- + p \rightarrow \Lambda + \omega$

Berkeley, California

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Research and Development

UCRL-10723  
UC-34 Physics  
TID-4500 (19th Ed.)

UNIVERSITY OF CALIFORNIA  
Lawrence Radiation Laboratory  
Berkeley, California

Contract No. W-7405-eng-48

PROPERTIES OF THE  $\omega$  MESON FROM  $K^- + p \rightarrow \Lambda + \omega$

Joseph J. Murray, Jr., Massimiliano Ferro-Luzzi,  
Darrell O. Huwe, Janice B. Shafer, Frank T. Solmitz,  
and M. Lynn Stevenson

March 14, 1963

Printed in USA. Price 50 cents. Available from the  
Office of Technical Services  
U. S. Department of Commerce  
Washington 25, D.C.

PROPERTIES OF THE  $\omega$  MESON FROM  $K^- + p \rightarrow \Lambda + \omega$

Joseph J. Murray, Jr., Massimiliano Ferro-Luzzi,  
Darrell O. Huwe, Janice B. Shafer, Frank T. Solmitz,  
and M. Lynn Stevenson

Lawrence Radiation Laboratory and Department of Physics  
University of California, Berkeley, California

March 14, 1963

ABSTRACT

Studies of the reaction  $K^- + p \rightarrow \Lambda + \omega$  at 1.51 BeV/c are reported. On the basis of 650  $\omega \rightarrow 3\pi$  decays and 40  $\omega \rightarrow$  neutral decays, the branching ratio of neutral to charged decay is found to be  $0.11 \pm 0.03$ . The width of the  $\omega \rightarrow 3\pi$  decay is determined to be  $4 \pm 4$  MeV by the use of an extrapolation method. Evidence for the rare two-pion decay mode is presented. The production angular distribution for the  $\omega$  peaks slightly forward, as does that of the  $\rho$  meson. Cross sections for  $\omega$  production and subsequent decay into its charged and neutral modes are  $800 \pm 40$  and  $83 \pm 22$   $\mu\text{b}$ , respectively.

PROPERTIES OF THE  $\omega$  MESON FROM  $K^- + p \rightarrow \Lambda + \omega$  †

Joseph J. Murray, Jr., Massimiliano Ferro-Luzzi,  
Darrell O. Huwe, Janice B. Shafer, Frank T. Solmitz,  
and M. Lynn Stevenson

Lawrence Radiation Laboratory and Department of Physics  
University of California, Berkeley, California

March 14, 1963

Events of the type  $K^- + p \rightarrow \Lambda + \omega$  were studied in film obtained with the 72-inch hydrogen bubble chamber utilizing a beam of 1.51-BeV/c  $K^-$  mesons extracted from the Bevatron.<sup>1, 2</sup> Approximately 650 charged  $\omega$  decays and about 40 neutral decays were analyzed.


The following results are reported here:

- a) the branching ratio of neutral to charged (three-pion) decay
- b) a limit on the natural width of the predominant three-pion decay
- c) evidence for the two-pion decay mode.

At an incident momentum of 1.51 BeV/c (total c. m. energy of 2.025 BeV), approximately 1300 events of the type



and a similar number of the type



were analyzed. In addition, some 1400 events of the type




were processed. At present a systematic study is being made of the reaction  $K^- + p \rightarrow \Lambda + \omega$  as a function of  $K^-$  momentum; results will appear in a future publication.

For reactions (1), (2), and (3), an upper limit of 30 was imposed for the  $\chi^2$  value in kinematic fitting of the lambda decay. In ambiguous cases, where the vee decay fitted both  $K^0$  and lambda, a decision was made on the basis of either the goodness of fit of alternate production hypotheses or the ionization of the decay tracks. A production hypothesis [for reaction (1) or (2)] was accepted when its  $\chi^2$  value was less than ten times the number of constraints and the  $\chi^2$  value for any other hypothesis was higher relative to the number of constraints. The separation of  $\Sigma^0 \pi^+ \pi^-$  from the  $\Lambda \pi^+ \pi^-$  events removed about 7% of the  $\Lambda \pi^+ \pi^-$  events, but left a residual of  $\Sigma^0 \pi^+ \pi^-$  events which was only 5% of the  $\Lambda \pi^+ \pi^-$  sample. (For more details, see the earlier report.<sup>1</sup>)

## THREE-PION DECAY MODE

The effective-mass spectrum of the three pions in reaction (2) is shown in Fig. 1. It is apparent from this spectrum that a large fraction of the events result from the two-step process:




For  $\omega$  mass limits of 760 to 810 MeV, about 650  $\omega$ 's are observed above background. The background can be estimated as approximately 15% of the events within these mass limits.

The c. m. angular distribution of the recoil lambda for the events satisfying  $K^- + p \rightarrow \Lambda + \omega$  kinematics is given in Fig. 2. The  $\omega$ 's are produced predominantly in the forward direction.

The cross section for the two-step process (4) has been found by comparing the number of three-pion  $\omega$  decays with the number of  $\tau$  decays in the same film sample. After correction for scanning efficiency (about 90%) and for measurement failures (about 15% loss, mostly in kinematic fitting), this cross section is  $800 \pm 40 \mu\text{b}$ . The process  $K^- + p \rightarrow \Sigma^0 + \omega$  is thought to account for an enhancement on the high-mass side of the  $\omega$  peak in the mass spectrum of Fig. 1. Measurement errors may allow events of this type to fit hypothesis (2). Detailed study of the  $\Sigma^0\omega$  reaction is difficult, since it is not kinematically overdetermined. The number of events of the latter type which give spurious fits to the  $\Lambda\omega$  hypothesis is estimated to be less than 5% of the  $\Lambda\omega$  events.

## NEUTRAL DECAY MODE

The square of the missing neutral mass (or recoil mass), of reaction (3) is shown in Fig. 3. Here also the presence of  $\omega$  production and decay,




is apparent in the enhancement occurring approximately at the square of the  $\omega$  mass. (Instead of the linear mass scale of Fig. 1, the square of the missing mass is used as the basic scale of Fig. 3; the latter quantity is approximately normally distributed, even in the vicinity of zero mass.) Some 40 events are seen above background.

The cross section for the process given in (5) has also been obtained by comparing these events with the number of  $\tau$  decays in the same film. With corrections for scanning efficiency (about 90%) and for measurement failures (about 10% loss in track reconstruction, largely from poor measurement, and about 13% loss in kinematic fitting) the value is  $83 \pm 22 \mu\text{b}$ .

From the above cross sections, the branching ratio of neutral to three-pion  $\omega$  decays (i. e., presumably of  $\pi^0\gamma$  to  $\pi^+\pi^-\pi^0$  final states) is


$$\Gamma(\text{neutral})/\Gamma(3\pi) = 0.11 \pm 0.03. \quad (6)$$





MU-28528

Fig. 1. Three-pion effective mass spectrum in the reaction  $K^- + p \rightarrow \Lambda + \pi^+ + \pi^- + \pi^0$  at 1.51 BeV/c. About 650  $\omega$ 's are seen above background. The width of the  $\omega$  peak is due chiefly to measurement error.



MU-30022

Fig. 2. Production angular distribution for  $K^- + p \rightarrow \Lambda + \omega$ . The slight backward peaking of the  $\Lambda$  indicates that the  $\omega$  is produced preferentially in the forward direction.




Fig. 3. Mass of missing neutrals in the reaction  $K^- + p \rightarrow \Lambda + \text{neutrals}$  at 1.51 BeV/c. About 40 neutral  $\omega$  decays (presumably  $\pi^0 + \gamma$ ) are seen above background.

WIDTH OF THE  $\omega$  MESON

Determination of the true width of the three-pion  $\omega$  decay is difficult. The mass spread due to measurement error alone, called the "resolution function," is determined from the effective-mass errors calculated by our IBM program EXAMIN; these calculations are based on momentum and angle errors determined by our program PACKAGE, which effects the track reconstruction and kinematic fitting of events. The errors on kinematic quantities determined in track reconstruction appear to be underestimated for the particular film sample considered here. The  $\chi^2$  distributions obtained in fitting the events have approximately the correct form, but have median values that are about twice those expected. Underestimation of fitted errors yields a resolution function with too small a width. Thus unfolding the resolution function from the observed mass spectrum gives too large a value for the width of the  $\omega$ . (Without correction of the resolution function, a sample of the best-measured  $\Lambda\omega$  events yields a value of 14 MeV for the over-estimated width.)

In the kinematic fitting of the events, the approximations made in the calculation of momentum and angle errors by the track-reconstruction program appear to cause a scaling of the  $\chi^2$  distributions; that is, all errors seem to have been underestimated by approximately the same amount. Without a good understanding of the causes of these inaccuracies in error estimation, correction cannot be made in a consistent way. In the analysis of the three-pion decays of the omega, it was observed that the experimental  $\chi^2$  values for  $\Lambda\pi^+\pi^-\pi^0$  production compared well with the theoretical distribution if the latter were scaled by a factor of approximately 2.0.

The distribution of errors calculated in the effective mass of any particle system is generally assumed to be Gaussian. This was checked for the  $\omega \rightarrow 3\pi$  events by assuming the natural width of the  $\omega$  to be much smaller than the program-estimated mass errors, which ranged from 3 to more than 15 MeV. A dimensionless or normalized mass spectrum showing the spread of values of the  $3\pi$  effective mass  $M$  about the  $\omega$  mass was obtained by histogramming the events as a function of  $(M-M_\omega)/dM$ . This histogram was found to have a Gaussian form and further to have a variance  $(M-M_\omega)^2/dM^2$  of about 3.0 rather than the value 1.0 expected for correctly estimated errors. The mass spectrum thus was consistent with the  $\chi^2$  distribution in that the errors appeared to be underestimated by a factor of  $\leq \sqrt{3}$  from the former, and by a factor of  $\sqrt{2}$  from the latter.

One possible method of estimating the true  $\omega$  width would be to utilize only events with small mass errors and to correct the resolution function for these events by the " $\chi^2$  scale factor." Because of the complexity of the error calculations, it is not clear that this  $\chi^2$  scale factor is directly applicable to the resolution function.

An alternative method is to divide the data into several samples in accordance with the size of the  $3\pi$  mass errors and to extrapolate to zero measurement error the width values obtained for the several samples. This method does not necessitate quantitative knowledge of the resolution function nor the " $\chi^2$  scale factor", but does demand the use of the relative variation of this scale factor to correct relative values of the resolution functions. It

has the advantage of using more of the available data than the first method and also of providing determination of the absolute value of the scale factor demanded by the resolution function.

The values of effective three-pion mass for the 1361 events of reaction (2) were histogrammed for particular ranges of program-estimated error in the effective mass. The ranges chosen--3 to 5 MeV, 5 to 7 MeV, and 7 to 10 MeV--were narrow enough to permit the assumption that the values of effective mass were Gaussianly distributed with variance equal to the mean value of the squared error in each of the three groups. Events above background in each of the three error-classified samples were fitted with a Gaussian curve. The error on the variance of each of the three Gaussian curves was obtained by using the fact that the variance is distributed like  $\chi^2$  with its degrees of freedom equal to the number of events in the peak less one. This error was increased slightly to account for the uncertainty in background estimation.

The three observed widths result from folding a Breit-Wigner resonance curve with a Gaussian measurement-error distribution. These three experimental widths are shown as a function of the mean measurement error in Fig. 4. Figure 4a is a plot of the squared quantities, and Fig. 4b is a linear plot. If we assume that the scale factor required to correct the measurement errors varies inappreciably over the range of errors represented by the three samples of events, the appropriate form of the curve used for width extrapolation to zero measurement error can be taken from Fig. 5. The latter presents widths for the folding of Breit-Wigner (S-wave form)<sup>4</sup> and Gaussian distribution curves, with various Breit-Wigner widths assumed. (The folded distribution is given by


$$N(t) \propto \int_{-\infty}^{\infty} \left[ x^2 + (\Gamma/2)^2 \right]^{-1} \exp^{-t(x)^2/2(dM)^2} dx.$$

It is evident that the curves of Fig. 5a or 5b have similar slopes, which become greater (Fig. 5a) or smaller (Fig. 5b) in approaching zero.

The experimental-error scale factor could not be assumed to be constant. Examination of the  $\chi^2$  distributions for the three samples indicated that the relative scale factors were 1.0, 0.98, and 0.82 for the squared error  $(dM)^2$  for points (1), (2), and (3), respectively. Corrections for this variation brought the three points closer to a straight line. The slope of this line indicated again that the program-estimated errors are about a factor of  $\sqrt{2}$  smaller than they should be.


Because of the difficulty in representing the curves of Fig. 5 analytically, straight lines have been drawn through the data in Fig. 4a and 4b after correction for scale-factor differences. The intercepts have values of  $\Gamma^2 = 50^{+70}_{-50}$  MeV<sup>2</sup> and  $\Gamma = 2.5^{+3.5}_{-2.5}$  MeV for Fig. 4a and 4b, respectively. Straight-line extrapolations of the theoretical curves were also made in Fig. 5a and 5b; the intercepts of the latter for  $\Gamma = 4.0$  agreed well with the experimental (Fig. 4) values. Thus the value of three-pion decay width afforded by the extrapolation method is

$$\Gamma(\omega \rightarrow 3\pi) = 4 \pm 4 \text{ MeV.} \quad (7)$$



MU-30023

Fig. 4. (a) Square of observed width vs square of estimated mass error for the peak in three samples of events. The dashed curve represents a simple curve through the data. The solid line is a least-squares fit to the data after correction of estimated errors for relative variation in the error "scale factor" (see text for definition). The intercept of the straight-line fit gives an upper limit on the true width of the  $\omega$ . (The intercept would equal the true width if the measurement error width and the true width could be combined Gaussianly; then  $\Gamma^2$  (observed) would equal  $\Gamma^2$  (true) +  $K\Gamma^2$  (meas.) =  $\Gamma^2$  (true) +  $K \langle dM \rangle^2$ , where K represents the measurement-error scale factor. See Fig. 5 for the calculated results of folding Breit-Wigner and Gaussian curves.) (b) Observed width versus r. m. s. measurement error for the  $\omega$  peak in three samples of data. The dashed curve is drawn through the data points. The straight line is a least-squares fit to the data after correction for the error "scale factor." The intercept of the straight line represents a lower limit on the true width of the  $\omega$ . The estimated mass errors of the samples of events used for these figures are (1) 3 to 5 MeV; (2) 5 to 7 MeV; and (3) 7 to 10 MeV.



MU-30024

Fig. 5. (a) Square of observed width vs square of mass error, as determined from many calculated curves representing the folding of a Breit-Wigner resonance curve (of specified width  $\Gamma$ ) with Gaussian mass error curves of various  $\langle (dM)^2 \rangle$  values. (b) Observed width vs mass error for the same calculated curves resulting from the folding of a Breit-Wigner distribution with Gaussian distributions. The Breit-Wigner width appears most significant in the change of slope of the curves as the mass errors approach zero; the slope increases in (a) and decreases in (b) as  $dM$  goes to zero, the change being greater for larger Breit-Wigner widths.

## TWO-PION DECAY

The events of reaction (1) might be expected to show evidence of the well-known resonances,  $Y_1^{*+}$ ,  $Y_1^{*-}$ , and  $\rho$ . The  $Y_1^{*+}$  and  $Y_1^{*-}$  are clearly seen on the Dalitz plot of the square of  $\Lambda\pi^+$  effective-mass of the  $\Lambda\pi^+$  versus that of the  $\Lambda\pi^-$  (see Fig. 6). Diagonal lines at  $M_{\pi\pi} = 700$  and  $800$  MeV on the Dalitz plot indicate the expected peak of the two-pion  $\rho$  resonance; the  $\rho$  is difficult to see because it is a broad resonance, but interference between  $\rho$  and  $Y_1^*$  is evident in the nonuniform population of the  $Y_1^*$  mass bands.

In an attempt to distinguish the  $\rho$ , a portion of the Dalitz plot just inside the  $Y_1^*$  bands was examined. Two strips with  $1430 \text{ MeV} \leq (M_{\Lambda\pi^+} \text{ or } M_{\Lambda\pi^-}) \leq 1510 \text{ MeV}$  (shaded in Fig. 6) were projected onto the 45-degree line which is the two-pion mass axis. Bands rather than the entire sector between the  $Y_1^*$ 's were used to make the contribution of the  $Y_1^*$  tails homogeneous along the pion-pion axis outside the region of  $\rho$ - $Y_1^*$  interference.


The pion-pion mass spectrum resulting from this projection is shown in Fig. 7. A narrow peak is observed near  $M^2(2\pi) = 0.6 \text{ BeV}^2$ , which is very close to the square of the  $\omega$  mass. The number of events in the peak is about 15, while the number of  $\rho$  events plus background events in the same mass interval is also about 15. Hence the sharp spike is about a three-standard-deviation effect. Its central value is  $781 \text{ MeV}$ , and its width is about  $15 \text{ MeV}$  (approximately equal to the resolution function width if corrected for underestimation of measurement errors). It seems very likely that the sharp spike is the result of  $\omega$  decay into two pions, a mode resulting from the electromagnetic mixing of the  $\rho^0$  and  $\omega$  states. (See the discussion of theory below.)

The data have been fitted with a constant background, a P-wave Breit-Wigner resonance curve for the  $\rho$  meson (with  $E_0 = 750 \text{ MeV}$  and with  $\Gamma = 100 \text{ MeV}$ ), and a Gaussian curve for the " $\omega \rightarrow 2\pi$  decay". A good fit was obtained with parameters for the  $\omega$  decay of  $E_0 = 781 \text{ MeV}$  and  $\Gamma = 11 \text{ MeV}$  at half-maximum (mostly measurement error). The total numbers of events assigned to the various hypotheses by this fit are

$$\begin{aligned} N(\omega) &= 15.5 \pm 5.6 \\ N(\rho) &= 93 \pm 19 \\ N(\text{background}) &= 224 \pm 22. \end{aligned} \tag{8}$$


The possibility that the narrow peak may be a statistical fluctuation due entirely to the  $\rho$  decay cannot be completely ruled out. The  $\pi$ - $\pi$  mass spectrum has been fitted between  $300$  and  $900 \text{ MeV}$  with just constant background and the  $\rho$  resonance. The fit gives a  $\chi^2$  value of about  $47$  when  $35$  is expected, in contrast to the value of  $38$  for the above-mentioned fit with  $\omega$  included. A fit between  $660$  and  $850 \text{ MeV}$  gives  $\chi^2 = 12.8$  for  $12$  degrees of freedom without the  $\omega$ , and gives  $\chi^2 = 5.4$  for  $11$  degrees of freedom with the  $\omega$  included (see Fig. 8). However, the peak might conceivably be better explained as a  $\rho$  mass fluctuation if a mass spectrum enhanced at high mass by  $\rho$ - $Y_1^*$  interference were considered. (The narrowness of the peak, if clearly substantiated, would of course rule out this last objection. Calculations indicate that a peak caused by  $\rho$ - $Y_1^*$  interference would be at least  $50$ -MeV wide.)





MU-28877

Fig. 6. Dalitz plot of 1852 events representing the reaction  $K^- + p \rightarrow \Lambda + \pi^+ + \pi^-$  at 1.51 BeV/c. The 45-deg line is the axis for the square of the mass in the  $\pi^+ - \pi^-$  system; lines orthogonal to this axis are shown to indicate 700 and 800 MeV for the  $\pi - \pi$  mass. Bands between 1340 and 1430 MeV are outlined for the  $Y^{*+}$  and  $Y^{*-}$  systems; it can be seen that most of the events cluster within these bands. The  $\pi - \pi$  system was studied by considering two bands parallel to the  $Y^*$  bands, between  $\Lambda - \pi$  masses of 1430 and 1510 MeV. Events within these limits were projected onto the  $\pi - \pi$  mass axis, with the intention of keeping the  $Y^*$  contribution constant as a function of the  $\pi - \pi$  mass.



MU-28529

Fig. 7. Pion-pion mass spectrum resulting from the projection described in Fig. 6. A narrow peak (or a statistical fluctuation) appears superimposed on a broader bump (the  $\rho$  meson) in the region of 780 MeV. The width is comparable with the measurement resolution.




Fig. 8. A portion of the pion-pion mass spectrum shown in Fig. 7. One of the curves represents a fit to the data with a P-wave  $\rho$  meson and constant background; the other curve represents a fit with a P-wave  $\rho$  plus background and also a narrow Gaussian centered at 780 MeV to represent the  $\omega$  meson.

Another problem is that the 1660-MeV resonance<sup>5</sup> has begun to appear in the  $\Lambda\pi^+$  mass spectrum. This distorts somewhat the constant background assumed for the  $\pi$ - $\pi$  study, but it can hardly be expected to cause a narrow peaking.

The 15.5 events that are possible  $\omega \rightarrow 2\pi$  decays must be increased to 43 to correct for the fact that area selection on the Dalitz plot used 0.36 of the mass band in which  $\omega$ 's could be expected. The density of  $\omega \rightarrow 2\pi$  events in this mass band is taken to be uniform, since study of the  $\omega \rightarrow 3\pi$  events in the same film sample indicated isotropic decay.<sup>6</sup> On comparison with the 950 three-pion decays expected in the same sample, a branching ratio for two-pion to three-pion decay is estimated to be:

$$\Gamma(\omega \rightarrow 2\pi)/\Gamma(\omega \rightarrow 3\pi) = 0.045 \pm 0.016. \quad (9)$$

Simple checks have been made to ascertain that there are no spurious  $\Lambda\pi^+\pi^-$  fits of actual  $\Lambda\pi^+\pi^-\pi^0$  events appearing in the 780-MeV peak. Values of  $\chi^2$  used for selection of events were examined more closely. For events under the 780-MeV peak, no  $\chi^2$  for  $\Lambda\pi^+\pi^-\pi^0$  was lower than 10, and the three events with the  $\Lambda 3\pi$   $\chi^2$  between 10 and 20 had  $\chi^2$  values for the  $\Lambda\pi^+\pi^-$  hypothesis that were much better.


Additional checks have been made. The  $\omega \rightarrow 2\pi$  decay was not discernible in the complete  $\pi$ - $\pi$  mass spectrum. However, the entire sector between the  $Y^*$  bands was compared with the  $\pi$ - $\pi$  spectrum obtained by the incoherent combination of amplitudes for the  $\rho$ , the  $Y_1^*$  (1385 MeV), and the  $Y_1^*$  (1660 MeV) with weighting factors obtained from the fitting of the total sample. The  $2\pi$  peak at the  $\omega$  mass was still prominent, again as an approximate three-standard-deviation effect. Further, the angular distribution of the lambda produced with the two pions in the events under the " $\omega$  peak" (with about 50% background) was found to have a form closer to the  $\Lambda$  distribution in  $K^- + p \rightarrow \Lambda + \omega \rightarrow \Lambda + 3\pi$  than to the  $\Lambda$  distribution in the other  $K^- + p \rightarrow \Lambda + 2\pi$  events. Figure 9 shows the  $\pi$ - $\pi$  mass spectrum for various production angles; the " $\omega \rightarrow 2\pi$ " peak is most prominent at forward  $\omega$  angles.

#### EARLIER STUDY OF THE TWO-PION DECAY MODE

A number of theoretical discussions of the  $\omega \rightarrow 2\pi$  decay have appeared in the literature.<sup>7</sup> Two groups of experimentalists have indicated that their data on the  $\rho$  mass spectrum showed some evidence for the  $\omega \rightarrow 2\pi$  decay.<sup>8</sup>

Theory suggests that the identity of the  $\rho^0$  and  $\omega$  mesons in all quantum numbers except isotopic spin permits electromagnetic transitions such that an  $\omega$  meson can become a  $\rho$  meson (or vice versa) by passing through a virtual photon state. Another way of stating this is to say that an  $\omega$  meson (or a  $\rho^0$  meson) can be coupled to any charged pair ( $e^+e^-$ ,  $\mu^+\mu^-$ , or  $\pi^+\pi^-$ ) by an intermediate photon, and by virtue of the  $\rho$  resonance in the  $\pi^+\pi^-$  final state, this mode can be considerably enhanced.

The ratio of the anomalous amplitude to the isotopic-spin-conserving amplitude in  $\omega$  (or  $\rho^0$  decay) is estimated to be about 1/50; the square of this ratio times the width of the  $\rho$  meson gives the rate of  $\omega \rightarrow 2\pi$  decay. If the width of the three-pion decay of the  $\omega$  is about 1 MeV, then the two-pion



MU-29074

Fig. 9. Pion-pion mass spectra at various production angles for the recoil  $\Lambda$ . The same events are used as in Fig. 7. Although the  $\rho$  meson also appears to peak forward so that the  $\omega \rightarrow 2\pi$  decay cannot be separated, it is evident that this peak behaves rather differently from the general  $\Lambda - \pi^+ - \pi^-$  "background." (Note: Since the bin of interest, at 780 MeV, is one-fifth as wide as the other bins, the observed number of events and its error were multiplied by five for this bin.)

decay can be expected to be approximately 5% of the dominant three-pion decay. Whether this small rate is detectable in a given experiment and whether it appears as a peak or a dip superimposed on the mass spectrum depends on the relative amplitudes for  $\rho$  and for  $\omega$  production.

### OTHER RARE DECAY MODES

We have studied the decay of the  $\omega$  into  $e^+ + e^-$ . All vee-two-prong events were fitted for the hypotheses  $K^- + p \rightarrow \Lambda + e^+ + e^-$  and also for  $K^- + p \rightarrow \Lambda + \omega$ ,  $\omega \rightarrow e^+ + e^-$ . As an additional check, colinearity of the charged secondaries was examined after transformation into the  $\omega$  rest frame (determined solely from the measured  $K^-$  momentum and the  $\Lambda$  momentum fitted in decay) under the assumptions that they were  $e^+e^-$  and also that they were  $\pi^+\pi^-$ . On the basis of  $\chi^2$  selection criteria and the colinearity check, two good examples and three more possible examples of  $\omega \rightarrow e^+ + e^-$  were found in a sample of 4892  $K^- + p \rightarrow \Lambda + 2$ -prong events. (The likely  $e^+e^-$  candidates were examined for delta rays, but none was found.) The branching ratio thus obtained is

$$\Gamma(\omega \rightarrow e^+e^-)/\Gamma(\omega \rightarrow 3\pi) \leq 0.01. \quad (10)$$

We are studying the rare decay mode  $\omega \rightarrow \pi^+\pi^-\gamma$  by examining missing neutral mass. No definite results have yet been obtained.

### CONCLUSIONS

The experiment outlined above has given results for three properties of the  $\omega$  meson. The neutral-to charged branching ratio of  $0.11 \pm 0.03$  agrees approximately with the estimate of Gell-Mann, Sharp, and Wagner<sup>7</sup> of 0.17 (considered perhaps the most reliable branching ratio given by their model). The width for three-pion decay of the  $\omega$  is consistent with the estimate of approximately 0.5 MeV by Gell-Mann, Sharp, and Wagner<sup>7</sup> and by Feld,<sup>9</sup> and with an estimate of the order of 2 MeV by Zemach and Zachariasen.<sup>10</sup> The experimental result does not disagree with smaller estimates, such as 0.05 MeV of Nambu and Sakurai<sup>7</sup> and 0.15 MeV of Feinberg.<sup>11</sup> It does differ somewhat from the 10- to 20-MeV width calculated by Frazer and Wong.<sup>12</sup> The experimentally determined  $2\pi/3\pi$  decay ratio is in very good agreement with the 5% value given by most theoretical estimates; however, since the estimates depend on the absolute  $3\pi$  rate, this does not support any theoretical model without a more nearly exact determination of the  $3\pi$  width.

It seems evident that the experiment  $K^- + p \rightarrow \Lambda + 2\pi$  or  $\Lambda + 3\pi$  reported here is particularly well-suited to the study of the  $2\pi$  (or two-lepton) mode of  $\omega$  decay, as the number of  $\rho$  mesons produced was considerably less than the number of  $\omega$  mesons (50% of all  $3\pi$  final states). The situation was much less favorable in the antiproton experiment,<sup>13</sup> and it is somewhat less favorable in the experiments mentioned above.<sup>8</sup>

### ACKNOWLEDGMENTS

Discussions with Prof. Charles Zemach have been very helpful. Also the interest of Prof. Sheldon Glashow in the evidence for two-pion decay was much appreciated. Thanks are also due Prof. Arthur Rosenfeld for his interest. The assistance of Mr. Daniel Siegel in the  $\omega \rightarrow 3\pi$  analysis was greatly appreciated.

Drs. Stanley Wojcicki and Margaret Alston gave assistance in data reduction which was very valuable in this experiment. Discussions with Dr. Ulrich Kruse were helpful in work on rare decay modes. Finally, the encouragement of Prof. Luis Alvarez was very much appreciated.

## FOOTNOTES AND REFERENCES

† This work was done under the auspices of the U. S. Atomic Energy Commission.

1. A preliminary report was made on these data in J. Button-Shafer, M. Ferro-Luzzi, J. J. Murray, M. L. Stevenson, and F. T. Solmitz, Proceedings of the 1962 International Conference on High-Energy Physics at CERN, (CERN, Geneva, 1962), p. 307.
2. Qualitatively similar results were obtained from the same reaction by W. A. Cooper, H. Courant, H. Filthuth, E. I. Malamud, A. Minguizzi-Ranzi, H. Schneider, A. M. Segar, G. A. Snow, W. Willis, E. S. Gelsema, J. C. Kluyver, A. G. Tenner, K. Browning, I. S. Hughes, R. Turnbull, in Proceedings of the 1962 International Conference on High-Energy Physics at CERN, (CERN, Geneva, 1962), p. 298. The  $\omega$  meson was studied in other reactions by B. C. Maglič, L. W. Alvarez, A. H. Rosenfeld, M. L. Stevenson, *Phys. Rev. Letters* 7, 178 (1961); A. Pevsner, R. Kraemer, M. Nussbaum, C. Richardson, P. Schlein, R. Strand, T. Toohig, M. Block, A. Engler, R. Gessaroli, and C. Meltzer, *Phys. Rev. Letters* 7, 421 (1961); M. L. Stevenson, L. W. Alvarez, B. C. Maglič, and A. H. Rosenfeld, *Phys. Rev.* 125, 687 (1962); N. H. Xuong and G. R. Lynch, *Phys. Rev. Letters* 7, 327 (1961); and by the following authors in Proceedings of the 1962 International Conference on High-Energy Physics at CERN, (CERN, Geneva, 1962): W. D. Walker, E. West, A. R. Erwin, and R. H. March, p. 42; C. Alff, D. Colley, N. Gelfaud, V. Nauenberg, D. Riller, J. Steinberger, T. H. Tan, H. Brugger, P. Kramer, and R. Plano, p. 50; G. B. Chadwick, W. T. Davies, M. Derrick, C. J. B. Hawkins, P. B. Jones, J. H. Mulvey, D. Radojicic, C. A. Wilkinson, M. Cresti, A. Grigoletto, S. Limentani, A. Loria, L. Peruzzo, and R. Santangelo, p. 73; R. Armenteros, R. Budde, L. Montanet, D. R. O. Morrison, S. Nilsson, A. Shapira, J. Vandermeulen, C. d'Andlau, A. Astier, C. Ghesquière, B. Gregory, D. Rahm, P. Rivet, and F. Solmitz, p. 90; T. Toohig, R. Kraemer, L. Madansky, M. Meer, M. Nussbaum, A. Pevsner, C. Richardson, R. Strand, M. Block, p. 99; and M. Meer, R. Strand, R. Kraemer, L. Madansky, M. Nussbaum, A. Pevsner, C. Richardson, T. Toohig, M. Block, S. Orenstein, and T. Fields, p. 103. More recent work was done by E. L. Hart, R. I. Louttit, and T. W. Morris, *Phys. Rev. Letters* 9, 133 (1962) and C. Alff, D. Berley, D. Colley, N. Gelfaud, V. Nauenberg, D. Miller, J. Schultz, J. Steinberger, T. H. Tan, H. Brugger, P. Kramer, R. Plano, *Phys. Rev. Letters* 9, 322 (1962) and *Phys. Rev. Letters* 9, 325 (1962).
3. This method was applied to  $\overline{K}^0 \pi^- p$  events to find the ( $\sim$  zero) width of the  $\overline{K}^0$ ; it was found to give the correct width within statistical errors.
4. For the small width,  $\Gamma$ , of interest here, the  $J = 1^-$  resonance shape differs negligibly from the generally used  $J = 0$  resonance curve.



5. L. W. Alvarez, M. H. Alston, M. Ferro-Luzzi, D. O. Huwe, G. R. Kalbfleisch, D. H. Miller, J. J. Murray, A. H. Rosenfeld, J. B. Shafer, F. T. Solmitz, and S. G. Wojcicki, Phys. Rev. Letters 10, 184 (1963); P. L. Bastien and J. P. Berge, Phys. Rev. Letters 10, 188 (1963).
6. Isotropy is assumed to be undistorted by  $\rho$ - $\omega$  mixing. Populations of different portions of the  $\omega \rightarrow 2\pi$  strip of the Dalitz plot are compatible with isotropy; however, even with some anisotropy in the  $\omega \rightarrow 2\pi$  decay, the density of events in the two strips utilized above appears to be a good average value.
7. S. Fubini, Phys. Rev. Letters 7, 466 (1961); S. Glashow, Phys. Rev. Letters 7, 469 (1961); Y. Nambu and J. Sakurai, Phys. Rev. Letters 8, 79 (1962); G. Feinberg, Phys. Rev. Letters 8, 151 (1962); M. Gell-Mann, D. Sharp, and W. G. Wagner, Phys. Rev. Letters 8, 261 (1962); and J. Bernstein and G. Feinberg, Il Nuovo Cimento 25, No. 6, 1343 (1962).
8. E. Pickup, D. K. Robinson, and E. O. Salant, Phys. Rev. Letters 9, 170; C. Alff et al. Phys. Rev. Letters 9, 322 (1962).
9. B. T. Feld, Phys. Rev. Letters 8, 181 (1962).
10. F. Zachariasen and C. Zemach, Phys. Rev. 128, 849 (1962).
11. G. Feinberg, Phys. Rev. Letters 8, 151 (1962).
12. W. R. Frazer and D. Y. Wong, in Proceedings of the 1962 International Conference on High-Energy Physics at CERN, (CERN, Geneva, 1962), p. 144.
13. J. Button, G. Kalbfleisch, G. Lynch, B. Maglič, A. H. Rosenfeld, and M. L. Stevenson, Phys. Rev. 126, 1858 (1962).

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

