UC Irvine

UC Irvine Previously Published Works

Title

Unique Clinical Aspects of Nasal Scarring.

Permalink

<https://escholarship.org/uc/item/4jj517q1>

Journal

Facial plastic surgery clinics of North America, 25(1)

ISSN

1064-7406

Authors

Caughlin, Benjamin P
Barnes, Christian
Nelson, J Stuart
et al.

Publication Date

2017-02-01

DOI

10.1016/j.fsc.2016.08.004

Peer reviewed

Unique Clinical Aspects of Nasal Scarring

Benjamin P. Caughlin, MD^{a,b}, Christian Barnes, MD^a, J. Stuart Nelson, MD, PhD^a, Brian J.F. Wong, MD, PhD^{a,*}

KEY POINTS

- Various methods are available for refining scars of the external nose and optimal scar revision frequently requires the utilization of multiple techniques.
- Differing anatomy of nasal subunits and their underlying structural framework limit surgical options in nasal scar revision compared with other areas of the face.
- An understanding of a variety of laser technologies and their specific applications can vastly aid in fine, controlled scar revision.
- Achieving optimal scar reduction regularly requires multiple stages of intervention, close follow-up, and repeat procedures.

A main focus of facial aesthetic surgery is to draw attention to the eyes. Any imperfection in nasal contour that leads to asymmetry will be noticed, and this emphasizes the importance of treating nasal injury and scar formation aggressively. The flexible cartilaginous framework of the nose is enveloped by a skin soft tissue envelope (SSTE) of variable thickness. It is this framework particularly over the nasal tip that must counter the forces generated by wound healing and contracture. In contrast, over the superior nasal dorsum (supported by bone), and the middle vault (by the relatively firm fusion of the dorsal quadrangular and upper lateral cartilages) deformation of the SSTE is less dramatic after injury or defect closure. The varying thickness of the skin in combination with the varying stability of the structural framework (eg, tip vs dorsum) leads to differences in response in contrast to scar and injury management in most other regions of the face. This is amplified by the ability of the eye to discern even small changes in facial symmetry and structure. For example, the smallest amount of contracture at the alar rim or soft tissue triangle can result in asymmetry and thus noticeable deformation. In contrast, injury and subsequent scar formation over the osseous nasal framework can be treated conservatively, as the surface is firm and adjacent skin is linked to bone via periosteal attachments.

Finally, the nose is a functional organ that supports air flow and mechanically filters, warms, and humidifies inspired air. Significant scarring can lead to collapse of the internal and external nasal valves. All nasal surgeons must appreciate this when planning wound management or reconstruction and when deciding how to best treat nasal scars and defects.

MANAGEMENT AND TREATMENT

The Art of Doing Nothing

Healing by secondary intention can produce stable outcomes that are cosmetically acceptable, when applied in a measured and judicious manner. This is important to consider especially with early scars produced by thermal injury or ballistic trauma in which the true extent of injury evolves over time. Aggressive local wound care is important, and this can be time and labor intensive, requiring frequent office visits and constant patient reassurance. The key is to strike a balance between doing the least-invasive treatment possible and obtaining the best results (Fig. 1). As no reconstructive surgery is performed, injury to normal adjacent tissue is minimized and other treatment options are preserved for future reconstruction if required. Adjunctive therapies, such as hyperbaric therapy,¹ local steroid injections,² silicone dressings,³ massage,⁴ pulsed dye laser,⁵ fractionated laser,⁶ and microneedling⁷ may have value for modulating the evolving scar process. This works best over a concave surface overlying a firm bony framework, such as in the nasal dorsum adjacent to the canthal region.

Fig. 1. The nasal tip of a 28-year-old Hispanic man who underwent open rhinoplasty with defatting of the tip SSTE and aggressive postoperative tape application (elsewhere). This led to postoperative vascular compromise followed by infection and tissue breakdown. The wound was treated with local wound care including antibiotic ointment (mupirocin) and frequent dressing changes during the acute phases of wound healing by the senior author (BJFW). The patient additionally received 6 rounds of hyperbaric oxygen therapy, frequent visits for wound debridement, comedone extractions, and 1 episode of free hand microneedling to release the scar. Images were obtained at presentation (A–C), 5 days (D–F), 1 week (G–I), 2 weeks (J–L), 4 weeks (M–O), 8 weeks (P–R), 14 weeks (S–U), and 27 weeks (V–X). The final images show the defect at 46 weeks and after 1 treatment with injectable hyaluronic acid (Y–AA) (performed elsewhere).

Fig. 1. (continued)

Laser and Resurfacing

In our practice, the laser has largely replaced dermabrasion. Although many surgeons achieve outstanding results with dermabrasion, we focus on laser resurfacing for this section. The risk of laser devices to both physician and patient must be appreciated, and proper eye protection is imperative. A comprehensive understanding of the physiology of laser treatments for scars is evolving as technology advancement exceeds the rate of clinical adoption and use. In our practice, laser technology is used both early and later in treatment protocols.^{8,9} We often start scar treatments with laser resurfacing and remodeling to varying degrees. This timing then allows us to better define the scar, to soften it, and to build the local collagen and tissues for any future reconstructions. If surgical scar revision is required, the reconstruction is followed again by more laser treatments to assist with tissue remodeling.

Laser fundamentals are beyond the scope of this article, but the advantage of laser therapy is that specific epidermal and dermal structures can be selectively targeted through precise selection of dosimetry, and as a noncontact technique there is no mechanical trauma to the skin.^{8,10,11} In contrast, one limitation of dermabrasion (using sandpaper or rotary burrs) is that force is exerted on the tissue, and this may distract wounds. Hence, dermabrasion is seldom used during the acute phases of wound healing, and frequently it is used at 8 weeks.¹⁰ Our laser treatment protocols begin on the day of suture removal, during the acute phase of wound healing.^{8,9} We then select the appropriate wavelength and dosimetry parameters depending on what each injury requires and how it is likely to respond to treatment. A typical approach would start with pulsed dye laser therapy at suture removal with consideration of fractional ablative resurfacing 2 weeks later. Pulsed dye lasers target the scar vasculature. The fractional ablative resurfacing can be every 4 weeks for up to 5 treatments. At approximately 5 to 6 weeks, nonablative lasers maybe used at 2-week to 4-week intervals to support collagen remodeling.⁸⁻¹⁰

For early scar redness and erythema, a “vascular” laser, such as the 595-nm pulsed dye laser is used. Light from this laser is preferentially absorbed by hemoglobin.^{8,9,12} Being nonablative, it creates no eschar for the patients to care for following treatment.¹³ Scars will have many different-sized blood vessels and the laser settings can be adjusted to target each specifically. The pulse width can be set shorter for smaller vessels and longer for larger vessels. For general redness and erythema, we use settings similar to rosacea treatments: an energy density of 7.5 J/cm² with a 6-ms pulse width using a 10-mm spot size. We often make 2 passes to complete a single treatment. It is valuable to have a built-in cooling device or cooling technique to allow the epidermis to be cooled selectively.¹¹ The laser we use is the V-Beam Perfecta (Syneron/Candela, Wayland, MA) with a 30/30 cooling setting (30 ms spurt of cryogen with a 30-ms delay until laser burst). For larger vessels and telangiectasias, the laser can be set to an energy density of 12.5 J/cm², 20-ms pulse width, and 30/30 cooling using a 7-mm spot size. Often a specific scar will get different treatment protocols at the same time depending on what is causing the erythema. It is important to allow the tissue to cool and to not be overly aggressive, as burns and blisters can occur.

For dermal collagen remodeling, the nonablative fractional 1550-nm wavelength lasers can be used¹⁴ with multiple passes. This is often used in a fractional manner with multiple

passes. Typically this laser is reserved for depressed scars that require multiple treatments to restore collagen over time. These lasers can generate significant temperature elevations, and without the traditional visual feedback of pinpoint bleeding seen with ablative lasers, burns and blisters are possible. It is important to wait at least 1 minute between each successive pass to allow the skin to thermally relax. A common setting used in our practice for nasal tip or dorsal scars is an energy of 45 mJ, treatment coverage area of 26% delivered with 6 passes. Again, it must be stressed to cool the superficial skin between each pass for extended periods (at least 1 minute with a cooling technique) to prevent burns and blistering.

Ablative lasers have the highest water absorption peaks and thus lead to destruction and removal of the superficial epidermal and dermal tissue.^{8,9} In our practice, we use the 2940-nm wavelength Er:YAG laser with a tunable pulse duration. Er:YAG has high absorption in water, and thus a vertically shallow optical penetration depth. The 10,600-nm wavelength CO₂ laser is also frequently used in this regard as well by many surgeons with excellent results, although thermal injury and depth of ablation are substantially deeper.^{8,9,11} These ablative lasers are used in a fractional pattern of 11% to 22% treatment coverage area with multiple passes as desired.

A standard protocol for a hypertrophic nasal scar is 500 mm of depth of ablation, 11% of fractional treatment coverage area delivered with 3 passes. Pinpoint bleeding will be obtained similar to dermabrasion. This protocol will require posttreatment wound care and prophylaxis at the discretion of the surgeon (Fig. 2).

Surgical Scar Revision

Surgical revision of nasal scars is indicated when the scar distorts the natural contours of the nose. A tight scar band can pull the ala medially for example. This can be released with a Z-plasty to reorient the scar and then followed by laser treatments as indicated to blend the incisions.

Depressed scars with resulting soft tissue defect can be replaced with a full-thickness skin graft (FTSG) to release the pull and straining created from contraction. Fig. 3 shows an example of when an FTSG can be used to fill the soft tissue defect. Grafting can be followed by laser as stated previously to blend the areas. This patient elected to have no further treatment beyond the FTSG.

The soft tissue triangle area of the nose is a particular area of interest to the nasal surgeon. This web of tissue has the propensity to be easily distorted by scar. It is a classic scenario that the soft tissue triangle is scarred leading to retraction and notching of the ala. This can be repaired with an auricular composite graft to supply the needed rigid framework in conjunction with soft tissue simultaneously. Alar retraction is a common consequence of nasal reconstruction and trauma as well, and can be treated with composite grafts to the vestibular inner lining or by using structural rhinoplasty techniques, as seen in Fig. 4.

Injectable Steroids and Fillers

It is not our practice to use injectable fillers in the nose for scars, although many surgeons do report excellent results with such intervention. This topic is addressed in other sections of

treating facial scars, as there are roles for such treatments in specific locations and instances. Steroid injections are a commonly accepted method to settle inflammation and hypertrophy of scars. Our typical protocol is triamcinolone 40 mg/mL and volume is determined by the scar size. For 1 to 2 cm² 5 20 to 40 mg/course, 2 to 6 cm² 5 40 to 80 mg/course, 6 to 12 cm² 5 80 to 120 mg/course. We inject at 1-week to 4-week intervals (2–3 injections total). The patient in **Fig. 5** demonstrates the utility of local steroid injections. As discussed within this article, most patients undergo a combination of treatments to maximize results.¹⁵

Total Subunit Reconstruction

Essentially after the previously discussed more conservative options have been considered or attempted, the final stage is total nasal subunit reconstruction. If the scar is confined within one nasal subunit, often replacement of the entire subunit can lead to better results than having visible scars within the boundaries of a specific nasal subunit. Interpolated flaps such as a paramedian forehead or a melolabial flap are often used to replace the soft tissue when a significant volume is required. This is well established in the facial plastic surgery literature and is discussed with a patient example here. The patient in Fig. 6 is a 53-year-old woman who underwent total nasal reconstruction after roughly 9 rhinoplasty operations (performed elsewhere).

PEARLS AND PITFALLS

Pearl: Let the tissue and scar define themselves before intervention.

Pitfall: Treating scars before allowing the tissue to demarcate will delay and distort the nasal Structures and may mislead reconstructive efforts.

Pearl: With laser energy, always be conservative. Slow and steady progress allows you to appreciate how the scar and the patient will respond to laser treatments.

Pitfall: Lasers are powerful devices. Being overly aggressive with the laser can worsen and create new scars.

^a UC Irvine Department of Head & Neck Surgery, 101 The City Drive South, Orange, CA 92868, USA; ^b Facial Plastic and Reconstructive Surgery, 50 E Washington, Chicago, IL 60602, USA

* Corresponding author.

E-mail address: bjwong@uci.edu

Fig. 2. A 55-year-old woman with a history of melanoma in situ of the nasal dorsum resected by Mohs micrographic surgery (MMS) and reconstructed with nasal dorsum (Rieger) flap 8 months before presentation; note glabellar, dorsal, and sidewall scarring (A) (cancer resection and primary reconstruction performed elsewhere). Her recovery was complicated by wound infection that resolved with oral antibiotics. The patient used silicone strips to reduce scarring postoperatively and noted increased rotation of the tip with flap contracture (A). Unfortunately, she developed recurrent disease at the columella and right nasal ala (B). MMS was again used to remove the lesions, which resulted in a 9-mm columellar defect and a 5-mm right alar rim defect that were reconstructed with an FTSG from the left preauricular area. Scarring and telangiectasias were treated with combined pulsed dye laser and ablative Er:YAG fractional lasers with results seen 5 months after skin graft placement and 2 rounds of laser treatments (C, D) as well as 8 months postoperatively after 4 laser treatments (E, F). The patient chose not to undergo any additional intervention.

Fig. 3. A 44-year-old woman presents with morpheaform basal cell carcinoma of the right nasal ala (A). (B) Final defect after wide local excision of tumor. Surgical defect was covered with a hydrophilic dressing (MeroGel, Medtronic Xomed, Inc., Minneapolis, MN). Two weeks later, after pathologic confirmation of clear margins on permanent sections, the patient was taken back to the operating room for reconstruction with postauricular FTSG. The lesion as seen immediately before grafting (C) and 4 weeks after FTSG placement (D). The patient elected for no further treatment of the graft such as dermabrasion or laser resurfacing. Follow-up at 4 months (E), 10 months (F), 14 months (G), 23 months (H), 35 months (I), and 45 months after placement of skin graft (J).

Fig. 4. A 54-year-old woman with early rhinophyma presenting 5 years after Mohs excision of a right lateral nasal basal cell cancer with local reconstruction (performed elsewhere) and 3 years after subsequent rhinoplasty for persistent cosmetic defect (performed elsewhere). Note twisting of the tip and right alar rim notching from soft tissue triangle retraction (A), excessive columellar show (B), and nostril asymmetry with rightward deviation of the columella (C) and redness and telangiectasias in photographs at consultation. Her contour irregularities were treated with revision rhinoplasty with costal cartilage (septal extension graft, articulated alar rim grafts, lateral crural tensioning). Eight weeks postoperative enhanced telangiectasias of the nasal tip are seen (D–F). Telangiectasias were treated with a pulsed dye laser and the skin textural changes were treated with an ablative Erb:YAG fractional laser with results shown 3 weeks after the first treatment and 4 months postoperatively (G–I), as well as 11 weeks after the fourth treatment and 10 months postoperatively (J–L).

Fig. 5. A 52-year-old woman presents 3 months after excision of a right nasal sidewall basal cell carcinoma and reconstruction with a fascial-musculocutaneous glabellar (Banner) flap (performed elsewhere). One month after her surgery she noted thickening of the skin at the incision, was referred to the senior author (BJFW) for management, and underwent local injection of triamcinolone. (A) The patient at presentation after 1 treatment with injected triamcinolone. A second dose of triamcinolone (0.2 mL of Kenalog-40) was injected within the scar at this time. A third injection of triamcinolone was delivered 4 weeks later. (B) Two months after the third steroid injection and 1 month after her first session of laser skin resurfacing (by JSN). (C) Eleven months after initial reconstruction followed by 3 rounds of local steroid injections and multiple sessions of laser resurfacing.

Fig. 6. A 53-year-old woman who presented approximately 9 rhinoplasty operations (performed elsewhere). She elected to undergo total nasal reconstruction with a paramedian forehead flap. Her procedure was performed in 3 stages with 5 minor additional refinement and defatting procedures performed under local.16 Laser therapy was also provided. Images show presentation (A–C), 4 weeks after final staging of the paramedian forehead flap (D–F), and 4 years after final staging of paramedian forehead flap with 4 refinement procedures and 1 treatment of combination pulsed dye and fractional laser resurfacing (G–I).

REFERENCES

1. Dauwe PB, Pulikkottil BJ, Lavery L, et al. Does hyperbaric oxygen therapy work in facilitating acute wound healing: a systematic review. *Plast Reconstr Surg* 2014;133(2):208e–15e.
2. Manuskiatti W, Fitzpatrick RE. Treatment response of keloidal and hypertrophic sternotomy scars: comparison among intralesional corticosteroid, 5-fluorouracil, and 585-nm flashlamp-pumped pulsed-dye laser treatments. *Arch Dermatol* 2002; 138(9):1149–55.
3. Sidgwick GP, McGeorge D, Bayat A. A comprehensive evidence-based review on the role of topicals and dressings in the management of skin scarring. *Arch Dermatol Res* 2015; 307:461–77.
4. Shin TM, Bordeaux JS. The role of massage in scar management: a literature review. *Dermatol Surg* 2012;38:414–23.
5. Hultman CS, Friedstat JS, Edkins RE, et al. Laser resurfacing and remodeling of hypertrophic burn scars: the results of a large, prospective, before-after cohort study, with long-term follow-up. *Ann Surg* 2014;260(3):519–29.
6. Carniol PJ, Hamilton MM, Carniol ET. Current status of fractional laser resurfacing. *JAMA Facial Plast Surg* 2015;17(5):360–6.
7. El-Domyati M, Barakat M, Awad S, et al. Microneedling therapy for atrophic acne scars: an objective evaluation. *J Clin Aesthet Dermatol* 2015;8(7):36–42.
8. Oliaei S, Nelson JS, Fitzpatrick R, et al. Laser treatment of scars. *Facial Plast Surg* 2012;28(5):518–24.
9. Oliaei S, Nelson JS, Fitzpatrick R, et al. Use of lasers in acute management of surgical and traumatic incisions on the face. *Facial Plast Surg Clin North Am* 2011;19(3):543–50.
10. Leake DS, Baker SR. Scar revision and local flap refinement. In: Baker SR, editor. *Local flaps in facial reconstruction*. Philadelphia: Elsevier Inc; 2007. p. 727–66.
11. Wu EC, Wong BJF. Lasers and optical technology in facial plastic surgery. *Arch Facial Plast Surg* 2008; 10:381–409.
12. Cohen BE, Brauer JA, Geronemus RG. Acne scarring: a review of available therapeutic lasers. *Lasers Surg Med* 2016;48(2):95–115.
13. Finney R, Torbeck R, Saedi N. Non-ablative fractional resurfacing in the treatment of scar contracture. *Lasers Surg Med* 2016;48(2):170–3.
14. Friedmann DP, Tzu JE, Kauvar AN, et al. Treatment of facial photodamage and rhytides using a novel 1,565nm non-ablative fractional erbium-doped fiber laser. *Lasers Surg Med* 2016;48(2):174–80.
15. On HR, Lee SH, Lee YS, et al. Evaluating hypertrophic thyroidectomy scar outcomes after treatment with triamcinolone injections and copper bromide therapy. *Lasers Surg Med* 2015;47(6): 479–84.
16. Menick FJ. Aesthetic refinements in use of forehead for nasal reconstruction: the paramedian forehead flap. *Clin Plast Surg* 1990;17(4):607–22.