

# UC Berkeley

## UC Berkeley Previously Published Works

### Title

Measurement of the branching fraction and CP content for the decay  $B^0 \rightarrow D^{*+} D^{*-}$ .

### Permalink

<https://escholarship.org/uc/item/4jk8r8wd>

### Journal

Physical review letters, 89(6)

### ISSN

0031-9007

### Authors

Aubert, B  
Boutigny, D  
Gaillard, J-M  
[et al.](#)

### Publication Date

2002-08-01

### DOI

10.1103/physrevlett.89.061801

### License

<https://creativecommons.org/licenses/by/4.0/> 4.0

Peer reviewed

## Measurement of the Branching Fraction and $CP$ Content for the Decay $B^0 \rightarrow D^{*+} D^{*-}$

B. Aubert,<sup>1</sup> D. Boutigny,<sup>1</sup> J.-M. Gaillard,<sup>1</sup> A. Hicheur,<sup>1</sup> Y. Karyotakis,<sup>1</sup> J. P. Lees,<sup>1</sup> P. Robbe,<sup>1</sup> V. Tisserand,<sup>1</sup> A. Zghiche,<sup>1</sup> A. Palano,<sup>2</sup> A. Pompili,<sup>2</sup> G. P. Chen,<sup>3</sup> J. C. Chen,<sup>3</sup> N. D. Qi,<sup>3</sup> G. Rong,<sup>3</sup> P. Wang,<sup>3</sup> Y. S. Zhu,<sup>3</sup> G. Eigen,<sup>4</sup> B. Stugu,<sup>4</sup> G. S. Abrams,<sup>5</sup> A. W. Borgland,<sup>5</sup> A. B. Breon,<sup>5</sup> D. N. Brown,<sup>5</sup> J. Button-Shafer,<sup>5</sup> R. N. Cahn,<sup>5</sup> M. S. Gill,<sup>5</sup> A. V. Gritsan,<sup>5</sup> Y. Groysman,<sup>5</sup> R. G. Jacobsen,<sup>5</sup> R. W. Kadel,<sup>5</sup> J. Kadyk,<sup>5</sup> L. T. Kerth,<sup>5</sup> Yu. G. Kolomensky,<sup>5</sup> J. F. Kral,<sup>5</sup> C. LeClerc,<sup>5</sup> M. E. Levi,<sup>5</sup> G. Lynch,<sup>5</sup> P. J. Oddone,<sup>5</sup> M. Pripstein,<sup>5</sup> N. A. Roe,<sup>5</sup> A. Romosan,<sup>5</sup> M. T. Ronan,<sup>5</sup> V. G. Shelkov,<sup>5</sup> A. V. Telnov,<sup>5</sup> W. A. Wenzel,<sup>5</sup> T. J. Harrison,<sup>6</sup> C. M. Hawkes,<sup>6</sup> D. J. Knowles,<sup>6</sup> S. W. O'Neale,<sup>6</sup> R. C. Penny,<sup>6</sup> A. T. Watson,<sup>6</sup> N. K. Watson,<sup>6</sup> T. Deppermann,<sup>7</sup> K. Goetzen,<sup>7</sup> H. Koch,<sup>7</sup> M. Kunze,<sup>7</sup> B. Lewandowski,<sup>7</sup> K. Peters,<sup>7</sup> H. Schmuecker,<sup>7</sup> M. Steinke,<sup>7</sup> N. R. Barlow,<sup>8</sup> W. Bhimji,<sup>8</sup> N. Chevalier,<sup>8</sup> P. J. Clark,<sup>8</sup> W. N. Cottingham,<sup>8</sup> B. Foster,<sup>8</sup> C. Mackay,<sup>8</sup> F. F. Wilson,<sup>8</sup> K. Abe,<sup>9</sup> C. Hearty,<sup>9</sup> T. S. Mattison,<sup>9</sup> J. A. McKenna,<sup>9</sup> D. Thiessen,<sup>9</sup> S. Jolly,<sup>10</sup> A. K. McKemey,<sup>10</sup> V. E. Blinov,<sup>11</sup> A. D. Bukin,<sup>11</sup> D. A. Bukin,<sup>11</sup> A. R. Buzykaev,<sup>11</sup> V. B. Golubev,<sup>11</sup> V. N. Ivanchenko,<sup>11</sup> A. A. Korol,<sup>11</sup> E. A. Kravchenko, A. P. Onuchin,<sup>11</sup> S. I. Serednyakov,<sup>11</sup> Yu. I. Skovpen,<sup>11</sup> V. I. Telnov,<sup>11</sup> A. N. Yushkov,<sup>11</sup> D. Best,<sup>12</sup> M. Chao,<sup>12</sup> D. Kirkby,<sup>12</sup> A. J. Lankford,<sup>12</sup> M. Mandelkern,<sup>12</sup> S. McMahon,<sup>12</sup> D. P. Stoker,<sup>12</sup> K. Arisaka,<sup>13</sup> C. Buchanan,<sup>13</sup> S. Chun,<sup>13</sup> D. B. MacFarlane,<sup>14</sup> S. Prell,<sup>14</sup> Sh. Rahatlou,<sup>14</sup> G. Raven,<sup>14</sup> V. Sharma,<sup>14</sup> C. Campagnari,<sup>15</sup> B. Dahmes,<sup>15</sup> P. A. Hart,<sup>15</sup> N. Kuznetsova,<sup>15</sup> S. L. Levy,<sup>15</sup> O. Long,<sup>15</sup> A. Lu,<sup>15</sup> M. A. Mazur,<sup>15</sup> J. D. Richman,<sup>15</sup> W. Verkerke,<sup>15</sup> J. Beringer,<sup>16</sup> A. M. Eisner,<sup>16</sup> M. Grothe,<sup>16</sup> C. A. Heusch,<sup>16</sup> W. S. Lockman,<sup>16</sup> T. Pulliam,<sup>16</sup> T. Schalk,<sup>16</sup> R. E. Schmitz,<sup>16</sup> B. A. Schumm,<sup>16</sup> A. Seiden,<sup>16</sup> M. Turri,<sup>16</sup> W. Walkowiak,<sup>16</sup> D. C. Williams,<sup>16</sup> M. G. Wilson,<sup>16</sup> E. Chen,<sup>17</sup> G. P. Dubois-Felsmann,<sup>17</sup> A. Dvoretzki,<sup>17</sup> D. G. Hitlin,<sup>17</sup> S. Metzler,<sup>17</sup> J. Oyang,<sup>17</sup> F. C. Porter,<sup>17</sup> A. Ryd,<sup>17</sup> A. Samuel,<sup>17</sup> S. Yang,<sup>17</sup> R. Y. Zhu,<sup>17</sup> S. Devmal,<sup>18</sup> S. Jayatilleke,<sup>18</sup> G. Mancinelli,<sup>18</sup> B. T. Meadows,<sup>18</sup> M. D. Sokoloff,<sup>18</sup> T. Barillari,<sup>19</sup> P. Bloom,<sup>19</sup> W. T. Ford,<sup>19</sup> U. Nauenberg,<sup>19</sup> A. Olivas,<sup>19</sup> P. Rankin,<sup>19</sup> J. Roy,<sup>19</sup> J. G. Smith,<sup>19</sup> W. C. van Hoek,<sup>19</sup> L. Zhang,<sup>19</sup> J. Blouw,<sup>20</sup> J. L. Harton,<sup>20</sup> M. Krishnamurthy,<sup>20</sup> A. Soffer,<sup>20</sup> W. H. Toki,<sup>20</sup> R. J. Wilson,<sup>20</sup> J. Zhang,<sup>20</sup> T. Brandt,<sup>21</sup> J. Brose,<sup>21</sup> T. Colberg,<sup>21</sup> M. Dickopp,<sup>21</sup> R. S. Dubitzky,<sup>21</sup> A. Hauke,<sup>21</sup> E. Maly,<sup>21</sup> R. Müller-Pfefferkorn,<sup>21</sup> S. Otto,<sup>21</sup> K. R. Schubert,<sup>21</sup> R. Schwierz,<sup>21</sup> B. Spaan,<sup>21</sup> L. Wilden,<sup>21</sup> D. Bernard,<sup>22</sup> G. R. Bonneaud,<sup>22</sup> F. Brochard,<sup>22</sup> J. Cohen-Tanugi,<sup>22</sup> S. Ferrag,<sup>22</sup> S. T'Jampens,<sup>22</sup> Ch. Thiebaut,<sup>22</sup> G. Vasileiadis,<sup>22</sup> M. Verderi,<sup>22</sup> A. Anjomshoaa,<sup>23</sup> R. Bernet,<sup>23</sup> A. Khan,<sup>23</sup> D. Lavin,<sup>23</sup> F. Muheim,<sup>23</sup> S. Playfer,<sup>23</sup> J. E. Swain,<sup>23</sup> J. Tinslay,<sup>23</sup> M. Falbo,<sup>24</sup> C. Borean,<sup>25</sup> C. Bozzi,<sup>25</sup> L. Piemontese,<sup>25</sup> E. Treadwell,<sup>26</sup> F. Anulli,<sup>27,\*</sup> R. Baldini-Ferroli,<sup>27</sup> A. Calcaterra,<sup>27</sup> R. de Sangro,<sup>27</sup> D. Falciari,<sup>27</sup> G. Finocchiaro,<sup>27</sup> P. Patteri,<sup>27</sup> I. M. Peruzzi,<sup>27,\*</sup> M. Piccolo,<sup>27</sup> Y. Xie,<sup>27</sup> A. Zallo,<sup>27</sup> S. Bagnasco,<sup>28</sup> A. Buzzo,<sup>28</sup> R. Contri,<sup>28</sup> G. Crosetti,<sup>28</sup> M. Lo Vetere,<sup>28</sup> M. Macri,<sup>28</sup> M. R. Monge,<sup>28</sup> S. Passaggio,<sup>28</sup> F. C. Pastore,<sup>28</sup> C. Patrignani,<sup>28</sup> E. Robutti,<sup>28</sup> A. Santroni,<sup>28</sup> S. Tosi,<sup>28</sup> M. Morii,<sup>29</sup> R. Bartoldus,<sup>30</sup> R. Hamilton,<sup>30</sup> U. Mallik,<sup>30</sup> J. Cochran,<sup>31</sup> H. B. Crawley,<sup>31</sup> P.-A. Fischer,<sup>31</sup> J. Lamsa,<sup>31</sup> W. T. Meyer,<sup>31</sup> E. I. Rosenberg,<sup>31</sup> J. Yi,<sup>31</sup> G. Grosdidier,<sup>32</sup> A. Höcker,<sup>32</sup> H. M. Lacker,<sup>32</sup> S. Laplace,<sup>32</sup> F. Le Diberder,<sup>32</sup> V. Lepeltier,<sup>32</sup> A. M. Lutz,<sup>32</sup> S. Plaszczynski,<sup>32</sup> M. H. Schune,<sup>32</sup> S. Trincaz-Duvoid,<sup>32</sup> G. Wormser,<sup>32</sup> R. M. Bionta,<sup>33</sup> V. Brigljević,<sup>33</sup> D. J. Lange,<sup>33</sup> M. Mugge,<sup>33</sup> K. van Bibber,<sup>33</sup> D. M. Wright,<sup>33</sup> A. J. Bevan,<sup>34</sup> J. R. Fry,<sup>34</sup> E. Gabathuler,<sup>34</sup> R. Gamet,<sup>34</sup> M. George,<sup>34</sup> M. Kay,<sup>34</sup> D. J. Payne,<sup>34</sup> R. J. Sloane,<sup>34</sup> C. Touramanis,<sup>34</sup> M. L. Aspinwall,<sup>35</sup> D. A. Bowerman,<sup>35</sup> P. D. Dauncey,<sup>35</sup> U. Egede,<sup>35</sup> I. Eschrich,<sup>35</sup> G. W. Morton,<sup>35</sup> J. A. Nash,<sup>35</sup> P. Sanders,<sup>35</sup> D. Smith,<sup>35</sup> J. J. Back,<sup>36</sup> G. Bellodi,<sup>36</sup> P. Dixon,<sup>36</sup> P. F. Harrison,<sup>36</sup> R. J. L. Potter,<sup>36</sup> H. W. Shorthouse,<sup>36</sup> P. Strother,<sup>36</sup> P. B. Vidal,<sup>36</sup> G. Cowan,<sup>37</sup> S. George,<sup>37</sup> M. G. Green,<sup>37</sup> A. Kurup,<sup>37</sup> C. E. Marker,<sup>37</sup> T. R. McMahon,<sup>37</sup> S. Ricciardi,<sup>37</sup> F. Salvatore,<sup>37</sup> G. Vaitsas,<sup>37</sup> D. Brown,<sup>38</sup> C. L. Davis,<sup>38</sup> J. Allison,<sup>39</sup> R. J. Barlow,<sup>39</sup> J. T. Boyd,<sup>39</sup> A. C. Forti,<sup>39</sup> F. Jackson,<sup>39</sup> G. D. Lafferty,<sup>39</sup> N. Savvas,<sup>39</sup> J. H. Weatherall,<sup>39</sup> J. C. Williams,<sup>39</sup> A. Farbin,<sup>40</sup> A. Jawahery,<sup>40</sup> V. Lillard,<sup>40</sup> J. Olsen,<sup>40</sup> D. A. Roberts,<sup>40</sup> J. R. Schieck,<sup>40</sup> G. Blaylock,<sup>41</sup> C. Dallapiccola,<sup>41</sup> K. T. Flood,<sup>41</sup> S. S. Hertzbach,<sup>41</sup> R. Kofler,<sup>41</sup> V. B. Koptchev,<sup>41</sup> T. B. Moore,<sup>41</sup> H. Staengle,<sup>41</sup> S. Willocq,<sup>41</sup> B. Brau,<sup>42</sup> R. Cowan,<sup>42</sup> G. Sciolla,<sup>42</sup> F. Taylor,<sup>42</sup> R. K. Yamamoto,<sup>42</sup> M. Milek,<sup>43</sup> P. M. Patel,<sup>43</sup> F. Palombo,<sup>44</sup> J. M. Bauer,<sup>45</sup> L. Cremaldi,<sup>45</sup> V. Eschenburg,<sup>45</sup> R. Kroeger,<sup>45</sup> J. Reidy,<sup>45</sup> D. A. Sanders,<sup>45</sup> D. J. Summers,<sup>45</sup> C. Hast,<sup>46</sup> J. Y. Nief,<sup>46</sup> P. Taras,<sup>46</sup> H. Nicholson,<sup>47</sup> C. Cartaro,<sup>48</sup> N. Cavallo,<sup>48,†</sup> G. De Nardo,<sup>48</sup> F. Fabozzi,<sup>48</sup> C. Gatto,<sup>48</sup> L. Lista,<sup>48</sup> P. Paolucci,<sup>48</sup> D. Piccolo,<sup>48</sup> C. Sciacca,<sup>48</sup> J. M. LoSecco,<sup>49</sup> J. R. G. Alsmiller,<sup>50</sup> T. A. Gabriel,<sup>50</sup> J. Brau,<sup>51</sup> R. Frey,<sup>51</sup> E. Grauges,<sup>51</sup> M. Iwasaki,<sup>51</sup> N. B. Sinev,<sup>51</sup> D. Strom,<sup>51</sup> F. Colecchia,<sup>52</sup> F. Dal Corso,<sup>52</sup> A. Dorigo,<sup>52</sup> F. Galeazzi,<sup>52</sup> M. Margoni,<sup>52</sup> G. Michelon,<sup>52</sup> M. Morandin,<sup>52</sup> M. Posocco,<sup>52</sup> M. Rotondo,<sup>52</sup> F. Simonetto,<sup>52</sup> R. Stroili,<sup>52</sup> E. Torassa,<sup>52</sup> C. Voci,<sup>52</sup> M. Benayoun,<sup>53</sup> H. Briand,<sup>53</sup> J. Chauveau,<sup>53</sup> P. David,<sup>53</sup> Ch. de Vaissière,<sup>53</sup> L. Del Buono,<sup>53</sup> O. Hamon,<sup>53</sup> Ph. Leruste,<sup>53</sup> J. Ocariz,<sup>53</sup> M. Pivk,<sup>53</sup> L. Roos,<sup>53</sup> J. Stark,<sup>53</sup> P. F. Manfredi,<sup>54</sup> V. Re,<sup>54</sup> V. Speziali,<sup>54</sup> E. D. Frank,<sup>55</sup> L. Gladney,<sup>55</sup> Q. H. Guo,<sup>55</sup> J. Panetta,<sup>55</sup> C. Angelini,<sup>56</sup> G. Batignani,<sup>56</sup> S. Bettarini,<sup>56</sup> M. Bondioli,<sup>56</sup> F. Bucci,<sup>56</sup> E. Campagna,<sup>56</sup> M. Carpinelli,<sup>56</sup> F. Forti,<sup>56</sup> M. A. Giorgi,<sup>56</sup> A. Lusiani,<sup>56</sup> G. Marchiori,<sup>56</sup> F. Martinez-Vidal,<sup>56</sup> M. Morganti,<sup>56</sup> N. Neri,<sup>56</sup>

E. Paoloni,<sup>56</sup> M. Rama,<sup>56</sup> G. Rizzo,<sup>56</sup> F. Sandrelli,<sup>56</sup> G. Simi,<sup>56</sup> G. Triggiani,<sup>56</sup> J. Walsh,<sup>56</sup> M. Haire,<sup>57</sup> D. Judd,<sup>57</sup> K. Paick,<sup>57</sup> L. Turnbull,<sup>57</sup> D. E. Wagoner,<sup>57</sup> J. Albert,<sup>58</sup> C. Lu,<sup>58</sup> V. Miftakov,<sup>58</sup> S. F. Schaffner,<sup>58</sup> A. J. S. Smith,<sup>58</sup> A. Tumanov,<sup>58</sup> E. W. Varnes,<sup>58</sup> G. Cavoto,<sup>59</sup> D. del Re,<sup>59</sup> R. Faccini,<sup>14,59</sup> F. Ferrarotto,<sup>59</sup> F. Ferroni,<sup>59</sup> M. A. Mazzoni,<sup>59</sup> S. Morganti,<sup>59</sup> G. Piredda,<sup>59</sup> M. Serra,<sup>59</sup> C. Voena,<sup>59</sup> S. Christ,<sup>60</sup> R. Waldi,<sup>60</sup> T. Adye,<sup>61</sup> N. De Groot,<sup>61</sup> B. Franek,<sup>61</sup> N. I. Geddes,<sup>61</sup> G. P. Gopal,<sup>61</sup> S. M. Xella,<sup>61</sup> R. Aleksan,<sup>62</sup> S. Emery,<sup>62</sup> A. Gaidot,<sup>62</sup> S. F. Ganzhur,<sup>62</sup> P.-F. Giraud,<sup>62</sup> G. Hamel Monchenault,<sup>62</sup> W. Kozanecki,<sup>62</sup> M. Langer,<sup>62</sup> G. W. London,<sup>62</sup> B. Mayer,<sup>62</sup> B. Serfass,<sup>62</sup> G. Vasseur,<sup>62</sup> Ch. Yèche,<sup>62</sup> M. Zito,<sup>62</sup> M. V. Purohit,<sup>63</sup> H. Singh,<sup>63</sup> A. W. Weidemann,<sup>63</sup> F. X. Yumiceva,<sup>63</sup> I. Adam,<sup>64</sup> D. Aston,<sup>64</sup> N. Berger,<sup>64</sup> A. M. Boyarski,<sup>64</sup> G. Calderini,<sup>64</sup> M. R. Convery,<sup>64</sup> D. P. Coupal,<sup>64</sup> D. Dong,<sup>64</sup> J. Dorfan,<sup>64</sup> W. Dunwoodie,<sup>64</sup> R. C. Field,<sup>64</sup> T. Glanzman,<sup>64</sup> S. J. Gowdy,<sup>64</sup> T. Haas,<sup>64</sup> V. Halyo,<sup>64</sup> T. Himel,<sup>64</sup> T. Hryn'ova,<sup>64</sup> M. E. Huffer,<sup>64</sup> W. R. Innes,<sup>64</sup> C. P. Jessop,<sup>64</sup> M. H. Kelsey,<sup>64</sup> P. Kim,<sup>64</sup> M. L. Kocian,<sup>64</sup> U. Langenegger,<sup>64</sup> D. W. G. S. Leith,<sup>64</sup> S. Luitz,<sup>64</sup> V. Luth,<sup>64</sup> H. L. Lynch,<sup>64</sup> H. Marsiske,<sup>64</sup> S. Menke,<sup>64</sup> R. Messner,<sup>64</sup> D. R. Muller,<sup>64</sup> C. P. O'Grady,<sup>64</sup> V. E. Ozcan,<sup>64</sup> A. Perazzo,<sup>64</sup> M. Perl,<sup>64</sup> S. Petrak,<sup>64</sup> H. Quinn,<sup>64</sup> B. N. Ratcliff,<sup>64</sup> S. H. Robertson,<sup>64</sup> A. Roodman,<sup>64</sup> A. A. Salnikov,<sup>64</sup> T. Schietinger,<sup>64</sup> R. H. Schindler,<sup>64</sup> J. Schwiening,<sup>64</sup> A. Snyder,<sup>64</sup> A. Soha,<sup>64</sup> S. M. Spanier,<sup>64</sup> J. Stelzer,<sup>64</sup> D. Su,<sup>64</sup> M. K. Sullivan,<sup>64</sup> H. A. Tanaka,<sup>64</sup> J. Va'vra,<sup>64</sup> S. R. Wagner,<sup>64</sup> M. Weaver,<sup>64</sup> A. J. R. Weinstein,<sup>64</sup> W. J. Wisniewski,<sup>64</sup> D. H. Wright,<sup>64</sup> C. C. Young,<sup>64</sup> P. R. Burchat,<sup>65</sup> C. H. Cheng,<sup>65</sup> T. I. Meyer,<sup>65</sup> C. Roat,<sup>65</sup> R. Henderson,<sup>66</sup> W. Bugg,<sup>67</sup> H. Cohn,<sup>67</sup> J. M. Izen,<sup>68</sup> I. Kitayama,<sup>68</sup> X. C. Lou,<sup>68</sup> F. Bianchi,<sup>69</sup> M. Bona,<sup>69</sup> D. Gamba,<sup>69</sup> L. Bosisio,<sup>70</sup> G. Della Ricca,<sup>70</sup> S. Dittongo,<sup>70</sup> L. Lanceri,<sup>70</sup> P. Poropat,<sup>70</sup> G. Vuagnin,<sup>70</sup> R. S. Panvini,<sup>71</sup> C. M. Brown,<sup>72</sup> P. D. Jackson,<sup>72</sup> R. Kowalewski,<sup>72</sup> J. M. Roney,<sup>72</sup> H. R. Band,<sup>73</sup> E. Charles,<sup>73</sup> S. Dasu,<sup>73</sup> M. Datta,<sup>73</sup> A. M. Eichenbaum,<sup>73</sup> H. Hu,<sup>73</sup> J. R. Johnson,<sup>73</sup> R. Liu,<sup>73</sup> F. Di Lodovico,<sup>73</sup> Y. Pan,<sup>73</sup> R. Prepost,<sup>73</sup> I. J. Scott,<sup>73</sup> S. J. Sekula,<sup>73</sup> J. H. von Wimmersperg-Toeller,<sup>73</sup> S. L. Wu,<sup>73</sup> Z. Yu,<sup>73</sup> T. M. B. Kordich,<sup>74</sup> and H. Neal<sup>74</sup>

## (BABAR Collaboration)

<sup>1</sup>Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France<sup>2</sup>Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy<sup>3</sup>Institute of High Energy Physics, Beijing 100039, China<sup>4</sup>University of Bergen, Institute of Physics, N-5007 Bergen, Norway<sup>5</sup>Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720<sup>6</sup>University of Birmingham, Birmingham, B15 2TT, United Kingdom<sup>7</sup>Ruhr Universität Bochum, Institut für Experimentalphysik I, D-44780 Bochum, Germany<sup>8</sup>University of Bristol, Bristol BS8 1TL, United Kingdom<sup>9</sup>University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1<sup>10</sup>Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom<sup>11</sup>Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia<sup>12</sup>University of California at Irvine, Irvine, California 92697<sup>13</sup>University of California at Los Angeles, Los Angeles, California 90024<sup>14</sup>University of California at San Diego, La Jolla, California 92093<sup>15</sup>University of California at Santa Barbara, Santa Barbara, California 93106<sup>16</sup>University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064<sup>17</sup>California Institute of Technology, Pasadena, California 91125<sup>18</sup>University of Cincinnati, Cincinnati, Ohio 45221<sup>19</sup>University of Colorado, Boulder, Colorado 80309<sup>20</sup>Colorado State University, Fort Collins, Colorado 80523<sup>21</sup>Technische Universität Dresden, Institut für Kern und Teilchenphysik, D-01062 Dresden, Germany<sup>22</sup>Ecole Polytechnique, F-91128 Palaiseau, France<sup>23</sup>University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom<sup>24</sup>Elon University, Elon University, North Carolina 27244-2010<sup>25</sup>Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy<sup>26</sup>Florida A&M University, Tallahassee, Florida 32307<sup>27</sup>Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy<sup>28</sup>Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy<sup>29</sup>Harvard University, Cambridge, Massachusetts 02138<sup>30</sup>University of Iowa, Iowa City, Iowa 52242<sup>31</sup>Iowa State University, Ames, Iowa 50011-3160<sup>32</sup>Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France<sup>33</sup>Lawrence Livermore National Laboratory, Livermore, California 94550<sup>34</sup>University of Liverpool, Liverpool L69 3BX, United Kingdom

- <sup>35</sup>University of London, Imperial College, London, SW7 2BW, United Kingdom  
<sup>36</sup>Queen Mary, University of London, E1 4NS, United Kingdom  
<sup>37</sup>University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom  
<sup>38</sup>University of Louisville, Louisville, Kentucky 40292  
<sup>39</sup>University of Manchester, Manchester M13 9PL, United Kingdom  
<sup>40</sup>University of Maryland, College Park, Maryland 20742  
<sup>41</sup>University of Massachusetts, Amherst, Massachusetts 01003  
<sup>42</sup>Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139  
<sup>43</sup>McGill University, Montréal, Quebec, Canada H3A 2T8  
<sup>44</sup>Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy  
<sup>45</sup>University of Mississippi, University, Mississippi 38677  
<sup>46</sup>Université de Montréal, Laboratoire René J. A. Lévesque, Montréal, Quebec, Canada H3C 3J7  
<sup>47</sup>Mount Holyoke College, South Hadley, Massachusetts 01075  
<sup>48</sup>Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126 Napoli, Italy  
<sup>49</sup>University of Notre Dame, Notre Dame, Indiana 46556  
<sup>50</sup>Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831  
<sup>51</sup>University of Oregon, Eugene, Oregon 97403  
<sup>52</sup>Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy  
<sup>53</sup>Universités Paris VI et VII, Lab de Physique Nucléaire H.E., F-75252 Paris, France  
<sup>54</sup>Università di Pavia, Dipartimento di Elettronica and INFN, I-27100 Pavia, Italy  
<sup>55</sup>University of Pennsylvania, Philadelphia, Pennsylvania 19104  
<sup>56</sup>Università di Pisa, Scuola Normale Superiore and INFN, I-56010 Pisa, Italy  
<sup>57</sup>Prairie View A&M University, Prairie View, Texas 77446  
<sup>58</sup>Princeton University, Princeton, New Jersey 08544  
<sup>59</sup>Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy  
<sup>60</sup>Universität Rostock, D-18051 Rostock, Germany  
<sup>61</sup>Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom  
<sup>62</sup>DAPNIA, Commissariat à l'Energie Atomique/Saclay, F-91191 Gif-sur-Yvette, France  
<sup>63</sup>University of South Carolina, Columbia, South Carolina 29208  
<sup>64</sup>Stanford Linear Accelerator Center, Stanford, California 94309  
<sup>65</sup>Stanford University, Stanford, California 94305-4060  
<sup>66</sup>TRIUMF, Vancouver, British Columbia, Canada V6T 2A3  
<sup>67</sup>University of Tennessee, Knoxville, Tennessee 37996  
<sup>68</sup>University of Texas at Dallas, Richardson, Texas 75083  
<sup>69</sup>Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy  
<sup>70</sup>Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy  
<sup>71</sup>Vanderbilt University, Nashville, Tennessee 37235  
<sup>72</sup>University of Victoria, Victoria, British Columbia, Canada V8W 3P6  
<sup>73</sup>University of Wisconsin, Madison, Wisconsin 53706  
<sup>74</sup>Yale University, New Haven, Connecticut 06511

(Received 5 March 2002; published 19 July 2002)

We report a measurement of the branching fraction of the decay  $B^0 \rightarrow D^{*+} D^{*-}$  and of the  $CP$ -odd component of its final state using the BABAR detector. With data corresponding to an integrated luminosity of  $20.4 \text{ fb}^{-1}$  collected at the  $Y(4S)$  resonance during 1999–2000, we have reconstructed 38 candidate signal events in the mode  $B^0 \rightarrow D^{*+} D^{*-}$  with an estimated background of  $6.2 \pm 0.5$  events. From these events, we determine the branching fraction to be  $\mathcal{B}(B^0 \rightarrow D^{*+} D^{*-}) = [8.3 \pm 1.6(\text{stat}) \pm 1.2(\text{syst})] \times 10^{-4}$ . The measured  $CP$ -odd fraction of the final state is  $0.22 \pm 0.18(\text{stat}) \pm 0.03(\text{syst})$ .

DOI: 10.1103/PhysRevLett.89.061801

PACS numbers: 13.25.Hw, 11.30.Er, 12.15.Hh

After the observation of time-dependent  $CP$ -violating asymmetries in the decays of neutral  $B$  mesons to  $CP$  eigenstates containing charmonium [1,2], it is interesting to extend the search for  $CP$ -violating effects to Cabibbo-suppressed double charm modes, such as  $B^0 \rightarrow D^{(*)+} D^{(*)-}$  [3]. The interference of the dominant tree amplitude with the mixing diagram is sensitive to the angle  $\beta$  of the unitarity triangle in this case as well; however, the theoretically uncertain contribution of penguin amplitudes with

different weak phases is potentially significant and may shift the observed asymmetry by an amount that depends on the ratio of the penguin and tree contributions and their relative weak phases. The  $B^0 \rightarrow D^{*+} D^{*-}$  vector-vector final state has very clear experimental signatures that make it an interesting candidate for  $CP$ -violation measurements. However, it is not a pure  $CP$  eigenstate and a dilution of the measured asymmetry can be produced by a  $P$ -wave,  $CP$ -odd, component. A time-dependent angular

analysis of the decay products [4] can remove the dilution by resolving the  $CP$ -even and  $CP$ -odd components. As a precursor to measuring time-dependent  $CP$ -violating asymmetries using the decay  $B^0 \rightarrow D^{*+}D^{*-}$ , we report in this letter a measurement of the  $B^0 \rightarrow D^{*+}D^{*-}$  branching fraction and a measurement of the  $CP$ -odd component,  $R_{\perp}$ , of the final state. These measurements represent significant improvements over the previous measurements  $\mathcal{B}(B^0 \rightarrow D^{*+}D^{*-}) = [9.9_{-3.3}^{+4.2}(\text{stat}) \pm 1.2(\text{syst})] \times 10^{-4}$  and  $(1 - R_{\perp}) < 0.11$  at 90% C.L. [5].

The data used in this analysis were collected with the BABAR detector [6] at the PEP-II storage ring [7] located at the Stanford Linear Accelerator Center. This data sample represents an integrated luminosity of  $20.4 \text{ fb}^{-1}$  collected on the  $Y(4S)$  resonance. Assuming 50% of the  $Y(4S)$  decays give  $B^0\bar{B}^0$ , the number of neutral  $B$  mesons in this sample is  $(22.8 \pm 0.4) \times 10^6$ .

Charged particles are detected and their momenta measured with the combination of a 40-layer drift chamber (DCH) and a 5-layer silicon vertex tracker (SVT), both operating in a 1.5 T solenoidal magnetic field. The charged particle tracking system allows particles with low momentum in the laboratory frame to be reconstructed efficiently, a property that is very important for this analysis. This efficiency begins to turn on at a momentum of  $\sim 60 \text{ MeV}/c$  and reaches its maximum value at about  $200 \text{ MeV}/c$ . Photons are detected by a CsI(Tl) electromagnetic calorimeter (EMC) that provides high detection efficiency for energies above 20 MeV, with typical energy and angular resolutions of 3% and 4 mrad, respectively, for 1 GeV photons. Charged particle identification is provided by the ionization loss measurements in the SVT and DCH, and by an internally reflecting ring-imaging Cherenkov detector (DIRC) covering the central region of the detector.

Events are selected by requiring three or more charged tracks and a normalized second Fox-Wolfram moment [8] of the event of less than 0.6. We also require that the cosine of the angle between the reconstructed  $B$  direction and the thrust axis of the rest of the event, calculated in the  $Y(4S)$  rest frame, be less than 0.9. These criteria discriminate  $Y(4S)$  events from nonresonant back-ground events.

$B^0$  mesons are exclusively reconstructed by combining two charged  $D^*$  candidates, using a number of  $D^{*+}$  and  $D$  decay modes. The  $D^{*+}$  mesons are reconstructed in their decays  $D^{*+} \rightarrow D^0\pi^+$  and  $D^{*+} \rightarrow D^+\pi^0$ . We include in this analysis the decay combinations  $D^{*+}D^{*-}$  decaying to  $(D^0\pi^+, \bar{D}^0\pi^-)$  or  $(D^0\pi^+, D^-\pi^0)$ , but not  $(D^+\pi^0, D^-\pi^0)$  due to the smaller branching fraction and larger expected backgrounds.  $D^0$  and  $D^+$  candidates are subjected to a mass-constrained fit to provide an improved measurement of the  $D$  meson's momentum. They are combined with pion candidates, referred to as "soft" pions due to their low ( $< 200 \text{ MeV}/c$ ) transverse momentum, to form  $D^{*+}$  candidates. A topological vertex fit is performed that includes the mean position of the  $e^+e^-$  interaction point to improve the angular resolution of the soft pion.

The decay modes of  $D^0$  and  $D^+$  are selected by an optimization of  $S^2/(S+B)$  based on Monte Carlo simulations, where  $S$  and  $B$  are the expected number of signal and background events, respectively. We first determine, based on Monte Carlo simulations, the expected  $S-B$  for each of the decay mode combinations individually. Then, we successively add modes in order of decreasing  $S-B$  to compute an overall  $S^2/(S+B)$  value until  $S^2/(S+B)$  no longer increases. The decay modes used are  $D^0 \rightarrow K^-\pi^+$ ,  $D^0 \rightarrow K^-\pi^+\pi^0$ ,  $D^0 \rightarrow K^-\pi^+\pi^-\pi^+$ ,  $D^0 \rightarrow K_s^0\pi^-\pi^+$ ,  $D^+ \rightarrow K^-\pi^+\pi^+$ ,  $D^+ \rightarrow K_s^0\pi^+$ , and  $D^+ \rightarrow K^-K^+\pi^+$ .  $D^0$  ( $D^+$ ) meson candidates are required to have a reconstructed invariant mass within  $20 \text{ MeV}/c^2$  of the nominal  $D^0$  ( $D^+$ ) mass [9].

Charged kaon candidates are required to be inconsistent with the pion hypothesis, as inferred from the Cherenkov angle measured by the DIRC and the ionizations measured by the SVT and DCH. No particle identification is required for the kaon from the decay  $D^0 \rightarrow K^-\pi^+$ .

$K_s^0 \rightarrow \pi^+\pi^-$  candidates are required to have an invariant mass within  $15 \text{ MeV}/c^2$  of the nominal  $K_s^0$  mass. The angle between the flight direction and the momentum vector of the  $K_s^0$  candidate is required to be less than 200 mrad, and the transverse flight distance from the primary event vertex, obtained from the remaining charged tracks in the event, must be greater than 2 mm.

Neutral pion candidates are formed from pairs of photons in the EMC with energy above 30 MeV, an invariant mass within  $35 \text{ MeV}/c^2$  of the nominal  $\pi^0$  mass, and a summed energy greater than 200 MeV. A mass-constrained fit is applied to these  $\pi^0$  candidates. In the case of the soft  $\pi^0$  from  $D^{*+} \rightarrow D^+\pi^0$  decays, the energy cut is replaced by a momentum cut, in the  $Y(4S)$  frame, of  $70 < p^* < 450 \text{ MeV}/c$ .

To select  $B^0$  candidates with well-reconstructed  $D^{*+}$  and  $D$  mesons, we form a  $\chi^2$  that includes all measured  $D^{*+}$  and  $D$  masses:

$$\chi_{\text{mass}}^2 = \left( \frac{m_D - \hat{m}_D}{\sigma_{m_D}} \right)^2 + \left( \frac{m_{\bar{D}} - \hat{m}_{\bar{D}}}{\sigma_{m_{\bar{D}}}} \right)^2 + \left( \frac{\Delta m_{D^*} - \Delta \hat{m}_{D^*}}{\sigma_{\Delta m_{D^*}}} \right)^2 + \left( \frac{\Delta m_{\bar{D}^*} - \Delta \hat{m}_{\bar{D}^*}}{\sigma_{\Delta m_{\bar{D}^*}}} \right)^2, \quad (1)$$

where the caret over a value refers to the nominal value, and  $\Delta m_{D^*}$  is the  $D^{*+} - D$  mass difference. For  $\sigma_{m_D}$  we use values computed for each  $D$  candidate, while for  $\sigma_{\Delta m_{D^*}}$  we use fixed values of  $0.83 \text{ MeV}/c^2$  for  $D^{*+} \rightarrow D^0\pi^+$  and  $1.18 \text{ MeV}/c^2$  for  $D^{*+} \rightarrow D^+\pi^0$ . A requirement that  $\chi_{\text{mass}}^2 < 20$  is applied to all  $B^0$  candidates. In events with more than one  $B^0$  candidate, we choose the candidate with the lowest value of  $\chi_{\text{mass}}^2$ .

A  $B$  meson candidate is characterized by two kinematic variables: the energy-substituted mass,

$$m_{\text{ES}} \equiv \sqrt{E_{\text{beam}}^{*2} - p_B^{*2}}, \quad (2)$$

and the difference of the  $B$  candidate's energy from the beam energy,

$$\Delta E \equiv E_B^* - E_{\text{beam}}^*. \quad (3)$$

$E_B^*$  ( $p_B^*$ ) are the energy (momentum) of the  $B$  candidate in the center-of-mass frame and  $E_{\text{beam}}^*$  is one-half of the total center-of-mass energy. The signal region in the  $\Delta E$  vs  $m_{\text{ES}}$  plane is defined to be  $|\Delta E| < 25$  MeV and  $5.273 < m_{\text{ES}} < 5.285$  GeV/ $c^2$ . Based on Monte Carlo simulations, the width of this region corresponds to approximately  $\pm 2.5\sigma$  in both  $\Delta E$  and  $m_{\text{ES}}$ .

To determine the expected contribution from background in the signal region, we scale the number of events seen in a sideband in the  $\Delta E$  vs  $m_{\text{ES}}$  plane defined as  $|\Delta E| < 200$  MeV,  $5.20$  GeV/ $c^2 < m_{\text{ES}} < 5.26$  GeV/ $c^2$  and  $50$  MeV  $< |\Delta E| < 200$  MeV,  $5.26 < m_{\text{ES}} < 5.29$  GeV/ $c^2$ . The scaling factor is calculated by parametrizing the shape of the background in the  $\Delta E$  vs  $m_{\text{ES}}$  plane as the product of an ARGUS function [10] in  $m_{\text{ES}}$  and a first-order polynomial in  $\Delta E$ . Based on this parametrization we estimate that the ratio of the number of background events in the signal region to the number of events in the sideband region is  $(1.72 \pm 0.10) \times 10^{-2}$ . The uncertainty is derived from the observed variation of this ratio under alternative assumptions for the background shape in  $m_{\text{ES}}$  and  $\Delta E$  using Monte Carlo simulations. The simulations also indicate that there are no significant sources of background appearing in the signal region beyond that indicated by the sideband extrapolation.

After all selection criteria, 38 events are located in the signal region, with 363 events in the sideband region. The latter, together with the scaling factor determined above, implies an expected number of background events in the signal region of  $6.24 \pm 0.33(\text{stat}) \pm 0.36(\text{syst})$ . The systematic uncertainty comes from the background shape variation mentioned previously. Figure 1 shows a projection of the data on the  $m_{\text{ES}}$  axis after requiring  $|\Delta E| < 25$  MeV.

We use a Monte Carlo simulation of the *BABAR* detector to determine the efficiency for reconstructing the signal. The efficiencies range from 17.4% to 2.7%, depending on the  $D$  decay modes. This, together with the total number of neutral  $B$  mesons produced during data collection, allows us to determine the branching fraction for  $B^0 \rightarrow D^{*+}D^{*-}$  to be

$$\mathcal{B}(B^0 \rightarrow D^{*+}D^{*-}) = [8.3 \pm 1.6(\text{stat}) \pm 1.2(\text{syst})] \times 10^{-4}.$$

The high charged particle multiplicity makes this measurement particularly sensitive to the tracking system. Therefore the dominant systematic uncertainty comes from our level of understanding of the charged particle tracking efficiency. Systematic errors are assigned on a per track basis for  $\pi$ ,  $K$ , and soft  $\pi$ , and are added linearly




FIG. 1. The  $m_{\text{ES}}$  distribution of  $B^0 \rightarrow D^{*+}D^{*-}$  events with  $|\Delta E| < 25$  MeV. The curve represents a fit with the sum of a Gaussian to model the signal, and an ARGUS function [10] to model the background shape.

(9.9%). The effect on acceptance due to the imprecisely known partial-wave content of the  $B^0 \rightarrow D^{*+}D^{*-}$  final state is another source of potential systematic bias (6.6%). Other significant potential systematic biases arise due to the uncertainties on the branching fractions [9] of the  $D^{*+}$ ,  $D^0$ , and  $D^+$  (5.6%) and the uncertainties in mass resolutions of reconstructed mesons (4.1%). The total systematic uncertainty from all considered sources is 15%.

In addition to the branching fraction quoted above, we have also measured the  $CP$ -odd fraction of the final state. This fraction,  $R_{\perp}$ , is determined from the angular distribution of the soft pions in the decay, analyzed in the transversity basis [4]. In this reference frame, three decay amplitudes determine the distribution of three decay angles. Integrating over time,  $B$  flavor, and two of these three angles yields the following expression:

$$\frac{1}{\Gamma} \frac{d\Gamma}{d\cos\theta_{\text{tr}}} = \frac{3}{4}(1 - R_{\perp}) \sin^2\theta_{\text{tr}} + \frac{3}{2}R_{\perp} \cos^2\theta_{\text{tr}}. \quad (4)$$

Here  $\Gamma$  is the decay rate and  $\theta_{\text{tr}}$  is the angle between the normal to the  $D^{*-}$  decay plane and the line of flight of the soft pion from the  $D^{*+}$  evaluated in the  $D^{*+}$  rest frame.

We perform an unbinned maximum likelihood (ML) fit to the 38 events in the signal region described previously. The fit takes into account the presence of background, whose properties are derived from the sideband sample, and the angular resolution  $\sigma_{\theta}$  estimated from Monte Carlo

simulations. We define the likelihood function to be

$$\begin{aligned} \mathcal{L} &= \prod_{i=1,n} \mathcal{L}_i \\ &= \prod_{i=1,n} [p \times \mathcal{F}(\theta_{tr,i}, \sigma_{\theta,i}, R_{\perp}^{\text{sig}}) + (1-p) \\ &\quad \times \mathcal{F}(\theta_{tr,i}, \sigma_{\theta,i}, R_{\perp}^{\text{bkg}})], \end{aligned} \quad (5)$$

where  $n$  is the number of selected events. The contribution to the total likelihood from the  $i$ th event,  $\mathcal{L}_i$ , is defined in terms of the purity,  $p$ , of the sample and the probability density functions  $\mathcal{F}(\theta_{tr,i}, \sigma_{\theta,i}, R_{\perp})$  for the signal and background.  $R_{\perp}^{\text{sig}}$  and  $R_{\perp}^{\text{bkg}}$  are the parameters describing the shapes of the signal and background angular distributions, respectively, and  $\theta_{tr,i}$  is the measured transversity angle in event  $i$ . The probability density functions  $\mathcal{F}$  are obtained from the convolution of the angular distribution [Eq. (4)] with Gaussian resolution functions describing the measurement uncertainties  $\sigma_{\theta,i}$ . From studies of simulated data,  $\sigma_{\theta}$  was measured to be 0.11 (0.12) radians for charged (neutral) slow pions. A 10% uncertainty of these values is considered when estimating the corresponding systematic error.

The value of  $R_{\perp}^{\text{bkg}}$  is evaluated by fitting the 363 events in the sideband region and setting  $p = 0$  in Eq. (5). The result of this fit is  $R_{\perp}^{\text{bkg}} = 0.29 \pm 0.04$ , compatible with the value expected for a uniform distribution ( $R_{\perp} = 1/3$ ). To determine  $R_{\perp}^{\text{sig}}$ , we fit the 38 events in the signal region with  $R_{\perp}^{\text{bkg}}$  fixed to 0.29 and  $p$  fixed at 83.6%. The result of the fit to the signal region, without the correction for angular acceptance bias described below, is  $R_{\perp}^{\text{sig}} = 0.25 \pm 0.18(\text{stat})$ , and is shown in Fig. 2. The probability of obtaining a lower likelihood, evaluated using a Monte Carlo technique, is 66%.

It should be noted that Eq. (4) is the differential decay rate  $\Gamma$  integrated over the full ranges of the other two decay angles in the transversity basis, neglecting any bias in the projected  $\theta_{tr}$  distribution introduced by detector acceptance effects. A detailed study of the kinematics of the decay shows that the incomplete detector coverage of the polar angle with respect to the beam axis does not introduce any bias in the distributions of the decay angles in the transversity basis. However, an inefficiency in detecting soft pions below a threshold in transverse momentum may indeed introduce such a bias due to the correlations between the decay angles and particle momenta in the laboratory frame. The amount of these acceptance losses depends on the population of phase space, determined by the values of the decay amplitudes.

An accurate correction for these acceptance effects requires the complete determination of the decay amplitudes using a full angular analysis on a sufficiently large data sample. To estimate the size of the acceptance bias on  $R_{\perp}$  without knowing the decay amplitudes, the fit procedure was tested on several samples of  $B^0 \rightarrow D^{*+} D^{*-}$  simulated




FIG. 2. The  $\cos\theta_{tr}$  distribution from the unbinned ML fit, superimposed on the histogram of the  $B^0 \rightarrow D^{*+} D^{*-}$  candidates in the signal region. The solid line represents the  $\cos\theta_{tr}$  distribution from the unbinned ML fit for the selected events. The dotted and dashed lines represent the fitted  $CP$ -odd and  $CP$ -even components, respectively, for the signal. The dot-dashed curve represents the fitted background component.

events generated with different sets of decay amplitudes. The different amplitudes affect, to varying extents, the correlated soft pions' transverse momenta and angular distributions. The fitted  $R_{\perp}$  values were found to be consistent with the generated values in the limit of negligible soft pion inefficiency. Depending on the mix of decay amplitudes, they did reveal a bias once the pion-detection threshold was taken into account. Considering the full possible range of decay amplitudes, the calculated bias on  $R_{\perp}$  ranged from  $-0.048$  to  $+0.004$ . The central value of this interval is taken as a correction to the fitted  $R_{\perp}^{\text{sig}}$ , while its half width is taken as an estimate of the corresponding systematic uncertainty (0.026). Additional, smaller systematic uncertainties affecting the  $R_{\perp}$  measurement arise from the imperfect knowledge of the resolution in the transversity angle  $\theta_{tr}$  (0.006), the angular distribution of the background (0.008), and the purity of the signal sample (0.0003). The total systematic uncertainty on  $R_{\perp}$  is determined to be 0.028, giving the final corrected result:

$$R_{\perp} = 0.22 \pm 0.18(\text{stat}) \pm 0.03(\text{syst}).$$

In summary, we have observed a signal of  $31.8 \pm 6.2(\text{stat}) \pm 0.4(\text{syst})$  events in the decay  $B^0 \rightarrow D^{*+} D^{*-}$ . Our measurement of the branching ratio is

$$\mathcal{B}(B^0 \rightarrow D^{*+} D^{*-}) = [8.3 \pm 1.6(\text{stat}) \pm 1.2(\text{syst})] \times 10^{-4}.$$

From the transversity angular distribution of these events, we have also measured the  $CP$ -odd fraction,  $R_{\perp}$ , of the final state. These measurements provide a starting point for measuring time-dependent  $CP$ -violating asymmetries in these decays when more data become available.

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF (Germany), INFN (Italy), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from the A.P. Sloan Foundation, Research Corporation, and Alexander von Humboldt Foundation.

\*Also with Università di Perugia, Perugia, Italy.

†Also with Università della Basilicata, Potenza, Italy.

- [1] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **87**, 091801 (2001).
- [2] Belle Collaboration, K. Abe *et al.* Phys. Rev. Lett. **87**, 091802 (2001).
- [3] Charge-conjugate states are implied throughout this paper and the symbol  $D^{(*)}$  refers to either  $D$  or  $D^*$ .
- [4] I. Dunietz *et al.*, Phys. Rev. D **43**, 2193 (1991).
- [5] CLEO Collaboration, E. Lipeles *et al.*, Phys. Rev. D **62**, 032005 (2000).
- [6] *BABAR* Collaboration, B. Aubert *et al.* Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 1 (2002), hep-ex/0105044 (2001).
- [7] PEP-II Conceptual Design Report No. SLAC-R-418 (1993).
- [8] G. C. Fox and S. Wolfram, Phys. Rev. Lett. **41**, 1581 (1978).
- [9] Particle Data Group, D. E. Groom *et al.*, Eur. Phys. J. C **15**, 1 (2000).
- [10] ARGUS Collaboration, H. Albrecht *et al.*, **48**, 543 (1990).