

Lawrence Berkeley National Laboratory

Recent Work

Title

THE SOURCE OF OXYGEN IN RHODOPSEUDOMONAS SPHEROIDES CAROTENOID PIGMENT CONVERSION

Permalink

<https://escholarship.org/uc/item/4m05g54z>

Author

Shneour, Elie A.

Publication Date

1962-07-01

University of California

**Ernest O. Lawrence
Radiation Laboratory**


TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.



UCRL-10387

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

THE SOURCE OF OXYGEN IN RHODOPSEUDOMONAS SPHEROIDES
CAROTENOID PIGMENT CONVERSION

Elie A. Shneour

July 1962

U.S. GOVERNMENT PRINTING OFFICE

THE SOURCE OF OXYGEN IN RHODOPSEUDOMONAS SPHEROIDES CAROTENOID

PIGMENT CONVERSION

by Elie A. Shneour^a

Bio-Organic Chemistry Group, Lawrence Radiation Laboratory, University of
California, Berkeley 4, California^b

ABSTRACT

When an anaerobic culture of Rhodopseudomonas spheroides was exposed to O_2^{18} , 94.2% (dark) and 89.6% (light) of the theoretical oxygen incorporation into the pigment spheroidenone was observed. These results are consistent with the conclusion that the sole source of the ketone oxygen in this carotenoid pigment is molecular oxygen.

When anaerobic cultures of Rhodopseudomonas spheroides are exposed to oxygen, the yellow carotenoid pigment spheroidene is rapidly converted to the red carotenoid pigment spheroidenone (Table I). This reaction involves the specific oxidation of the methylene group at carbon 2 to a ketone^{1,2}.

It has long been considered likely that molecular oxygen contributes one atom to the spheroidenone molecule, but no direct evidence of this was available. A complicating factor was the observation that the rate and extent of the reaction are enhanced by exposure to light^{2,3}. The availability of nearly pure O_2^{18} and of improved instrumentation and techniques have made a precise mass spectrographic analysis of the product of this reaction possible. This is the object of this report.

Wild type Rhodopseudomonas spheroides, strain 2.4.1. originally obtained from the collection of Professor C.B. van Niel, was grown anaerobically in the manner described earlier² and in amounts sufficient to yield approximately 1 mg of spheroidenone. Pellets of the bacteria, obtained by centrifugation, were resuspended in 10 ml of the modified Hutner's resting cell medium², and transferred to 20 ml flasks connected to a Pyrex glass vacuum manifold and evacuated to 20 mm Hg. A gaseous mixture at S.T.P. consisting of 20% O_2^{18} (98.4 atom %)^c and 80% argon by volume was next introduced into the manifold and the whole system was shaken for three hours at 20°C. One flask was kept in complete darkness while the other was exposed to saturating incandescent illumination during the shaking phase². At the end of this period, the crude pigments were quantitatively extracted according to the method described earlier². Isolation and purification of spheroidenone was obtained by repeated chromatography on 32 x 150 mm columns of Micro-Cel C^d with 1% acetone in light petroleum. Carrier spheroidenone was prepared by the same procedure.

Spheroidenone 482 μ $E_{1\text{ cm}}^{1\%}$ 2120 (acetone-methanol 7:2).

Samples of approximately 5 mg spheroidenone, including carrier, were pyrolysed with Hg_2Cl_2 at 500°C for an hour, to yield CO_2 according to the method of Rittenberg and Pontecorvo⁴. The $^{18}\text{O}/^{16}\text{O}$ ratio in CO_2 was determined on a Consolidated Electroynamics Co. Mass Spectrometer Model 21-130, a cycloid tube type which is capable of good measurements with relatively small samples. Results are shown in Table II. Since there are two oxygen atoms per molecule of spheroidenone, a 100% contribution from molecular oxygen to the ketone would result in a 50% overall incorporation of the isotope in the CO_2 . The observed experimental value of 47.1% for the dark culture sample is consistent, within experimental error, with this assumption. Under the conditions of these experiments, synthesis of spheroidene, with incorporation of label into the methoxyl oxygen is improbable. While the incorporation of the label into the light saturated culture sample was lower by 1.5%, this difference is certainly insufficient to suggest that a contribution from oxygen resulting from photolysis of water was involved. On the basis of the evidence presented here, it is most likely that the sole source of the ketone oxygen in spheroidenone is molecular oxygen.

FOOTNOTES



- a. This work was done during the tenure of an Advanced Research Fellowship of the American Heart Association.
- b. The work described in this paper was sponsored, in part, by the United States Atomic Energy Commission.
- c. Molecular Oxygen-18, 98.4 atom % purity, was obtained from the Isotope Division, Weitzman Institute of Science, Rehovot, Israel.
- d. Micro-Cel C is a synthetic calcium silicate manufactured by Johns Manville Co. For best results, it was necessary to pyrolyse it in an oven at 400°C for one hour before use.

REFERENCES

1. Barber, M.S., L.M. Jackman, B.C.L. Weedon and H. Yokoyama, Proc. Chem. Soc., 1962 (in press).
2. Shneour, E.A., Biochim. Biophys. Acta 62, 534 (1962).
3. van Niel, C.E., Antonie van Leeuwenhoek J. Microbiol. Serol. 12, 156 (1947).
4. Rittenberg, D. and L. Pontecorvo, J. Appl. Radiation and Isotopes 1, 208 (1956).

Table I

Structure of Carotenoid Pigments in R. spheroides.



MU-25232

Table II

¹⁸O Analysis of Spheroidenone

INCUBATION	ATOM %	ATOM % EXCESS	DILUTION FACTOR			ATOM % EXCESS X DILUTION	% INCORPORATION**
			Initial*	Carrier	Total		
Dark	6.77	6.57	1.64	5.41	7.05	46.3	47.1
Light	8.64	8.44	1.64	3.67	5.31	44.8	45.6

Tank CO₂ control: 0.202 ± 0.003 atoms per cent ¹⁸O

* Dilution caused by spheroidenone present before incubation with enriched oxygen.

** Molecular oxygen-18 - 98.2 atoms per cent excess.

[Faint, illegible text covering the majority of the page]

