

# UCLA

## UCLA Previously Published Works

### Title

Effect of State Immunization Information System Based Reminder/Recall for Influenza Vaccinations: A Randomized Trial of Autodialer, Text, and Mailed Messages

### Permalink

<https://escholarship.org/uc/item/4mc0g3zc>

### Authors

Szilagyi, Peter G  
Albertin, Christina S  
Saville, Alison W  
[et al.](#)

### Publication Date

2020-06-01

### DOI

10.1016/j.jpeds.2020.02.020

Peer reviewed



Published in final edited form as:

*J Pediatr.* 2020 June ; 221: 123–131.e4. doi:10.1016/j.jpeds.2020.02.020.

## Effect of State Immunization Information System Based Reminder/Recall for Influenza Vaccinations: A Randomized Trial of Autodialer, Text, and Mailed Messages

Peter Szilagyi, MD MPH<sup>1</sup>, Christina Albertin, BSN, MSPH<sup>1</sup>, Alison W. Saville, MSPH, MSW<sup>2,3</sup>, Rebecca Valderrama, MPH<sup>1</sup>, Abigail Breck, BS<sup>1</sup>, Laura Helmkamp, MS<sup>2,3</sup>, Xinkai Zhou, MS<sup>4</sup>, Sitaram Vangala, MS<sup>4</sup>, L. Miriam Dickinson, PhD<sup>2,3</sup>, Chi-Hong Tseng, PhD<sup>4</sup>, Jonathan D. Campbell, PhD<sup>5</sup>, Melanie Whittington, PhD<sup>4</sup>, Heather Roth, MA<sup>6</sup>, Cynthia Rand, MD MPH<sup>7</sup>, Sharon G Humiston, MD MPH<sup>8</sup>, Dina Hoefer, PhD<sup>9</sup>, Allison Kempe, MD MPH<sup>2,3</sup>

<sup>1</sup>Department of Pediatrics, UCLA Mattel Children's Hospital, University of California at Los Angeles, Los Angeles, California

<sup>2</sup>Adult and Child Consortium for Health Outcomes Research and Delivery Science (ACCORDS), University of Colorado School of Medicine and Children's Hospital Colorado, Aurora, CO

<sup>3</sup>Department of Pediatrics, University of Colorado School of Medicine, Aurora, CO

<sup>4</sup>Department of Medicine, Statistics Core, David Geffen School of Medicine, University of California, Los Angeles, California

<sup>5</sup>Department of Clinical Pharmacy, University of Colorado Anschutz Medical Campus, Aurora, CO

<sup>6</sup>Colorado Immunization Information System, Colorado Department of Public Health and Environment

<sup>7</sup>Department of Pediatrics, University of Rochester School of Medicine and Dentistry, Rochester, NY

<sup>8</sup>Department of Pediatrics, Children's Mercy, Kansas City, Missouri

<sup>9</sup>New York State Immunization Information System, New York State Department of Health

### Abstract

**Objective:** To evaluate the effect of different modalities of centralized reminder/recall (C-R/R) (autodialer, text, mailed reminders) on increasing childhood influenza vaccination.

**Study design:** Two simultaneous randomized clinical trials (RCTs) conducted 10/2017–4/1/2018 in New York State and Colorado. 61,931 children in New York (136 practices) and 23,845 children in Colorado (42 practices) were randomized to different C-R/R modalities—4

---

**Corresponding author:** Peter Szilagyi; Professor and Executive Vice-Chair, Department of Pediatrics, UCLA. 10833 LeConte. MC 175217. Los Angeles, CA 90095. Phone 310-206-6328. FAX 310 206-4855. pszilagyi@mednet.ucla.edu.

The other authors declare no conflicts of interest.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

arms in NY (autodialer, text, mailed, no-reminder control) and 3 arms in Colorado (autodialer, mailed, no-reminder control). Message content was similar across modalities. Up to three reminders were sent for intervention arms. The main outcome measure was receipt of at least one influenza vaccine.

**Results:** In New York, compared with the control arm (26.6%), post-intervention influenza vaccination rates in the autodialer arm (28.0%) were 1.4 percentage points higher; adjusted risk ratio (ARR) 1.06 (1.02, 1.10)], but rates for text (27.6%) and mail (26.8%) arms were not different from controls. In Colorado, compared with the control arm (29.9%), post-intervention influenza vaccination rates for autodialer (32.9%) and mail (31.5%) arms were 3.0 percentage points and 1.6 percentage points higher, respectively [ARRs 1.08 (1.03, 1.12) and 1.06 (1.02, 1.10), respectively]. Compared with the control arm, the incremental cost-per-additional vaccine delivered was \$20 (New York) and \$16 (Colorado) for autodialer messages.

**Conclusions:** Centralized reminder/recall for childhood influenza vaccine was most effective via autodialer, less effective via mail, and not effective via text messages. The impact of each modality was modest. Compared with no reminders, the incremental cost-per-additional vaccine delivered was also modest for autodialer messages.

**Trial registration**—[ClinicalTrials.gov: NCT03294473](https://clinicaltrials.gov/ct2/show/study/NCT03294473) and [NCT03246100](https://clinicaltrials.gov/ct2/show/study/NCT03246100)

Seasonal influenza causes substantial illnesses, hospitalizations, ambulatory visits, and deaths throughout the U.S among both children and adults.<sup>1,2</sup> Despite longstanding national recommendations for influenza vaccination of all children over 6 months of age and Healthy People 2020 goals for >80% childhood influenza vaccination rates,<sup>3</sup> national influenza vaccination rates using the 2017–2018 National Immunization Survey (which does not include provider record checks for influenza vaccine), were only 58% for children 6 months to 17 years of age.<sup>4</sup>

Patient reminder/recall (R/R) can raise influenza vaccination coverage.<sup>5,6</sup> Most published studies of R/R for influenza vaccination involved practice-based reminders, and many of these targeted high-risk subgroups of patients and not an entire population of children.<sup>5</sup> The Task Force on Community Preventive Services recommends sending patients R/R messages for any vaccine,<sup>7</sup> yet only 20–33% of primary care practices send any R/R messages for any vaccinations including influenza vaccination.<sup>8–10</sup> A frequently cited barrier is limited resources.<sup>9,11,12</sup>

Some experts have begun focusing upon centralized R/R (C-R/R) as a potential scalable R/R strategy. Centralized systems such as state immunization information systems (IISs) or health systems can use economies of scale to send reminders to patients. Several studies of IIS-based C-R/R have noted improvements in routine childhood vaccination rates,<sup>13–15</sup> but few investigated the impact of C-R/R for raising *influenza vaccination rates among an entire population of children*. The few prior studies using C-R/R for influenza vaccine have focused upon children with chronic diseases,<sup>8</sup> subsets of the population,<sup>16,17</sup> senior adults,<sup>18,19</sup> pregnant women,<sup>20</sup> or small numbers of practices.<sup>20,21</sup> High vaccine hesitancy<sup>22</sup> for influenza vaccination,<sup>23–25</sup> potential confusion among parents about locations for influenza vaccinations<sup>26</sup> (ie, practices, schools, pharmacies), and general barriers for vaccination<sup>27,28</sup> may reduce the impact of C-R/R for influenza vaccination for the entire child population. On

the other hand, even a small increase in vaccination rates, if scaled up widely, may reduce influenza-related vaccine-related morbidity.<sup>29</sup>

Patient R/R messages can use a variety of modalities- e.g., phone, mail, and text.<sup>5</sup> Phone reminders can involve live person calls or autodialers; autodialers are more scalable.

## Methods

This study evaluated: (1) the effectiveness of C-R/R upon influenza vaccination of an entire population of children, (2) the relative effectiveness of autodialer (phone), text message, or mailed C-R/R, and (3) the cost-effectiveness of different reminder modalities. We simultaneously conducted two multi-arm RCTs of IIS-based C-R/R across New York (NY) and Colorado (CO). We hypothesized *a priori* that C-R/R would raise influenza vaccination rates overall, but did not have *a priori* hypotheses about relative effectiveness by reminder modality.

The Institutional Review Boards (IRB) at UCLA, New York State Department of Health, University of Colorado, and the Colorado Department of Public Health and the Environment all approved this study. The C-R/R messages were sent between October 2017 and December 31, 2017 (Figure 1; available at [www.jpeds.com](http://www.jpeds.com)). In NY, we evaluated a 4-arm RCT (autodialer, text, mail, or no-C-R/R control arm). In CO, we evaluated a 3-arm RCT (autodialer, mail, or no-C-R/R control arm), without a text message arm due to state limitations for sending text messages without consent from recipients.

These two states had different IIS reporting laws, compliance regulations for community vaccinators, and involvement with C-R/R. NY had mandatory IIS reporting for vaccines given to persons <19 years of age; CO did not. CO allowed vaccination of children in pharmacies; NY did not until late in the study data collection. CO had performed IIS-based C-R/R for routine childhood vaccinations; NY had not.

In NY, the setting included 57 counties (2.3 million children) outside of the five New York City boroughs. We excluded New York City which has a different immunization registry. In CO the setting included ten urban counties encompassing Denver Metro (total child population of about 660,000) plus several urban counties in northern and southern Colorado.

### New York State and Colorado Immunization Information Systems and Policies

In both states, practices routinely sent vaccination and demographic information to the state IIS via electronic transfers from practice EHRs or via direct data entry into a web-enabled application.

New York State's IIS (NYSIIS) and the New York City IIS exchange immunization updates found on children who have a home address in the other's jurisdiction. New York mandates that all vaccinators send vaccination data to the IIS. CDC has a standard method for state IISs to report on completeness of IIS data, by comparing the number of unique individuals contained in the IISs with census denominators. In NY the number of children <6 years with 2 vaccination records in NYSIIS was about equal to the actual census count of children <6 years, and the number of adolescents who had 2 vaccination records in NYSIIS was 97%

of the census count of adolescents. Prior to January 25, 2018, NY pharmacies could only administer adult influenza vaccines. Starting 1/25/2018 (after all reminders had been sent) NY began allowing pharmacies to vaccinate children ages  $\geq 2$  without a physician order; however few children were vaccinated at pharmacies. NYSIIS had performed an HPV C-R/R mailed reminder study<sup>30</sup> but no prior C-R/R studies for influenza vaccine.

Colorado's IIS (CIIS), did not have mandated reporting, yet >99% of <6 year-olds, 95% of 6 to 10 year-olds, and 80% of 11 to 17 year-olds had at least two immunization records in CIIS. In CO, pharmacists, local public health agencies, and primary care practices vaccinate children. About half of pharmacies and all local health clinics submitted vaccination data to CIIS, and pharmacies can vaccinate children of any age (although they rarely vaccinate children less than 2 years of age). CIIS had a previous history of C-R/R for childhood vaccination<sup>13-15</sup> but not for influenza vaccine.

### Study Populations

Using a stratified, two-stage cluster sampling approach, with practice as the primary sampling unit and rural/urban location as the strata, we randomly selected practices from the pool of all eligible practices in each IIS, sampling practices proportional to practice size. We then randomly selected children per practice and then within each practice randomly assigned them to autodialer message, text message (NY only), mailed message, or no-reminder groups. In NY we allocated 30% of subjects to autodialer, text message, or no-reminder arms and a smaller number (10%, due to costs) to the mailed C-R/R arm. In CO, a smaller overall sample size was selected, and patients were assigned evenly between the three arms. A statistician for each state performed the randomizations using SAS version 9.4. The sample sizes in both states were sufficient to provide >80% power to detect improvements of 2.5 percentage points in each intervention arm versus the control arm. This conservatively assumes a 50% control arm vaccination rate, a 2-fold Bonferroni correction in CO and a 3-fold correction in NY for the multiple intervention arms, and an overall significance level of 0.05.

Figure 1 outlines practice and patient inclusion/exclusion criteria. We included children 6 months to 17.9 years old, grouped siblings by common phone or address, and randomly selected one child per family in families with  $\leq 1$  child. We excluded children who had been vaccinated prior to 10/18/2017 in NY or 10/2/2017 in CO (start of the reminder fieldwork).

### Centralized Reminder (C-R/R) Fieldwork

**Message Content (Figure 3; available at [www.jpeds.com](http://www.jpeds.com)):** the content of the message was similar in both states and across all intervention modalities. In NY, irrespective of modality (autodialer, text, mail), messages stated that the message was from the NY State Health Department and named the primary care practice. In CO, messages stated that the reminder was on behalf of the primary care practice and also included practice names. Reminders in both states had a different educational message each month.

**Phone Messages:** A commercial telephone company (<http://www.teletask.com>) sent autodialer calls for both states, in English and Spanish (recipients pressed "2" for Spanish).

We used the phone number listed in the IIS as the primary contact number, whether cell or landline. Based upon state preferences and also requirements of the telephone company, both states used an “800” number rather than the practice’s phone number; thus, the families using caller identification methods did not see the child’s primary care practice’s phone number. Voicemails were left if a call went unanswered. Recipients could opt out of receiving further calls by calling a toll-free number, or pressing “9”.

**Text messages:** In NY, Teletask also sent text messages, in English and Spanish. Recipients could reply, “SPAN” to receive the text message in Spanish or “STOP” to opt out.

**Mailed Messages:** We sent postcard messages in English and Spanish. In NY, to comply with privacy concerns, the postcard was folded and sealed with the main message (including the primary care practice name and office phone number) on the interior, and in CO a regular postcard was used. Recipients could call or email to be removed from the reminder list.

**Protocols:** Up to three messages were sent to eligible patients approximately every 4 to 6 weeks. Patients who received an influenza vaccine according to the IIS were removed from the calling list between reminder rounds, and phone numbers and addresses were updated if they were updated in the IIS.

## Outcome Measures

The primary study outcome was IIS-based documentation of 1 influenza vaccine within six months of the start of the study (October 2017 through March 31, 2018 [Figure 1]). We assessed vaccinations from the IISs after April 1 (to allow for data to be uploaded into the ISSs.) Secondary outcomes included costs of the intervention stratified by personnel and other related costs, and cost-effectiveness. We estimated the total cost by summing the costs related to: consensus building and preliminary work; training; software costs; collaboration; implementation meetings; and reminders. We used the viewpoint of the state IIS when calculating cost effectiveness measures.

## Statistical Analyses

The primary analysis compared the effectiveness of IIS-based C-R/R, sent by autodialer, text (NY only), or mail, compared with no-reminder control, in increasing receipt of 1 influenza vaccine. We used generalized linear mixed modeling to assess the impact of C-R/R on receipt of 1 influenza vaccine. We used Poisson regression with a robust variance estimator to obtain risk ratios, and we adjusted for covariates including patient age, receipt of 1 influenza vaccine in the prior 2 years and type of practice. These covariates have been noted in prior studies to affect vaccination rates and could affect the response to our intervention. Possible within-practice correlation was accounted for using a random effect for practice. We also tested interactions between each predictor and study arm to determine whether there were differential intervention effects by covariates. We employed intention-to-treat analyses and used version 9.4, SAS Institute, Inc.

In addition, we performed two additional analyses to test whether the intervention resulted in earlier receipt of influenza vaccine; this is relevant because optimal immunity from the

vaccine is achieved a couple of weeks following vaccination and also influenza can arrive in December in some seasons. We repeated the unadjusted and adjusted Poisson regression to obtain risk ratios for vaccination but now with the endpoint being December 1 rather than March 31. Second, we used time-to-event analyses to assess differences in timeliness of vaccination by study arm, adjusted for clustering of patients within practices.

Cost effectiveness analyses took into account personnel time to plan and send reminders in addition to cost of autodialer, text, and mail messaging. We considered the cost differential to carry out C-R/R for autodialer, text (NY only), and mail vs no IIS-based C-R/R reminder. The total cost for each randomized arm and state was the sum of the cost activities related to personnel and other expenses relevant to a state IIS. We reported the cost per child randomized within each arm and state as well as the incremental cost per additional vaccine delivered for each active arm versus the control arm.

## Results

Enrollment occurred from September 2017 to April 2019. Table 1 shows patient and practice characteristics by state. There were large numbers of practices and children across the practice types, geographic regions, study arms, and age groups. Very few children in either state had opted out of the IIS or had missing phone numbers.

### Influenza Vaccination Rates

**New York:** Influenza vaccination rates (Table 2) were slightly higher (by 1.4% points) in the autodialer arm than in the control arm ( $p=0.007$ ) but not statistically different in the text or mail arms versus the control arm: autodialer-28.0%, text-27.6%, mail-26.8%, control-26.6%. In adjusted models, the probability of vaccination was significantly increased in the autodialer arm compared with the control arm [ARR = 1.6, 95% CI (1.02, 1.10)], but not for the text or mail arms. The intervention effect differed (not shown in tables) by age (interaction effect  $p=.04$ ), but not practice type ( $p=.83$ ), or prior vaccination ( $p=.07$ ). Children 6 months to 2 years were most likely to have received the vaccine and those <11 years were more likely than those 11 years to be vaccinated.

The costs per child randomized were \$0.28 for autodialer, \$0.24 for text and \$1.76 for mail (Table 3). Compared with the control arm, the incremental cost per additional vaccine delivered was \$20 for autodialer and \$24 for text arms, but with no observed statistically significant difference in vaccination rates between text and control arms. The incremental cost for the mail arm was \$869 (more costly but not more effective than the control arm).

**Colorado:** Vaccination rates were higher in the autodialer arm (by 3.0 percentage points,  $P < .001$  in adjusted model) and the mail arm (by 1.6 percentage points,  $p=0.01$  in adjusted model) than the control arm (autodialer-32.9%, mail-31.5%, control-29.9%). In adjusted models, the probability of vaccination was significantly increased compared with the control arm [ARR (95% CI) for autodialer 1.08 (1.03, 1.12), and for mail 1.06 (1.02, 1.10)]. The effect of the intervention did not differ by age (interaction effect  $p=.86$ ), practice type ( $p=.27$ ), region ( $p=.37$ ) or prior vaccination ( $p=.70$ ). Children <11 years (compared with



older subjects) and children who received an influenza vaccine in the prior year (compared with those who did not) were more likely to receive a vaccine during the study season.

The costs per child randomized were \$0.49 for autodialer and \$1.72 for mail (Table 3). Compared with the control arm, the incremental cost per additional vaccine delivered was \$16 for autodialer and \$108 for mail arms.

**Both States:** We performed two post-hoc analyses to assess whether the intervention improved the timeliness of influenza vaccination. Table 4 (available at [www.jpeds.com](http://www.jpeds.com)) shows unadjusted and adjusted risk ratios vaccination by 8 weeks after the start of the study in each state. Compared with the findings for end-of-season vaccination, findings were identical in NY but showed a slightly greater impact in CO. Figure 2 shows time-to-event analyses including Kaplan-Meier curves and smoothed hazard functions to assess differences in timeliness of vaccination by study arm. In NY, the autodialer arm had higher likelihood of vaccination in the first 8 weeks than the control arm. In CO both the autodialer and mailed arms had higher vaccination rates in the first 8 weeks than the control arm.

Finally, we performed a post-hoc analysis utilizing cell phone-scrubbing software to identify landline vs. cell numbers; this did not change the study findings.

## Discussion

Our study tested IIS-based C-R/R for influenza vaccine across large populations and also in comparing different modalities of C-R/R (autodialer, text messaging, and mailed reminders versus controls). We found that IIS-based C-R/R messages sent by autodialer calls had a modest impact on raising influenza vaccination coverage in both NY and CO, and that the impact of autodialer calls in both states occurred largely during the first 8 weeks after the initial call. IIS-based C-R/R sent by text message did not raise influenza vaccination rates in NY. Mailed C-R/R messages had no effect in NY and had a small effect in CO, again primarily during the first 8 weeks.

A Cochrane systematic review noted that R/R was effective in increasing influenza vaccination rates in children.<sup>5</sup> However, only one pediatric study focused on healthy children rather than children with high risk conditions, and this study noted only a 4.4% point improvement for children 6 to 23 months of age.<sup>16</sup> One trial of text reminders for healthy children of all age groups conducted in 2012 across four New York City practices noted an absolute difference of 3.7% points between text message and no-reminder control arms.<sup>31</sup> Thus, the expected effect size for C-R/R for influenza vaccine for the entire child population, whether practice-based or centralized, might be small. Importantly, a recent CDC modeling study of adult plus child influenza vaccination noted that a 5% point improvement in vaccination rates would substantially reduce population-wide influenza morbidity.<sup>29</sup> If an intervention could be scaled up widely, even an impact of 2 to 3 percentage points would have a significant public health benefit.

We evaluated 3 common R/R modalities. Autodialer R/Rs can be scaled-up widely and are low-cost. Text messages might have greater sense of urgency or importance than phone calls and are scalable and low-cost. Mailed R/Rs might be perceived as more important and can



remain in patients' homes as a continuous reminder. One prior study noted that mailed reminders were more effective though also more costly than autodialer reminders;<sup>32</sup> however, that study focused only on low-income adolescents. Because mailed reminders are more costly, they are less scalable than phone or text reminders.

Unlike some other studies,<sup>21,31,33</sup> text messages had no benefit in NY whereas autodialer reminders had a small benefit. Perhaps frequent texting in everyday life diminished the impact of text message reminders. In addition, leaving messages on answering machines might cause autodialer messages to have greater impact than text messages. More study is needed to compare the relative impact of different reminder modalities.

For any R/R to be effective, phone or mail contact information must be accurate. The accuracy of IIS-based contact information is unclear, and the relative accuracy of phone numbers versus mailing addresses is unknown; both depend on data uploads and corrections from immunization providers. Our trials were pragmatic because we did not improve on contact information but rather used existing IIS-based data. Optimizing IIS-based contact information might improve the impact of IIS reminders.

One important issue that has not been well studied is whether R/R, either practice-based or C-R/R, might be less effective today than in the past. The Cochrane review includes R/R studies since 1974, and did not formally test time-trends in effectiveness of R/R. We speculate that there may be some current challenges to IIS-based C-R/R for influenza vaccination. Because of state requests and also requirements of the telephone company, our autodialer calls emanated from an 800 number rather than from the child's pediatric practice phone number. Some parents might automatically ignore or delete calls from unfamiliar sources. It is also possible that the flood of telemarketing calls,<sup>34</sup> now experienced by many individuals, might diminish the potential impact of health-related autodialer reminders. One report noted that there were 3.4 billion robocalls made during one month in 2018, representing >10 calls per US resident that month.<sup>35</sup> Our autodialer calls might have been ignored by many parents due to these concerns.

Underreporting to IISs of childhood influenza vaccination might limit our ability to observe the full impact of the intervention. Underreporting by practices might occur if they do not upload electronic data directly to the IIS or offer cash-only flu vaccine clinics without entering the vaccine into the electronic medical record. Our vaccination coverage was markedly lower than levels reported by the National Immunization Survey and Behavioral Risk Factor Surveillance System (BRFSS) combined data<sup>2</sup> which showed >60% coverage in both states (based on self-report without verification for influenza vaccine) but which might overestimate rates.<sup>36,37</sup> One reason is that we excluded children who had been vaccinated prior to the start of the intervention (October 18 in NY, October 2 in CO). Also, we included both NY, a mandatory reporting state, and CO, a non-mandatory reporting state in which many children receive flu vaccination in pharmacies (where IIS underreporting may be more common). Yet the impact of IIS-based C-R/R was similar in both states. Unfortunately, we cannot quantify the degree of underreporting or its impact on our study. Linkages with insurance/Medicaid databases or billing exchanges, and encouraging providers to use CMS meaningful use standards might improve reporting to state IISs.

Another factor that may have limited the impact of our intervention is influenza vaccine hesitancy due to parent concerns about sub-optimal influenza vaccine effectiveness, vaccine safety, or general vaccine hesitancy. Vaccine hesitancy for influenza is well described<sup>23,25,38</sup> although the prevalence is unknown in either these states or nationally. More intensive Interventions are needed to address vaccine hesitancy.

As expected, we found that younger children and those previously vaccinated were more likely to be vaccinated during our study, but the effect of the intervention didn't vary consistently by these two factors. For older children, and those not previously vaccinated, practice-based interventions or other options such as school-located influenza vaccinations<sup>26</sup> or vaccination in community settings such as pharmacies<sup>9</sup> should be considered.

Most of the effect of the autodialer and (for CO) mail reminders occurred during the first 8 weeks; for these study arms the intervention raised overall vaccination rates and also shifted vaccinations earlier. This might be beneficial because the benefit of the vaccine is optimal >2 weeks after delivery, and because the onset of influenza epidemics is variable across years. Also, the unexpected bump in vaccination rates around January for all groups (including controls) may have been due to media reports of influenza disease.<sup>39-41</sup>

The cost of centralized R/R in this study was approximately \$0.28 to \$0.49 per child reminded by autodialer or text. Our estimates are most similar to a previous study that calculated the mean cost of IIS-based C-R/R using an autodialed method as \$0.53 per contact.<sup>14</sup> A study of C-R/R for a relatively small adolescent population reported \$0.78 per adolescent sent a reminder.<sup>42</sup> The incremental cost per additional vaccine delivered comparing the autodialer arms to the control arm were relatively consistent across both NY and CO, i.e., an estimated additional cost of \$20 (NY) or \$16 (CO) to deliver an additional influenza vaccine in the autodialer arm. Our cost-effectiveness finding is similar to the cost per child for any preschool immunization for children 19–35 months of age using IIS-based C-R/R.<sup>14</sup> These cost-effectiveness estimates could help decision makers who face competing implementation alternatives and fixed budgets. Some health systems or practices, particularly those receiving additional reimbursement for influenza vaccination rates, might consider such costs worthwhile.

Our study has both strengths and limitations. The use of pragmatic trials across 2 states enhance generalizability. The study design (randomization within practices and large numbers of practices and patients) enhances internal validity by allowing us to control for multiple potential confounders. Limitations include an inability to use practice telephone numbers that might have been recognized by parents in the autodialer or text message arms, a potential underreporting to the IISs of updated contact information and influenza vaccination that would blunt our ability to detect an intervention effect, and data from only two states. In addition, some other R/R studies<sup>31,43,44</sup> have employed more reminder messages than the three we sent; we limited to three reminders based upon prior input from state leaders and parents. We received very few calls or letters from parents objecting to the reminders, although a small number opted out of future autodialer, text, or mail reminders in NY (1%, 2%, and <1%) or CO (3%, 1%, and 1%). Of note, there were no adverse outcomes requiring reporting to the IRBs.

In conclusion, our pragmatic trials lend support to centralized reminder-recall for influenza vaccination among children, particularly using autodialer or mailed reminders although autodialer reminders are more cost-effective. Our findings do not support text message reminders for IIS-based C-R/R. Because of its scalability and potential low cost, and evidence that even small improvements in influenza vaccination rates at the population level could prevent substantial morbidity, IIS-based C-R/R remains a viable option for raising influenza vaccination rates. However, more intensive interventions are needed to increase substantially the U.S.'s rates of childhood influenza vaccination.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

We thank Megan D Meldrum at the New York State Department of Health for New York State data curation and contributing largely to the project.

Supported by the National Institute of Allergy and Infectious Diseases of the National Institutes of Health (R01AI114903 [to P.S.]). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. The funding organization was not involved in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; or the decision to submit the manuscript for publication. The authors had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. S.H. serves as a consultant to Sanofi Pasteur.


Portions of this study were presented at the Pediatric Academic Societies Annual Meeting, April 24-May 1, 2019, Baltimore, Maryland.

## REFERENCES

1. Grohskopf LA, Sokolow LZ, Broder KR, et al. Prevention and Control of Seasonal Influenza with Vaccines. *MMWR Recomm Rep.* 2016;65(5):1–54.
2. Centers for Disease Control and Prevention. FluVaxView: Influenza vaccination coverage estimates for persons 6 months and older by State, HHS Region, and the United States, National Immunization Survey-Flu (NIS-Flu) and Behavioral Risk Factor Surveillance System (BRFSS), 2015–16 influenza season. <https://www.cdc.gov/flu/fluvoxview/reportshtml/reporti1516/reportii/index.html> . Accessed January 21, 2020.
3. Office of Disease Prevention and Health Promotion. Healthy People 2020. Increase the percentage of children aged 6 months through 17 years who are vaccinated annually against seasonal influenza Web site. [https://www.healthypeople.gov/node/6359/data\\_details#revision\\_history\\_header](https://www.healthypeople.gov/node/6359/data_details#revision_history_header). Published 2018. Accessed January 21, 2020.
4. Centers for Disease Control and Prevention, National Center for Immunization and Respiratory Diseases (NCIRD), Estimates of Flu Vaccination Coverage among Children — United States, 2017–18 Flu Season. <https://www.cdc.gov/flu/fluvoxview/coverage-1718estimates-children.htm>. Accessed January 21, 2020.
5. Jacobson Vann JC, Jacobson RM, Coyne-Beasley T, Asafu-Adjei JK, Szilagyi PG. Patient reminder and recall interventions to improve immunization rates. *Cochrane Database Syst Rev.* 2018;1:CD003941.
6. Szilagyi PG, Bordley C, Vann JC, et al. Effect of patient reminder/recall interventions on immunization rates: A review. *JAMA.* 2000;284(14):1820–1827. [PubMed: 11025835]
7. Briss PA, Rodewald LE, Hinman AR, et al. Reviews of evidence regarding interventions to improve vaccination coverage in children, adolescents, and adults. The Task Force on Community Preventive Services. *Am J Prev Med.* 2000;18(1 Suppl):97–140.

8. Dombkowski KJ, Harrington LB, Dong S, Clark SJ. Seasonal influenza vaccination reminders for children with high-risk conditions: a registry-based randomized trial. *Am J Prev Med*. 2012;42(1):71–75. [PubMed: 22176850]
9. Saville AW, Szilagyi P, Helmkamp L, et al. Potential Strategies to Achieve Universal Influenza Vaccination for Children: Provider Attitudes in Two States. *Acad Pediatr*. 2018;18(8):873–881. [PubMed: 30031132]
10. Tierney CD, Yusuf H, McMahon SR, et al. Adoption of reminder and recall messages for immunizations by pediatricians and public health clinics. *Pediatrics*. 2003;112(5):1076–1082. [PubMed: 14595049]
11. Saville AW, Albright K, Nowels C, et al. Getting under the hood: exploring issues that affect provider-based recall using an immunization information system. *Acad Pediatr*. 2011;11(1):44–49. [PubMed: 21272823]
12. Kempe A, Wortley P, O’Leary S, et al. Pediatricians’ attitudes about collaborations with other community vaccinators in the delivery of seasonal influenza vaccine. *Acad Pediatr*. 2012;12(1):26–35. [PubMed: 21900066]
13. Kempe A, Saville A, Dickinson LM, et al. Population-based versus practice-based recall for childhood immunizations: a randomized controlled comparative effectiveness trial. *Am J Public Health*. 2013;103(6):1116–1123. [PubMed: 23237154]
14. Kempe A, Saville AW, Dickinson LM, et al. Collaborative centralized reminder/recall notification to increase immunization rates among young children: a comparative effectiveness trial. *JAMA Pediatr*. 2015;169(4):365–373. [PubMed: 25706340]
15. Kempe A, Saville AW, Beaty B, et al. Centralized Reminder/Recall to Increase Immunization Rates in Young Children: How Much Bang for the Buck? *Acad Pediatr*. 2017;17(3):330–338. [PubMed: 27913163]
16. Kempe A, Daley MF, Barrow J, et al. Implementation of universal influenza immunization recommendations for healthy young children: results of a randomized, controlled trial with registry-based recall. *Pediatrics*. 2005;115(1):146–154. [PubMed: 15629993]
17. Baker AM, McCarthy B, Gurley VF, Yood MU. Influenza immunization in a managed care organization. *J Gen Intern Med*. 1998;13(7):469–475. [PubMed: 9686713]
18. Mullooly JP. Increasing influenza vaccination among high-risk elderly: a randomized controlled trial of a mail cue in an HMO setting. *Am J Public Health*. 1987;77(5):626–627. [PubMed: 3565663]
19. Nexoe J, Kragstrup J, Ronne T. Impact of postal invitations and user fee on influenza vaccination rates among the elderly. A randomized controlled trial in general practice. *Scand J Prim Health Care*. 1997;15(2):109–112. [PubMed: 9232713]
20. Stockwell MS, Westhoff C, Kharbanda EO, et al. Influenza Vaccine Text Message Reminders for Urban, Low-Income Pregnant Women: A Randomized Controlled Trial. *Am J Public Health*. 2013.
21. Stockwell MS, Hofstetter AM, DuRivage N, et al. Text message reminders for second dose of influenza vaccine: a randomized controlled trial. *Pediatrics*. 2015;135(1):e83–91. [PubMed: 25548329]
22. Edwards KM, Hackell JM, Committee On Infectious Diseases TCOP, Ambulatory M. Countering Vaccine Hesitancy. *Pediatrics*. 2016;138(3).
23. Cunningham RM, Minard CG, Guffey D, Swaim LS, Opel DJ, Boom JA. Prevalence of Vaccine Hesitancy Among Expectant Mothers in Houston, Texas. *Acad Pediatr*. 2018;18(2):154–160. [PubMed: 28826731]
24. Real FJ, DeBlasio D, Beck AF, et al. A Virtual Reality Curriculum for Pediatric Residents Decreases Rates of Influenza Vaccine Refusal. *Acad Pediatr*. 2017;17(4):431–435. [PubMed: 28126612]
25. Strelitz B, Gritton J, Klein EJ, et al. Parental vaccine hesitancy and acceptance of seasonal influenza vaccine in the pediatric emergency department. *Vaccine*. 2015;33(15):1802–1807. [PubMed: 25744225]
26. Szilagyi PG, Schaffer S, Rand CM, et al. Impact of elementary school-located influenza vaccinations: A stepped wedge trial across a community. *Vaccine*. 2018;36(20):2861–2869. [PubMed: 29678459]

27. Rand CM, Goldstein NPN. Patterns of Primary Care Physician Visits for US Adolescents in 2014: Implications for Vaccination. *Acad Pediatr*. 2018;18(2S):S72–S78. [PubMed: 29502641]
28. Allison MA, O’Leary ST, Lindley MC, et al. Financing of Vaccine Delivery in Primary Care Practices. *Acad Pediatr*. 2017;17(7):770–777. [PubMed: 28600199]
29. Hughes MM, Reed C, Flannery B, et al. Projected Population Benefit of Increased Effectiveness and Coverage of Influenza Vaccination on Influenza Burden - United States. *Clin Infect Dis*. 2019.
30. Coley S, Hoefler D, Rausch-Phung E. A population-based reminder intervention to improve human papillomavirus vaccination rates among adolescents at routine vaccination age. *Vaccine*. 2018;36(32 Pt B):4904–4909. [PubMed: 30037480]
31. Stockwell MS, Kharbanda EO, Martinez RA, Vargas CY, Vawdrey DK, Camargo S. Effect of a text messaging intervention on influenza vaccination in an urban, low-income pediatric and adolescent population: a randomized controlled trial. *JAMA*. 2012;307(16):1702–1708. [PubMed: 22535855]
32. Szilagyi PG, Albertin C, Humiston SG, et al. A randomized trial of the effect of centralized reminder/recall on immunizations and preventive care visits for adolescents. *Acad Pediatr*. 2013;13(3):204–213. [PubMed: 23510607]
33. Stockwell MS, Kharbanda EO, Martinez RA, et al. Text4Health: impact of text message reminder-recalls for pediatric and adolescent immunizations. *Am J Public Health*. 2012;102(2):e15–21.
34. Fazzini K. CNBC. Robocalls jumped 60 percent in the U.S. last year and scammers are finding more ways to make money. 1 4, 2019 <https://www.cnn.com/2019/01/02/as-robo-calling-ramps-up-consumers-increasingly-wonder-why-carriers-cant-stop-scammers-from-spoofing-their-phone-numbers.html>. Accessed January 21, 2020.
35. Johnson TP. Presidential Address Legitimacy, Wicked Problems, and Public Opinion Research. *Public Opin Quart*. 2018;82(3):614–621.
36. Centers for Disease Control and Prevention. . <https://www.cdc.gov/mmwr/preview/mmwrhtml/mm5916a1.htm>. Published 2010. Accessed December 4 Interim Results: State-Specific Seasonal Influenza Vaccination Coverage --- United States, August 2009--January 2010.
37. Santibanez TA, Grohskopf LA, Zhai Y, Kahn KE. Complete Influenza Vaccination Trends for Children Six to Twenty-Three Months. *Pediatrics*. 2016;137(3):e20153280.
38. Healy CM, Rench MA, Montesinos DP, Ng N, Swaim LS. Knowledge and attitudes of pregnant women and their providers towards recommendations for immunization during pregnancy. *Vaccine*. 2015;33(41):5445–5451. [PubMed: 26307234]
39. Centers for Disease Control and Prevention. CDC: Flu Season Ongoing with Tens of Thousands Hospitalized So Far. 1 11, 2018 <https://www.cdc.gov/flu/spotlights/flu-season-updates-2018.htm>. Accessed January 16, 2020.
40. New York Times. The Flu Is Widespread in the U.S., and It’s Not Too Late to Get Vaccinated. 1 11, 2018 <https://www.nytimes.com/2019/01/11/health/flu-widespread-cdc-.html> . Accessed January 16, 2020.
41. Yoo BK, Frick K. Determinants of influenza vaccination timing. *Health Econ*. 2005;14(8):777–791. [PubMed: 15700301]
42. Whittington MD, Gurfinkel D, I L, Lockhart L, Beatty B, Dickinson M, Roth H. & Kempe A. Cost of Centralized and Decentralized Reminder/Recall for Accountable Care Organizations. *American Journal of Accountable Care*. Forthcoming 2018.
43. Rand CM, Brill H, Albertin C, et al. Effectiveness of centralized text message reminders on human papillomavirus immunization coverage for publicly insured adolescents. *J Adolesc Health*. 2015;56(5 Suppl):S17–20. [PubMed: 25863549]
44. Dini EF, Linkins RW, Sigafos J. The impact of computer-generated messages on childhood immunization coverage(2)(2). *Am J Prev Med*. 2000;19(1):68–70. [PubMed: 10865167]



|                  | Eligibility criteria by state   |  |
|------------------|---|--|
|                  | New York  | Colorado   |
| <b>Practices</b> | Eligibility: <ul style="list-style-type: none"> <li>Pediatric, family medicine, or community health centers in 57 counties outside of New York City</li> </ul> Ineligibility: <ul style="list-style-type: none"> <li>Large HMOs with own R/R systems</li> </ul> | Eligibility: <ul style="list-style-type: none"> <li>Pediatric, family medicine, or community health centers located in 10 urban counties</li> <li>Serve minimum of 200 children</li> <li>Sent immunization data via HL7 data exchange to CIIS</li> <li>Had &gt;20% flu vaccination recorded at CIIS 2016-2017</li> <li>Uploaded immunizations into IIS within 3 mon prior to study</li> </ul> Ineligibility: <ul style="list-style-type: none"> <li>Large HMOs with own R/R systems (e.g. Kaiser)</li> </ul> |

- Practices that opted out

Opt out of study:

- Possible


Opt out of study:

- Not possible since R/R by the Public Health Department is permitted by legislation

|              | New York   | Colorado  |
|--------------|--|---|
| <b>Child</b> | <p>Eligibility:</p> <ul style="list-style-type: none"> <li>• Age 6 months – 17.9 years</li> </ul> <p>Ineligibility:</p> <ul style="list-style-type: none"> <li>• Sibling/household members (identifiable by duplicate phone numbers)</li> <li>• Patients unlikely to be residing in the state- if:                             <ul style="list-style-type: none"> <li>○ Only demographic information from a birth certificate or the birth Hepatitis B dose</li> <li>○ Ages 13 – 17.0 with no immunizations in IIS since 9th birthday (since school Tdap requirements)</li> <li>○ Ages 6.0 – 12.9 with no immunizations in IIS since 4th birthday (since school vaccine requirements)</li> <li>○ Ages 0 – 17 with no immunizations aside from a single dose of Hepatitis B</li> </ul> </li> </ul> <p>Excluded children who were vaccinated prior to October 18, 2017 – i.e., before the first reminder</p> <p>Patients who opted out of IIS (&lt;2%)</p> | <p>Eligibility:</p> <ul style="list-style-type: none"> <li>• Age 6 months – 17.9 years</li> </ul> <p>Ineligibility:</p> <ul style="list-style-type: none"> <li>• Sibling/household members (identifiable by duplicate phone numbers)</li> <li>• Patients unlikely to be residing in the state- if:                             <ul style="list-style-type: none"> <li>○ Only demographic information from a birth certificate or the birth Hepatitis B dose</li> <li>○ Ages 13 – 17.0 with no immunizations in IIS since 9th birthday (since school Tdap requirements)</li> <li>○ Ages 6.0 – 12.9 with no immunizations in IIS since 4th birthday (since school vaccine requirements)</li> <li>○ Ages 0 – 17 with no immunizations aside from a single dose of Hepatitis B</li> </ul> </li> </ul> <p>Excluded children who were vaccinated prior to October 2, 2017 – i.e., before the first reminder</p> |

**Figure 1:**  
Consort Diagram plus practice and patient eligibility criteria





**Figure 2:**

Time-to-event analysis of influenza vaccinations

Cumulative incidence curves **A** and **C** above show the cumulative percent of study subjects vaccinated over time in NY ( $p=0.004$ ) and CO ( $p<0.001$ ), with final vaccination rates among children not already vaccinated at the start of the study ranging from 26–33%. Smoothed hazard plots **B** and **D** show the probability of an unvaccinated subject receiving a vaccination at any given time through the study period. The peak vaccination time for both states occurred shortly after the start of the intervention in October, November, and early December, with instantaneous vaccination rates decreasing through the remainder of the season apart from a small uptick around late January. Differences between study arms were observed primarily during peak vaccination time in the first 8 weeks of the study; at the end of 8 weeks adjusted RR in NY for autodial, text, and mail were 1.06 [1.01, 1.11], 1.04 [0.99, 1.09], 0.90 [0.83, 0.98] respectively and in CO for autodial and mail arms were 1.13 (1.08, 1.19) and 1.10 (1.05, 1.15) respectively.

| <b>October</b>  |
|---|
| <p><b>Autodialer Script</b></p> <p>This is a call regarding your child’s health. Press 1 for English; or presiona 2 para Español. Hello! This is an important message on behalf of [Practice Name] and the [State] Department of Health. Our records show that your child is due for a flu vaccine. All children over 6 months old need the flu shot to keep them healthy. Please call [practice name] at [practice phone number] to schedule a vaccine visit. If you would like to be removed from this list, please press 9. Please press 0 to repeat this message.*</p>              |
| <p><b>Text Message</b></p> <p>[Practice Name/StateDeptofHealth]: Our records show that your child is due for the flu vaccine. All children over 6 months old need the flu shot to keep them healthy. Call [Practice Name] at [practice phone number] to schedule a vaccine visit. Responde SPAN para Español. Reply STOP to opt out</p>   |
| <b>November</b>   |
| <p><b>Autodialer Script</b></p> <p>This is a call regarding your child’s health. Press 1 for English; or presiona 2 para Español. Hello! This is an important message on behalf of [Practice Name] and the [State] Department of Health. Our records show that your child is due for a flu vaccine. The flu vaccine is safe and people cannot get the flu from the shot. Please call [practice name] at [practice phone number] to schedule a vaccine visit. If you would like to be removed from this list, please press 9. Please press 0 to repeat this message.</p>                 |
| <p><b>Text Message</b></p> <p>[Practice Name/StateDeptofHealth]: Our records show that your child is due for the flu vaccine. The flu vaccine is safe and people CANNOT get the flu from the shot. Call [Practice Name] at [practice phone number] to schedule a vaccine visit. Responde SPAN para Español. Reply STOP to opt out</p>   |
| <b>December</b>   |
| <p><b>Autodialer Script</b></p> <p>This is a call regarding your child’s health. Press 1 for English; or presiona 2 para Español. Hello! This is an important message on behalf of [Practice Name] and the [State] Department of Health. Our records show that your child is due for a flu vaccine. Since the flu virus changes every year, it’s important to get vaccinated every year. Please call [practice name] at [practice phone number] to schedule a vaccine visit. If you would like to be removed from this list, please press 9. Please press 0 to repeat this message.</p> |
| <p><b>Text Message</b></p> <p>[Practice Name/StateDeptofHealth]: Our records show that your child is due for the flu vaccine. Since the flu virus changes every year, it’s important to get vaccinated every year. Call [Practice Name] at [practice phone number] to schedule a vaccine visit. Responde SPAN para Español. Reply STOP to opt out</p>   |

**Online Figure.**

Autodialer and text message content in the two states.

\*Note: Colorado did not include “state health department” in scripts.

**Table 1:**

Characteristics of study practices and patients in each state (intervention and control subjects combined).

| Characteristics                   |   | New York n (%) | Colorado n (%) |
|-----------------------------------|---|----------------|----------------|
| <b>Practices</b>                  |   |                |                |
| <b>Number</b>                     |   | 136            | 42             |
| <b>Practice Type</b>              | Pediatric                                   | 95 (69.9)      | 19 (45.2)      |
|                                   | Family Medicine                             | 31 (22.8)      | 19 (45.2)      |
|                                   | Community Health Center (CHC)               | 10 (7.4)       | 4 (9.5)        |
| <b>Geography</b>                  | <u>New York</u>                             |                |                |
|                                   | -Downstate (mostly urban)                   | 57 (41.9)      |                |
|                                   | -Upstate urban                              | 22 (16.2)      |                |
|                                   | -Upstate rural                              | 57 (41.9)      |                |
|                                   | <u>Colorado</u>                             |                |                |
|                                   | -Metro (Denver and surrounds)               |                | 21 (50.0)      |
|                                   | -North of Denver                            |                | 12 (28.6)      |
| -South of Denver                  |   | 9 (21.4)       |                |
| <b>Children</b>                   |   |                |                |
| <b>Number</b>                     | All study arms                              | 61931          | 23,845         |
| <b>Study Arm</b>                  | Autodialer                                  | 19003 (30.7)   | 7,910 (33.2)   |
|                                   | Mail  | 4779 (7.7)     | 8,007 (33.6)   |
|                                   | Text  | 19090 (30.8)   | N/A            |
|                                   | Control                                     | 19059 (30.8)   | 7,928 (33.2)   |
| <b>Age Group</b>                  | 6 months to <2 years                        | 3967 (6.4)     | 1,952 (8.2)    |
|                                   | 2–5 years                                   | 13620 (22.0)   | 5,810 (24.4)   |
|                                   | 6–10 years                                  | 18728 (30.2)   | 6,534 (27.4)   |
|                                   | 11–17 years                                 | 25616 (41.4)   | 9,549 (40.0)   |
| <b>Vaccinated in last 2 years</b> | Vaccinated in 2015–2016 or 2016–2017 season | 24957 (40.3)   | 10,457 (43.9)  |
| <b>Miscellaneous</b>              | Total patients that opted-out* of R/R       | 1199(1.9)      | NA             |
|                                   | Missing phone numbers                       | 129(0.2)       | 910 (3.8)      |

\* Opted-out by calling to have their name removed from the recall list or pressing 9 during autodial message (this was only possible in New York).

**Table 2:**

Vaccination rates at the end of the study and unadjusted and adjusted risk ratios for influenza vaccination by patient characteristics and study arm.

|                                   | Category             | n by category | Vaccinated per category, (%) | Unadjusted Risk Ratio    | Adjusted Risk Ratio      |
|-----------------------------------|----------------------|---------------|------------------------------|--------------------------|--------------------------|
| <b>New York</b>                   |                      |               |                              |                          |                          |
| <b>Age category</b>               | 6 months – 1.9 years | 3967          | 52.3                         | <b>2.40 [2.28, 2.53]</b> | <b>2.28 [2.16, 2.40]</b> |
|                                   | 2 – 5.9 years        | 13620         | 30.8                         | <b>1.41 [1.35, 1.47]</b> | <b>1.09 [1.05, 1.14]</b> |
|                                   | 6 – 10.9 years       | 18728         | 25.8                         | <b>1.17 [1.12, 1.21]</b> | <b>1.10 [1.06, 1.14]</b> |
|                                   | 11 – 17.9 years      | 25616         | 22.4                         | -Reference-              | -Reference-              |
| <b>Practice type</b>              | Family Medicine      | 7273          | 20.9                         | -Reference-              | -Reference-              |
|                                   | Pediatrics           | 49390         | 28.1                         | <b>1.38 [1.17, 1.62]</b> | <b>1.17 [1.04, 1.33]</b> |
|                                   | CHC/RHC              | 5268          | 23.5                         | 1.20 [0.90, 1.59]        | 0.98 [0.80, 1.20]        |
| <b>Rurality</b>                   | Downstate            | 27456         | 26.9                         | -Reference-              | -Reference-              |
|                                   | Upstate rural        | 8540          | 24.8                         | <b>0.82 [0.67, 0.99]</b> | 0.93 [0.82, 1.07]        |
|                                   | Upstate urban        | 25935         | 29.1                         | 1.02 [0.88, 1.17]        | 1.05 [0.95, 1.16]        |
| <b>Vaccinated in last 2 years</b> | Unvaccinated         | 36974         | 10.6                         | -Reference-              | -Reference-              |
|                                   | Vaccinated           | 24957         | 51.3                         | <b>4.68 [4.51, 4.85]</b> | <b>4.27 [4.13, 4.40]</b> |
| <b>Study Arm</b>                  | Autodialer           | 19003         | 28.0                         | <b>1.06 [1.02, 1.10]</b> | <b>1.06 [1.02, 1.10]</b> |
|                                   | Text                 | 19090         | 27.6                         | 1.04 [1.00, 1.08]        | 1.03 [0.99, 1.07]        |
|                                   | Mail                 | 4779          | 26.8                         | 1.01 [0.95, 1.07]        | 1.00 [0.94, 1.06]        |
|                                   | Usual care           | 19059         | 26.6                         | -Reference-              | -Reference-              |
| <b>Colorado</b>                   |                      |               |                              |                          |                          |
| <b>Age category</b>               | 6 months – 1.9 years | 1,952         | 59.5                         | <b>2.43 (2.24, 2.63)</b> | <b>2.15 (2.02, 2.30)</b> |
|                                   | 2 – 5.9 years        | 5,810         | 35.9                         | <b>1.47 (1.32, 1.63)</b> | <b>1.16 (1.10, 1.23)</b> |
|                                   | 6 – 10.9 years       | 6,534         | 29.3                         | <b>1.20 (1.11, 1.30)</b> | <b>1.15 (1.10, 1.20)</b> |
|                                   | 11 – 17.9 years      | 9,549         | 24.5                         | -Reference-              | -Reference-              |
| <b>Practice type</b>              | Family Medicine      | 6,451         | 27.7                         | -Reference-              | -Reference-              |
|                                   | Pediatrics           | 15,071        | 33.0                         | 1.19 (0.99, 1.44)        | <b>1.17 (1.06, 1.28)</b> |
|                                   | CHC/RHC              | 2,323         | 31.6                         | 1.14 (0.92, 1.42)        | <b>1.17 (1.02, 1.34)</b> |
| <b>Rurality</b>                   | Metro                | 11,083        | 35.3                         | -Reference-              | -Reference-              |
|                                   | North                | 5,431         | 30.5                         | 0.86 (0.72, 1.03)        | 1.02 (0.92, 1.13)        |
|                                   | South                | 7,331         | 26.3                         | <b>0.75 (0.63, 0.88)</b> | <b>0.88 (0.79, 0.99)</b> |
| <b>Vaccinated in last 2 years</b> | Unvaccinated         | 13,388        | 13.8                         | -Reference-              | -Reference-              |
|                                   | Vaccinated           | 10,457        | 54.0                         | <b>3.91 (3.56, 4.29)</b> | <b>3.71 (3.40, 4.05)</b> |
| <b>Study Arm</b>                  | Autodialer           | 7,910         | 32.9                         | <b>1.10 (1.05, 1.15)</b> | <b>1.08 (1.03, 1.12)</b> |
|                                   | Mail                 | 8,007         | 31.5                         | <b>1.05 (1.01, 1.10)</b> | <b>1.06 (1.02, 1.10)</b> |
|                                   | Usual care           | 7,928         | 29.9                         | -Reference-              | -Reference-              |

vaccination by patient characteristics and study arm.

Overall Interaction p-values:

New York: Age \* study arm p=0.03; Practice type \* study arm p=0.92; Region \* study arm p=0.34; Prior 2yr vaccination \* study arm p= 0.10

Colorado: Age \* study arm p=0.83; Practice type \* study arm p=0.01; Region \* study arm p=0.63; Prior 2yr vaccination \* study arm p= 0.85

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Table 3:**

Cost of centralized reminder-recall

| Type of Cost                                | New York                 |                    |                   |                       | Colorado                |                   |                      |
|---|--------------------------|--------------------|-------------------|-----------------------|-------------------------|-------------------|----------------------|
|   | Autodialer<br>(n=19,003) | Text<br>(n=19,090) | Mail<br>(n=4,779) | Control<br>(n=19,059) | Autodialer<br>(n=7,910) | Mail<br>(n=8,007) | Control<br>(n=7,928) |
| Personnel                                   | \$2,473                  | \$2,213            | \$2,133           | \$0                   | \$1,683                 | \$1,670           | \$0                  |
| Other                                       | \$2,753                  | \$2,395            | \$6,265           | \$0                   | \$2,195                 | \$12,131          | \$0                  |
| <b>TOTAL</b>                                | <b>\$5,226</b>           | <b>\$4,608</b>     | <b>\$8,398</b>    | <b>\$0</b>            | <b>\$3,878</b>          | <b>\$13,801</b>   | <b>\$0</b>           |
| Cost per child randomized                   | \$0.28                   | \$0.24             | \$1.76            | \$0                   | \$0.49                  | \$1.72            | \$0                  |
| Percent vaccinated                          | 28.0%                    | 27.6%              | 26.8%             | 26.6%                 | 32.9%                   | 31.5%             | 29.9%                |
| Incremental cost per additional vaccination | \$20                     | \$24               | \$879             | -                     | \$16                    | \$108             | -                    |

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript