

**University of California**

**Ernest O. Lawrence  
Radiation Laboratory**

**TWO-WEEK LOAN COPY**

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545*

**A NEW REGION OF ALPHA RADIOACTIVITY**

**Berkeley, California**

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Submitted to Phys. Rev. Letters

UCRL-11765

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory  
Berkeley, California

AEC Contract No. W-7405-eng-48

A NEW REGION OF ALPHA RADIOACTIVITY

Ronald D. Macfarlane and Antti Siivola

November 1964

A NEW REGION OF ALPHA RADIOACTIVITY\*

Ronald D. Macfarlane  
Department of Chemistry  
McMaster University  
Hamilton, Canada

and

Antti Siivola  
Lawrence Radiation Laboratory  
University of California  
Berkeley, California

Alpha radioactivity is a mode of decay commonly associated with the heavier elements. In this paper we report on a discovery of alpha decay from nuclides in the region of mass 100.

It has long been recognized from atomic mass data that an enhancement of the alpha decay energies of the tellurium isotopes is present due to the effect of the 50-proton closed shell. The enhancement, however, is not large enough to produce a detectable instability toward alpha particle emission among the tellurium isotopes presently known. Some unsuccessful attempts to produce neutron-deficient tellurium alpha emitters by high energy proton spallation have been previously reported<sup>1,2</sup>. We have obtained results on the alpha radioactivity of the very light tellurium isotopes whose alpha decay energies are additionally enhanced by the  $N = 50$  closed shell.

We bombarded a 90% enriched  $\text{Ru}^{96}$  target with high energy  $\text{O}^{16}$  ions from the Berkeley heavy-ion linear accelerator (Hilac), and obtained alpha particle spectra of the products using techniques described in an earlier paper<sup>3</sup>. Two weak alpha groups (established by  $dE/dx$  experiments) were observed at 3.28 MeV and 3.08 MeV which decayed with half-lives of 2.2 sec and 5.3 sec. respectively. An alpha particle spectrum of these activities is shown in Fig. 1. We established that these activities were due to isotopes of tellurium when we observed that they could not be produced in  $\text{Ru}^{96} + \text{N}^{14}$  bombardments.

We obtained mass assignments for these activities from excitation function measurements using the results of Black on heavy-ion reactions in this mass region to interpret our data<sup>5</sup>. As shown in Fig. 2, the peak cross section for the production of the 3.28 MeV alpha activity falls at a bombarding energy of 104 MeV. This energy compares most favourably with that expected for the reaction  $\text{Ru}^{96}(\text{O}^{16}, 5n)\text{Te}^{107}$ . The excitation function for the 3.08 MeV group peaks at a bombarding energy of 87 MeV which is close to the value expected for the reaction  $\text{Ru}^{96}(\text{O}^{16}, 4n)\text{Te}^{108}$ . The results are summarized in Table 1.

Table 1

Summary of Results

<u>Nuclide</u>	<u><math>E_{\alpha}</math> (MeV)</u>	<u><math>t_{1/2}</math> (sec)</u>
$\text{Te}^{107}$	$3.28 \pm 0.03$	$2.2 \pm 0.2$
$\text{Te}^{108}$	$3.08 \pm 0.03$	$5.3 \pm 0.4$

These nuclides represent the first opportunity to study alpha decay from nuclei where the valence neutrons and protons are in the same single particle level, in this case, the  $1g_{7/2}$  level. This may give rise to a kind of "super-allowed" alpha decay resulting in large reduced alpha widths. At present, we cannot give any estimates of the alpha reduced widths for  $\text{Te}^{107}$  and  $\text{Te}^{108}$  because the alpha branching ratios are not known.

The most alpha labile tellurium isotope is expected to be  $\text{Te}^{104}$  since it can alpha decay to the double closed shell nuclide  $\text{Sn}^{100}$ . Our first experiments were directed toward producing this nuclide by the reaction  $\text{Ru}^{96}(\text{O}^{16}, 8n)\text{Te}^{104}$ . Apparently, however, the peak cross section for this reaction as well as for the  $\text{O}^{16}, 7n$  and  $\text{O}^{16}, 6n$  reactions is extremely small due to the high neutron binding energies in this region.

We expect that it will also be possible to observe alpha decay from

the very neutron-deficient isotopes of iodine and xenon.

We would like to thank Professor I. Perlman, Professor J. O. Rasmussen, Dr. E. K. Hyde, and A. Ghiorso for their interest in this work. One of us (R.D.M.) would like to express his appreciation to the Chemistry Division of the Lawrence Radiation Laboratory for their kind hospitality.


\* Research performed under the auspices of the U.S. Atomic Energy Commission.

1. J. O. Rasmussen, S. G. Thompson, and A. Ghiorso, Phys. Rev. 89, 33 (1953).
2. M. Karras, G. Andersson, and M. J. Nurmi, Arkiv Fysik 23, 57 (1962).
3. R. D. Macfarlane and R. D. Griffioen, Nucl. Instr. Methods 24, 461 (1963).
4. R. P. Black, "Heavy Ion Reactions in the Medium Mass Region" (thesis) Massachusetts Institute of Technology, (1964).

#### Figure Captions


Figure 1. Alpha particle spectrum of the products of the reaction  $Ru^{96} + O^{16}$   
MUB-4579 at a bombarding energy of 95 MeV (lab).

Figure 2. Excitation functions for the production of the new Te alpha emitters.  
MUB-4580



MUB-4579

Fig. 1



MUB-4580

Fig. 2



This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

