

Lawrence Berkeley National Laboratory

Recent Work

Title

TUNNELING CORRECTIONS TO UNIMOLECULAR RATE CONSTANTS WITH APPLICATION TO FORMALDEHYDE

Permalink

<https://escholarship.org/uc/item/4nj3c5s9>

Author

Miller, W.H.

Publication Date

1979

Submitted to American Chemical Society

LBL-8649
Preprint

e-2

TUNNELING CORRECTIONS TO UNIMOLECULAR RATE CONSTANTS
WITH APPLICATION TO FORMALDEHYDE

RECEIVED
LAWRENCE
BERKELEY LABORATORY

William H. Miller

MAR 23 1979


January 1979

LIBRARY AND
DOCUMENTS SECTION

Prepared for the U. S. Department of Energy
under Contract W-7405-ENG-48

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 6782*



LBL-8649e-2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Tunneling Corrections to Unimolecular Rate Constants,
with Application to Formaldehyde

By

William H. Miller[‡]

Department of Chemistry, and Materials and Molecular Research Division,
of the Lawrence Berkeley Laboratory, University of California,
Berkeley, CA 94720

Abstract

Tunneling corrections to the rate constant for unimolecular reactions in an isolated molecule are treated within the standard transition state (i.e., RRKM) theory of such processes. The microcanonical distribution relevant to the unimolecular case causes tunneling effects to enter in a somewhat more complicated fashion than in the analogous transition state theory for thermally averaged bimolecular rate constants; e.g., even within the separable approximation they do not enter as simply a multiplicative correction factor. Application of the theoretical expressions to some unimolecular processes ($\text{H}_2\text{CO} \rightarrow \text{H}_2 + \text{CO}$, trans $\text{HCOH} \rightarrow \text{H}_2\text{CO}$) of interest in the collisionless photochemistry of formaldehyde indicate that tunneling effects are quite significant for rates of 10^9 sec^{-1} or slower. Isotope effects are also considered and seen to be quite interesting.

Much has been written over the years about tunneling corrections to transition state theory¹ for thermal rate constants of bimolecular reactions, but there has been little discussion about the effect of tunneling in the analogous transition state (i.e., RRKM) theory² of unimolecular reactions. The purpose of this paper is to consider such effects and to illustrate them by application to some processes of current interest in the photochemistry of formaldehyde.

Brief Summary of the Standard Transition State
(i.e. RRKM) Theory for Unimolecular Processes

To simplify the presentation, rotational degrees of freedom will be ignored here; the Appendix shows how the formulae are modified to take proper account of total angular momentum conservation and other aspects of the rotational degrees of freedom. With this proviso, the standard expression² for the unimolecular rate constant (units sec^{-1}) of an isolated molecule with total energy E is

$$k(E) = \frac{N(E)}{2\pi h N_0(E)} \quad , \quad (1)$$

where $N(E)$ and $N_0(E)$ are the integral densities of states for the transition state and for the reactant molecule, respectively. Specifically,

$$N(E) = \sum_{\tilde{n}} h(E - \epsilon_{\tilde{n}}^{\ddagger}) \quad (2a)$$

$$N_0(E) = \sum_{\tilde{n}} h(E - \epsilon_{\tilde{n}}) \quad , \quad (2b)$$

where $h(x)$ is the usual step-function,

$$h(x) = \begin{cases} 0, & x < 0 \\ 1, & x > 0 \end{cases}$$

and ϵ_n^\ddagger and ϵ_n are the vibrational energy levels of the transition state and the reactant molecule. In practice the vibrational energy levels are almost always assumed to be given by a separable harmonic oscillator approximation, so that

$$\epsilon_n \approx \sum_{i=1}^s \hbar \omega_i (n_i + \frac{1}{2}) \quad (3a)$$

$$\epsilon_n^\ddagger \approx V_0 + \sum_{i=1}^{s-1} \hbar \omega_i^\ddagger (n_i + \frac{1}{2}) \quad (3b)$$

where s is the number of vibrational degrees of freedom of the stable molecule, $\{\omega_i\}$ and $\{\omega_i^\ddagger\}$ are the normal mode frequencies of the reactant molecule and transition state, and V_0 is the "bare" barrier height, i.e., the energy of the saddle point of the potential energy surface (i.e., the transition state) relative to the minimum of the potential energy surface which corresponds to the reactant molecule. The total energy E is also defined relative to the minimum of the potential energy surface.

Because of the large sums involved in computing the densities of states, it is customary in practice to approximate them by closed form expressions. The simplest such approximation is the classical expression² which gives (if the Marcus-Rice "semiclassical" modification² is included)

$$N_0(E) = \frac{E^s}{s! \prod_{i=1}^s (\hbar\omega_i)} \quad (4a)$$

$$N(E) = \frac{(E-V_0)^{s-1}}{(s-1)! \prod_{i=1}^{s-1} (\hbar\omega_i^\ddagger)} \quad (4b)$$

With these approximations Eq. (1) gives the simple classical rate expression²

$$k(E) = A \left(\frac{E-V_0}{E} \right)^{s-1} \quad (5a)$$

where A is a frequency factor (units sec⁻¹)

$$A = \left(\prod_{i=1}^s \omega_i \right) / 2\pi \left(\prod_{i=1}^{s-1} \omega_i^\ddagger \right) \quad (5b)$$

Modification to Include Tunneling

The only simple way to include the effect of tunneling along the reaction coordinate in transition state theory is to assume that this degree of freedom--i.e., motion along the reaction coordinate--is separable from the other degrees of freedom; this approximation is also consistent with the use of Eq. (3) for the energy levels. Within the separable approximation tunneling is accounted for by replacing N(E) in Eq. (1) by $N_{QM}(E)$,¹

$$N_{QM}(E) = \sum_{\tilde{n}} P(E - \epsilon_{\tilde{n}}^\ddagger) \quad (6)$$

where $P(E_1)$ is the one-dimensional tunneling probability as a function of the energy E_1 in the reaction coordinate; in the classical limit of no tunneling $P(E_1) \rightarrow h(E_1)$, and $N_{QM}(E) \rightarrow N(E)$. The expression for the unimolecular rate constant which incorporates tunneling is thus

$$k_{QM}(E) = \frac{\sum_n P(E - \epsilon_n^\ddagger)}{2\pi\hbar N_0'(E)} \quad (7)$$

If the barrier along the reaction coordinate is approximated as an inverted parabola, then the tunneling probability is given by

$$P(E_1) = e^{-\epsilon} / (1 + e^{-\epsilon})$$

with

$$\epsilon = \frac{2\pi E_1}{\hbar\omega_b}$$

where ω_b is the magnitude of the imaginary frequency related to the barrier. The generalized Eckart potential^{1a} in general provides a more accurate representation of the barrier, and in this case the tunneling probability is given by

$$P(E_1) = \frac{\sinh(a) \sinh(b)}{\sinh^2\left(\frac{a+b}{2}\right) + \cosh^2(c)} \quad (8)$$

where

$$a = \frac{4\pi}{\hbar\omega_b} \sqrt{E_1 + V_0} (V_0^{-1/2} + V_1^{-1/2})^{-1}$$

$$b = \frac{4\pi}{\hbar\omega_b} \sqrt{E_1 + V_1} (V_0^{-1/2} + V_1^{-1/2})^{-1}$$

$$c = 2\pi \sqrt{\frac{V_0 V_1}{(\hbar\omega_b)^2} - \frac{1}{16}} \quad ;$$

V_0 is, as before, the barrier height relative to reactants, and V_1 is the barrier height relative to products; $V_1 - V_0$ is the exoergicity of the reaction (neglecting zero point energies).

Eq. (6) can be written in another form by the following manipulations:

$$\begin{aligned} N_{QM}(E) &\equiv \sum_{\tilde{n}} P(E - \epsilon_{\tilde{n}}^{\ddagger}) = \int dE_1 P(E_1) \sum_{\tilde{n}} \delta(E - E_1 - \epsilon_{\tilde{n}}^{\ddagger}) \\ &= \int_{-V_0}^{E-V_0} dE_1 P(E_1) N'(E - E_1) \\ &= \int_{-V_0}^{E-V_0} dE_1 P'(E_1) N(E - E_1) \quad , \end{aligned} \quad (9)$$

where $N(E - E_1)$ is the density of states defined by Eq. (2a); i.e., $N_{QM}(E)$ is given by a convolution of the classical approximation to it, namely $N(E)$, and the tunneling probability. [For comparison, it is interesting to note that the tunneling correction factor Γ for a thermally averaged rate constant is given in terms of the tunneling probability by

$$\begin{aligned} \Gamma &= \int_{-V_0}^{\infty} dE_1 P(E_1) \frac{e^{-E_1/kT}}{kT} \\ &= \int_{-V_0}^{\infty} dE_1 P'(E_1) e^{-E_1/kT} \quad .] \end{aligned}$$

One is tempted to use Eq. (9) with the approximation to $N(E-E_1)$ given by Eq. (4b), thereby obtaining the following simple expression for the tunneling rate constant:

$$k_{QM}(E) = A \int_{-V_0}^{E-V_0} dE_1 P'(E_1) \left(\frac{E-V_0-E_1}{E} \right)^{s-1}, \quad (10)$$

where A is the same frequency factor as above. It should be emphasized, however, that this is not a good thing to do. Although approximating $N_0(E)$ by Eq. (4a) is valid (at least for the applications below³), in the threshold region where tunneling is important there are so few terms that contribute to the sum in Eq. (6) that Eq. (4b) is a poor approximation to $N(E-E_1)$ in the integrand of Eq. (9). Eq. (10) thus gives values much too large, and one must retain the discrete sum; since only a few terms do contribute to the sum, this causes no computational difficulties. The final expression we use (except for the modification due to rotation discussed in the Appendix) is thus

$$k_{QM}(E) = \frac{(s-1)! \prod_{i=1}^s \hbar \omega_i}{2\pi \hbar E^{s-1}} \sum_{\tilde{n}} P[E-V_0 - \hbar \tilde{\omega}^\ddagger \cdot (\tilde{n} + \frac{1}{2})], \quad (11)$$

where

$$\tilde{n} = n_1, n_2, \dots, n_{s-1}$$

$$\tilde{\omega}^\ddagger \cdot (\tilde{n} + \frac{1}{2}) = \sum_{i=1}^{s-1} \omega_i^\ddagger (n_i + \frac{1}{2}) \quad ;$$

for the applications below the tunneling probability for the Eckart barrier [Eq. (8)] was used.

Applications to Unimolecular Processes in the Ground

Electronic State (S_0) of Formaldehyde

The motivation for this work has been the current interest in the photochemistry of formaldehyde,⁴ for which there are several potentially relevant unimolecular processes that can take place on the potential energy surface of the ground electronic state (S_0). (The excited electronic state S_1 which is initially produced by laser excitation is assumed to undergo a radiationless transition back to S_0 .) Figure 1 shows a schematic of the potential energy surface⁵ for S_0 , and we consider the unimolecular decomposition of formaldehyde to molecular products



and the isomerization of the metastable species trans-hydroxymethylene-- which has been suggested⁶ as the species first formed from S_1 --to formaldehyde,



Goddard and Schaefer⁵ have recently carried out extensive self-consistent field and configuration interaction calculations on formaldehyde and have determined all the parameters needed to evaluate the rate expressions given above. The energies of the two stable species (i.e., H_2CO and HCOH) and of the two transition states are shown in Figure 1,

and Table I gives the six vibrational frequencies and three rotation constants for all four species. Unless stated otherwise, all rates constants given below were calculated from Eq. (A.8), which for $J = 0$ is identical to Eq. (11), with the tunneling probability of Eq. (8).

Figure 2 shows the unimolecular rate constant for reaction (R1) as a function of total energy E (relative to the bottom of the potential energy surface of H_2CO), and for comparison the "semiclassically" modified classical rate constant of Eq. (5) is also shown (broken line). The arrow on the energy scale indicates the value \tilde{V}_0 , the "bare" barrier height plus the zero point energy of the transition state,

$$\tilde{V}_0 = V_0 + \sum_{i=1}^{s-1} \frac{1}{2} \hbar \omega_i^\ddagger, \quad (12)$$

which would be the threshold of the reaction in a completely classical approximation; one sees that tunneling allows a significant rate ($> 10^9 \text{ sec}^{-1}$) at this threshold energy. The rate has fallen only to 10^6 sec^{-1} at an energy $\sim 8 \text{ kcal/mole}$ below \tilde{V}_0 . The exponential energy dependence of $k(E)$ (i.e., the linearity of the semi-log plot) for $E < \tilde{V}_0$ also indicates that the process is dominated by tunneling in this region.

Figure 2 also shows that the "semiclassically" modified classical rate expression [Eq. (5)] provides a crude description of the tunneling region; it is at least better than the purely classical result which goes to zero at \tilde{V}_0 . This partial success is the result of a cancellation of errors: the "semiclassical" approximation to $N(E)$ [Eq. (4b)] is known² to become much too large as E decreases toward threshold, which tends to

make the rate too large, but neglect of tunneling tends to make it too small, thus the partially off-setting errors. If, on the other hand, one includes tunneling via Eq. (9) and uses the "semiclassical" approximation to $N(E-E_1)$, then the rate obtained [Eq. (10)] is much (over an order of magnitude) too large in the tunneling region.

Figure 3 shows similar results for the rate constant of reaction (R2). The arrow shows the classical threshold for the reaction, \tilde{V}_0 , and one again sees that tunneling is substantial. In this case the "semiclassically" modified classical rate is rather poor in the threshold region, probably a more typical situation than the better agreement in Figure 2.

To assess the effect of rotation, calculations were also carried out for total angular momentum $J > 0$. The rate constant in general decreases with increasing J , but the effect is not large for the present examples: for $J = 10$, as large a value as is probably of interest, the rate constant for both reactions is decreased by a factor of ~ 2.5 at $E = 90$ kcal/mole, and the factor decreases approximately uniformly to ~ 1.2 at $E = 120$ kcal/mole.

Isotope Effects

Tunneling is significant for these reactions because they primarily involve the motion of hydrogen atoms, as evidenced by the large imaginary barrier frequencies $i\omega_b$ for the transition states in Table I. One thus expects large isotope effects in the tunneling region if H-atoms are replaced by D atoms. The relevant frequencies (and rotation constants) for the deuterated species have also been determined by Goddard and Schaefer⁵ and are given in Table I.

Figure 4 shows the isotope effect, i.e., the ratio of the hydrogen to the deuterium rate constant for the two reactions. It is easy to see that the classical rate expression [Eq. (5)] gives an energy-independent isotope ratio, and one sees in Figure 4 that this limit is approached for energies above the classical thresholds. In the threshold region and below, however, the isotope ratio is strongly dependent on energy: for energies significantly below the classical threshold there is an exponential energy dependence (i.e., the semi-log plot is linear), but near the classical threshold itself the energy dependence is quite complicated, showing a pronounced minimum, in the vicinity of which the deuterium versions of the reactions are actually faster than the hydrogen versions.

Although comparing different isotopes at the same total energy (as in Figure 4) is the most meaningful comparison from a theoretical point of view, the experimental situation often dictates otherwise. Thus for formaldehyde the excitation energy from the ground vibrational state of S_0 to the ground vibrational state of S_1 is⁴ 80.6 kcal/mole for H_2CO and 80.9 kcal/mole for D_2CO , and since the zero point energies of H_2CO and D_2CO are 16.1 kcal/mole and 12.8 kcal/mole, respectively, the total energies resulting from these vibrationless excitations are 96.7 kcal/mole for H_2CO and 93.7 kcal/mole for D_2CO . At these energies the present calculations give

$$k_1^H = 5.8 \times 10^6 \text{ sec}^{-1} \quad , \quad k_1^D = 1.4 \times 10^5 \text{ sec}^{-1} \quad (13a)$$

$$k_2^H = 1.9 \times 10^7 \text{ sec}^{-1} \quad , \quad k_2^D = 5.7 \times 10^5 \text{ sec}^{-1} \quad , \quad (13b)$$

where k_1 and k_2 refer to reactions (R1) and (R2), respectively. (The values for k_1^H and k_1^D in Eq. (13a) include an extra factor of 2 due to symmetry, which has heretofore been omitted; this is because H_2CO and D_2CO have a two-fold rotation axis and thus a symmetry number of 2.) These rates are for total angular momentum $J = 0$; for $J = 10$ they are all about a factor of 2 smaller.

Similar calculations have been carried out for the mixed isotope HDCO and the rates are, perhaps not unexpectedly, intermediate between those for H_2CO and D_2CO . For the vibrationless $S_0 \rightarrow S_1$ excitation, for example, the total energy is 95.3 kcal/mole and the rate of reaction (R1) is

$$k_1^{HD} = 9.5 \times 10^5 \text{ sec}^{-1} \quad (14)$$

Concluding Remarks

The main purpose of this paper has been to show how tunneling can be incorporated in the transition state (i.e., RRKM) theory for unimolecular reactions in a manner analogous to the way it is included in transition state theory for thermal bimolecular reactions. Because the unimolecular case corresponds to a fixed energy rather than a fixed temperature, the effect of tunneling is somewhat more complicated, i.e., it does not enter as simply a multiplicative correction factor but rather in a more convoluted manner.

With regard to the applications to formaldehyde, one must be somewhat cautious regarding the specific values obtained for the rate constants because it is known that when tunneling effects are substantial, the separable approximation for tunneling can be poor.^{1c} Nevertheless, the

results obtained for the rates do indicate that on the time scale of interest in the collisionless photochemistry of formaldehyde, i.e., 10^{-5} - 10^{-6} secs, tunneling is likely to play a significant role.

Acknowledgments

I would like to thank Professors C. B. Moore and H. F. Schaefer for a number of stimulating discussions regarding the photochemistry of formaldehyde and their suggestion to consider the role of tunneling in unimolecular processes. Thanks are also due Dr. J. D. Goddard and Professor Schaefer for providing the results of their extensive calculations on formaldehyde prior to publication. This work has been supported by the Division of Chemical Sciences, Office of Basic Energy Sciences, U. S. Department of Energy under contract No. W-7405-Eng-48 and the Miller Institute of Sciences.

Appendix: Effect of Rotational Degrees of Freedom

There are three degrees of freedom associated with the overall rotation of a non-linear molecule, the quantum numbers for which we designate J , M_J , K . J , the total angular momentum quantum number, and M_J , its projection onto a space-fixed axis, are always conserved, while K is in general not conserved. (For a rigid symmetric top, K is also conserved.) One thus needs to define the unimolecular rate constant $k(E, J)$ which corresponds to a fixed value of total angular momentum J as well as total energy E ; because of the isotropy of space, the rate is independent of M_J .

We assume that K is a statistical degree of freedom, i.e., that it interchanges energy statistically with all the vibrational degrees of freedom. The unimolecular rate constant is then given by

$$k(E, J) = \frac{N(E, J)}{2\pi\hbar \frac{\partial N_0(E, J)}{\partial E}}, \quad (\text{A.1})$$

where

$$N(E, J) = \sum_{K=-J}^J \sum_{\tilde{n}} h(E - \epsilon_{\tilde{n}, J, K}^\ddagger) \quad (\text{A.2a})$$

$$N_0(E, J) = \sum_{K=-J}^J \sum_{\tilde{n}} h(E - \epsilon_{\tilde{n}, J, K}) \quad (\text{A.2b})$$

$\{\epsilon_{\tilde{n}, J, K}^\ddagger\}$ and $\{\epsilon_{\tilde{n}, J, K}\}$ being the rotational-vibrational energy levels of the transition state and of the reactant molecule, respectively. In practice these energy levels are obtained by assuming a rigid rotor-harmonic

oscillator approximation,

$$\epsilon_{\tilde{n},J,K} = W_{J,K} + \sum_{i=1}^s \hbar\omega_i \left(n_i + \frac{1}{2}\right) \quad (\text{A.3a})$$

$$\epsilon_{\tilde{n},J,K}^{\ddagger} = V_0 + W_{J,K}^{\ddagger} + \sum_{i=1}^{s-1} \hbar\omega_i^{\ddagger} \left(n_i + \frac{1}{2}\right) \quad , \quad (\text{A.3b})$$

where $W_{J,K}^{\ddagger}$ and $W_{J,K}$ are the rotational energies of the transition state and of the reactant molecule. Since the energy levels of most asymmetric rotors are reasonably well approximated by assuming an "almost symmetric top", we invoke that approximation here, so that⁷

$$W_{J,K} = \frac{1}{2} (A+B) [J(J+1) - K^2] + CK^2 \quad (\text{A.4a})$$

$$W_{J,K}^{\ddagger} = \frac{1}{2} (A^{\ddagger} + B^{\ddagger}) [J(J+1) - K^2] + C^{\ddagger}K \quad , \quad (\text{A.4b})$$

where (A,B,C) and $(A^{\ddagger}, B^{\ddagger}, C^{\ddagger})$ are the three rotation constants of the molecule and of the transition state. (A and B are chosen as the two most nearly equal rotation constants of the three $A, B,$ and $C,$ and A^{\ddagger} and B^{\ddagger} similarly.)

The sums over vibrational quantum number \tilde{n} may be approximated as before,² but because we consider cases of small $J,$ the discrete sum over K is retained. Analogous to Eq. (4) the classical approximation to the sum over \tilde{n} in Eq. (4.2) thus gives

$$N(E,J) = \sum_{K=-J}^J \frac{(E - V_0 - W_{J,K}^{\ddagger})^{s-1}}{(s-1)! \prod_{i=1}^{s-1} (\hbar\omega_i^{\ddagger})} \quad (\text{A.5a})$$

$$N_0(E, J) = \sum_{K=-J}^J \frac{(E - W_{J,K})^s}{s! \prod_{i=1}^s \hbar \omega_i} , \quad (\text{A.5b})$$

so that the classical rate expression which replaces Eq. (5a) is

$$k(E, J) = A \frac{\sum_{K=-J}^J (E - V_0 - W_{J,K}^\ddagger)^{s-1}}{\sum_{K=-J}^J (E - W_{J,K})^{s-1}} , \quad (\text{A.6})$$

A being the frequency factor of Eq. (5b). Note that for $J = 0$, Eq. (A.6) reduces to Eq. (5a).

The effect of tunneling along the reaction coordinate is included in the same manner as before, by replacing $N(E, J)$ by $N_{QM}(E, J)$,

$$N_{QM}(E, J) = \sum_{K=-J}^J \sum_{\tilde{n}} P(E - \epsilon_{\tilde{n}, J, K}^\ddagger) , \quad (\text{A.7})$$

where again $P(E_1)$ is the one-dimensional tunneling probability. The expression for the tunneling rate constant which generalizes Eq. (11) by including rotation is thus

$$k_{QM}(E, J) = \frac{(s-1)! \prod_{i=1}^s \hbar \omega_i \sum_{K=-J}^J \sum_{\tilde{n}} P[E - V_0 - W_{J,K}^\ddagger - \hbar \omega^\ddagger \cdot (\tilde{n} + \frac{1}{2})]}{2\pi\hbar \sum_{K=-J}^J (E - W_{J,K})^{s-1}} , \quad (\text{A.8})$$

with the rotational energies $W_{J,K}$ and $W_{J,K}^\ddagger$ given by Eq. (A.4). One notes that for the case $J = 0$ Eq. (A.8) reduces to the result in the text [Eq. (11)] that ignores rotation altogether.

References

- † Camille and Henry Dreyfus Teacher-Scholar.
1. See, for example, (a) H. S. Johnston, Gas Phase Reaction Rate Theory, Ronald Press, N.Y., 1966, pp. 37-47; (b) P. Pechukas, in Dynamics of Molecular Collisions, Part B, Vol. 2 of Modern Theoretical Chemistry, ed. W. H. Miller, Plenum, N.Y., 1976, pp. 269-322; (c) W. H. Miller, Accts. Chem. Res. 9, 306 (1976).
 2. (a) P. J. Robinson and K. A. Holbrook, Unimolecular Reactions, Wiley, N.Y., 1972; (b) Wendell Forst, Theory of Unimolecular Reactions, Academic Press, N.Y., 1973.
 3. $N_0'(E)$ was actually computed using the more accurate Whitton-Rabinovitch formula (cf. ref. 2a, pp. 131-137), but this makes only a few percent difference in the present applications.
 4. See (a) E. S. Yeung and C. B. Moore, J. Chem. Phys. 58, 3988 (1973); (b) P. L. Houston and C. B. Moore, J. Chem. Phys. 65, 757 (1976); and many other references cited therein.
 5. J. D. Goddard and H. F. Schaefer, "The Photodissociation of Formaldehyde: Potential Energy Surface Features", preprint.
 6. M. J. H. Kemper, J. M. F. Dijk, and H. M. Buck, J. Amer. Chem. Soc. 100, 7841 (1978).
 7. C. H. Townes and A. L. Schalow, Microwave Spectroscopy, McGraw-Hill, N.Y., 1955, pp. 84-86.

Table I. Vibrational Frequencies and Rotation Constants^a

Hydrogen Species			
<u>H₂CO</u>	<u>tr-HCOH</u>	<u>TS-M</u>	<u>TS-R</u>
2843	3634	2760	3675
2766	2684	1654	2803
1746	1595	1137	2339
1501	1264	941	1568
1247	1101	697	1221
1164	1093	2288 i	2299 i


1.13	1.04	1.05	1.09
1.30	1.16	1.18	1.25
9.41	10.52	9.06	8.86
Deuterated Species			
<u>D₂CO</u>	<u>tr-DCOD</u>	<u>TS-M</u>	<u>TS-R</u>
2160	2525	2186	2759
2056	1979	1503	2134
1700	1430	820	1735
1106	933	724	1408
990	921	505	974
938	799	1900 i	1675 i

0.88	0.84	0.76	0.89
1.07	0.97	0.89	1.10
4.70	6.28	5.45	4.75

^aThe six values above the dotted lines are the vibrational frequencies, and the three values below it are the rotation constants. Units for all are cm^{-1} . TS-M and TS-R are the molecular and rearrangement transition states indicated in Figure 1.


Figure Captions

1. Schematic of the potential energy surface for the ground electronic state (S_0) of formaldehyde. Units of energy are kcal/mole, and the values shown are from the work in reference 5.
2. Unimolecular rate constant for the reaction indicated, as a function of total energy, for total angular momentum $J = 0$. The solid curve includes tunneling effects and is computed from Eq. (11). The broken curve is from the "semiclassically" modified classical expression in Eq. (5). The arrow indicates the classical threshold for the reaction, as defined by Eq. (12).
3. Same as Figure 2.
4. Isotope effects. Plotted is the log of the ratio of hydrogen to deuterium rate constants for the reactions indicated, as a function of total energy.




XBL 791-8083

Figure 1




XBL 791-7797

Figure 2



XBL 791-7798

Figure 3



XBL 791-7799

Figure 4

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720