# **UCSF**

# **UC San Francisco Previously Published Works**

# **Title**

First Complete Genome Sequence of Corynebacterium riegelii

## **Permalink**

https://escholarship.org/uc/item/4p01r2ds

# **Journal**

Microbiology Resource Announcements, 5(13)

## **ISSN**

2576-098X

# **Authors**

Greninger, Alexander L Streithorst, Jessica Chiu, Charles Y et al.

## **Publication Date**

2017-03-30

# DOI

10.1128/genomea.00084-17

Peer reviewed







# First Complete Genome Sequence of Corynebacterium riegelii

**ABSTRACT** Here, we report the first complete genome sequence of *Corynebacterium riegelii* strain PUDD\_83A45, isolated from the urine of a patient with urinary tract infection. The genome measured 2.56 Mb and contained no plasmid.

Corynebacterium riegelii is a newly described species of corynebacteria originally isolated in 1997 from pure cultures of urinary tract infections in females and is named after corynebacteriologist Philippe Riegel (1, 2). Since its discovery, it has been uniformly associated with urinary tract infections and twice from urosepsis (3, 4). Biochemically, it is distinguished by its strong urease positivity and ability to ferment maltose but not glucose (1). We sequenced the first complete genome of *C. riegelii* strain PUDD\_83A45, isolated from a 2003 urine sample from a 9-year-old male with congenital hypospadias and hydronephrosis who presented with symptoms of urinary tract infection. The patient had multiple surgical repairs as a child to correct anatomical and obstructive urinary tract defects and had multiple urinary tract infections prior to this presentation. The isolate was identified biochemically and determined to be susceptible to penicillin and vancomycin by disk diffusion testing.

DNA from *C. riegelii* strain PUDD\_83A45 was extracted using the Qiagen EZ1 kit. Nextera XT paired-end and Nextera mate-pair libraries were sequenced  $2 \times 300$  bp and  $2 \times 80$  bp, respectively, on an Illumina MiSeq platform. DNA was also sequenced using approximately half of a flow cell of an Oxford Nanopore Minion sequencer. Illumina sequences were adapter and quality (Q30) trimmed using Cutadapt or NxTrim, *de novo* assembled using SPAdes version 3.5, metagenomically screened for contaminating sequences with SURPI, and annotated via Prokka version 1.1 (5–10). A total of 2,265,156 paired-end reads and 4,092,796 mate-pair reads were recovered after trimming, while 1,381 "pass" and 26,418 "fail" reads were recovered from the Nanopore sequencer. *De novo* assembly combined with manual integration of Nanopore data via LASTZ alignments yielded a single scaffold of 2,563,736 bp, an average coverage of 225 $\times$ , and a total of 2,395 coding sequences. Two gaps of <1.5 kb remained in the genome due to a transposase that was also present at 96% nucleotide identity in *Corynebacterium imitans* strain DSM 44264, as well as a highly repetitive sequence in a Rib surface antigen family protein.

BLASTn analysis of the complete 16S sequence from strain PUDD\_83A45 showed 99.8% identity to *C. riegelii* strain CTF08-1967 (EU848548) and *C. riegelii* isolate 99-0185 (AF537602). BLASTn analysis of the complete *rpoB* gene demonstrated 98.8% identity to *C. riegelii* strain CIP 103310 (AY492278). The genome contained a 5.1-kb locus that included *ureA-G* genes that aligned 71 to 80% by nucleotide to known urealytic corynebacteria such as *C. pseudotuberculosis*, *C. ulcerans*, *C. urealyticum*, and *C. ureicele-rivorans*. Annotated maltose-related genes included a *malF* maltose permease that aligned with 76% nucleotide identity to *C. ureicelerivorans* strain IMMIB RIV-2301 (CP009215) and a *malE* maltose-binding periplasmic protein that aligned with 76% nucleotide identity to *C. glutamicum* strain B253 (CP010451). No high-confidence antibiotic resistance genes were identified by Comprehensive Antibiotic Resistance

**Received** 25 January 2017 **Accepted** 27 January 2017 **Published** 30 March 2017

Citation Greninger AL, Streithorst J, Chiu CY, Miller S. 2017. First complete genome sequence of *Corynebacterium riegelii*. Genome Announc 5:e00084-17. https://doi.org/10.1128/ genomeA.00084-17.

**Copyright** © 2017 Greninger et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

Address correspondence to Steve Miller, steve.miller@ucsf.edu.

Greninger et al. genameAnnouncements™

Database analysis, although distant matches included two multidrug MFS transporters, one closest to *Actinomyces neuii* (63% amino acid identity to WP\_024330734) and one closest to *Propionibacterium* sp. KPL1838 (73% amino acid identity to ERS38161) (11).

**Accession number(s).** This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession number CP012342.

#### **ACKNOWLEDGMENT**

This work was supported by the University of California, San Francisco, Department of Laboratory Medicine.

#### **REFERENCES**

- Funke G, Lawson PA, Collins MD. 1998. Corynebacterium riegelii sp. nov., an unusual species isolated from female patients with urinary tract infections. J Clin Microbiol 36:624–627.
- Riegel P, Liégeois P, Chenard MP, Mathelin C, Monteil H. 2004. Isolations of Corynebacterium kroppenstedtii from a breast abscess. Int J Med Microbiol 294:413–416. https://doi.org/10.1016/j.ijmm.2004.07.013.
- Aygun G, Midilli K, Cilingir H, Yilmaz M, Kutukcu A, Eker E. 2013. A fatal case of urosepsis due to Corynebacterium riegelii. Braz J Microbiol 44: 475–476. https://doi.org/10.1590/S1517-83822013000200022.
- Matusnami M, Sogi M, Kitazono H, Hosokawa N, Otsuka Y, Ohkusu K. 2012. Urosepsis caused by *Globicatella sanguinis* and *Corynebacterium riegelii* in an adult: case report and literature review. J Infect Chemother 18:552–554. https://doi.org/10.1007/s10156-011-0335-x.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics 30:2068–2069. https://doi.org/10.1093/bioinformatics/btu153.
- Naccache SN, Federman S, Veeraraghavan N, Zaharia M, Lee D, Samayoa E, Bouquet J, Greninger AL, Luk KC, Enge B, Wadford DA, Messenger SL, Genrich GL, Pellegrino K, Grard G, Leroy E, Schneider BS, Fair JN, Martínez MA, Isa P, Crump JA, DeRisi JL, Sittler T, Hackett J, Miller S, Chiu CY. 2014. A cloud-compatible bioinformatics pipeline for ultrarapid pathogen identification from next-generation sequencing of clinical samples. Genome Res 24:1180–1192. https://doi.org/10.1101/gr.171934.113.
- 7. Martin M. 2011. Cutadapt removes adapter sequences from high-

- throughput sequencing reads. EMBnetJ 17:10–12. https://doi.org/10.14806/ej.17.1.200.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol 19:455–477. https://doi.org/10.1089/cmb.2012.0021.
- Greninger AL, Langelier C, Cunningham G, Keh C, Melgar M, Chiu CY, Miller S. 2015. Two rapidly growing mycobacterial species isolated from a brain abscess: first whole genome sequence of *Mycobacterium immu-nogenum* and *Mycobacterium Ilatzerense*. J Clin Microbiol 53:2374–2377. https://doi.org/10.1128/JCM.00402-15.
- Greninger AL, Chorny I, Knowles S, Ng VL, Chaturvedi V. 2015. Draft genome sequences of four NDM-1-producing Klebsiella pneumoniae strains from a health care facility in northern California. Genome Announc 3(3):e00421-15. https://doi.org/10.1128/genomeA.00421-15.
- McArthur AG, Waglechner N, Nizam F, Yan A, Azad MA, Baylay AJ, Bhullar K, Canova MJ, De Pascale GD, Ejim L, Kalan L, King AM, Koteva K, Morar M, Mulvey MR, O'Brien JS, Pawlowski AC, Piddock LJV, Spanogiannopoulos P, Sutherland AD, Tang I, Taylor PL, Thaker M, Wang W, Yan M, Yu T, Wright GD. 2013. The Comprehensive Antibiotic Resistance Database. Antimicrob Agents Chemother 57:3348–3357. https://doi.org/10.1128/AAC.00419-13.

Volume 5 Issue 13 e00084-17 genomea.asm.org **2**