

Lawrence Berkeley National Laboratory

Recent Work

Title

MEASUREMENT OF THE Z -ASYMMETRY PARAMETER

Permalink

<https://escholarship.org/uc/item/4pj4f71b>

Authors

Tripp, Robert D.
Watson, Mason B.
Ferro-Luzzi, M.

Publication Date

1962-05-15

University of California

**Ernest O. Lawrence
Radiation Laboratory**

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UCRL-10238

UNIVERSITY OF CALIFORNIA
Lawrence Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

MEASUREMENT OF THE Σ^- ASYMMETRY PARAMETER

Robert D. Tripp, Mason B. Watson, and Maccimiliano Ferro-Luzzi

May 15, 1962

MEASUREMENT OF THE Σ^- ASYMMETRY PARAMETER^oRobert D. Tripp, Mason B. Watson, and Massimiliano Ferro-Luzzi[†]Lawrence Radiation Laboratory
University of California
Berkeley, California

May 15, 1962

In a previous letter we reported a hydrogen bubble-chamber experiment in which the existence of an excited 1520-Mev hyperon was established.¹ Moreover, a study of the angular distributions and polarizations of Σ hyperons from the reaction $K^-p \rightarrow \Sigma v$ in the vicinity of this resonance showed the $K\Sigma$ parity to be negative.² A further study of this reaction permits us to determine the asymmetry parameters a in Σ^\pm decay by means of their decay asymmetries. The a for $\Sigma_+^+ \rightarrow n\pi^+$ (called a_+) and $\Sigma_0^+ \rightarrow p\pi^0$ (called a_0) have already been well established, and our measurements are in good agreement with the published values, although statistically less accurate. The a_- for $\Sigma_-^- \rightarrow n\pi^-$ has however resisted measurement via the Σ^- decay asymmetry due to uncertainty in Σ^- polarization. Since the procedure described in reference (2) yields a relatively precisely specified and rather large value for the $\sin\theta \cos\theta$ term in Σ^- polarization over the K^- momentum range 350 to 450 Mev/c, we have been able to determine the helicity $a_- = -0.16 \pm 0.21$.³

Figure 1 shows the measurements as a function of momentum, of $(a\bar{P})_{\sin\theta \cos\theta}$ for the reaction $K^-p \rightarrow \Sigma^- v^+$, where θ is the c. m. angle between the K^- and v^+ , and \bar{P} is the average value of the Σ^- polarization over the angular interval $0.95 > |\cos\theta| > 0.30$ as described in reference 2. The solid curve shows the value of \bar{P} for this reaction, calculated using a least-squares fitting program to determine the S-, P-, and D-wave amplitudes.⁴ These

amplitudes are essentially those of reference 2, except that the full width of the $D_{3/2}$ resonance is now $\Gamma \approx 25$ Mev. The dashed curve shows a maximum-likelihood fit to $a_{\bar{P}}$ over the previously noted momentum range, resulting in the value of $a_{\bar{P}}$ quoted above. Similar procedures applied to Σ_{+}^{+} and Σ_{0}^{+} decays yield the values shown in column 2 of Table 1. The uncertainties are statistical and correspond to standard deviations for a gaussian likelihood function ($e^{-1/2}$ points on the likelihood curve). The uncertainties in the calculated Σ polarization are small relative to this. From our experiment, a_{+} , a_{-} , and a_{0} all have the same sign. The overall sign is determined by the sign of a_{0} as measured by Beall et al.⁵ Column 3 gives a as measured by other experimenters.^{6,7}

We shall now describe these asymmetries in terms of the $|\Delta T| = 1/2$ triangular relationship of Gell-Mann and Rosenfeld.⁸ In their notation, $\underline{N}_{+} + \sqrt{2} \underline{N}_{0} = \underline{N}_{-}$, where $\underline{N}_{\pm, 0}$ signifies the decay amplitudes for $\Sigma_{\pm, 0}^{\pm}$. For spin-1/2 hyperons decaying into S and P states with amplitudes S and P, we have $\underline{N} = \underline{S} + \underline{P}$ and

$$a = \frac{2 \operatorname{Re} S^{*} P}{|S|^2 + |P|^2} = \frac{2 SP}{S^2 + P^2} \quad (1)$$

Here we have used time-reversal invariance to relate the phases of these amplitudes to the πN -scattering phase shifts. These phase shifts are listed in reference 8 and are seen to be small. Taking the amplitudes to be real (the resulting error is small relative to experimental uncertainties), one then describes the various decay amplitudes as vectors in an S-P plane. The magnitudes of the vectors are determined from the three decay rates.⁹

The directions are obtained from Eq. (1) by expressing $a = \sin 2v$, where v is the angle with respect to the coordinate axis. Two ambiguities remain--the labeling of the S and P axes.

Using the combined values for the helicities a listed in Table I, we construct Fig. 2. The two directions for N_0 arise from $|a_0|$ being less than one, corresponding to S/P greater or less than one. For $a_0 = -1$, the triangle would close well within experimental errors, and the triangular relationship given by the $|\underline{\Delta T}| = 1/2$ rule would hold. The inconsistency with the $|\underline{\Delta T}| = 1/2$ rule lies between two and three standard deviations.

REFERENCES

^oWork supported by the U. S. Atomic Energy Commission.

[†]National Academy of Sciences Fellow.

1. M. Ferro-Luzzi, R. D. Tripp, and M. B. Watson, *Phys. Rev. Letters* 8, 28 (1962).
2. R. D. Tripp, M. B. Watson, and M. Ferro-Luzzi, *Phys. Rev. Letters* 8, 175 (1962).
3. The decay asymmetry is written as $1 + aP_{\Sigma} \cos\phi$, where ϕ is the angle between the hyperon polarization, P_{Σ} , and the nucleon direction. This convention has the merit of having the nucleon helicity equal to the decay asymmetry parameter. The more usual, but not universal, convention of following the pion leads to an annoying minus sign relating the helicity to the asymmetry parameter.
4. M. B. Watson, M. Ferro-Luzzi, and R. D. Tripp (to be submitted to *Phys. Rev.*).
5. E. F. Beall, Bruce Cork, D. Keefe, W. C. Murphy, and W. A. Wenzel, *Phys. Rev. Letters* 9, 75 (1962).
6. Bruce Cork, L. T. Kerth, W. A. Wenzel, J. W. Cronin, and R. L. Cool, *Phys. Rev.* 120, 1000 (1960).
7. See reference 5. The value we quote uses the $|\Delta T| = 1/2$ rule, but in only a very weak way involving the πN phase shifts. We have adjusted their uncertainties to correspond to standard deviations.
8. M. Gell-Mann and A. H. Rosenfeld, *Ann. Rev. Nuclear Sci.* 7, 454 (1957).

9. W. H. Humphrey and R. R. Ross, *Low-Energy Interactions of K^- Mesons in Hydrogen*, Lawrence Radiation Laboratory Report UCRL-10018, January 12, 1962 (to be submitted to *Phys. Rev.*). The mass difference has been accounted for to first order by dividing the decay rates by the phase-space factor P/E to obtain $|N|^2$. If we assume the $|\Delta T| = 1/2$ rule, these decay amplitudes alone show that the triangle is nearly a right triangle (94 ± 5 deg).

Table I. Asymmetry parameters (helicities) in Σ decay

	<u>This experiment</u>	<u>Other experiments</u>	<u>Combined</u>
a_-	$-0.16 \pm .21$		$-0.16 \pm .21$
a_+	$-0.20 \pm .24$	$-0.03 \pm .08^a$	$-0.05 \pm .08$
a_0	$-0.90 \pm .25$	$-0.70 \begin{matrix} -.08^b \\ +.09 \end{matrix}$	$-0.79 \begin{matrix} -.08 \\ +.09 \end{matrix}$


^a Reference 6.


^b Reference 7.

FIGURE LEGENDS

Fig. 1. Experimental values of $(\sigma P)_{\sin\theta \cos\theta}$ as a function of K^- laboratory momentum for the reaction $K^- + p \rightarrow \Sigma^- + \pi^+$. The dashed curve is the maximum likelihood fit to the data from 350 to 450 Mev/c; the solid line is the calculated behavior of $(\sigma P)_{\sin\theta \cos\theta}$.

Fig. 2. The Σ decay amplitudes in the S-P plane. The $|\Delta T| = 1/2$ rule requires that the three amplitudes form a triangle.





MU-26847