

# UC Riverside

## Journal of Citrus Pathology

### Title

Dispersal Behavior of *Diaphorina citri* Kuwayama (Hemiptera: Psyllidae) Under Laboratory Conditions

### Permalink

<https://escholarship.org/uc/item/4q36c4zv>

### Journal

Journal of Citrus Pathology, 1(1)

### Authors

Ruan, Chuanqing  
Liu, Bo  
Wu, Zhenquan  
et al.

### Publication Date

2014

### DOI

10.5070/C411024832

### Copyright Information

Copyright 2014 by the author(s). This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

**4.6 P**

**Dispersal Behavior of *Diaphorina citri* Kuwayama (Hemiptera: Psyllidae) Under Laboratory Conditions**

Ruan, C.<sup>1</sup>, Liu, B.<sup>1</sup>, Wu, Z.<sup>2</sup>, Li, T.<sup>2</sup>, Hu, H.<sup>1</sup>, Fan, G.<sup>1</sup>, Duan, Y.<sup>3</sup>, and Hall, D.G.<sup>3</sup>

<sup>1</sup>Fujian Academy of Agricultural Sciences, Fuzhou, Fujian 350003, China

<sup>2</sup>Fujian Agriculture and Forestry University, Fuzhou, Fujian 350002, China

<sup>3</sup>USDA-ARS, 2001 South Rock Road, Fort Pierce, FL 34945 USA

Asian citrus psyllid (ACP), *Diaphorina citri* Kuwayama, is the vector of huanglongbing (HLB), the most devastating disease of citrus worldwide. Knowledge of ACP dispersal behavior in locating host plants may contribute to our understanding of the spread of HLB within and between citrus trees. We conducted research in laboratory to evaluate ACP host plant finding behavior. In a free-choice situation, ACP adults initially settled at equal rates among seedlings of *Rhododendron simsii* (non host plant for ACP), *Murraya paniculata* and “Lugan” *Citrus reticulata*. However, at 18 and 42 h after ACP were released, the mean number of adults per plant on *R. simsii* was significantly lower than on *C. reticulata* and *M. paniculata*, respectively. Numbers of adults on *C. reticulata* and *M. paniculata* remained statistically equivalent until 90 h after releases. Within a plant, the ratio of ACP adults on new flush shoots did not significantly increased in comparison to other parts of the tree during 1-4 d after the psyllids were released. Even on 7<sup>th</sup> day after ACP release, about 30 % adults were observed on plant locations other than flushing shoots. The results indicated that ACP adults differentiated between host and non-host plants faster than they differentiated between different parts of a host plant.