Lawrence Berkeley National Laboratory

Recent Work

Title SUPERHEAVY ELEMENTS

Permalink https://escholarship.org/uc/item/4qh151mc

Author Thompson, S.G.

Publication Date 1972-05-01

1

LBL-665 Preprint

LBL-665 ,

SUPERHEAVY ELEMENTS


S.G. Thompson and C.F. Tsang

May 1972

AEC Contract No. W-7405-eng-48

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545



DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

SUPERHEAVY ELEMENTS

S. G. Thompson and C. F. Tsang

Nuclear Chemistry Division Lawrence Berkeley Laboratory University of California Berkeley, California 94720

LBL-665

Introduction

Since 1965 there has been considerable interest among nuclear physicists and chemists in the possibility of discovering superheavy elements. Extensive experimental efforts have been made in the past four years to detect them in nature, but up to this time the results are negative. Now various groups are attempting to make these superheavy elements in heavy-ion nuclear reactions, but results so far are inconclusive. With the new SuperHilac (Super Heavy Ion Linear Accelerator) being completed at Berkeley, a further major effort will soon be made in creating these superheavy elements.

In this article we will summarize recent efforts and indicate some future possibilities. Thus we shall begin with a general discussion of the developments leading to the expectations concerning superheavy elements. The theoretical predictions of the nuclear and chemical properties will be presented. The search for these elements in nature by many groups in the U.S. and Europe will then be summarized. The possibility of producing these elements by neutron-capture and heavy-ion reactions will then be discussed and some yetunanswered problems involving these methods will be pointed out. In the concluding section, some current thoughts will be given on various aspects of a new field of research in which "superheavies" are only a part. We have attempted to impart some feelings regarding the significance of the push into the previously inaccessible domains. Highly technical details will not be discussed, and complete and unbiased referencing has not been attempted. For those who wish to make a study in greater depth, a number of excellent review articles (1) are currently available.

LBL-665

Background

Superheavy elements are those elements that lie somewhat beyond the end of the present periodic table (2). Interest now is focused on a region of isotopes - centered at atomic number 114 and neutron number 184 - expected to have special stability. The nucleus 114_{184}^{298} (signifying atomic number 114, mass number 298 and neutron number 184) is expected to be extremely stable due to the closing of both a proton shell and a neutron shell at this location. Such shell closures, sometimes referred to as magic numbers, are somewhat analogous to the closing of electronic shells in atoms that give extra chemical stability to certain elements such as the rare gases.

About 90 natural elements are found in nature, and 15 more have been made artificially in the past 30 years. The element with the largest atomic number is hahnium, number 105. Isotope 105Ha²⁶² is so unstable that it can only be produced in extremely small amounts (3), and it disappears in a few minutes by radioactive decay. These known elements form a peninsula in a plane of proton and neutron numbers (Fig. 1), surrounded on three sides by a "sea of instability."

In trying to extend the periodic table still further it is well to understand the basic reason for the limited number of elements (4): Why are there now 105 elements rather than two or three, say, or two or three thousand? The underlying physics responsible for the limited extent of the periodic table is the competition between the cohesive nuclear forces and the disruptive electrostatic forces due to the protons. The limit of the periodic table at $Z \approx 105$ is set by the process of nuclear fission, which takes place when electrostatic repulsion between protons overcomes nuclear cohesion.

100.00

LBL-665

It has been recognized for some time that this limit to the periodic table, set by electrostatic repulsion, could be extended somewhat by nuclear shell effects. Thus the presence of a closed shell of protons or neutrons or preferably both - beyond the end of the periodic table would provide extra binding and extra stability to the nucleus (5). With suitable techniques it might then be possible to reach an island of superheavy nuclei, centered about this magic nucleus, with relatively long half-lives against fission.

No progress was made in this direction for several years, principally because it was assumed that the next closed proton shell, i.e., the next proton magic number, would be at atomic number Z = 126, in analogy with the known neutron magic number N = 126. Proton number Z = 126 was too far beyond the present periodic table to be reached with any kind of nuclear reaction available at that time.

The picture began to change as a result of a suggestion by H. W. Meldner (6) in 1965 that Z = 114, rather than 126, was the next magic number. The reason the neutron and proton closed shells do not occur at the same numbers for heavy elements may be traced to the influence of the electrostatic energy, which is beginning to play an increasingly important role towards the end of the periodic table.

A second factor that changed the outlook for superheavy elements was an improved insight into the relation between magic numbers and the height of the potential energy barrier against fission, which was achieved by W. D. Myers and W. J. Swiatecki (7) at about the same time that Meldner was finding evidence for a closed shell at Z = 114.

LBL-665

The result of the work of Myers and Swiatecki was the rather startling estimate that the stability against fission for a hypothetical nucleus with closed neutron and proton shells might be as high as - or even higher than that of many heavy elements. This result stimulated a considerable amount of theoretical and experimental work on the possible existence of superheavy nuclei.

The prediction of a doubly closed shell at Z = 114 and N = 184 together with the understanding of how shell effects increase stability against fission was still not sufficient to make detailed quantitative calculations. A way was needed of making rather precise calculations of nuclear masses and of deformation energies -- then decay modes and half-lives could be estimated. Microscopic models e.g., Nilsson's shell model (8), while giving local <u>changes</u> in nuclear masses very well, are subject to large errors in predicting the absolute values or general trends. At the other extreme the liquid-drop model (9), gives the absolute magnitude and general trends very well, but is unable to reproduce local fluctuations caused by shell effects.

A significant advance in the calculation of masses was obtained by merging the shell model with the liquid-drop model. The shell effects or local fluctuations (of the order of a few MeV) are extracted from the results of shell model calculations and combined with the liquid-drop binding energies (about 2000 MeV for a heavy nucleus). The philosophy of this two-part approach was proposed by Swiatecki (10) and others, but the prescription for merging was developed by V. Strutinsky of the USSR about 1966 (11), and came to be known as the Strutinsky method. The physical basis of the method is still under extensive investigation by various groups (12), but the method itself has been successfully applied to a host of phenomena. Besides reproducing experimental

LBL-665

nuclear masses (13) to within 2 MeV, the method also makes it possible to explain the existence of a large number of fission isomers (14), and provides a basis for the quantitative understanding of asymmetric fission (15) (i.e., the tendency of a heavy nucleus to split into two unequal rather than equal parts), which has been one of the outstanding problems in fission for more than 30 years.

Strutinski's method was then employed by S. G. Nilsson and co-workers (16) to make the first comprehensive predictions of the properties of superheavy nuclei. These results, which became available in 1968, indicated that some "superheavies" might have half-lives long enough for them to exist in nature and immediately "triggered" experimental searches for them at Berkeley and elsewhere, as will be discussed later. Detailed calculations were also made by several other groups (17), the most recent of which are those of J. R. Nix <u>et al</u>. (18) which will be discussed in more detail below. Great caution should be exercised in considering the theoretical results. These calculations involve great uncertainties. Thus the prediction of a half-life of 10^9 years may be uncertain by a factor of 10^6 either way; i.e., the half-life may well be anything between 10^3 and 10^{15} years.

Theoretical Predictions of Superheavy Nuclei

The above-mentioned calculations (18) indicate that the region centered around Z = 114 and N = 184 should be very stable. These nuclei form an island somewhat beyond the tip of the peninsula of known elements in a plane of proton and neutron numbers as shown in Fig. 1. Contours of total half-lives involving all major modes of decay, namely, spontaneous fission, alpha decay, and beta decay are shown in Fig. 2. Note that the island centers around proton number 110

LBL-665

(instead of 114) and neutron number 184. The shift from proton number 114 to 110 is mainly due to the competition between spontaneous fission (where 114 is the most stable) and alpha decay (where nuclei with lower proton numbers are more stable).

The island may be divided into four regions: the top, where the dominant mode of decay is alpha particle emission; the lower where the dominant decay is beta decay; and the two regions on the sides, where the dominant decay is spontaneous fission. In the alpha decay region the half-lives increase as atomic number, Z, decreases. These contours show a characteristic kink near the magic number. In the spontaneous-fission regions, the contours have a "diamond-like" shape centered around the magic numbers. (In this calculation, the neutron magic number is 184, but the proton magic region extends from Z = 114 to Z = 120). As regards beta decay, the half-lives decrease with decreasing proton numbers and distance from the island of stability. The beta-stable nuclei are marked. They form a belt extending diagonally across the island through the nucleus Z = 110 and N = 184. The longest-lived nucleus in the island appears to be 110_{18h}^{294} (or could be an adjacent odd A or odd Z nucleus), which has half-life as long as 10⁹ years. Such a half-life is nearly as long as the age of the solar system. If one considers only nuclei with half-lives 1 minute or longer, one is confined to an island with proton numbers between 106 and 116, and neutron numbers between 174 and 192. These are the nuclei experimentalists are attempting to produce.

The results of the estimates concerning the properties of superheavy nuclei raised a number of new questions. Do these elements exist in nature? Could they have been formed by astrophysical processes during the formation of

the solar system? Can they be produced in nuclear reactions -- for example, by neutron capture or by heavy ion reactions? What are their chemical properties? We shall first discuss their predicted chemical properties.

Predicted Chemical Properties

If the superheavy elements do become available, experimental studies of their chemical and physical properties will be a subject of considerable interest. The study of such superheavy nuclei and atoms will present a new frontier to the nuclear and inorganic chemists. For example, the study of the chemical properties of the superheavy elements should give some indication of how far the periodic system of the elements can be extended and, at the same time, should shed new light on the underlying electronic properties that allow the periodic system to exist.

The expected properties of these elements had already received significant attention as early as 1958 when G. T. Seaborg (19) predicted properties of the elements up to atomic number 118, eka-radon. These predictions in general have been borne out by more recent studies.

As a starting point for discussion of the chemistry, let us consider some experimental information concerning the most recently discovered elements. Element 103 was confirmed (20) as being the last member of the actinide series in which the 5f electron shell is filled. Element 104, rutherfordium, was found (21) to resemble hafnium in its chemical properties, confirming the expectation that the 6d electron subshell is filled. Experimental studies of the chemistry of element 105, hahnium, have not been carried out to date.

LBL-665

Detailed predictions concerning the chemical properties of superheavy elements are not easy to make. Nevertheless some progress in this direction has been made by using two different methods. The first is an extension of Mendeleev's method, in which the behavior of the well known elements as a function of their chemical group and period is extrapolated into unknown regions. Second, the order in which the electrons fill their orbits is studied by doing self-consistent calculations for the electrons surrounding the nuclei, e.g., by relativistic Hartree-Fock-Slater calculations. Such calculations have been performed by several groups (22) at Los Alamos, Oak Ridge, Northwestern University, Frankfurt, and elsewhere.

Certain effects that are negligible in the light elements are predicted to become very important in superheavy atoms, e.g., relativistic effects will be quite large and the spin-orbit splitting of levels becomes a dominant feature. These relativistic effects are likely to produce unexpected chemistry in certain superheavy elements. The physical limit of the periodic system, as we know it, may occur approximately at atomic number 170 (23). At about this point the inner electron shells might undergo a critical change because of the very large electrostatic field that exists in the atom.

The expected positions in the periodic table of the superheavy elements is indicated in Table I. The locations of elements 104 to 112 are the result of filling of the 6d electron subshell, which makes them homologs of the elements hafnium through mercury. These elements are expected to have their 7s electrons more strongly bound than the homologous elements hafnium to mercury, but the atomic radii of the "superheavies" should be slightly larger. The elements up to the middle of the series should tend to reach their maximum oxidation states,

but near the end of the series the ionization energies should be very large and these elements should be good noble metals. The metallic state should be predominant.

Elements 113 through 118 are characterized by the filling of the 7p subshell and are thus homologous with the elements thallium through radon. Element 114 can be called eka-lead, using Mendeleev's terminology. These elements will tend to prefer lower oxidation states than their homologs. The calculations (22) indicate that element 115 will be monovalent, thus precluding the extrapolation of its properties from those of As, Sb, and Bi, which only show oxidation states of 3 and 5. Element 115 therefore might exhibit a radically new chemistry differing from that of other monovalent elements as much as the chemistry of rubidium and silver or cesium and thallium differ from each other.

Elements 119 and 120 should be very similar to their homologs, francium and radium.

The elements in the vicinity of atomic number 120 to 125 present another interesting problem. At about this point a new inner 5g transition series with a maximum of 18 electrons is expected to begin. However, it seems possible that the 5g and 6f shells may be filled more or less simultaneously and it might be impossible to distinguish between the two shells. If this occurred, it would give rise to a series of 32 elements, for which Seaborg proposes the name "superactinides."

The predicted properties of some of the superheavy elements are shown in Table II.

LBL-665

The magnitude of the effort on the chemistry of superheavy elements will depend to a large extent on the number of elements produced and the range of half-lives and decay modes of the various isotopes. At Berkeley there will be collaboration with groups at Lawrence Livermore Laboratory, the Argonne National Laboratory, the Oak Ridge National Laboratory, and the Los Alamos Scientific Laboratory. Preparations are under way to carry out many different experiments. For example, very efficient extraction chromatographic separations based on the behavior of homologous elements have been developed already by Horwitz and his associates at Argonne (24).

It is not within the scope of this article to discuss the detailed procedures which might be used to make separations and study chemical properties. However, many different types of separations based on homologous elements have been considered, some of which, for example, come from an excellent series of monographs on radiochemical separations of the elements published by the National Academy of Sciences (Nuclear Science Series). A variety of separations are included, based on ion exchange, volatility, oxidation-reduction, solvent extraction, and precipitation methods.

The chemists who look forward to investigations of the chemical properties of the "superheavies" are faced with many difficulties. First of all, when these new elements are produced, it is likely to come about as a result of the heavy-ion reactions (discussed later). In this case the yields are likely to be small and many of the half-lives would be very short. Therefore, deductions of chemical properties are likely to be made by application of tracer methods involving rapid separations of very small amounts (even a few atoms) of radioactive materials and comparisons with homologous tracer elements.

LBL-665

Finally, it should be emphasized that it may be essential to apply chemical methods to the identification of the atomic numbers of some of the new elements. Though observation of alpha particle and spontaneous-fission events having higher energies than those associated with the decay of previously known isotopes would be an almost certain indicator of "superheavies" (25), such an indication, however, would not be sufficient to reveal the atomic number and identify the element. Nor would measurements of the mass numbers of the nuclei in question be sufficient. An element assignment could be made on the basis of measurements of the energies of the characteristic x-rays, but if the yields of the products formed in the reactions are small it may be difficult or even impossible to employ this method successfully. Another method commonly used to make element assignments (e.g., elements 102, 103, 104, and 105 (26)) is the observation of decay to daughters having well known characteristics. But this method may not be useful for "superheavies" because, as indicated in Fig. 2, the decay chains are expected to be terminated by spontaneous fission in the "sea of stability." Thus the chemical separation methods would be necessary. Even in this case proof of the atomic number may not be simple and straightforward. It might be necessary to decide on the basis of separations of tracer amounts of the element, without the benefit of prior knowledge whether element 110, for example, is more like platinum than it is like iridium. On the other hand, if it were similar to mercury with respect to the volatility of its metallic state, the identification might not be so difficult. Such problems will be very challenging, especially if some of the elements exhibit unexpected behavior.

LBL-665

Early Experimental Work

The first attempts (27) to produce superheavy elements by means of heavy ion reactions were carried out at Berkeley (28) in 1967 in response to the early suggestions by Myers and Swiatecki and by Meldner. Bombardments of 248 Cm with 40 Ar were carried out at the Berkeley Heavy Ion Linear Accelerator, using very sensitive apparatus for detecting spontaneous-fission events. The results were negative, in agreement with the results of the first comprehensive calculations by Nilsson <u>et al</u>. (16) which became available in 1968. The results of these calculations, however, stimulated an extensive search for superheavy elements in nature, as discussed in the following section.

Search for Superheavy Elements in Nature

The prediction by Nilsson <u>et al</u>. that the half-life of the nucleus 110_{184}^{294} (eka-platinum) should be in the neighborhood of 10^8 years suggested that small amounts of superheavy elements might be present in nature. The presence of these elements on the earth could have resulted from their formation along with the other elements at the time the earth was formed. If some of the nuclei have half-lives near 2 × 10^8 years, small fractions could have survived the period ($\sim 4.5 \times 10^9$ years) since the earth was formed.

Since large uncertainties are possible in these estimates of halflives and since odd A and odd Z-odd N nuclides generally exhibit retardation in decay compared with even Z-even N nuclei, the surviving nuclides could extend over several atomic numbers, e.g., 108-115. In addition, a search for very heavy particles in cosmic rays at high altitudes by P. H. Fowler and others (29) indicates the presence of uranium and adjacent elements and even suggests a

LBL-665

possible contribution from elements with atomic number greater than 100. Thus the possibility exists that superheavy nuclei might have been produced in more recent cosmic events (30), and shorter-half-life nuclei $(10^4-10^5 \text{ years})$ could conceivably have been deposited in small amounts on the surface of the earth.

A search for the new elements on the earth depends on suitable choices of the most promising minerals and ores containing elements having chemical properties most closely resembling those of the elements being sought. Therefore, certain assumptions had to be made about the chemical properties of the new elements (see the section on chemistry). The search for eka-platinum in nature was undertaken (31, 32) on natural platinum ore containing significant amounts of the neighboring elements. This selection was made on the basis that the most prominent chemical characteristic of elements 108 to 114 is expected to be their predicted nobility, and one might expect them to be found with the noble metals. Even so it is not certain that the superheavy elements will exhibit completely analogous chemical behavior to their homologs. Obviously a pure or purified metal might not be an ideal source since a high degree of selectivity in refining might eliminate homologous elements if they are not almost identical. At Berkeley the search for element number 110 in natural platinum ores utilized a variety of techniques (31), for example, the low-background counting of neutrons, gamma rays, and spontaneous fissions. The Berkeley group also used x-ray fluorescence, mass spectrometric, and activation-analysis techniques. The results were negative and corresponded to a limit of less than 10^{-11} gram of superheavy element per gram of platinum.

The most sensitive methods employed to search for the new elements are those involving the detection of spontaneous-fission events. In this case the assumption is made that the elements are not completely stable and that

spontaneous fission always occurs in the termination of a decay chain. This condition seems to hold for all the theoretical calculations and is well illustrated by the results in Fig. 2. The detection of spontaneous-fission events can be made either by observing fission tracks or by counting the neutrons emitted when spontaneous fission occurs.

The latter method has been employed by Grimm, Herrmann, and Schüssler (33), using a He³-filled proportional-type neutron counter. They examined multi-kilogram samples of pure and ore-grade minerals. Samples were selected on the basis of both homologous chemical behavior and geochemical rules, and covered the range from eka-osmium to eka-bismuth elements. They found no evidence for superheavy nuclei.

The method of observing fission tracks has been used by G. N. Flerov and co-workers (34), who reported results from spontaneous-fission measurements on lead-bearing samples -- in particular, lead glass -- which they felt could be explained as due to the presence of superheavy nuclei. These measurements were made by scanning plastic track detectors (about one square meter of Mylar foils) that had been in contact with lead foils for 100 days. Further investigations on lead glass samples, including a fragment from an 18th century glass vase, showed excess spontaneous-fission events above those expected from the small amounts of uranium and thorium present in the samples. However, similar experiments on other lead-bearing minerals were inconclusive. The results on lead glass appear to be confirmed by fission counting samples in a large area (1.96 square meters) proportional counters (35). Assuming a halflife of 10^9 years for the spontaneously fissioning nuclei, the concentration would be $v10^{-12}$ gram per gram of lead.

LBL-665

P. B. Price <u>et al</u>. (36) used a very sensitive method to obtain results which seem to be in conflict with those of Flerov <u>et al</u>. They searched for fission tracks accumulated over millions of years in ancient minerals (more than 10^8 years old). Their results for lead (Hardystone) and gold (quartz) bearing samples gave no evidence for the presence of superheavy elements, and concentration limits of less than 10^{-15} gram per gram of lead and less than 10^{-17} gram per gram of gold were assigned. However, since the samples were of different composition and origin, it might be possible for superheavy elements to be present in the lead samples of Flerov and not in the lead minerals of Price. Price (37) and D. Lal <u>et al</u>. (38) studied fission tracks in moon rock in search of the fission of superheavy nuclei, but the results were inconclusive.

Advantage can also be taken of the fact that the spontaneous fission of superheavy elements is expected to be different from the fission of wellknown elements. Rather simple theoretical considerations strongly indicate that when these new elements undergo fission the fragments should have significantly higher energies and should involve the emission of a larger number of neutrons per fission - probably about ten, rather than about two as in the case of uranium (39). By measuring the number of neutrons emitted in each fission event one should be able to distinguish superheavy elements from other elements. This approach was employed by the Berkeley group (40), who have recently concluded an extensive search for superheavy elements in nature. They used a large liquid scintillator to measure the number of neutrons per fission event in large samples of minerals and ores. Their counter was located in a tunnel about 850 feet below the surface of the earth to minimize the influence of interfering cosmic radiation. More than 40 samples of ores, minerals, and rocks were examined, including manganese nodules, moon rocks, large samples of gold, and platinum in their natural states. The selection of samples was also made to include the range of elements from eka-platinum to eka-bismuth. No evidence for superheavy elements was found in any of the samples. Groups at Oak Ridge (41) and in the U.S.S.R. (42) are also employing He³ counter systems to detect events in which large numbers of neutrons are emitted.

A number of other searches for superheavy elements have been made, but none have given conclusive evidence of their presence. Although the results up to now do not definitely rule out the presence of these elements in nature, the weight of evidence is such as to suggest strongly that they do not exist on the earth. Thus the question arises, how can we account for their absence? If the half-lives are much less than 2×10^8 years they would have disappeared by radioactive decay during the 4.5×10^9 years since the earth was formed. On the other hand if the half-lives are long enough, the conclusion would be that they probably were not formed during nucleogenesis in which the other elements were formed. Although some papers have been published suggesting that there should be no difficulty in producing the new elements in the neutroncapture process, an examination of some of the most reliable mass formulas indicates that there may be difficulty in binding neutrons to nuclei with such necessarily large neutron excess. Even if there were no difficulty in binding neutrons, neutron-induced fission or spontaneous fission might be able to cut off formation of the products of interest. Stability against fission is certain to be very small in the case of the intermediate nuclei in the formation process.

Production of Superheavy Elements in Neutron-Capture Reactions

Possible difficulties in producing the new elements in reactions associated with nucleogenesis have been indicated above, but these difficulties do not necessarily rule out their production by irradiation of high-atomic-number targets with neutrons on a slow time scale in, for example, the High Flux Isotope Reactor at Oak Ridge. However, the heaviest nucleus produced so far by this means is Fm^{257} , and this same isotope is also the heaviest one produced to date in thermonuclear explosions on a faster time scale. In this case the Fm isotopes were produced by the rapid successive capture of about 20 neutrons in targets such as U^{238} . Therefore these methods do not appear at all promising.

A possible means of circumventing the difficulties inherent in the production by neutron-capture processes in the extreme cases of fast or slow time scales has been suggested by H. W. Meldner (43). His proposal is to utilize capture of neutrons from thermonuclear explosions that would be controlled to allow some intermediate beta decays. This approach would tend to minimize some of the difficulties associated with the attempts described above but might be extremely difficult from a technical point of view.

Production with Heavy Ions

Probably the most promising approach to the production of superheavy nuclei involves the use of heavy ions. One of the major differences between the processes used to make elements up to 105 and those necessary for producing superheavy elements is that in the latter case there is a gap of very unstable nuclei between the island and the peninsula (see Fig. 1). Thus it is impossible to go step by step to the island; it is apparently essential to make a big jump

LBL-665

to the island by means of relatively heavier projectiles (such as Ca^{48} , Ge^{76} , and Kr^{86}). In this process the heavy ions are accelerated to a high energy and used to bombard a target nucleus such as Th^{232} or U^{238} . Hopefully, the projectile and target will fuse together, forming compound nuclei within the island of stability. Until recently only those projectiles up to Ar^{40} were available with sufficient energies to fuse with heavy targets, and it was not possible to make a jump close to the center of the island by using Ar^{40} . But with the newer heavy-ion accelerators at Dubna and Orsay, heavier projectiles are available. Even more intense beams of very heavy projectiles should become available soon at Berkeley, and also near the end of 1974 at the "Universal Linear Accelerator" in Darmstadt, Germany.

In Fig. 3 we show some of the projectile-and-target combinations that may make it possible to land close to the island of stability. However, even with the available long-lived elements as projectiles the center of the island cannot be reached unless we use special methods such as those suggested below. Furthermore, it does not seem advisable to use too heavy a projectile, because such a heavy projectile would overshoot the island and land in a region where the products are very unstable.

One of the most favorable target-projectile combinations, advocated by Swiatecki (44), after consideration of various effects that enter into heavy ion reactions, is

$$Ge^{76} + Th^{232} = 122_{182}^{304} + 4n$$

= $120_{181}^{301} + He^{4} + 3n$

Even with this reaction, certain difficulties are likely to be encountered as discussed in a later section.

LBL-665

In a discussion of heavy-ion reactions it is necessary to recognize the need to accelerate heavy ions to a sufficiently high energy. Why not just mix two elements together and extract a product of much higher atomic number? The electrostatic energy - which, as we have seen before, becomes increasingly important at the end of the periodic table - prevents fusion. The very large positive charges in the target and projectile nuclei prevent them from coming within the very small distances required to make them touch and fuse together. These distances are about 10^{-12} cm. In order to make even a relatively light ion, such as argon, fuse with uranium the argon must have its energy raised to about 200 MeV. It is rather difficult to achieve such high energies. One reason is that all accelerators require projectile atoms that have a net charge (ions). Thus electrons must be removed from the atoms in "ion sources." The heavier the atom, the larger is the number of electrons that have to be removed for acceleration to high energies. This stripping of electrons presents considerable difficulties and, in some cases, a procedure of stripping in successive stages is used.

One of the main difficulties in reaching the island of stability by heavy-ion reactions is that neither the projectiles nor targets have sufficient number of neutrons to form a compound nucleus near the center of the island. Four methods may be considered for dealing with this problem.

The first is by secondary reactions after multi-nucleon transfer. It has been observed that in the heavy ion collision, a few neutrons may be transferred from the target to the projectile or protons may be transferred from the projectile to the target (45). In both cases the projectile ends up being more neutron-rich. These neutron-enriched projectiles may then make a second reaction with the target, to form superheavy nuclei. Estimates of the yields to be

LBL-665

expected from such a process indicate there is some chance of success (46).

It is well known that the fission of heavy nuclei such as Cf^{252} yield neutron-rich fission fragments which are then available as projectiles with enough neutrons to make favorable heavy-ion reactions. The difficulty in this method is the acceleration of these fission fragments to a high enough energy to cause compound-nucleus formation when bombarding a target (47). The intensity of the beam of accelerated fission fragments may be too small to cause observable heavy-ion reaction products. One way to get over this difficulty is to accelerate, for example, the nucleus U^{238} to high energies first and then to allow the U^{238} particles to collide with deuterons or helium, in which case sufficiently energetic fission fragments would be produced.

Another method suggested by A. Marinov <u>et al.</u> (48) for producing veryneutron-rich projectiles involves the use of very-high-energy protons (\sim 24 GeV) which have about 100 times the energy required for ordinary nuclear reactions. The high-energy protons colliding with heavy target nuclei cause the target to fragment into pieces some of which would be very neutron rich and might have sufficient energies to produce heavy-ion reactions. So far this method has not been successful despite some initially encouraging results.

Another suggested method involves accelerating very heavy nuclei such as uranium and using them to bombard another uranium nucleus to give a total atomic number of 184 and mass number of 476 (49). This combination is expected to be very unstable and to divide into smaller pieces. One of the pieces might be right in the island of stability. However, very little is known about this process, and possibly, in such a violent reaction, only many small pieces would be produced.

LBL-665

The above methods are necessarily very speculative at this time; further studies and quantitative estimates of production probabilities are needed. However, they may all be technically possible in the next few years.

Physical Methods of Detection of Products in Heavy-Ion Reactions

The predicted half-lives of nuclei near the center of the island of stability are very long. Thus radiochemical methods might be useful for their separation and identification. However, one must be prepared to allow for the large uncertainties inherent in these predictions. It may be that the theory overestimates half-lives by a large factor and one is confronted with the detection of products with very short half-lives. Further, most combinations of target and projectile (see Fig. 3) will yield products some distance away from the center of the stability island, with very short half-lives. When the halflives are below about ten seconds it becomes difficult to apply radiochemical methods despite great improvements in recent years, and the application of physical methods becomes necessary.

Fortunately, a large number of rapid and sensitive methods of detecting and identifying the products of heavy-ion reactions were already developed in extending the periodic table upward through the transuranium elements 103, 104, and 105. Recent developments in fast electronics, solid state counters, and reliable high-speed computers have been able to meet the stringent technical requirements for studying very short half-lives.

In many of the physical methods of detection, advantage is taken of one of the characteristics of the heavy-ion reactions themselves: When an energetic heavy ion strikes a target nucleus the resulting nucleus is driven forward with

Thompson 23

relatively high energy as a consequence of momentum conservation. Such a highenergy product is able to escape from the target and can be transported rapidly by various means to detectors that record the properties of the subsequent radioactive decay (26). Commonly used methods of transporting such "recoil atoms" to counters involve rapidly flowing gases such as helium. Rapidly moving drums or tapes are also used to collect and transport the recoil atoms to locations near counters that can measure their radiations. Other methods involve bending the recoil atoms in a magnetic field (to separate out the products of interest) and measuring their velocities and energies simultaneously. By such means one can determine the masses of the recoil nuclei. Means are also available for observing the properties of daughters of the radioactive decay and for measuring the energies associated with the decay. Most of the superheavy nuclei are predicted to decay by successive alpha decay before undergoing spontaneous fission (18). The alpha energies are expected to be much larger than those from the decay of previously known nuclei. Thus a measurement of these successive high-energy alpha decays will be characteristic of the presence of superheavy nuclei. These particular methods are only a few examples to demonstrate some approaches being taken to the problem.

Problems Associated with Heavy-Ion Reactions

Even with the most favorable target-projectile combinations there are formidable difficulties, which are summarized as follows:

<u>High Excitation Energies</u>. When a heavy-ion projectile fuses with the target nucleus the resulting compound nucleus has a large excitation energy. Calculations by

LBL-665

L. G. Moretto (50) show that the very shell effects responsible for the stability of the superheavy nuclei (and in fact for the existence of the island of stability) will be destroyed as the excitation energy is increased. Then the stability against fission that comes from the shell effect is lost. The excitation energy can only be reduced by the emission of several neutrons or charged particles. Thus, survival at such high excitation energies becomes a question of the competition between fission and neutron or charged-particle emission. If neutron emission is sufficiently probable, then the superheavy nuclei (minus the number of neutrons emitted) may survive. If fission is too probable, then superheavy nuclei will not survive. In typical reactions the emission of four neutrons should be sufficient to reduce the excitation energy to a safe level. In Fig. 3 we show rough estimates for the survival probabilities due to these competing effects and indicate the best landing areas in the island of stability. The optimum combination, $Ge^{76} + Th^{232}$, that Swiatecki has proposed should have a survival probability approximately 1:10,000. That is to say, the yield of the final product is expected to be reduced by the factor 10^{-4} because of fission competition.

<u>Angular Momentum Effects</u>. When a heavy ion collides with a target nucleus, rotational angular momentum is necessarily introduced. The centrifugal forces which arise make the system less stable. Some estimates of the result of this effect have been made which indicate that the expected yields of superheavy products should be further reduced by factors ranging from 10^{-1} to 10^{-3} . <u>Fusion Probability</u>. The largest uncertainty of all has to do with the probabilities for the projectiles and targets to fuse together (44). It is not

sufficient for the projectile and the target nuclei to merely come in contact with each other: an extra push is necessary to force them to fuse together. The probability that they can be made to fuse together into a final spherical compound nucleus involves not only the nuclear inertia which acts against the push, but also the viscosity of the flow of nuclear matter, which is a dissipative effect converting the pushing energy into useless excitation energy. Good estimates of nuclear inertias and viscosities have not been made so far; these are important gaps in our knowledge of nuclear properties.

Present Status

Small beams of energetic Kr ions have been available at Orsay in France, and Zn and Xe projectiles have been available at Dubna in the U.S.S.R. Attempts by both the French (51) and the Soviet (52,53) groups (summarized in Table III together with previous efforts) to produce observable amounts of superheavy nuclei have not been successful at the time of this writing. However a new heavy-ion linear accelerator is near completion in Berkeley that is expected to produce heavy-ion beams with much larger intensities than either the French or the Soviet group. A major assault on the production of superheavy elements will be made. It is a completely open question whether these elements will indeed be made by heavy-ion reactions, and scientists all over the world are anxiously waiting for the outcome. Extensive searches for these elements in nature have been carried out, and it seems that the prospects for finding them in terrestrial materials are greatly diminished by the negative results obtained so far. There

LBL-665

is still a possibility that superheavy elements might be found in cosmic radiation. The results of recent searches have not been conclusive.

If the superheavy elements are discovered, it will be the crowning success of an international effort of many years with contributions being made by many scientific workers and groups from many parts of the world. We would be facing the happy prospect of confronting the comprehensive theoretical map of the island of superheavy elements with new experimental data. The confrontation will provide a new testing ground for our understanding of the chemistry of the elements and the physics of the nucleus. The possible existence of superheavy elements on the earth in some time past or in supernovae would also have impact on other fields such as geology and astrophysics. Practical and useful applications would be forthcoming eventually, as is always the case with basic research, although in most cases definite predictions of the direction and nature of the applications cannot be made.

On the other hand if we were not able to produce the superheavy nuclei either because their half-lives are too short or because of difficulties associated with fusion of projectile and target, there are still many exciting studies that can be made. We will study the collisions between target and projectile nuclei and obtain information on the fragments and radiations emitted as a result of the collisions. This information will be able to tell us about the conditions prevailing during the very brief time when the target and projectile nuclei are together. A study of such transient systems may extend our present knowledge of the nucleus in two ways. First, these systems may have a wide variety of shapes, such as asymmetric dumbells, triaxial ellipsoids and

Thompson 27

cylindrical shapes. Effects of large centrifugal forces on these shapes may also be studied by observing off-center collisions. Secondly, these systems have atomic numbers up to about 200 and mass numbers up to about 500, well beyond our present periodic table. We will be confronted with the most intense electric fields occurring anywhere in the universe, specifically, the fields in the close vicinity of such heavy systems. In the neighborhood of atomic number 170, a critical condition might occur. In this case quantum electrodynamics has to be applied to the understanding of the observed phenomena. The result would be an unusually intimate interaction between nuclear physics and quantum electrodynamics in which the theory of matter and radiation will undergo testing under unique conditions.

References and Notes

 J. R. Nix, Ann. Rev. Nucl. Sci., <u>22</u> (1972); Physics Today (April 1972), p. 30. Thomas Johansson, Sven Göste Nilsson and Zdzisław Szymański, Ann. Phys. <u>5</u>, 377 (1970). G. T. Seaborg, Ann. Rev. Nucl. Sci. <u>18</u>, 53 (1968). G. T. Seaborg and J. L. Bloom, Sci. Am. <u>220</u>, No. 4, 56-67 (1969).

Thompson

28

- 2. The term "Superheavy elements" was first introduced as far back as 1958 by F. G. Werner and J. A. Wheeler (Phys. Rev. <u>109</u>, 126 (1958)), who considered nuclei far beyond the known regions based on the liquid drop model without any shell closure effects. In the present context the superheavy nuclei are predicted to be particularly stable due to shell closure effects.
- 3. A. Ghiorso, M. Nurmia, K. Eskola, J. Harris and P. Eskola, Phys. Rev. Letters <u>24</u>, 1498 (1970).
- 4. W. J. Swiatecki, Talk given at the Research Progress Meeting, Lawrence Berkeley Laboratory, University of California, Berkeley, February 1968, unpublished.
- 5. The known closed shells (magic numbers) for protons and neutrons are at 2, 8, 28, 50, 82, and, for neutrons, also at 126.
- H. W. Meldner, private communication to W. J. Swiatecki (1965), (see Ref. 7);
 Ark. Fys. <u>36</u>, 593 (1967).
- 7. W. D. Myers and W. J. Swiatecki, Nucl. Phys. <u>81</u>, 1 (1966).
- G. Gustafson, I. L. Lamm, B. Nilsson and S. G. Nilsson, Ark. Fys <u>36</u>, 613 (1967).
- The model is more appropriately called "the leptodermous model," see
 C. F. Tsang, Ph.D. Thesis, University of California, Berkeley, Lawrence
 Berkeley Laboratory Report UCRL-18899 (1969).

- W. J. Swiatecki, Proc. Int. Conf. Nuclidic Masses, Vienna, 2nd, July 1963 (1964), pp. 58-66. (Vienna:Springer-Verlag).
- 11. V. M. Strutinsky, Yad. Fiz. <u>3</u>, 614 (1966); Sov. J. Nucl. Phys. <u>3</u>, 449 (1966); Nucl. Phys. <u>A95</u>, 420 (1967); <u>A122</u>, 1 (1968).
- 12. See for example: W. H. Bassichis, A. K. Kerman, C. F. Tsang, D. R. Tuerpe and L. Wilets, UCRL-73044 (Jan. 1971); to be published in "Magic Without Magic (Wheeler Festschrift) (San Francisco:Freeman); also Bull. Am. Phys. Soc. (Series II) <u>17</u>, 507 (1972). S. J. Krieger and C. Y. Wong, Phys. Rev. Letters <u>28</u>, 690 (1972). See also the first reference in Ref. 17.
- 13. See for example: Ref. 16. And also, P. A. Seeger, Proceedings of Int. Conf. on the Properties of Nuclei Far from the Region of Beta-Stability, Leysin, Switzerland, August 31-September 4, 1970, p. 217.
- 14. D. D. Clark, Phys. Today <u>24</u>, No. 12, 23 (Dec. 1971).
- 15. Peter Möller and S. G. Nilsson, Phys. Letters <u>31B</u>, 283 (1970).
 - J. B. Wilhemy and C. F. Tsang, Nucl. Phys. <u>A184</u>, 417 (1972).
- 16. S. G. Nilsson, C. F. Tsang, A. Sobiczewski, Z. Szymański, and S. Wycech,
 C. Gustafson, I. L. Lamm, P. Möller, and B. Nilsson, Nucl. Phys. <u>A131</u>, 1 (1969).
- M. Brack, J. Damgaard, A. Stenholm-Jensen, H. C. Pauli, V. M. Strutinsky, and C. Y. Wong, Rev. Mod. of Phys. <u>44</u>, 320 (1972). J. Grumann, U. Mosel,
 B. Fink, and W. Greiner, Z. Physik <u>228</u>, 371 (1969). Also Ref. 16 and Ref. 18. It is interesting to note the international nature of the theoretical efforts by listing the nationalities of the many investigators: American, Chinese, Danish, German, Polish, Soviet, Swedish and Swiss.

- M. Bolsterli, E. O. Fiset, J. R. Nix, J. L. Norton, Phys. Rev. Letters <u>27</u>, 681 (1971); Phys. Rev. <u>C5</u>, 1050 (1972).
- 19. G. T. Seaborg, <u>The Transuranium Elements</u>, New Haven Yale University Press, 1958, page 277.
- 20. R. Silva, T. Sikkeland, M. Nurmia, and A. Ghiorso, Inorg. Nucl. Chem. Letters <u>6</u>, 733 (1970).
- 21. R. Silva, J. Harris, M. Nurmia, K. Eskola, and A. Ghiorso, Inorg. Nucl. Chem. Letters 6, 871 (1970).
- 22. B. Fricke and J. T. Waber, Actinides Reviews <u>1</u>, 433 (1971); and references therein.
- I. Pomeranchuk and Ya. Smorodinsky, J. Phys. USSR <u>9</u>, 97 (1945). V. S. Popov,
 Sov. J. Nucl. Phys. <u>14</u>, 257 (1972).
- 24. E. T. Horwitz, Private communication (1972).
- 25. See for example Ref. 18.
- 26. A. Ghiorso, Lawrence Berkeley Laboratory Report LBL-213, (July 1971), to be published in <u>Handbook of Experimental Pharmacology</u>.
- 27. An earlier experiment was done by T. Sikkeland (Phys. Letters <u>27B</u>, 277 (1968)) to study fission fragments produced in the reaction ${}_{18}Ar^{40} + {}_{92}U^{238}$. Some of the properties of the fragments were consistent with those expected from the fission of the compound system 110^{278}_{168} , but the data were far from conclusive. The recent calculations show that 110^{278}_{168} is far from the island of stability.
- 28. H. R. Bowman, R. C. Gatti, R. C. Jared, L. G. Moretto, W. J. Swiatecki, and S. G. Thompson, Lawrence Berkeley Laboratory, Nuclear Chemistry Annual Report 1967.(UCRL-17989) Stanley G. Thompson, Talk given at the APS

LBL-665

Meeting, Miami, Nov. 25-27, 1968. M. Nurmia, T. Sikkeland, R. Silva, and A. Ghiorso, Phys. Letters <u>26B</u>, 78 (1967).

- 29. P. H. Fowler, R. A. Adams, V. G. Cowan, J. M. Kidd, Proc. Roy. Soc. (London) <u>A301</u>, 39 (1967).
- 30. Some attention has been given to the question of how superheavy nuclei might be synthesized in astrophysical processes (D. N. Schramm and W. A. Fowler, Nature 231, 103 (1971); M. Kowalski and Br. Kuchowicz, Phys. Letters 30B, 79 (1969)). The proposal has been made that isotopes might be accelerated by staying in phase with the electromagnetic fields of stellar plasmas and reach energies sufficient to undergo fusion reactions with other nuclei.
- 31. S. G. Thompson, R. C. Gatti, L. G. Moretto, H. R. Bowman, and M. C. Michel, Lawrence Berkeley Laboratory, Nuclear Chemistry Annual Report 1968 (UCRL-18667) p. 277; Stanley G. Thompson, talk given at the APS meeting, Miami, Nov. 25-27, 1968; S. G. Nilsson, S. G. Thompson, and C. F. Tsang, Phys. Letters 28B, 458 (1969).
- 32. J. J. Wesolowski, W. John, R. Jewell, and F. Guy, Phys. Letters <u>28B</u>, 544 (1969).
- W. Grimm, G. Herrmann, and H. D. Schüssler, Phys. Rev. Letters <u>26</u>, 1040 (1971).
- 34. G. N. Flerov and V. P. Perelygin, Atomnaya Energiya <u>26</u>, 520 (1969);
 E. Cieslak, Joint Institute of Nuclear Research (Dubna USSR) Report P15-4738, 1969.
- 35. G. N. Flerov, N. K. Skobelev, G. M. Ter-Akopyan, V. G. Subbotin,
 B. A. Gvozdev, and M. P. Ivanov, Joint Institute of Nuclear Research
 (Dubna USSR) Report D6-4554, 1969.

- 36. P. B. Price, R. L. Fleischer, and R. T. Woods, Phys. Rev. <u>1</u>, 1819 (1970).
 37. P. B. Price, private communications, 1972.
- 38. N. Bhandari, S. Bhat, D. Lal, G. Rajagopalan, A. S. Tamhane, and
 V. S. Venkatararadan, Apollo 12 Lunar Science Conference (1971), to be published; Nature 230, 219 (1971).
- 39. J. R. Nix, Phys. Letters <u>30B</u>, 1 (1969).
- 40. E. Cheifetz, R. C. Jared, E. R. Giusti, and S. G. Thompson, "Search for Superheavy Elements in Nature," Lawrence Berkeley Laboratory Report LBL-613 (January 1972), to be published in Phys. Rev.
- 41. R. L. Macklin, F. M. Glass, J. Halperin, R. T. Roseberry, H. W. Schmitt,
 R. W. Stoughton, and M. Tobias, "Neutron Multiplicity Counter," to be
 published in Nucl. Instr. and Methods.
- 42. G. N. Flerov, private communication (Leysin, Switzerland, 1970); see also <u>Proceedings of Int. Conf. on the Properties of Nuclei Far from the Region</u> <u>of Beta-Stability</u>, Leysin, Switzerland, August 31-September 4, 1970, p. 771.
 43. H. W. Meldner, Phys: Rev. Letters 28, 975 (1972).
- 44. W. J. Swiatecki, talk given at Ebeltoft Conference, Ebeltoft, Denmark, May 1971, Lawrence Berkeley Laboratory Report, LBL-549 (Dec. 1971).
- 45. A. G. Artukh, V. V. Avdeichikov, G. F. Gridnev, V. L. Mikheev, V. V. Volkov, and J. Wilczynski, Nucl. Phys. <u>A168</u>, 321 (1971); also Nucl. Phys. <u>A160</u>, 511 (1971); A126, 1 (1969).
- 46. Unpublished work by K. Eskola; W. J. Swiatecki, and C. F. Tsang.
- 47. E. Cheifetz, R. C. Gatti, R. C. Jared, and S. G. Thompson, Phys. Rev. Letters 24, 148 (1970).

LBL-665

- 48. A. Marinov, C. J. Batty, A. I. Kilvington, G. W. A. Newton, V. J. Robinson, and J. D. Hemingway, Nature <u>229</u>, 464 (1971).
- 49. G. N. Flerov, talk given at the Paris Conference dedicated to the centenary of the Mendeleev Periodic Table, Joint Institute of Nuclear Research (Dubna, USSR) report E15-4651 (August 1969).
- 50. L. G. Moretto, Nucl. Phys. <u>A182</u>, 641 (1972); <u>A180</u>, 337 (1972).
- 51. R. Bimbot, C. Deprun, D. Gardès, H. Gauvin, Y. Le Beyec, M. Lefort,
 J. Pèter, Nature <u>234</u>, 215 (1971).
- 52. Yu. Ts. Oganessian, O. A. Orlova, Yu. E. Penionshkevich, K. A. Gavrilov, and Kim De En, Joint Institute of Nuclear Research (Dubna USSR) report P7-6300, 1972.
- 53. G. N. Flerov, S. A. Karamian, Yu. E. Penionshkevich, S. P. Tretiakova, and I. A. Shelaev, Joint Institute of Nuclear Research (Dubna USSR) report P7-6262, 1972.
- 54. We have greatly profited from numerous discussions with W. J. Swiatecki; many of the ideas mentioned in the article originated with him. We also thank S. Kataria, L. G. Moretto, W. D. Myers, C. P. Pezzotti and

G. T. Seaborg for critical comments on the manuscript.

55. Work was performed under the auspices of the U. S. Atomic Energy Commission.

CONVENTIONAL FORM OF PERIODIC TABLE SHOWING PREDICTED LOCATIONS OF NEW ELEMENTS

H												· .					He
Li	Be											В 5	С 6	N 7	8 8	F 9	Ne 10
Na -	Mg						•					Al 13	Si 14	P 15	S 16	CI 17	Ar 18
K 19	Ca 20	Sc 21	Ti 22	¥ 23	Cr 24	M n 25	Fe 26	C0 21	Ni 28	. Си 29	Zn 30	Ga 31	Ge 32	As 33	Se 34	B r 35	Kr 36
Rb 37	- Sr 38	Y 39	Zr 40	Nb 41	M0 42	TC 43	Ru 44	Rh 45	Pd	Ag 47	Cd 48	In 49	Sn 50	51 51	Te	53	Xe 54
C S 55	Ba 56	La 57	H1 72	та 73	₩ 14	Re 75	0s	, lr	Pt	Au 79	Hg 80	T1 81	Pb 82	Bi 83	P0 84	At 85	R n 86
Fr 87	R a 88	AC 89	Rf 104	Ha 105	(106)	(107)	(108)	(109)	(110)	(111)	(112)	(113)	(114)	(115)	(116)	(117)	(118)
(119)	(120)		(154)	(155)	(156)	(157)	(158)	(159)	(160)	(161)	(162)	(163)	(164)	(165)	(166)	(167)	(168)
ANT	HANI	DES	Ce 58	Pr 59	N d 60	Pm 61	Sm 62	Eu 63	Gd 64	Tb 65	Dy 66	H0 67	Er 68	Tm 69	Yb 70	Lu]
ACT	INID	ES	Th 90	Pa	U 92	Np 93	Pu 94	Am 95	Cm 96	Bk 97	Cf 98	99 59	Fm 100	- M d 101	N 0 102	Lr 103	
	IPER INID		(122)	(123)	(124)							/ i i /	/ \ }				
												.*					

Table I

LBL-665

Thompson

μ

Table 11. Some Physical and Chemical P						
Element	110	111	112	113	114	115
Electronic Ground-State Configuration	6a ⁴ 6a ⁴ 7s ²	6a ⁴ 6a ⁵ 7s ²	6a ⁴ 6d ⁶ 7s ²	7p ¹ 7s ²	$7p^27s^2$	$7p^27p^17s^2$
Chemical Group	VIII	IB	IIB	IIIA	IVA	VA
Most Probable Oxidation States	+6	+1,3	+1,2	+1	+2	+1,3
First Ionization Potential	9.4	10.3	11.2	7.4	8.5	5.9
Metallic Radius, Å	1.4	1.5	1.6	1.7	1.8	1.9
Ionic Radius, Å		-	 —	1.5	1.3	· -
Density g/cm ³	27.4	24.4	16.8	16	14	13.5
Melting Point ^o C	-	-	-	430	67	- ·
Boiling Point °C	-	-	-	1130	147	. – .
K _α x-ray Energy (keV) [*]	157	161	165	169	173	177

1

* From Ref. 31.

Table II. Some Physical and Chemical Properties of Elements 110 - 115 According to Fricke and Waber (22)

Thompson

ω S

LBL-665

×


Investigators	Reference	Reaction	Results
Sikkeland	27	$18^{Ar^{40}} + 92^{U^{238}} \rightarrow \text{fission fragments}$	fission fragment from 110 ²⁷⁸ (?)
Thompson, Ghiorso <u>et al</u> .	28	$18^{\text{Ar}^{40}} + 96^{\text{Cm}^{248}} \rightarrow 114^{288-x} + xn$	$\sigma < 5 \times 10^{-32} \text{ cm}^2$ for $\tau > 10^{-9} \text{ s}$
Oganesyan <u>et</u> <u>al</u> .	52	$54^{\text{Xe}^{136}} + 92^{238} \rightarrow \text{fission fragments}$	fission fragments from 146^{374} (?)
Flerov <u>et al</u> .	53	30^{2n} ⁶⁶ + $92^{238} \rightarrow 122^{304-x} + xn$	$\sigma \lesssim 5 \times 10^{-30} \text{ cm}^2$ for $\tau > 10^{-8} \text{ sec}$
Bimbot <u>et al</u> .	51	$36^{\text{Kr}^{84}} + 90^{\text{Th}^{232}} \rightarrow 126^{316-x} + xn$	High energy α from 126 ^{316-x} (?)
	• * · ·		
	en e		

Table III. Previous Attempts to make Superheavy Elements by Heavy Ion Reactions

Figure Captions


Fig. 1. Nuclear stability is illustrated in a scheme that shows a peninsula of known elements and an island of predicted stability (nuclei around proton number 11⁴ and neutron number 18⁴) in a "sea of instability." Grid lines show magic numbers of protons and neutrons giving rise to exceptional stability. Magic regions on the mainland peninsula are represented by mountains or ridges.

- Fig. 2. Total half-lives for decay of even Z-even N superheavy nuclei given as contours, which are labeled by the logarithms (to the base 10) of the half-lives in years. The points indicate nuclei that are calculated to be beta-stable. Taken from Nix (1).
- Fig. 3. Available landing places near the island of stability in heavy-ion reactions. These are designated by $\sqrt{}$, \times , or o. The latter two symbols denote landing places that cannot be reached without Cm^{250} or Cf^{252} targets respectively. The longer curve marks out the region where the probability of the compound nucleus surviving four successive neutron-fission competitions is expected to be in excess of 10^{-3} . This region will be decreased to the area indicated by the shorter curve if the calculated fission barrier is arbitrarily cut down by 2 MeV.




٤.

¢.



XBL725-2955



Э

Fig. 3

-LEGAL NOTICE-

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

0

TECHNICAL INFORMATION DIVISION LAWRENCE BERKELEY LABORATORY UNIVERSITY OF CALIFORNIA BERKELEY, CALIFORNIA 94720

٠

.

N