UC Berkeley

UC Berkeley Previously Published Works

Title

MICA-Expressing Monocytes Enhance Natural Killer Cell Fc Receptor-Mediated Antitumor Functions.

Permalink

<https://escholarship.org/uc/item/4qp41942>

Journal

Cancer immunology research, 5(9)

ISSN

2326-6066

Authors

Campbell, Amanda R
Duggan, Megan C
Suarez-Kelly, Lorena P
et al.

Publication Date

2017-09-01

DOI

10.1158/2326-6066.cir-16-0005

Peer reviewed

MICA-Expressing Monocytes Enhance Natural Killer Cell Fc Receptor-Mediated Antitumor Functions

Amanda R. Campbell^{1,2}, Megan C. Duggan^{1,2}, Lorena P. Suarez-Kelly¹, Neela Bhawe¹, Kallan S. Opheim¹, Elizabeth L. McMichael^{1,2}, Prashant Trikha¹, Robin Parihar¹, Eric Luedke^{1,3}, Adrian Lewis¹, Bryant Yung⁴, Robert Lee⁴, David Raulet⁵, Susheela Tridandapani⁶, Veronika Groh⁷, Lianbo Yu⁸, Vedat Yildiz⁸, John C. Byrd^{1,9}, Michael A. Caligiuri^{1,9}, and William E. Carson III^{1,3}

Abstract

Natural killer (NK) cells are large granular lymphocytes that promote the antitumor response via communication with other cell types in the tumor microenvironment. Previously, we have shown that NK cells secrete a profile of immune stimulatory factors (e.g., IFN γ , MIP-1 α , and TNF α) in response to dual stimulation with the combination of antibody (Ab)-coated tumor cells and cytokines, such as IL12. We now demonstrate that this response is enhanced in the presence of autologous monocytes. Monocyte enhancement of NK cell activity was dependent on cell-to-cell contact as determined by a Transwell assay. It was hypothesized that NK cell effector functions against Ab-coated tumor

cells were enhanced via binding of MICA on monocytes to NK cell NKG2D receptors. Strategies to block MICA–NKG2D interactions resulted in reductions in IFN γ production. Depletion of monocytes *in vivo* resulted in decreased IFN γ production by murine NK cells upon exposure to Ab-coated tumor cells. In mice receiving trastuzumab and IL12 therapy, monocyte depletion resulted in significantly greater tumor growth in comparison to mock-depleted controls ($P < 0.05$). These data suggest that NK cell–monocyte interactions enhance NK cell antitumor activity in the setting of monoclonal Ab therapy for cancer. *Cancer Immunol Res*; 5(9); 778–89. ©2017 AACR.

Introduction

Natural killer (NK) cells are innate immune cells that target and lyse virally infected cells as well as malignant cells. NK cells contain cytolytic granules and express an array of cellular adhesion molecules and cytokine receptors (1–4). They produce a variety of immunostimulatory cytokines, including IFN γ , TNF α , MIP-1 α , RANTES, and GM-CSF (5, 6). Constitutive expression of multiple cytokine receptors facilitates their response to inflammatory events (7, 8). NK cell effector functions are regulated by a

balance between activating and inhibitory receptors, as well as the expression of target cell MHC class I antigens (8). NK cells express an activating receptor for the Fc portion of immunoglobulin G (Fc γ RIIIa) that enables recognition of antibody (Ab)-coated targets. Support for the concept that antitumor activity of therapeutic monoclonal antibodies (mAb) is in part dependent on FcR-bearing cells stems from experiments conducted in FcR-deficient mice (9) and the observation that polymorphisms in the human FcR gene influence the clinical efficacy of therapeutic mAbs (10, 11). Studies from our group have shown that cytokine stimulation of NK cells enhances the immune response to Ab-coated tumor cells *in vitro*, in murine models, and in phase I clinical trials (12–15). Costimulation via the IL12 receptor (IL12R) and FcR has been shown to activate multiple cellular signaling pathways (e.g., JAK/STAT and PI3-kinase/Akt) and lead to maximal stimulation of NK cell effector functions (15).

NK cells express an activating, lectin-like receptor known as NKG2D (16). Human NKG2D binds to ligands MICA, MICB, and ULBPs that are expressed by a variety of tumor cell types (17, 18). Engagement of NKG2D on NK cells results in activation of the adapter protein DAP10, recruitment of the DAP10 binding partner Grb2 and signaling through Vav1, and also activation of PI3 kinase and the downstream serine threonine kinase Akt (19, 20). NK-cell interactions with NKG2D ligand-expressing tumor cells lead to NK cell–mediated cytotoxicity (21).

The role of reciprocal activation between monocytes and NK cells in immune responses is gaining increased recognition (22–25). NK cells function as one of the early sources of immunoregulatory cytokines (1). Upon activation, NK cells produce

¹The Ohio State University Comprehensive Cancer Center, Columbus, Ohio.

²Medical Scientist Training Program and Biomedical Sciences Graduate Program, The Ohio State University, Columbus, Ohio. ³Department of Surgery, The Ohio State University, Columbus, Ohio. ⁴Department of Pharmacy, The Ohio State University, Columbus, Ohio. ⁵Department of Molecular and Cell Biology, University of California, Berkeley, California. ⁶Division of Pulmonary, Allergy, Critical Care and Sleep Medicine, The Ohio State University, Columbus, Ohio. ⁷Clinical Research Division, Fred Hutchinson Cancer Research Center, Seattle, Washington. ⁸Center for Biostatistics, The Ohio State University, Columbus, Ohio. ⁹Division of Hematology, The Ohio State University, Columbus, Ohio.

Note: Supplementary data for this article are available at Cancer Immunology Research Online (<http://cancerimmunolres.aacrjournals.org/>).

A.R. Campbell and M.C. Duggan contributed equally to this article.

Corresponding Author: William E. Carson III, The Ohio State University, N924 Doan Hall, 410 West 12th Avenue, Columbus, OH 43210. Phone: 614-293-6306; Fax: 614-293-3465; E-mail: william.carson@osumc.edu

doi: 10.1158/2326-6066.CIR-16-0005

©2017 American Association for Cancer Research.

IFN γ , the prototypic macrophage activating factor, which optimizes monocyte/macrophage function (26, 27). Activated monocytes then produce cytokines like IL12 and IL15 that stimulate NK cells (28, 29). This positive feedback system supports the immune response against invading organisms and enables the rapid and effective clearance of infectious agents by monocytes/macrophages. We hypothesized that NK cell effector functions against Ab-coated tumor cells would be enhanced upon interactions with monocytes. We now report the ability of MICA-expressing monocytes to engage NKG2D on NK cells and promote NK-cell activity against Ab-coated tumor targets.

Materials and Methods

Cytokines and antibodies

Recombinant human IL12 (rhIL12) was provided by Genetics Institute Inc. Recombinant human IL15 and IL18 were purchased from Peprotech, Inc. The anti-HER2 mAb trastuzumab was provided by Genentech Inc. Antibodies utilized for flow cytometry include APC anti-hNKG2D (BD Biosciences, 558071), PE anti-hIFN γ (BD Biosciences, 559326), PE anti-hMICA (R&D Systems, FAB1300P), and PE anti-hMICA/B (BioLegend, 320906). The anti-MICA human F(ab')₂ Ab and human IgG F(ab')₂ control utilized for neutralization studies were generated as described (30). sMICA used in monocyte coculture experiments was generated as described (31).

Cell lines

The SK-BR-3 and MDA-MB-468 human breast adenocarcinoma cell lines were obtained from American Type Culture Collection. The C1R and C1R-MICA cell lines were provided by Dr. Veronika Groh. The identity of these cell lines was not authenticated. Cells were maintained in RPMI-1640 medium containing 10% FBS, 100 U/mL penicillin, 100 μ g/mL streptomycin, and C1R-MICA cultures were supplemented with G418 sulfate (Life Technologies Inc.). All cell lines used in this study were routinely tested for *mycoplasma* infection using the MycoAlert Mycoplasma Detection Kit (Lonza) and Plasmocin prophylactic (InvivoGen) was added to all growth media at a concentration of 2.5 μ g/mL. Cell lines were maintained in culture for no more than 10 passages.

Isolation of human NK cells and monocytes

PBMC and NK cells were isolated from donor leukopacks (American Red Cross, Columbus, OH) as described (15). CD14⁺ monocytes were isolated using anti-CD14 magnetic beads (Miltenyi Biotec, Auburn, CA). Immune cells were cultured in 10% HAB medium.

In vitro coculture with tumor cells

SK-BR-3 and MDA-MB-468 cells were cultured in 96-well plates. Cells were treated the following day with 100 μ g/mL trastuzumab for 1 hour. Immune cells were added to wells (2×10^5 NK cells/well) in 10% HAB medium supplemented with IL12 (10 ng/mL). Cocultures were plated at a 2:1 NK cell to monocyte/PBMC ratio. Supernatants were harvested and analyzed for cytokines by ELISA (R&D Systems) as described (12).

Intracellular staining for IFN γ

NK-cell IFN γ production was analyzed using an IFN γ PE-conjugated mAb and a CD56 APC-conjugated mAb as described (13).

Real-time PCR

Cellular RNA was isolated using RNeasy Mini-kits (Qiagen). cDNA was generated with random hexamer primers and MMLV-RT according to manufacturer's recommendations (Life Technologies Inc.). Real-time PCR for huIFN γ transcript was performed using an ABI prism 7700 sequence detector (Applied Biosystems) with an 18S rRNA internal control (PE Applied Biosystems).

Transwell cocultures

SK-BR-3 cells were cultured in 24-well plates and treated the next day with 100 μ g/mL trastuzumab for 1 hour. A Transwell insert with a 0.4- μ m filter was placed in each well, and NK cells and/or monocytes were added in media $\pm$ IL12 (10 ng/mL). Supernatants were analyzed for IFN γ content by ELISA.

MICA siRNA and monocyte transfection

Monocytes were transfected using the Amaxa Nucleofector apparatus (Amaxa Biosystems). Cells were resuspended in Nucleofector Solution T and nucleofected with scrambled or human MICA siRNA from Ambion. Cells were transferred to RPMI media supplemented with 10% FBS and M-CSF (20 ng/mL). Cy3-labeled GAPDH siRNA was used to analyze transfection efficiency.

Murine studies

Mice received two intraperitoneal (i.p.) injections of F4/80 Ab (250 μ g), isotype control Ab (250 μ g), or PBS. CT-26^{HER2/neu} cells were incubated in PBS plus 10% FBS with 4D5 (100 μ g/mL) (6). A total of 4×10^6 cells in PBS containing 1 μ g muIL12 were injected i.p. Serum was harvested at 24 hours and analyzed by ELISA. Splenocytes were collected to assess monocyte depletion by flow cytometry. For *ex vivo* studies, wild-type BALB/c splenocytes were cocultured with CT-26^{HER2/neu} tumor cells. Culture supernatants were analyzed for muIFN γ by ELISA. NKG2D knockout mice were provided by Dr. David Raulet.

For the murine tumor study, mice received i.p. injections of control or clodronate-containing liposomes (1 mg/kg in 100 μ L PBS) on days 0 and 4 with respect to tumor inoculation and every fourth day thereafter (32). On day 0, mice were inoculated with 8×10^5 EMT6^{HER2/neu} cells in the mammary fat pad (33). On day 7 and every third day, mice received i.p. injections of trastuzumab and IL12 (10 mg/kg and 2.5 ng, respectively). Tumor volume = $0.5 \times [(large\ diameter)^2 \times (small\ diameter)]$. Upon completion of the study, mice were sacrificed and tumors were collected. Cells were labeled with F4/80 PE-conjugated Ab and CD11b APC-conjugated Ab to evaluate monocyte depletion.

Statistical analysis

Statistical analyses of cytokine levels were performed using Student *t* tests. Changes in tumor volume over time were assessed via a longitudinal model. Tumor data was log transformed, and a linear mixed effects model was applied to account for correlations of observations from the same mouse.

Results

NK cells secrete immune stimulatory cytokines in response to IL12 and tumor cells

To investigate the ability of monocytes to enhance NK-cell interactions with Ab-coated tumor cells, we first evaluated NK-cell cytokine production and lytic activity *in vitro* in response to a stimulatory strategy: a therapeutic mAb coupled with the cytokine originally referred to as NK-cell stimulatory factor, IL12. The

HER2 overexpressing breast cancer cell line SK-BR-3 or the HER2 negative MDA-MB-468 cell line were cocultured with NK cells in the presence or absence of trastuzumab and IL12. NK-cell production of IFN γ in response to trastuzumab-coated SK-BR-3 cells was enhanced in the presence of IL12, as compared to control conditions; however, this was not observed with the HER2-negative cell line (Fig. 1A). The capacity of total peripheral blood mononuclear cells (PBMC) to respond to IL12 and Ab-coated tumor cells was also tested. PBMC (plated at the same cell density as pure NK cells) secreted more IFN γ compared to NK cells alone in response to dual stimulation via Ab-coated tumor cells and IL12. This relationship held true for other NK cell-derived cytokines including TNF α and MIP-1 α (Fig. 1B). On average, PBMC IFN γ production was 200% higher than that of pure NK cells. The number of NK cells within the PBMC population added to each well was approximately 4-fold less than the number of pure NK

cells plated (2×10^5 cells per well). Next, PBMC IFN γ production in response to various stimuli (e.g., IL1 β , IL12, IL15, IL18, and/or trastuzumab-coated cells) was compared to that of NK cells. PBMC IFN γ production was greater than that of purified NK cells for all conditions tested (Fig. 1C). The most potent individual cytokine stimulus for NK-cell IFN γ production in response to Ab-coated tumor cells was IL12. Subsequent investigation sought to uncover which cellular compartment within total PBMC could be responsible for providing a stimulatory signal to increase NK cell antitumor activity in the presence of what was considered to be a strong stimulatory strategy, namely, FcR and IL12R coactivation.

Monocytes within a PBMC population enhance the NK-cell cytokine response
The impact of different immune cell populations on the NK-cell IFN γ response was investigated. Depletion of monocytes from

Figure 1. Cytokine production is enhanced in the presence of PBMC. **A**, NK cells were cocultured with SK-BR-3 or MDA-MB-468 tumor cells in the presence of medium, rhuIL12 (IL12), trastuzumab (Tras), or the combination. **B**, NK cells or PBMC were cultured in medium alone or with trastuzumab-coated SK-BR-3 cells and IL12. **C**, NK cells or PBMCs were cultured in the presence of rhuIL1 β , IL12, IL15, or IL18 and/or trastuzumab-coated SK-BR-3 cells. Bars represent the mean concentration $\pm$ SD of cytokine (IFN γ for **A**, **B**, **C**, MIP-1 α for **B**, or TNF α for **B**) content from culture supernatants analyzed by ELISA. Data are representative of three independent experiments with similar results (**A**) or two experiments (**B** and **C**). *, $P < 0.05$.

PBMCs, but not B cells or T cells, resulted in decreased IFN γ production following exposure to IL12 and trastuzumab-coated tumor cells (Fig. 2A, Supplementary Fig. S1). The ability of monocyte-depleted PBMC to induce IFN γ production in response to IL12 and Ab-coated tumor cells was restored in a dose-dependent fashion by the reintroduction of autologous monocytes (Fig. 2B). To confirm the effect of monocytes on NK-cell

cytokine production, NK cells were cocultured with purified autologous monocytes. NK cells cocultured with monocytes in the presence of Ab-coated tumor cells and IL12 produced more IFN γ than NK cells alone (Fig. 2C). Monocytes produced no measureable amounts of IFN γ . Similar results were obtained with IL15 as the cytokine stimulus (Fig. 2D). Coculture with suppressive monocyte-derived cells such as tumor-associated

Figure 2.

Monocytes promote NK-cell IFN γ production. **A**, PBMC (~14% CD14⁺ cells), PBMC depleted of monocytes (<0.03% CD14⁺ cells), or PBMC depleted of B cells (>98% CD19⁺ cells) were cocultured with SK-BR-3 cells in medium, IL12, trastuzumab, or the combination. **B**, Monocyte-depleted PBMCs were mixed with increasing numbers of autologous monocytes and cultured with SK-BR-3 cells in the presence of medium, IL12, trastuzumab, or the combination. **C**, NK cells, or NK cells plus monocytes were cultured with SK-BR-3 cells in the presence of medium, IL12, trastuzumab, or the combination. **D**, NK cells were mixed with monocytes and cocultured with SK-BR-3 cells in the presence of medium, IL15, trastuzumab, or the combination. **E**, NK cells were cultured in the presence of trastuzumab-coated SK-BR-3 cells and IL12, and monocytes and/or *in vitro*-generated autologous TAM or MDSC at a 2:1 ratio were added to the coculture (47, 48). **F**, NK cells were cultured with varying numbers of autologous monocytes in the presence of trastuzumab-coated SK-BR-3 cells and IL12. **G**, NK cells alone or in the presence of monocytes were cocultured with trastuzumab-coated SK-BR-3 cells and IL12. Cells were harvested and intracellular IFN γ was examined by flow cytometry. **H**, NK cells, monocytes, or both were cultured in the presence of trastuzumab-coated SK-BR-3 cells and IL12. Bars represent the mean concentration $\pm$ SD of IFN γ content from culture supernatants analyzed by ELISA (**A-F**) or IFN γ transcript as measured by PCR (**H**). Data shown are representative of two (**A**, **B**, **D**, **F**, **G**) or three (**E**) independent experiments with similar results, or the mean concentration $\pm$ SD of $n = 8$ donors (**C**). *, $P < 0.05$.

macrophages (TAM) or myeloid-derived suppressor cells (MDSC) generated *in vitro* from autologous monocytes led to a reduction in NK-cell IFN γ production in this system (Fig. 2E). Thus, it appeared that monocytes, although not the direct source of IFN γ , were important for enhancement of NK-cell IFN γ production. The effect of monocyte addition was dose dependent, as the introduction of increasing numbers of monocytes led to NK-cell IFN γ production that was equivalent to the IFN γ output of unmanipulated PBMC (Fig. 2F). The effect of monocytes on NK-cell IFN γ production was confirmed by intracellular flow cytometry (Fig. 2G). Also, IFN γ transcript levels were evaluated in dual-stimulated NK cells in the presence and absence of monocytes. Addition of monocytes to coculture enhanced NK-cell IFN γ transcript levels (Fig. 2H). These results suggest that monocytes within the PBMC population are capable of enhancing NK-cell cytokine production in response to Ab-coated tumor cells and IL12.

Monocytes enhance the lytic activity of NK cells

The effect of monocytes on the ability of IL12-activated NK cells to lyse Ab-coated tumor cells was examined (Fig. 3). At the 50:1 effector to target ratio, IL12-activated NK cells mediated approximately 48% lysis of trastuzumab-coated SK-BR-3 target cells (Fig. 3, top). Monocytes alone exhibited negligible cytotoxic activity (Fig. 3, middle). It has been demonstrated that monocytes are capable of ADCC 8 to 14 hours after priming with IFN γ ; however, this was not observed in our standard 4-hour chromium release assay conditions. Exposure of NK cells to monocytes (Fig. 3, bottom) prior to the addition of Ab-coated target cells and IL12 facilitated an increase in NK cell-mediated cytotoxicity to nearly 98% at the 50:1 effector to target ratio ($P < 0.05$).

NK cell activation is enhanced by direct contact between NK cells and monocytes

Increased NK-cell IFN γ production upon interaction with monocytes could be due to direct cell-to-cell contact or the diffusion of soluble mediators. To determine whether cell-to-cell contact was required for stimulation, a Transwell coculture experiment was performed. When NK cells and monocytes were separated by a cell-impermeable membrane, NK cells exposed to trastuzumab-coated tumor cells and IL12 produced normal amounts of IFN γ . However, coculture of NK cells and monocytes in the same compartment (permitting contact) led to significantly greater IFN γ production (Fig. 4A, $P = 0.03$). Figure 4B demonstrates the same experiment with a full set of controls, which was challenging to perform due to the requirement for a large number of NK cells from a single donor. The effect of monocytes on NK-cell cytokine production in Fig. 4B, although substantial and reproducible, was less than in previous experiments possibly due to the physical characteristics of larger Transwell cultures. In summary, monocyte enhancement of NK-cell IFN γ production in response to Ab-coated tumor cells and IL12 requires cell-to-cell contact, and this effect is lost when monocytes are separated from NK cells by a cell-impermeable membrane.

Ligand interactions essential for monocyte enhancement of NK-cell IFN γ production

Bauer and colleagues reported that interactions between NK cell NKG2D and its ligands MICA/B or ULBPs, expressed by tumor cells, enhances NK cell cytokine production (18). However, NKG2D ligands may also be expressed by cells of the

Figure 3.

Monocytes enhance NK cell ADCC. NK cells, monocytes, or both were incubated overnight in medium alone or medium containing IL12. NK cell ADCC was assessed in a standard 4-hour chromium release assay with trastuzumab-coated SK-BR-3 target cells. Symbols represent target cell mean percent lysis $\pm$ SD for three independent experiments.

innate immune system (34, 35). Thus, we hypothesized that monocytes could activate NK cells via expression of NKG2D ligands. Monocyte MICA surface expression was measured by flow cytometry. Though variability exists in NKG2D ligand expression among healthy donors, substantial MICA expression was detected in freshly isolated monocytes (Supplementary Fig. S2). Further, it was shown that MICA expression could be upregulated following stimulation of monocytes with IFN α or LPS (Fig. 5A). LPS treatment of monocytes enhanced their ability to augment NK-cell IFN γ production (Fig. 5B). To further explore the interaction between NKG2D and MICA, the MICA overexpressing cell line C1R-MICA was used and was confirmed to selectively overexpress MICA but not other NKG2D ligands (Supplementary Fig. S3). Coculture of NK cells with C1R-MICA cells in the presence of trastuzumab-coated SK-BR-3 cells and IL12 led to increased NK-cell IFN γ production that was comparable to that observed with the addition of

monocytes. Coculture of NK cells with the parental cell line C1R (which lacks MICA expression) did not modify NK-cell IFN γ production (Fig. 5C). Introduction of an anti-MICA-neutralizing Ab abrogated the ability of C1R-MICA cells to enhance NK-cell IFN γ production (Fig. 5D).

Experiments involving an anti-MICA-neutralizing Ab were difficult to optimize with donor monocytes and NK cells due to FcR expression. An F(ab')₂ against MICA was designed and utilized in an attempt to block NKG2D-MICA interactions; however, unintended NK cell activation persisted. An anti-MICA siRNA was employed to study the interaction between NK cell NKG2D and monocyte MICA. MICA expression levels were downregulated in monocytes using this siRNA approach. Reduced MICA expression at the transcript level and on the surface of monocytes was confirmed by real-time PCR and flow cytometry (Fig. 5E and F). Down regulation of MICA impaired the ability of monocytes to augment NK-cell cytokine production and led to reduction in IFN γ levels in response to Ab-

coated tumor cells and IL12 (Fig. 5G). The interaction between NK-cell NKG2D and monocyte MICA was also disrupted by pretreating NK cells with soluble MICA (sMICA) to block NKG2D receptor availability prior to coculture with monocytes (31). This strategy resulted in a dose-dependent decrease in IFN γ production, supporting the hypothesis that direct NKG2D-MICA interactions between NK cells and monocytes promote NK-cell activity (Fig. 5H). No correlation was found between MICA expression levels on donor monocytes and IFN γ production from NK cells, indicating that although MICA expression on monocytes is necessary for enhanced NK cell activation, the level to which MICA is expressed is not as important (Supplementary Fig. S4).

Modulation of NKG2D expression on NK cells

It was confirmed that NK cells express NKG2D, and expression may be upregulated upon NK-cell stimulation via IL12 or IL15 (Fig. 6A; ref. 36). Pretreatment of NK cells with IL15 (followed by

Figure 5.

Monocyte expression of the NKG2D ligand MICA promotes NK cell antitumor activity. **A**, Monocyte MICA expression at baseline (medium) and following treatment with LPS or IFN α was evaluated by flow cytometry. **B**, Untreated and LPS-treated monocytes were cocultured with NK cells, trastuzumab-coated SK-BR-3 cells, and IL12. **C**, NK cells were cocultured with C1R or C1R-MICA cells in the presence of trastuzumab-coated SK-BR-3 cells and IL12. **D**, C1R-MICA cells were incubated with control IgG or anti-MICA Ab. C1R-MICA cells then were cocultured with NK cells, trastuzumab-coated SK-BR-3 cells, and IL12. **E**, Monocytes were transfected with MICA siRNA or scrambled control siRNA. Transfected cells were analyzed for MICA transcript by real-time PCR or (F) MICA/B surface expression was examined by flow cytometry. **G**, Monocytes with downregulated levels of MICA were cocultured with NK cells in the presence of trastuzumab-coated SK-BR-3 cells and IL12. **H**, NK cells were pretreated with soluble MICA (sMICA) prior to coculture with monocytes in the presence of trastuzumab-coated SK-BR-3 cells and IL12. Bars represent the mean concentration $\pm$ SD of IFN γ content from supernatants analyzed by ELISA (**B**, **C**, **G**, **H**). Data shown for each panel are representative of two (**B**, **C**, **G**), three (**A**, **H**) independent experiments with similar results or the mean concentration $\pm$ SD of $n = 3$ donors (**D**). *, $P < 0.05$; ***, $P < 0.001$; ns, not significant.

an overnight rest) increased NKG2D expression and enhanced NK-cell IFN γ production in the presence of monocytes (Fig. 6B). Additionally, NK cell NKG2D expression was not decreased following interactions with MICA expressing cells *in vitro* (Supplementary Fig. S5). Further, an anti-NKG2D-neutralizing Ab was used to block activation of this receptor (37). NKG2D blockade reduced NK-cell IFN γ production in response to Ab-coated tumor

cells and IL12 in the presence of C1R-MICA cells (Fig. 6C) as well as monocytes (Fig. 6D).

Murine NK-cell IFN γ production is enhanced in the presence of monocytes

Next, experiments were conducted using murine models. Though mice do not express MIC-derived molecules, they do

Figure 6.

NK cells interact with MICA on monocytes via the NKG2D receptor. **A**, NK cell NKG2D expression was evaluated at baseline (medium) and following stimulation via IL12 or IL15. **B**, NK cells were treated with IL15, rested overnight, then added to coculture with monocytes, trastuzumab-coated SK-BR-3 cells, and IL12. **C**, NK cells were incubated with a control IgG or anti-NKG2D Ab. NK cells then were cultured in the presence of trastuzumab-coated SK-BR-3 cells, IL12, and C1R-MICA cells, or (D) monocytes. Bars represent the mean concentration $\pm$ SD of IFN γ content from supernatants analyzed by ELISA. Data shown for each panel are representative of two independent experiments with similar results.

express similar MHC class I-like molecules, including RAE-1, H60, and MULT-1, that engage murine NKG2D and promote NK-cell activation (38). Guerra and colleagues showed in murine models that tumor immunosurveillance is inhibited in NKG2D-deficient mice (39). Given that *in vitro* experiments in our laboratory demonstrated a relationship between NK cells and monocytes, we sought to evaluate whether interruption of this interaction would impact the NK-cell response to mAb therapy in murine models as well.

Splenocytes from wild-type mice were cultured in the presence of IL12 and 4D5 (a murine anti-HER2 mAb)-coated HER2-positive murine tumor cells. Splenocytes depleted of NK cells produced less IFN γ , identifying NK cells as the source of IFN γ , as described by our group previously (Fig. 7A, ref. 6). Clodronate-containing liposomes were used as a means of depleting mice of monocytes *in vivo*. Splenocytes isolated from mice treated with control liposomes retained normal monocyte numbers and produced IFN γ *ex vivo* in response to IL12- and 4D5-coated tumor cells. In contrast, treatment with clodronate-containing liposomes led to a 60% reduction in monocyte numbers and a

reduction in NK-cell IFN γ production (Fig. 7B). An F4/80 antibody was used as an additional strategy to deplete monocytes. Control mice treated with an isotype Ab retained normal monocyte numbers, and the amount of IFN γ in the serum increased in response to i.p. administration of IL12 and 4D5-coated tumor cells. However, treatment with an F4/80 Ab led to a 61% reduction in monocytes in the spleen and resulted in a 2-fold reduction in serum IFN γ levels in response to dual stimulation (Fig. 7C). Next, mice rendered genetically deficient in NKG2D were used. Splenocytes were harvested from wild-type and NKG2D knockout mice and cultured in the presence of murine IL12- and 4D5-coated CT26^{HER2/neu} tumor cells. IFN γ production was inhibited in mice lacking the NKG2D receptor (Fig. 7D). Lastly, a study was conducted in which mice bearing HER2-positive tumors were treated with trastuzumab and IL12 along with systemic administration of clodronate-containing liposomes or control liposomes. Treatment of mice with clodronate-containing liposomes led to significantly larger tumors as compared to mice receiving control liposomes ($P < 0.05$; Fig. 7E).

Figure 7.

Monocytes enhance NK cell function *in vivo*. **A**, Murine splenocytes were depleted of NK cells and cocultured with CT26^{HER2/neu} tumor cells in the presence of medium, IL12, 4D5, or the combination. Bars represent the mean concentration $\pm$ SD of IFN γ content from supernatants analyzed by ELISA, $n = 2$ mice. **B**, Splenocytes from wild-type BALB/c mice treated with clodronate-containing or control liposomes were cocultured with 4D5-coated CT26^{HER2/neu} tumor cells and IL12. Bars represent the mean concentration $\pm$ SD of IFN γ content, $n = 2$ mice. **C**, BALB/c mice were treated with PBS, an isotype control Ab, or an F4/80 Ab and were injected i.p. with 4D5-coated CT26^{HER2/neu} tumor cells and IL12. Serum was collected 24 hours later and analyzed for IFN γ content by ELISA. The means $\pm$ SD are shown ($n = 5$ for each condition). **D**, Splenocytes were isolated from wild-type (WT) and NKG2D-deficient mice (NKG2D K.O.) and cocultured with 4D5-coated CT26^{HER2/neu} tumor cells and IL12. Bars represent the mean concentration $\pm$ SD of IFN γ content, $n = 3$. **E**, Trastuzumab and IL12 were administered to EMT6^{HER2/neu} tumor-bearing mice that were treated with clodronate-containing or control liposomes. Average tumor volume $\pm$ SEM is shown ($n = 5$ mice for each group). *, $P < 0.05$; **, $P < 0.02$; ***, $P < 0.001$.

Discussion

Here, we investigated the role of NK cell–monocyte interactions in the setting of mAb therapy for cancer. We have demonstrated previously that NK cell-derived IFN γ serves as a key mediator of the immune response to cancer. NK cells release IFN γ following costimulation via Ab-coated tumor cells and IL12, a therapeutic strategy proven to elicit NK cell activity *in vitro* and *in vivo*. Now, it has been shown that monocytes enhance NK-cell IFN γ production in response to a stimulus. This stimulatory interaction requires direct cell-to-cell contact between NK cells and monocytes and is mediated primarily by monocyte MICA engagement of the NKG2D receptor on NK cells. Lastly, it was demonstrated in murine models that the immune response to combination therapy with trastuzumab and IL12 is enhanced through NK cell–monocyte interactions. Taken together, these results suggest that NK cells and monocytes in the tumor microenvironment participate in crosstalk to optimize Ab-mediated antitumor activity, and mechanisms to promote these interactions may enhance the response to tumor-specific mAbs.

Interactions between NK cells and monocytes promote the immune response against invading pathogens and transformed cancer cells. However, the role of monocytes in modulating NK-cell activity in the setting of mAb therapy is not well understood. We noted that exposure of PBMC to various NK stimulatory conditions led to more IFN γ production than that of NK cells alone. Monocytes were the cells within the PBMC population responsible for this boost in NK-cell cytokine production. Others have demonstrated that reciprocal activation between NK cells and monocytes occurs via contact in the setting of infection as well as chronic arthritis (40, 41). However, the molecules responsible for this cross-talk were not determined. In the present study, a key interface between NK cells and monocytes, the NKG2D-MICA axis, was shown to be responsible for monocyte-mediated enhancement of NK cell antitumor activity. These results suggest that NK-cell NKG2D and monocyte MICA interactions play an important role in the immune response to mAb therapy.

Few studies have delved into the relationship between NK cells and monocytes in regard to cell-mediated cytotoxicity against tumor cells (42). We now show that interactions with monocytes promote NK cell-mediated cytotoxic activity. Bhatnagar and colleagues (43) described reciprocal cross-talk between NK cells and monocytes in the setting of viral infection, concluding that monocytes enhanced NK cell cytotoxicity against HIV-1–infected cells via direct contact in an Fc γ RII (CD32)–dependent manner. The role of NK cell recognition and lysis of NKG2D-ligand expressing tumor cells has been described previously (44). Pende and colleagues analyzed tumor expression levels of the NKG2D ligands MICA and ULBPs in melanoma, leukemia, neuroblastoma, and various carcinoma cell lines and correlated expression levels to increased tumor cell susceptibility to NK cell-mediated lysis. The impact of NKG2D ligand expression by circulating immune cells was not explored. Wang and colleagues evaluated IFN γ pretreated monocytes and determined that this strategy induced monocyte expression of MIC molecules and membrane-bound IL15 (45). These molecules promoted NK-cell IFN γ production and cytotoxicity against lymphoblastoid cells. This was observed only with IFN γ pretreated monocytes *in vitro*, and NK-cell activity was not enhanced in the presence of freshly isolated monocytes, as was the case in the present study. Also, Wang and colleagues utilized allogeneic NK cell–monocyte cocultures at a

5:1 NK cell to monocyte ratio, whereas the current study investigated autologous NK cells and monocytes cultured at a 2:1 ratio, modeling what might be found *in vivo*. Kloss and colleagues demonstrated that MICA-positive, LPS-activated monocytes enhanced NK cell cytokine production *in vitro*. The presence of monocytes led to increased NK cell proliferation, regardless of whether monocytes were untreated or primed with LPS (35). The current study demonstrates that in the setting of mAb therapy for cancer, interactions between MICA-positive monocytes and NKG2D-expressing NK cells facilitate NK-cell cytokine production and ADCC against Ab-coated tumor cells both *in vitro* and in murine models.

A recent study by Crane and colleagues described that myeloid cells, including monocytes isolated from glioblastoma multiforme (GBM) patients, expressed NKG2D ligands (46). It was shown that glioma cell line–derived lactate dehydrogenase-containing supernatants had the capacity to upregulate monocyte NKG2D ligand expression and promote NK-cell degranulation and cytokine production *in vitro*. The authors proposed that immune evasion in GBM patients may be due to NKG2D-dependent NK cell exhaustion. In contrast, this study demonstrated that interruption of NK cell–monocyte cross-talk *in vivo* inhibited the NK cell effector response against Ab-coated breast cancer cells. These contrasting results emphasize the need to further explore the possible effects of NKG2D ligand expression by myeloid cells.

This study has provided evidence that monocytes promote NK-cell cytokine production and tumor cell lysis in response to FcR and IL12R stimulation, an effective dual stimulation regimen with proven antitumor activity. The antitumor effects of mAb therapy may be enhanced by strategies aimed at maximizing monocyte expression of NK-cell activating molecules.

Disclosure of Potential Conflicts of Interest

D. Raulet is a consultant/advisory board member for Innate Pharma SAS. No potential conflicts of interest were disclosed by the other authors.

Authors' Contributions

Conception and design: A.R. Campbell, L.P. Suarez-Kelly, N. Bhavé, R. Lee, J.C. Byrd, W.E. Carson III

Development of methodology: A.R. Campbell, L.P. Suarez-Kelly, N. Bhavé, P. Trikha, R. Parihar, R. Lee, V. Groh, W.E. Carson III

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): A.R. Campbell, M.C. Duggan, L.P. Suarez-Kelly, N. Bhavé, E.L. McMichael, P. Trikha, R. Parihar, E. Luedke, A. Lewis, R. Lee, D. Raulet, S. Tridandapani, J.C. Byrd

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): A.R. Campbell, M.C. Duggan, L.P. Suarez-Kelly, N. Bhavé, K.S. Opheim, P. Trikha, R. Lee, S. Tridandapani, V. Groh, L. Yu, V. Yildiz, W.E. Carson III

Writing, review, and/or revision of the manuscript: A.R. Campbell, M.C. Duggan, L.P. Suarez-Kelly, N. Bhavé, K.S. Opheim, R. Lee, V. Groh, V. Yildiz, J.C. Byrd, M.A. Caligiuri, W.E. Carson III

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): M.C. Duggan, B. Yung

Study supervision: W.E. Carson III

Other (provided critical research materials): V. Groh

Acknowledgments

The authors thank The Ohio State University Comprehensive Cancer Center's Analytical Cytometry Shared Resource for providing equipment to facilitate flow cytometry analyses.

Grant Support

This work was supported by the Susan G. Komen Breast Cancer Foundation Dissertation Research Award (R. Parihar), the OSUCCC Pelotonia Graduate Fellowship (A.R. Campbell), and the NIH grants T32 GM068412 (M.C. Duggan) and P01 CA95426 (W.E. Carson).

References

- Bhave NS, Carson WE. The role of natural killer cells in innate immunity. Regulation of Innate Immune Function Editors: Clay Marsh, Susheela Tridandapani and Melissa Piper 2010:141–64.
- Caligiuri MA. Human natural killer cells. Blood 2008;112:461–9.
- Carson WE, Fehniger TA, Haldar S, Eckhert K, Lindemann MJ, Lai CF, et al. A potential role for interleukin-15 in the regulation of human natural killer cell survival. J Clin Invest 1997;99:937–43.
- Carson WE, Parihar R, Lindemann MJ, Personeni N, Dierksheide J, Meropol NJ, et al. Interleukin-2 enhances the natural killer cell response to Herceptin-coated Her2/neu-positive breast cancer cells. Eur J Immunol 2001;31:3016–25.
- Badgwell B, Parihar R, Magro C, Dierksheide J, Russo T, Carson WE 3rd. Natural killer cells contribute to the lethality of a murine model of Escherichia coli infection. Surgery 2002;132:205–12.
- Jaime-Ramirez AC, Mundy-Bosse BL, Kondadasula S, Jones NB, Roda JM, Mani A, et al. IL-12 enhances the antitumor actions of trastuzumab via NK-cell IFN- γ production. J Immunol 2011;186:3401–9.
- Carson WE, Ross ME, Baiocchi RA, Marien MJ, Boiani N, Grabstein K, et al. Endogenous production of interleukin 15 by activated human monocytes is critical for optimal production of interferon-gamma by natural killer cells in vitro. J Clin Invest 1995;96:2578–82.
- Vivier E, Nunès JA, Vély F. Natural killer cell signaling pathways. Science 2004;306:1517–9.
- Clynes RA, Towers TL, Presta LG, Ravetch JV. Inhibitory Fc receptors modulate in vivo cytotoxicity against tumor targets. Nat Med 2000;6:443–6.
- Carton G. Therapeutic activity of humanized anti-CD20 monoclonal antibody and polymorphism in IgG Fc receptor Fc γ RIIIa gene. Blood 2002;99:754–8.
- Weng WK, Levy R. Two immunoglobulin G fragment C receptor polymorphisms independently predict response to rituximab in patients with follicular lymphoma. J Clin Oncol 2003;21:3940–7.
- Parihar R, Dierksheide J, Hu Y, Carson WE. IL-12 enhances the natural killer cell cytokine response to Ab-coated tumor cells. J Clin Invest 2002;110:983–92.
- Parihar R, Nadella P, Lewis A, Jensen R, De Hoff C, Dierksheide JE, et al. A phase I study of interleukin 12 with trastuzumab in patients with human epidermal growth factor receptor-2-overexpressing malignancies. Clin Cancer Res 2004;10:5027–37.
- Roda JM, Parihar R, Magro C, Nuovo GJ, Tridandapani S, Carson WE 3rd. Natural killer cells produce T cell-recruiting chemokines in response to antibody-coated tumor cells. Cancer Res 2006;66:517–26.
- Kondadasula SV, Roda JM, Parihar R, Yu J, Lehman A, Caligiuri MA, et al. Colocalization of the IL-12 receptor and Fc γ RIIIa to natural killer cell lipid rafts leads to activation of ERK and enhanced production of interferon-gamma. Blood 2008;111:4173–83.
- Lanier LL. Up on the tightrope: natural killer cell activation and inhibition. Nat Immunol 2008;9:495–502.
- Groh V, Bahram S, Bauer S, Herman A, Beauchamp M, Spies T. Cell stress-regulated human major histocompatibility complex class I gene expressed in gastrointestinal epithelium. Proc Natl Acad Sci 1996;93:12445–50.
- Groh V, Groh V, Wu J, Steinle A, Phillips JH, Lanier LL, et al. Activation of NK cells and T cells by NKG2D, a receptor for stress-inducible MICA. Science 1999;285:727–9.
- Upshaw JL, Ameson LN, Schoon RA, Dick CJ, Billadeau DD, Leibson PJ. NKG2D-mediated signaling requires a DAP10-bound Grb2-Vav1 intermediate and phosphatidylinositol-3-kinase in human natural killer cells. Nat Immunol 2006;7:524–32.
- Wu J, Song Y, Bakker AB, Bauer S, Spies T, Lanier LL, et al. An activating immunoreceptor complex formed by NKG2D and DAP10. Science 1999;285:730–2.
- Cerwenka A, Lanier LL. Ligands for natural killer cell receptors: redundancy or specificity. Immunol Rev 2001;181:158–69.
- Amakata Y, Fujiyama Y, Andoh A, Hodohara K, Bamba T. Mechanism of NK cell activation induced by coculture with dendritic cells derived from peripheral blood monocytes. Clin Exp Immunol 2001;124:214–22.
- Fernandez NC, Flament C, Crepineau F, Angevin E, Vivier E, Zitvogel L. Dendritic cells (DC) promote natural killer (NK) cell functions: dynamics of the human DC/NK cell cross talk. Eur Cytokine Netw 2002;13:17–27.
- Cooper MA, Fehniger TA, Fuchs A, Colonna M, Caligiuri MA. NK cell and DC interactions. Trends Immunol 2004;25:47–52.
- Della Chiesa M, Romagnani C, Thiel A, Moretta L, Moretta A. Multidirectional interactions are bridging human NK cells with plasmacytoid and monocyte-derived dendritic cells during innate immune responses. Blood 2006;108:3851–8.
- Cooper MA, Fehniger TA, Turner SC, Chen KS, Ghaheri BA, Ghayur T, et al. Human natural killer cells: a unique innate immunoregulatory role for the CD56(bright) subset. Blood 2001;97:3146–51.
- DeMarco RA, Fink MP, Lotze MT. Monocytes promote natural killer cell interferon gamma production in response to the endogenous danger signal HMGB1. Mol Immunol 2005;42:433–44.
- Carson WE, Giri JG, Lindemann MJ, Linett ML, Ahdieh M, Paxton R, et al. Interleukin (IL) 15 is a novel cytokine that activates human natural killer cells via components of the IL-2 receptor. J Exp Med 1994;180:1395–403.
- Broudy VC, Kaushansky K, Segal GM, Harlan JM, Adamson JW. Tumor necrosis factor type alpha stimulates human endothelial cells to produce granulocyte/macrophage colony-stimulating factor. Proc Natl Acad Sci 1986;83:7467–71.
- Groh V, Steinle A, Bauer S, Spies T. Recognition of stress-induced MHC molecules by intestinal epithelial $\gamma\delta$ T cells. Science 1998;279:1737–40.
- Groh V, Wu J, Yee C, Spies T. Tumour-derived soluble MIC ligands impair expression of NKG2D and T-cell activation. Nature 2002;419:734–8.
- Bacci M, Capobianco A, Monno A, Cottone L, Di Puccio F, Camisa B, et al. Macrophages are alternatively activated in patients with endometriosis and required for growth and vascularization of lesions in a mouse model of disease. Am J Pathol 2009;175:547–56.
- Cho HM, Rosenblatt JD, Tolba K, Shin SJ, Shin DS, Calfa C, et al. Delivery of NKG2D ligand using an anti-HER2 antibody-NKG2D ligand fusion protein results in an enhanced innate and adaptive antitumor response. Cancer Res 2010;70:10121–30.
- Zwirner NW, Fernández-Viña MA, Stastny P. MICA, a new polymorphic HLA-related antigen, is expressed mainly by keratinocytes, endothelial cells, and monocytes. Immunogenetics 1998;47:139–48.
- Kloss M, Decker P, Baltz KM, Baessler T, Jung C, Rammensee HG, et al. Interaction of monocytes with NK cells upon Toll-like receptor-induced expression of the NKG2D ligand MICA. J Immunol 2008;181:6711–9.
- Roberts AL, Lee L, Schwarz E, Groh V, Spies T, Ebert EC, et al. NKG2D receptors induced by IL-15 costimulate CD28-negative effector CTL in the tissue microenvironment. J Immunol 2001;167:5527–30.
- Kwong KY, Baskar S, Zhang H, Mackall CL, Rader C. Generation, affinity maturation, and characterization of a human anti-human NKG2D monoclonal antibody with dual antagonistic and agonistic activity. J Mol Biol 2008;384:1143–56.
- Zafirova B, Mandarić S, Antulov R, Krmpotić A, Jonsson H, Yokoyama WM, et al. Altered NK cell development and enhanced NK cell-mediated resistance to mouse cytomegalovirus in NKG2D-deficient mice. Immunity 2009;31:270–82.
- Guerra N, Tan YX, Joncker NT, Choy A, Gallardo F, Xiong N, et al. NKG2D-deficient mice are defective in tumor surveillance in models of spontaneous malignancy. Immunity 2008;28:571–80.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received January 19, 2016; revised July 20, 2016; accepted July 5, 2017; published OnlineFirst July 19, 2017.

40. Newman KC, Korbel DS, Hafalla JC, Riley EM. Cross-talk with myeloid accessory cells regulates human natural killer cell interferon-gamma responses to malaria. *PLoS Pathog* 2006;2:e118.
41. Dalbeth N, Gundle R, Davies RJ, Lee YC, McMichael AJ, Callan MF. CD56bright NK cells are enriched at inflammatory sites and can engage with monocytes in a reciprocal program of activation. *J Immunol* 2004; 173:6418–26.
42. Deguine J, Breart B, Lemaitre F, Bousso P. Cutting edge: tumor-targeting antibodies enhance NKG2D-mediated NK cell cytotoxicity by stabilizing NK cell-tumor cell interactions. *J Immunol* 2012;189:5493–7.
43. Bhatnagar N, Ahmad F, Hong HS, Eberhard J, Lu IN, Ballmaier M, et al. FcγRIII (CD16)-mediated ADCC by NK cells is regulated by monocytes and FcγRII (CD32). *Eur J Immunol* 2014;44:3368–79.
44. Pende D, Rivera P, Marcenaro S, Chang CC, Biassoni R, Conte R, et al. Major histocompatibility complex class I-related chain A and UL16-binding protein expression on tumor cell lines of different histotypes: analysis of tumor susceptibility to NKG2D-dependent natural killer cell cytotoxicity. *Cancer Res* 2002;62: 6178–86.
45. Wang H, Ruan Z, Wang Y, Han J, Fu X, Zhao T, et al. MHC class I chain-related molecules induced on monocytes by IFN-gamma promote NK cell activation. *Mol Immunol* 2008;45:1548–56.
46. Crane CA, Austgen K, Haberthur K, Hofmann C, Moyes KW, Avanesyan L, et al. Immune evasion mediated by tumor-derived lactate dehydrogenase induction of NKG2D ligands on myeloid cells in glioblastoma patients. *Proc Natl Acad Sci USA* 2014;111:12823–8.
47. Lechner MG, Liebertz DJ, Epstein AL. Characterization of cytokine-induced myeloid-derived suppressor cells from normal human peripheral blood mononuclear cells. *J Immunol* 2010;185:2273–84.
48. Rey-Giraud F, Hafner M, Ries CH. In vitro generation of monocyte-derived macrophages under serum-free conditions improves their tumor promoting functions. *PLoS One* 2012;7:e42656.

Cancer Immunology Research

MICA-Expressing Monocytes Enhance Natural Killer Cell Fc Receptor-Mediated Antitumor Functions

Amanda R. Campbell, Megan C. Duggan, Lorena P. Suarez-Kelly, et al.

Cancer Immunol Res 2017;5:778-789. Published OnlineFirst July 19, 2017.

Updated version Access the most recent version of this article at:
doi:[10.1158/2326-6066.CIR-16-0005](https://doi.org/10.1158/2326-6066.CIR-16-0005)

Supplementary Material Access the most recent supplemental material at:
<http://cancerimmunolres.aacrjournals.org/content/suppl/2017/07/18/2326-6066.CIR-16-0005.DC1>

Cited articles This article cites 47 articles, 24 of which you can access for free at:
<http://cancerimmunolres.aacrjournals.org/content/5/9/778.full#ref-list-1>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, contact the AACR Publications Department at permissions@aacr.org.