

UC Santa Cruz

UC Santa Cruz Previously Published Works

Title

Probing the Activity Differences of Simple and Complex Brominated Aryl Compounds against 15-Soybean, 15-Human, and 12-Human Lipoxygenase §

Permalink

<https://escholarship.org/uc/item/4rj5c9kd>

Journal

Journal of Medicinal Chemistry, 47(16)

ISSN

0022-2623

Authors

Segraves, Erika N

Shah, Rachana R

Segraves, Nathaniel L

et al.

Publication Date

2004-07-01

DOI

10.1021/jm049872s

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Probing the Activity Differences of Simple and Complex Brominated Aryl compounds against 15-Soybean, 15-Human and 12-Human Lipoxygenase

Erika N. Seagraves,[†] Rachana R. Shah,[†] Nathaniel L. Seagraves,^{†,‡} Tyler A. Johnson,^{†,‡} Stephanie
Whitman,[†] Jennifer K. Sui,[†] Victor A. Kenyon,[†] Robert H. Cichewicz,^{†,‡} Phillip Crews,^{*,†,‡} and
Theodore R. Holman ^{*,‡}

Department of Chemistry and Biochemistry and Institute for Marine Sciences, University of California,
Santa Cruz, California 95064

phil@chemistry.ucsc.edu, tholman@chemistry.ucsc.edu


RECEIVED DATE (to be automatically inserted after your manuscript is accepted)

*To whom correspondence should be addressed. TRH, Tel: 831-459-5884, Fax: 831-459-2935,
tholman@chemistry.ucsc.edu; PC, Tel: 831-459-2603, Fax: 831-459-4197, phil@chemistry.ucsc.edu

[†]Department of Chemistry and Biochemistry, University of California, Santa Cruz.

[‡]Institute for Marine Sciences, University of California, Santa Cruz.

Table of Contents Graphic:



Abstract. Lipoxygenases (LO) have been implicated in asthma, immune disorders, and various cancers. As a consequence of these broad biological implications, there is great interest in understanding the effects of naturally occurring and environmental contaminants against its activity. On the basis of our earlier studies indicating that polybrominated diphenol ethers are potent inhibitors to mammalian 15-LO, we expanded our structure-activity study to include marine-derived brominated phenol ethers (including a newly discovered tribrominated diphenyl ether), dioxins, and bastadins, as well as the synthetic brominated fire-retardants, brominated bisphenol A (BBPA) and polybrominated diphenyl ethers (PBDEs). We report herein the effects of twenty-one simple and complex organobromine compounds against human platelet 12-LO, human reticulocyte 15-LO, and soybean LO-1.

Introduction.

Previously, we determined that marine-derived brominated diphenol ethers inhibited mammalian 15-lipoxygenase with good effectiveness ($IC_{50} \approx 1-2 \mu M$),¹ however it remained unclear if this class of compounds were selective against particular lipoxygenase (LO) isozymes. For the past several years, our labs have discovered over a dozen LO inhibitors from marine sponges with the goal of defining chemical scaffolds for targeted lipoxygenase therapeutics.^{2,3} LOs are a class of non-heme iron-containing enzymes that contribute to the eicosanoid pathway⁴ by the hydroperoxidation of arachidonic acid (AA). These enzyme products are precursors for the inflammatory mediators, leukotrienes and lipoxins, but are also involved in a variety of human diseases such as bronchial constriction,⁵ psoriasis,⁶ atherosclerosis,⁷ and cancer^{8,9} and thus are attractive pharmaceutical targets.

Over 1600 organobromine compounds have been isolated from natural sources and these substances possess a broad range of structural intricacy and biological activity.¹⁰ Some examples include the monocyclic areoplysinin-1, a relatively simple organobromine compound which is a potent anti-fouling agent. The bicyclic organobromine compound, pentabromopseulilin, was isolated from the marine bacterium *Alteromonas luteoviolaceus* and is the most active member in a group of more than 20 pyrrole antibiotics.¹¹ The more structurally intricate organobromine scaffold, bastadin, exhibits a plethora of activities, such as moderate antibacterial activity, cytotoxic activity against human tumor cell lines, anti-inflammatory activity and calcium channel activation.¹²

Polybrominated compounds are also synthesized commercially and used in large quantities as fire retardants,¹³ such as polybrominated diphenyl ethers (PBDE) and brominated bisphenol A (BBPA). These compounds have broad implications concerning human health due to the increasing levels of PBDEs detected in human blood plasma,¹⁴ adipose tissue,¹⁵ and breast milk.^{16,17} Their accumulation in human tissue raises concern because PBDEs have been found to induce murine phospholipase A₂ to release arachidonic acid,¹⁸ potentially affecting the eicosanoid signaling pathway and lipoxygenase in particular. In light of our previous discovery of marine-derived polybrominated LO inhibitors,

combined with the potential human health implications of organobromine compounds, we initiated the current investigation in order to examine this class of compounds in more detail and determine what structural features are required for inhibition against human platelet 12-lipoxygenase (12-hLO), human reticulocyte 15-lipoxygenase-1 (15-hLO), and soybean 15-lipoxygenase-1 (15-sLO).

Results and Discussion

The initial investigation of twenty-one organobromine compounds was to determine if their inhibitory activity was due to reduction of the active site ferric ion. Lipoxygenase inhibitors can be classified as either non-reductive or reductive inhibitors.¹⁹ The non-reductive inhibitors, such as puupehenone and jaspic acid, bind to the protein presumably at the catalytic and/or allosteric sites.³ The reductive inhibitors reduce the active ferric form of LO to the inactive, ferrous form and typically contain hydroquinone or catecholate moieties, as observed in jaspiquinol³ and nordihydroguaiaretic acid (NDGA),²⁰ respectively. These reductive inhibitors are less desirable as pharmaceutical targets due to their susceptibility to undergo chemical modification in the cell.⁸ The brominated diphenol ethers, which are one target of this publication, could potentially be reductive LO inhibitors due to their phenolic moieties. We probed this possibility and determined that on a kinetic timescale (less than 2 minutes), reduction of the ferric form of 15-sLO was not observed by fluorescence spectroscopy for any of the twenty-one organobromine compounds studied. However, fluorescence spectroscopy did reveal partial reduction of the active site ferric iron with many inhibitors (**1a-7b**) after 20 minutes of incubation, suggesting that these phenolic compounds could reduce the active site iron, but on a time scale that is kinetically irrelevant (i.e. greater than 2 minutes). This was further supported by EPR experiments of 15-sLO and compound **2a**,²¹ a micromolar inhibitor of both soybean and human lipoxygenases (IC_{50} (15-sLO) = $7 \pm 3 \mu\text{M}$, IC_{50} (12-hLO) = $0.7 \pm 0.2 \mu\text{M}$ and IC_{50} (15-hLO) = $1.8 \pm 0.4 \mu\text{M}$). Upon addition of 5 equivalents of **2a** to 15-sLO, the ferric signal was unchanged. The sample was then allowed to incubate for 20 minutes at 22 °C and subsequently showed a 40% decrease in the ferric EPR signal (data not shown). The EPR result, combined with the fluorescence data, lead us to

conclude that reductive inactivation of 15-sLO does occur slowly with some phenolic inhibitors, but it is not a relevant factor on a kinetic timescale for any of the compounds studied. Furthermore, there was no change in the EPR signal of 15-sLO upon addition of **1c**, **2a**, **3a**, and **5a** demonstrating that their phenolic moieties do not chelate to the active site iron atom.

The bioactivity studies of the three lipoxygenase enzymes began with seven simple brominated phenols (**1a-1g**), which in general showed no inhibition against 12-hLO, 15-hLO, and 15-sLO, except for two mild inhibitors (Table 1). The compound, 4-bromophenol (**1c**), weakly inhibited 15-sLO and 15-hLO, but did not inhibit 12-hLO, while 2,4-dibromophenol (**1f**) was a weak inhibitor of 15-hLO, but did not inhibit either 15-sLO or 12-hLO. These results suggest that *para*-bromination of simple phenols increases their potency. Nevertheless, these simple bromophenols are poor inhibitors against LO in general and more extended dicyclic structures are required for potent lipoxygenase activity (*vida infra*).

We next investigated the brominated diphenyl ethers (**2a-3c**),²¹⁻²³ which possess more complex structural features with regards to their degrees of hydroxylation, methoxylation, and bromination (Table 2). The compound, 3,4-dibromo-2-(5'-bromo-2'-hydroxyphenoxy)phenol (**3c**), is a novel marine natural product, with unique bromination relative to the phenol. These brominated diphenyl ethers (**2c-3c**), in general, were poor inhibitors of 15-sLO relative to the human enzymes except for the monohydroxylated diphenyl ethers, **2a** and **2b**, which affected moderate potency ($IC_{50} \approx 8 \mu\text{M}$ for both). Small structural changes to **2a** and **2b**, such as the addition of a hydroxide and/or the bromination position, abolish 15-sLO inhibition entirely, as seen for **2c**, **3a-3c**. Specifically, the positioning of the *para*-bromide for the simple phenols inhibition (*vida supra*) is not observed for the more complex brominated diphenyl ethers.

The diphenyl ethers (**2a-3c**) were markedly more potent against both human enzymes than 15-sLO, with increased potency against 12-hLO relative to 15-hLO. Interestingly, both the monohydroxy- and dihydroxy- diphenyl ethers manifest the same potency trends, in which increasing bromination increased potency. This increase in potency relative to increased bromination could be due to multiple

factors: an increase in hydrophobicity, a dependency on bromide position, or an increase in size. The hydrophobicity (i.e. cLog P) correlates well with IC₅₀ values within each sub-class (**2a-2c** and **3a-3c**), however, it does not cross-correlate between the two sub-classes: the mono- and diphenolic inhibitors. For example, **2a** and **3a** have similar IC₅₀ values but different cLog P values. The position of bromination could also be a factor, however, no clear trend is observed between the mono- and diphenolic compounds in this investigation. Increased size due to increased bromination is the strongest correlation with potency but it is unclear how size affects binding since both enzymes have previously been shown to accept far larger inhibitors, such as puerphenones.³ Since none of these factors completely explain the data, the inhibitor potency appears to be due to as yet undefined factors. Molecular docking studies are currently underway to model the sites in both human enzymes and assess possible binding scenarios for these inhibitors.

The next class of LO inhibitors are the monohydroxylated dioxins (**4a** and **4b**).^{24,25} These compounds were very potent against 15-hLO, similar to the monohydroxylated diphenyl ethers, however they were poor inhibitors towards 12-hLO (Table 3). The bromination position appears not to be a factor because **4b** has comparable bromination to **2b**, yet has a larger IC₅₀ value. Considering that their cLog P values are similar to the other potent brominated inhibitors against 12-hLO (such as **2c** & **3c**), this lowered inhibitor potency towards 12-hLO is most likely due to the rigid 3-ring structure of the dioxin scaffold.

The synthetic fire retardants are the next class of LO inhibitors to be investigated and can be divided into two subgroups, the phenolic (BBPAs) and the non-phenolic brominated diphenyls (PBDEs). The diphenolic fire retardants (**5a** & **5b**) were good inhibitors of both 12- and 15-hLO and have similar potency to **2b** and **3b**, indicating that the isopropylidene bridge and the position of the phenol relative to the bridge is not critical for micromolar inhibition (Table 4). BBPA fire-retardants (**5a** & **5b**) were not inhibitors to 15-sLO, which correlates well with the previous results of poor inhibition by non-brominated BPA against 15-sLO.²⁶ The brominated diphenyl ethers (**6a** & **6b**) are industrial mixtures of brominated isomers, typically used in toxicology studies.¹⁸ These fire retardant mixtures were poor

inhibitors with respect to both human enzymes ($IC_{50} > 100 \mu\text{M}$ for both 12-hLO and 15-hLO), which suggests the free phenols are required for inhibition. It should be noted that hydroxy-PBDE's are the primary metabolites of PBDEs in rodents,^{27,28} which could increase their potency towards lipoxygenase, as seen for the compounds of this study.

The marine-derived bastadins (**7a**, **7b**)²⁹ are the most structurally complex of the brominated natural products in this study. They were weak inhibitors towards 15-sLO, but potent inhibitors against the human enzymes (Table 5). The linear bastadin 2 (**7a**) was a submicromolar inhibitor of 12-hLO, while the macrocyclic bastadin 7 (**7b**) was ~ 6 -fold less potent. Significantly, 15-hLO was inhibited by both **7a** and **7b** with micromolar potency, and showed no difference between linear and macrocyclic bastadins. These data translate into a selectivity preference where the linear **7a** was ~ 5 -fold more selective for 12-hLO than for 15-hLO but the macrocycle **7b** demonstrated no selectivity. Interestingly, **7a** and **7b** are some of the largest LO inhibitors ($\text{FW} > 1000 \text{ g/mol}$) studied in our labs. This could account for their selectivity due to the fact that 12-hLO has a longer active site than 15-hLO,³⁰ and would require more molecular contacts, consistent with the modeled substrate binding in the two enzymes.³⁰ Even though compounds **7a** and **7b** are large, complex inhibitors, their structural intricacy does not enhance their potency relative to either **2a** or **3a**.

The overall selectivity of the 21 organobromine compounds against 12-hLO and 15-hLO tends toward favoring 15-hLO inhibition over that of 12-hLO, a common feature in the marine-derived compounds we have isolated previously^{2,3} (Table 6). The most noteworthy compounds in this regard are the dioxins (**4a**, **4b**), which inhibit 15-hLO more effectively than 12-hLO, 38-fold and 56-fold respectively. This selectivity is similar to that observed for the rigid, tetracyclic puerphenones derivatives from our previous publications and is comparable to the redox inhibitor, NDGA (12-hLO/15-hLO = 46).²⁰ Finally, the selectivity of these compounds with respect to 15-hLO versus 15-sLO indicates that there is little similarity between their inhibition activity, thus supporting the conclusion that 15-sLO should not be used as a 15-hLO mimic.

In summary, these SAR inhibitor results allow us three general conclusions. First, these brominated phenols can reduce the active site iron, however this property is not responsible for the inhibition potency because it is irrelevant on the kinetic timescale. Second, the brominated phenol ethers (**2a**, **3a**) are potent inhibitors to human LOs and increasing their rigidity, such as in the dioxins (**4a**, **4b**), can increase their potency and selectivity towards 15-hLO. Finally among the synthetic brominated fire-retardants, the phenolic BBPAs (**5a**, **5b**) are human LO inhibitors, while the non-phenolic PBDEs (**6a**, **6b**) are not, indicating a requirement of hydroxylation for potency.

Experimental.

Materials. Linoleic acid (LA) and AA were purchased from Aldrich Chemical Company (Milwaukee, WI). All other chemicals were reagent grade or better and were used without further purification.

Expression and Purification of Lipoxygenases. 12-hLO and 15-hLO were expressed and purified as described previously.³ Briefly, both His-tagged enzymes must be purified in one step and stored in 20% glycerol at $-80\text{ }^{\circ}\text{C}$, or significant inactivation occurs. 15-sLO was purified as previously published.³¹

Lipoxygenase Assay. The IC_{50} determination for all three enzymes was performed as previously described with the following modifications.² Under separate reaction conditions (12-hLO, 25 mM HEPES, pH 8; 15-hLO, 25 mM HEPES, pH 7.5; 15-sLO, 100 mM borate, pH 9.2), substrate was added (AA for 12-hLO and LA for 15-hLO/15-sLO). Reactions were initiated by adding the appropriate enzyme and reaction rates were monitored at 234 nm at room temperature (22-23 $^{\circ}\text{C}$) with constant stirring with a rotating magnetic stir-bar. Inhibitors were dissolved in MeOH to a concentration of 1 mg/mL. Control reactions were performed by adding an equivalent amount of solvent to the reaction. The data were fit to a simple hyperbolic curve with the program KaleidaGraph (Synergy) and IC_{50} calculated.

Purification of Marine-derived Compounds. Pure compounds **2a** and **2b** were obtained from a 100 g portion of *Psammocina* sp. extract (Papua New Guinea, pooled collections) that was partitioned with CH₂Cl₂, hexane, and MeOH. The CH₂Cl₂ partition (49 g) was fractionated using silica gel with serial elutions of 100% toluene, EtOAc, and finally MeOH. The bioactive fractions were combined and further fractionated using silica gel chromatography with hexane, CH₂Cl₂, and MeOH, then subjected to HPLC with a gradient of 70%-100% ACN in H₂O (with 0.1% formic acid in both solvents) yielding **2a** (1.5 mg) and **2b** (0.5 mg). Structures of **2a** & **2b** were determined based on comparison with published spectral data (¹H and ¹³C NMR).^{21,22} Compounds **2c**, **3a**, **4a**, **4b**, **7a**, **7b** were obtained from the Marine Natural Products Repository at UCSC, analytical data matches that in the literature.^{1,23-25,29}

3,4-dibromo-2-(5'-bromo-2'-hydroxyphenoxy)phenol (3c): Pure compounds **3b** and **3c** were obtained from the sponge identified in the field as belonging to the genus *Phyllospongia* (Solomon Islands, coll. no. 94034) following the published extraction scheme.³² A 0.5 g portion of the CH₂Cl₂ partition extract was fractionated using Sephadex LH-20 with 100% MeOH, then further separated using C₁₈ HPLC with a gradient of 50% to 100% MeOH in H₂O (0.1% trifluoroacetic acid in both solvents) to yield **3b** (8.8 mg) and **3c** (7.3 mg). The structure of **3b** was determined based on comparison with published spectral data (¹H and ¹³C NMR).²³ The structure of **3c** was determined by its physical properties: red solid. UV (MeOH) λ_{max} (log ε) 207 (4.7), 287 (3.7); ¹H and ¹³C NMR data see supporting information. TOFMS: *m/z* 435/437/439/441 [M -H] (calcd for C₁₂H₇Br₃O₂).

Purification of Synthetic Brominated Fire-Retardants. BBPA (BA-59P, Lot #2008JM17B) and PBDE mixtures (DE-69, Lot #21220B04A and DE-71, Lot #2525DI01A) were a gift from Great Lakes Chemical Corporation (West Lafayette, IN). A 0.5 g sample of BA-59P was purified by HPLC with a gradient of 75% to 100% MeOH in H₂O (0.1% formic acid in both solvents) to yield **5a** (4.8 mg) and **5b** (8.7 mg). Structures of **5a** and **5b** were verified by ¹H and ¹³C NMR, in comparison to predicted data derived from simulation using Advanced Chemical Dictionary, Inc. software. DE-69 and DE-71 were a mixture of pentabrominated (**6a**) and octabrominated (**6b**) PBDE isomers, respectively. Due to the low

solubility of pure compounds under assay conditions, these mixtures were used without further purification and are designated as compounds **6a** and **6b**.

Spectroscopic Characterization of the Ferric and Ferrous Soybean Lipoxygenase. The fluorescence of the resting, ferrous form of 15-sLO has been reported previously.^{33,34} Briefly, the ferrous signal was obtained by excitation at 280 nm and monitored at 328 nm on a Perkin-Elmer LS50B Spectrometer. Assays were 2 mL in volume and constantly stirred with a rotating magnetic bar. The ferrous signal was obtained by addition of 700 nM 15-sLO to an appropriate volume of 100 mM borate (pH 9.2) and was converted to the active ferric form by addition of one equivalent of 13-hydroperoxy-9(Z),11(E)-octadecadienoic acid (HPOD), quenching the fluorescence by 30%. The change in fluorescence was then monitored upon addition of 5-fold excess inhibitor dissolved in MeOH. Increasing signal was attributed to the reduction of the ferric enzyme to the ferrous form via the inhibitor, and verified by adding the known reductant, NDGA. A blank was performed by addition of MeOH and no increase in fluorescence was observed. The reduction of the active site was also examined by EPR in frozen solution at 4.3 K (data not shown). The procedure of Nelson³⁵ was followed with the following modifications. Addition of 1 equivalent of HPOD ($\approx 150 \mu\text{M}$) to the resting enzyme produced a signal at $g \approx 6$. The solution was thawed and 5 equivalents of inhibitor, dissolved in DMSO, were added. The solution was mixed and quickly refrozen in liquid nitrogen and the spectrum taken. If no reduction was seen, then the sample was thawed and allowed to incubate for 20 minutes at 22 °C, refrozen, and the spectrum re-taken. A decrease in the signal at $g \approx 6$ was ascribed to reduction of the ferrous iron.

cLog P determination. The cLog P was calculated using the atom fragmentation method,³⁶ available from Daylight Chemical Information Systems, Inc. via the world wide web.³⁷

ACKNOWLEDGMENT. The authors thank Great Lakes Chemical Corporation for providing samples of PBDE mixtures and Dr. M. Cristina Diaz (UCSC) for the identification of the sponge

taxonomy. Financial support was from the National Institute of Health grants R01 CA47135 (PC) and GM 56062-06 (TRH), American Cancer Society grant RPG-00-219-01-CDD (TRH). JKS was supported by the summer undergraduate research fellowship at UCSC by the National Science Foundation REU CHE-998784. Additional financial support was from equipment grants from the NSF BIR-94-19409 (NMR) and a supplement to NIH CA52955 (ESI-TOF-MS).

Table 1. LO inhibition activity ($\mu\text{M} \pm \text{SD}$) of simple bromophenols (**1a-1g**).

1

Compound d	R	X	cLog P	12-hLO	15-hLO	15-sLO
				IC ₅₀ (μM)	IC ₅₀ (μM)	IC ₅₀ (μM)
1a	OH	2-Br	2.35	>100	>100	>100
1b	OH	3-Br	2.63	>100	>100	>100
1c	OH	4-Br	2.59	>100	55 \pm 6	48 \pm 4
1d	CH ₂ OH	4-Br	1.97	>100	>100	>100
1e	OCH ₂ CH ₃	4-Br	3.59	>100	>100	>100
1f	OH	2,4- Br	3.22	>100	34 \pm 4	>100
1g	OH	2,6- Br	3.36	>100	>100	>100

Table 2. LO inhibition activity ($\mu\text{M} \pm \text{SD}$) of marine-derived polybrominated phenol ethers (**2a-2c**)^{1,21,22} and diphenol ethers (**3a-3c**).²³







Compound d	Structure	cLog P	12-hLO	15-hLO	15-sLO
			IC ₅₀ (μM)	IC ₅₀ (μM)	IC ₅₀ (μM)
2a		8.21	0.7 ± 0.2	1.8 ± 0.4	7 ± 3
2b		7.34	6 ± 2	5 ± 1	9 ± 3
2c		6.42	12 ± 2	10 ± 1	>100
3a		7.67	0.41 ± 0.03	0.79 ± 0.07	>100
3b		6.77	6.2 ± 0.8	2.2 ± 0.4	>100
3c		5.26	47 ± 8	11 ± 1	>100

Table 3. LO inhibition activity ($\mu\text{M} \pm \text{SD}$) of marine-derived polybrominated dioxins (**4a**, **4b**).^{24,25}




Compound	X	cLog P	12-hLO IC ₅₀ (μM)	15-hLO IC ₅₀ (μM)	15-sLO IC ₅₀ (μM)
4a	H	6.66	30 ± 11	0.8 ± 0.1	>100
4b	Br	7.42	50 ± 14	0.9 ± 0.1	>100

Table 4. LO inhibition activity ($\mu\text{M} \pm \text{SD}$) of synthetic brominated fire-retardants (BBPAs) (**5a**, **5b**).



Compound	X	cLog P	12-hLO IC ₅₀ (μM)	15-hLO IC ₅₀ (μM)	15-sLO IC ₅₀ (μM)
5a	H	6.12	7 ± 3	5 ± 2	>100
5b	Br	6.81	10 ± 5	4 ± 1	>100

Table 5. LO inhibition activity ($\mu\text{M} \pm \text{SD}$) of marine-derived bastadins (**7a**, **7b**).²⁹

Compound	cLog P	12-hLO IC ₅₀ (μM)	15-hLO IC ₅₀ (μM)	SLO-1 IC ₅₀ (μM)
7a	9.10	0.4 ± 0.1	2.0 ± 0.4	59 ± 45
7b	10.21	2.3 ± 0.6	4 ± 1	27 ± 12

Table 6. Selectivity of biologically active compounds (**1c**, **1f**, **2a-7b**).

Compound d	12-hLO/15- hLO	15-sLO/15-hLO	Concentration ^a -log(IC ₅₀) (M)
1c	> 2	1	
1f	> 3	> 3	
2a	0.4	4	
2b	1	2	
2c	1	> 10	
3a	1	> 130	
3b	3	> 45	
3c	4	> 10	
4a	38	> 130	
4b	56	> 110	
5a	1	> 20	
5b	2	> 20	
6a	≈ 1	≈ 1	
6b	≈ 1	0.3	
7a	0.2	30	
7b	1	7	

3.5 4 4.5 5 5.5 6 6.5

^aConcentration is expressed as – log (IC₅₀) in molar units. Open bars represent 12-hLO activity, solid bars represent 15-hLO activity, and hatched bars represent 15-sLO activity.

REFERENCES

- (1) Fu, X.; Schmitz, F. J.; Govindan, M.; Abbas, S. A.; Hanson, K. M.; Horton, P. A.; Crews, P.; Laney, M.; Schatzman, R. C. Enzyme inhibitors: New and known polybrominated phenols and diphenyl ethers from four indo-pacific *Dysidea* sponges. *J. Nat. Prod.* **1995**, *58*, 1384-1391.
- (2) Carroll, J.; Jonsson, E. N.; Ebel, R.; Hartman, M.; Holman, T. R.; Crews, P. Probing sponge-derived terpenoids for human 15-lipoxygenase inhibitors. *J. Org. Chem.* **2001**, *66*, 6847-6851.
- (3) Amagata, T.; Whitman, S.; Johnson, T. A.; Stessman, C. C.; Loo, C. P.; Lobkovsky, E.; Clardy, J.; Crews, P.; Holman, T. R. Exploring sponge-derived terpenoids for their potency and selectivity against 12-human, 15-human, and 15-soybean lipoxygenases. *J. Nat. Prod.* **2003**, *66*, 230-235.
- (4) Roberts, L. J. Introduction: lipids as regulators of cell function. *Cell. Mol. Life Sci.* **2002**, *59*, 727-728.
- (5) Nakano, H.; Inoue, T.; Kawasaki, N.; Miyataka, H.; Matsumoto, H.; Taguchi, T.; Inagaki, N.; Nagai, H.; Satoh, T. Synthesis and biological activities of novel antiallergic agents with 5-lipoxygenase inhibiting action. *Bioorg. Med. Chem.* **2000**, *8*, 373-380.
- (6) Hussain, H.; Shornick, L. P.; Shannon, V. R.; Wilson, J. D.; Funk, C. D.; Pentland, A. P.; Holtzman, M. J. Epidermis contains platelet-type 12-lipoxygenase that is overexpressed in germinal layer keratinocytes in psoriasis. *Am. J. Physiol.* **1994**, *266*, C243-C253.
- (7) Harats, D.; Shaish, A.; George, J.; Mulkins, M.; Kurihara, H.; Levkovitz, H.; Sigal, E. Overexpression of 15-lipoxygenase in vascular endothelium accelerates early atherosclerosis in LDL receptor-deficient mice. *Arterioscl. Throm. Vas. Biol.* **2000**, *20*, 2100-2105.

- (8) Steele, V. E.; Holmes, C. A.; Hawk, E. T.; Kopelovich, L.; Lubet, R. A.; Crowell, J. A.; Sigman, C. C.; Kelloff, G. J. Lipoxygenase inhibitors as potential cancer chemopreventives. *Cancer Epidemiol. Biomark. Prev.* **1999**, *8*, 467-483.
- (9) Avis, I.; Hong, S. H.; Martinez, A.; Moody, T.; Choi, Y. H.; Trepel, J.; Das, R.; Jett, M.; Mulshine, J. L. Five-lipoxygenase inhibitors can mediate apoptosis in human breast cancer cell lines through complex eicosanoid interactions. *FASEB J.* **2001**, *15*, U277-U306.
- (10) Gribble, G. W. The diversity of naturally produced organohalogens. *Chemosphere* **2003**, *52*, 289-297.
- (11) Laatsch, H.; Renneberg, B.; Hanefeld, U.; Kellner, M.; Pudleiner, H.; Hamprecht, G.; Kraemer, H. P.; Anke, H. Structure-activity relationships of phenyl- and benzoylpyrroles. *Chem. Pharm. Bull.* **1995**, *43*, 537-546.
- (12) Gribble, G. W. Naturally occurring organohalogen compounds. *Acc. Chem. Res.* **1998**, *31*, 141-152.
- (13) McDonald, T. A. A perspective on the potential health risks of PBDEs. *Chemosphere* **2002**, *46*, 745-755.
- (14) Hovander, L.; Malmberg, T.; Athanasiadou, M.; Athanassiadis, L.; Rahm, S.; Bergman, A.; Wehler, E. K. Identification of hydroxylated PCB metabolites and other phenolic halogenated pollutants in human blood plasma. *Arch. Environ. Contam. Tox.* **2002**, *42*, 105-117.
- (15) She, J. W.; Petreas, M.; Winkler, J.; Visita, P.; McKinney, M.; Kopec, D. PBDEs in the San Francisco Bay Area: measurements in harbor seal blubber and human breast adipose tissue. *Chemosphere* **2002**, *46*, 697-707.

- (16) Noren, K.; Meironyte, D. Certain organochlorine and organobromine contaminants in Swedish human milk in perspective of past 20-30 years. *Chemosphere* **2000**, *40*, 1111-1123.
- (17) Meironyte, D.; Noren, K.; Bergman, A. Analysis of polybrominated diphenyl ethers in Swedish human milk. A time-related trend study, 1972-1997. *J. Tox. Environ. Health* **1999**, *58*, 329-341.
- (18) Kodavanti, P. R. S.; Derr-Yellin, E. C. Differential effects of polybrominated diphenyl ethers and polychlorinated biphenyls on [H-3]arachidonic acid release in rat cerebellar granule neurons. *Tox. Sci.* **2002**, *68*, 451-457.
- (19) Nelson, M. J.; Batt, D. G.; Thompson, J. S.; Wright, S. W. Reduction of the active-site iron by potent inhibitors of lipoxygenases. *J. Biol. Chem.* **1991**, *266*, 8225-8229.
- (20) Whitman, S.; Gezginci, M.; Timmermann, B. N.; Holman, T. R. Structure-activity relationship studies of nordihydroguaiaretic acid inhibitors toward soybean, 12-human, and 15-human lipoxygenase. *J. Med. Chem.* **2002**, *45*, 2659-2661.
- (21) Salva, J.; Faulkner, D. J. A new brominated diphenyl ether from a Philippine *Dysidea* species. *J. Nat. Prod.* **1990**, *53*, 757-760.
- (22) Bowden, B. F.; Towerzey, L.; Junk, P. C. A new brominated diphenyl ether from the marine sponge *Dysidea herbacea*. *Aust. J. Chem.* **2000**, *53*, 299-301.
- (23) Norton, R. S.; Croft, K. D.; Wells, R. J. Polybrominated oxydiphenol derivatives from the sponge *Dysidea herbacea*. *Tetrahedron* **1981**, *37*, 2341-2349.
- (24) Utkina, N. K.; Denisenko, V. A.; Virovaya, M. V.; Scholokova, O. V.; Prokofeva, N. G. Two new minor polybrominated dibenzo-p-dioxins from the marine sponge *Dysidea dendyi*. *J. Nat. Prod.* **2002**, *65*, 1213-1215.

- (25) Utkina, N. K.; Denisenko, V. A.; Scholokova, O. V.; Virovaya, M. V.; Gerasimenko, A. V.; Popov, D. Y.; Krasokhin, V. B.; Popov, A. M. Spongiadioxins A and B, two new polybrominated dibenzo-p-dioxins from an Australian marine sponge *Dysidea dendyi*. *J. Nat. Prod.* **2001**, *64*, 151-153.
- (26) Rao, K. C. S.; Divakar, S.; Rao, A. G. A.; Karanth, N. G.; Sattur, A. P. A lipoxygenase inhibitor from *Aspergillus niger*. *Appl. Microbiol. Biotechnol.* **2002**, *58*, 539-542.
- (27) Orn, U.; Klasson-Wehler, E. Metabolism of 2,2',4,4'-tetrabromodiphenyl ether in rat and mouse. *Xenobiotica* **1998**, *28*, 199-211.
- (28) Hakk, H.; Larsen, G.; Klasson-Wehler, E. Tissue disposition, excretion and metabolism of 2,2',4,4',5-pentabromodiphenyl ether (BDE-99) in the male Sprague-Dawley rat. *Xenobiotica* **2002**, *32*, 369-382.
- (29) Kazlauskas, R.; Lidgard, R. O.; Murphy, P. T.; Wells, R. J.; Blount, J. F. Brominated tyrosine-derived metabolites from the sponge *Ianthella basta*. *Aust. J. Chem.* **1981**, *34*, 765-786.
- (30) Borngraber, S.; Browner, M.; Gillmor, S.; Gerth, C.; Anton, M.; Fletterick, R.; Kuhn, H. Shape and specificity in mammalian 15-lipoxygenase active site. *J. Biol. Chem.* **1999**, *274*, 37345-37350.
- (31) Holman, T. R.; Zhou, J.; Solomon, E. I. Spectroscopic and functional characterization of a ligand coordination mutant of soybean lipoxygenase: first coordination sphere analogue of human 15-lipoxygenase. *J. Am. Chem. Soc.* **1998**, *120*, 12564-12572.
- (32) Thale, Z.; Johnson, T.; Tenney, K.; Wenzel, P. J.; Lobkovsky, E.; Clardy, J.; Media, J.; Pietraszkiewicz, H.; Valeriote, F. A.; Crews, P. Structures and cytotoxic properties of sponge-derived bisannulated acridines. *J. Org. Chem.* **2002**, *67*, 9384-9391.

- (33) Finazzi-Agro, A.; Avigliano, L.; Veldink, G. A.; Vliegenhart, J. F. G.; Boldingh, J. The influence of oxygen on the fluorescence of lipoxygenase. *Biochim. Biophys. Acta* **1973**, *326*, 462-470.
- (34) Egmond, M. E.; Finazzi-Agro, A.; Fasella, P. M.; Veldink, G. A.; Vliegenhart, J. F. G. Changes in the fluorescence and absorbance of lipoxygenase-1 induced by 13-*L,S*-hydroperoxylinoleic acid and linoleic acid. *Biochim. Biophys. Acta* **1975**, *397*, 43-49.
- (35) Nelson, M. J. Catecholate complexes of ferric soybean lipoxygenase-1. *Biochemistry* **1988**, *27*, 4273-4278.
- (36) Ghose, A. K.; Pritchett, A.; Crippen, G. M. Atomic physicochemical parameters for 3-dimensional structure directed quantitative structure-activity relationships. *J. Comp.Chem.* **1988**, *9*, 80-90.
- (37) Daylight Chemical Information Systems, Inc. CLOGP: Calculation of hydrophobicity as Log P(o/w). Available from: <http://www.daylight.com/daycgi/clogp>, (accessed Sept. 2003).