

# UC Davis

## UC Davis Previously Published Works

### Title

TBX6 null variants and a common hypomorphic allele in congenital scoliosis.

### Permalink

<https://escholarship.org/uc/item/4v9775d7>

### Journal

The New England Journal of Medicine, 372(4)

### Authors

Shen, J  
Chen, Y  
Zhang, T  
et al.

### Publication Date

2015-01-22

### DOI

10.1056/NEJMoa1406829

Peer reviewed



Published in final edited form as:

*N Engl J Med.* 2015 January 22; 372(4): 341–350. doi:10.1056/NEJMoa1406829.

## ***TBX6* Null Variants and a Common Hypomorphic Allele in Congenital Scoliosis**

*A full list of authors and affiliations appears at the end of the article.*

### **Abstract**

**BACKGROUND**—Congenital scoliosis is a common type of vertebral malformation. Genetic susceptibility has been implicated in congenital scoliosis.

**METHODS**—We evaluated 161 Han Chinese persons with sporadic congenital scoliosis, 166 Han Chinese controls, and 2 pedigrees, family members of which had a 16p11.2 deletion, using comparative genomic hybridization, quantitative polymerase-chain-reaction analysis, and DNA sequencing. We carried out tests of replication using an additional series of 76 Han Chinese persons with congenital scoliosis and a multi-center series of 42 persons with 16p11.2 deletions.

**RESULTS**—We identified a total of 17 heterozygous *TBX6* null mutations in the 161 persons with sporadic congenital scoliosis (11%); we did not observe any null mutations in *TBX6* in 166 controls ( $P < 3.8 \times 10^{-6}$ ). These null alleles include copy-number variants (12 instances of a 16p11.2 deletion affecting *TBX6*) and single-nucleotide variants (1 nonsense and 4 frame-shift mutations). However, the discordant intrafamilial phenotypes of 16p11.2 deletion carriers suggest that heterozygous *TBX6* null mutation is insufficient to cause congenital scoliosis. We went on to identify a common *TBX6* haplotype as the second risk allele in all 17 carriers of *TBX6* null mutations ( $P < 1.1 \times 10^{-6}$ ). Replication studies involving additional persons with congenital scoliosis who carried a deletion affecting *TBX6* confirmed this compound inheritance model. In vitro functional assays suggested that the risk haplotype is a hypomorphic allele. Hemivertebrae are characteristic of *TBX6*-associated congenital scoliosis.

**CONCLUSIONS**—Compound inheritance of a rare null mutation and a hypomorphic allele of *TBX6* accounted for up to 11% of congenital scoliosis cases in the series that we analyzed.

CONGENITAL SCOLIOSIS, A FORM OF VERTEBRAL malformation, has an estimated prevalence of approximately 1 in 2000 live births.<sup>1</sup> It is manifested as a lateral curvature of the spine exceeding 10 degrees and results from defects in vertebra formation during embryogenesis.<sup>2,3</sup>

Previous evidence in animal models suggested that genetic factors contribute to vertebral malformations.<sup>3,4</sup> Genetic mutations have been implicated in human congenital scoliosis,

---

Copyright © 2015 Massachusetts Medical Society. All rights reserved

Address reprint requests to Dr. Feng Zhang at the State Key Laboratory of Genetic Engineering, School of Life Sciences, Fudan University, 2005 Song Hu Rd., Shanghai 200438, China, or at zhangfeng@fudan.edu.cn. Dr. Guixing Qiu at the Department of Orthopedic Surgery, Peking Union Medical College Hospital, Peking Union Medical College and Chinese Academy of Medical Sciences, No. 1 Shuaifuyuan, Beijing 100730, China, or at qiuguixingpumch@gmail.com.

Dr. N. Wu, Mr. X. Ming, Dr. J. Xiao, Dr. Z. Wu, and Dr. X. Chen contributed equally to this article.

Disclosure forms provided by the authors are available with the full text of this article at NEJM.org.

but their low penetrance highlights the complex molecular basis of the disorder and hinders molecular diagnosis of and genetic counseling for congenital scoliosis.<sup>4,5</sup>

The proximal 16p11.2 deletion in humans is rare but recurrent, with a population frequency of approximately 0.03% worldwide.<sup>6</sup> It is a likely cause of common diseases, including autism and obesity.<sup>7-9</sup> Vertebral malformations have been observed in a small proportion of persons with 16p11.2 copy-number variants (including deletion and duplication),<sup>10-12</sup> a finding that suggests the potential involvement of 16p11.2 copy-number variants (specifically, deletions)<sup>13</sup> in congenital scoliosis.

## METHODS

### STUDY PARTICIPANTS

We enrolled 237 consecutive Han Chinese persons who received a diagnosis of congenital scoliosis between October 2010 and June 2014 at Peking Union Medical College Hospital (PUMCH). Ancestry was determined by self-report (see the Methods section in the Supplementary Appendix, available with the full text of this article at NEJM.org). The discovery set consisted of 161 persons with sporadic congenital scoliosis (series 1), and we tested for replication in an additional series of 76 persons with the disorder (series 2). A clinical diagnosis of congenital scoliosis was confirmed by radiologic imaging. We excluded persons with syndromic diseases.

A total of 166 unrelated Han Chinese persons with no evidence of congenital scoliosis or other malformations served as population controls. We also evaluated 2 Han Chinese pedigrees, family members of which carried the 16p11.2 deletion and had discordant intrafamilial phenotypes.

For a further replication study, 42 unrelated persons with 16p11.2 deletion (series 3) were enrolled from multiple centers in the United States and China. These persons were initially referred for clinical chromosomal microarray testing owing to various medical problems.<sup>13-15</sup>

### CELL CULTURE AND INDUCED DIFFERENTIATION

The P19CL6 cell line (derived from a mouse embryonic carcinoma) was cultured and differentiated into cardiomyocytes, as described previously.<sup>16</sup> The cells were harvested every 24 hours after treatment with dimethyl sulfoxide.<sup>17</sup> We also used a human induced pluripotent stem cell (hiPSC) line that we derived from dermal fibroblasts by means of an efficient and integration-free method.<sup>18</sup> The culture of this hiPSC line and its induced differentiation to mesodermal-cell and cardiac-cell lineages has been described previously.<sup>19,20</sup> The cells were harvested 24 hours and 48 hours after differentiation with RPMI-B-27 medium without insulin.

### ANALYSIS OF GENOTYPE-PHENOTYPE CORRELATION

We reviewed the medical records and spinal radiographs of all 237 persons with sporadic congenital scoliosis. Vertebral malformations were classified as hemivertebra or hypoplasia, segmentation defect, or butterfly vertebra, as defined in a previous report.<sup>21</sup> Vertebral

malformations at more than one site in a patient were classified and counted independently according to location.

## STUDY OVERSIGHT

This study was approved by the institutional review boards of the PUMCH, Fudan University, the Capital Institute of Pediatrics, and the other participating institutions. We obtained written informed consent from the participants (those who were ≥18 years of age at the time of enrollment) or their guardians (for participants who were <18 years of age).

## STATISTICAL ANALYSIS

Fisher's exact test was used to assess the difference in *TBX6* mutation or variant frequencies between the persons with congenital scoliosis and controls. The unpaired t-test was used for statistical analysis of the luciferase reporter assays. Pearson's chi-square test was used to assess the difference in the prevalence of vertebral malformations between persons with *TBX6*-associated congenital scoliosis and those with congenital scoliosis but no *TBX6* nonsynonymous mutation. Odds ratios and 95% confidence intervals were calculated. The statistical analysis was performed with the use of SPSS software, version 15.0 (SPSS). A two-sided P value of less than 0.05 was considered to indicate statistical significance.

Further details of the study participants and of assays using comparative genomic hybridization (CGH), quantitative polymerase chain reaction, DNA sequencing of *TBX6*, haplotyping of *TBX6* variants, and luciferase assays can be found in the Supplementary Appendix.

## RESULTS

### PREVALENCE OF 16P11.2 DELETIONS IN PERSONS WITH CONGENITAL SCOLIOSIS

A genomewide analysis of copy-number variants was performed in 20 trios (consisting of a person with sporadic congenital scoliosis and two healthy parents) with the use of Agilent 1×1M CGH microarrays. We identified recurrent deletions in proximal 16p11.2 (Fig. 1A) in two persons (Patient XH004 and Patient XH042) (Fig. S1 in the Supplementary Appendix). Both of these heterozygous deletions were de novo (i.e., absent in the patient's parents).

Of the remaining 141 persons with sporadic congenital scoliosis in series 1, 10 had a heterozygous 16p11.2 deletion (Table 1), whereas none of the 166 Han Chinese controls had the deletion (12 of 161 persons with congenital scoliosis vs. 0 of 166 controls,  $P < 0.001$  by Fisher's exact test). These findings are consistent with the rarity of the 16p11.2 deletion in human populations (0.03%).<sup>22,23</sup> Remarkably, the prevalence of the deletion among persons with sporadic congenital scoliosis (12 of 161 persons, 7.5%) was much higher than its reported prevalence among persons with autism (0.4 to 0.8%)<sup>6,7</sup> or obesity (0.7%),<sup>8,9</sup> two phenotypes that it is known to cause. None of our patients with congenital scoliosis were obese. Their detailed neurodevelopmental records were not available for us to evaluate symptoms of autism.

## ASSOCIATION OF *TBX6* LOSS-OF-FUNCTION MUTATIONS WITH CONGENITAL SCOLIOSIS

The association of the 16p11.2 deletion with congenital scoliosis suggested that a gene related to the disorder may be located in the segment of DNA that is deleted: of these genes, *TBX6* was considered to be the best candidate (Fig. 1B).<sup>24-26</sup>

Accordingly, we hypothesized that other null or loss-of-function mutations (e.g., nonsense, frame-shift, and essential splice-site variants)<sup>27</sup> of *TBX6* cause congenital scoliosis. One heterozygous non-sense and four heterozygous frame-shift mutations of *TBX6* were identified among the patients with congenital scoliosis in series 1 (Table 1, and Fig. S2 in the Supplementary Appendix). The frame-shift mutation in Patient XH122 was de novo, whereas the status (i.e., de novo or inherited) of the other four null mutations is unknown. In contrast, we did not observe any null mutations in the controls (5 of 161 persons with congenital scoliosis vs. 0 of 166 controls,  $P=0.03$  by Fisher's exact test). These findings suggest an association of *TBX6* null mutations with congenital scoliosis.

Furthermore, we identified *TBX6* missense mutations in 7 of 161 persons with congenital scoliosis in series 1, a prevalence higher than that among the controls (3 of 166 persons) and among Han Chinese populations in Beijing and in southern China (CHB and CHS populations) in the 1000 Genomes Project (1 of 197 persons) (Fig. S3 and Tables S1 through S4 in the Supplementary Appendix).<sup>22</sup> In a combined analysis, the association of *TBX6* missense mutations with congenital scoliosis was significant (7 of 161 persons with congenital scoliosis vs. 4 of 363 controls,  $P=0.04$  by Fisher's exact test). Because missense mutations could be benign or even confer a gain of function, we did not include them in additional analyses, although we cannot exclude the possibility of a role for them in congenital scoliosis.

### A COMMON RISK HAPLOTYPE IN NULL MUTATION CARRIERS

Each person with congenital scoliosis in series 1 who carried a *TBX6* mutation was found to have only one null mutation (Table 1), which, at face value, suggests a simple dominant mode of inheritance. However, dominant inheritance is not supported by the low penetrance of congenital scoliosis in carriers of the 16p11.2 deletion affecting *TBX6*.<sup>10-12</sup> We therefore hypothesized that an additional genetic modifier acts in concert with *TBX6* null mutations to cause congenital scoliosis.

To test this hypothesis, we analyzed two pedigrees with an affected member and unaffected members who had the same 16p11.2 deletion. In Pedigree SE1 (Fig. 1C), the proband had congenital scoliosis with hemivertebra, whereas the father and sibling had the same deletion but did not have congenital scoliosis.<sup>12</sup> In Pedigree SE2 (Fig. 1D), the deletion in the proband was paternally inherited. The radiograph revealed a wedge-shaped vertebral malformation in the proband (Fig. 2); congenital scoliosis was not observed in the father.

Panel A shows the region affected by the proximal 16p11.2 deletion based on the human genome assembly hg19. A pair of long direct genomic repeats (shown in orange) can mediate 0.6-Mb recurrent deletions. The genes affected by this deletion, including *TBX6* (circled), are shown. Panel B shows three common single-nucleotide polymorphisms (SNPs)

in *TBX6*. Two SNPs, rs3809624 and rs3809627, are located in the 5' noncoding region, whereas rs2289292 is a synonymous SNP in the last exon. Panels C and D show two pedigrees with 16p11.2 deletions (SE1 and SE2). Squares denote male pedigree members, circles female pedigree members, solid symbols members with congenital scoliosis, and open symbols unaffected members; the probands are indicated by black arrows. The red bracket represents the deletion allele, and the nonreference alleles of rs2289292, rs3809624, and rs3809627 on the nondeletion chromosomes are shown in blue. The slash denotes a deceased family member. NA denotes not available.

We did not identify a rare mutation in *TBX6*, other than the deletion itself, in any of the family members. However, we did observe a common *TBX6* haplotype of T-C-A (defined by the nonreference alleles of three common single-nucleotide polymorphisms [SNPs] — rs2289292, rs3809624, and rs3809627, respectively) that cosegregated with congenital scoliosis in both pedigrees (Fig. 1). Therefore, we further evaluated the 17 persons with congenital scoliosis in series 1 who had the *TBX6* null mutations. All 17 persons carried the T-C-A haplotype at the second *TBX6* allele (Table 1). The probability of this coinheritance is  $P = (44\%)^{17}$ , which is equivalent to  $P < 1.1 \times 10^{-6}$ , where 44% is the frequency of the T-C-A haplotype among Han Chinese persons in the 1000 Genomes Project (Table S5 in the Supplementary Appendix).<sup>22</sup> Our observations suggest a genetic model of *TBX6* compound inheritance involving a combination of rare null and common risk alleles in the causation of congenital scoliosis. This model accounts for 17 of 161 cases of congenital scoliosis (11%) in series 1 (Table 2).

## REPLICATION STUDIES

We recruited an additional 76 persons with congenital scoliosis (series 2) and identified 6 *TBX6* null mutations (Table 1). Again, we observed the T-C-A risk haplotype at the second allele in all persons with these mutations. The *TBX6* compound inheritance model accounts for 6 of 76 cases of congenital scoliosis (8%) in series 2 (Table 2).

Furthermore, we enrolled 42 persons with the 16p11.2 deletion (series 3) from multiple centers (Table S6 in the Supplementary Appendix). These persons were categorized into three subgroups: those with congenital scoliosis (vertebral malformations), those who had scoliosis without vertebral malformations, and those with no evidence of scoliosis. This made it feasible to investigate whether the hemizygous T-C-A haplotype is associated with congenital scoliosis in 16p11.2 deletion carriers. In the congenital scoliosis subgroup, we identified 5 of 6 persons (83%) as having the hemizygous T-C-A haplotype (Table 2). The only exception (Patient PT03) was of African ancestry,<sup>13</sup> and the T-C-A haplotype is rare among such persons (prevalence, <1% among Africans, as compared with 44% among Asians and 33% among persons of European descent) (Table S5 in the Supplementary Appendix). Patient PT03 had the nonreference allele of SNP rs111939223 (present in 4% of Yoruba Africans but absent in Asians and Europeans), which is located at a predicted messenger RNA polyadenylation site of *TBX6*.<sup>28</sup> Therefore, we cannot exclude the possibility of compound inheritance of a *TBX6* null mutation and an alternative risk haplotype among persons of African ancestry in Patient PT03.

In the no-scoliosis subgroup, the T-C-A haplotype was present in only 5 of 30 persons (17%) (Table S7 in the Supplementary Appendix). The T-C-A risk haplotype was a significant risk factor for congenital scoliosis in series 3 (5 of 6 persons with congenital scoliosis vs. 5 of 30 persons without scoliosis,  $P=0.004$  by Fisher's exact test), which further supports the *TBX6* compound inheritance model of the disorder.

### THE HYPOMORPHIC RISK HAPLOTYPE IN VITRO

The *TBX6* SNPs defining the haplotype for risk of congenital scoliosis could be functional or, alternatively, could be a composite genetic marker that is linked with the true functional variants. We conducted a linkage-disequilibrium analysis using common human SNPs<sup>22</sup> and found that the linkage-disequilibrium block of *TBX6* is short (5 kb by the four-gamete rule) (Fig. S4 in the Supplementary Appendix). We identified no rare mutation in this block. The most parsimonious explanation for this observation is that the common *TBX6* SNPs defining the risk haplotype are functional.

The SNPs rs3809624 and rs3809627 are located in the 5' noncoding region of *TBX6* (Fig. 1B), potentially regulating *TBX6* expression.<sup>29</sup> Data from the Encyclopedia of DNA Elements (ENCODE) Project suggest the possible involvement of these two noncoding SNPs in the regulation of gene expression.<sup>30</sup> For example, both SNPs are located in the same binding site of upstream stimulatory factor 1 (USF1), a transcription factor potentially involved in somitogenesis.<sup>31,32</sup>

We used in vitro luciferase assays to study the effects of rs3809624 and rs3809627 on gene expression. Preliminary analyses in three human cell lines (HEK293, HepG2, and HeLa) showed significant reductions in the expression of the reporter gene when linked to the portion of *TBX6* harboring nonreference variants at both rs3809624 and rs3809627 (Fig. S5 and S6 in the Supplementary Appendix). We carried out the same experiment using the multipotent P19CL6 cells<sup>16,17</sup> and hiPSCs. *TBX6* was strongly expressed in these stem cells when they differentiated into mesodermal-cell lineages (Fig. S5 in the Supplementary Appendix). We observed that P19CL6 cells and hiPSCs carrying the luciferase reporter construct with nonreference alleles at both rs3809624 and rs3809627 expressed significantly less reporter protein on differentiation into mesodermal and cardiac cells (Fig. 3A). These findings consistently support the hypomorphic effects of the combination of nonreference alleles at rs3809624 and rs3809627 on *TBX6* expression.

### HEMIVERTEBRAE IN PERSONS WITH *TBX6*-ASSOCIATED CONGENITAL SCOLIOSIS

To assess the influence of *TBX6* disruption on the morphologic features of vertebral malformations, we evaluated the types of vertebral malformations in 23 persons with sporadic congenital scoliosis and *TBX6* null mutations and in 204 persons with sporadic congenital scoliosis and no detected *TBX6* nonsynonymous mutations. All 23 persons with *TBX6*-associated congenital scoliosis had one or more hemivertebrae (Table 1 and Fig. 2). The number of hemivertebrae was significantly higher in the persons with *TBX6*-associated congenital scoliosis than in those with congenital scoliosis but no apparent *TBX6* mutation ( $P<0.001$ ) (Table S8 in the Supplementary Appendix).



## DISCUSSION

The genetic basis for the risk of congenital scoliosis is poorly understood. The involvement of *Tbx6* in vertebral malformations in mice has been suggested.<sup>4,25,33</sup> However, a previous study involving 50 persons with congenital vertebral malformations identified no *TBX6* mutations,<sup>5</sup> and only a weak association between common *TBX6* SNPs and congenital scoliosis in the Han Chinese population was reported previously.<sup>26</sup> Our data show that *TBX6* null mutations are associated with sporadic congenital scoliosis, and in conjunction with the *TBX6* haplotype of T-C-A, these mutations accounted for 8 to 11% of congenital scoliosis cases in the study population with the disorder (Table 2). *TBX6* null mutations and noncoding variants therefore contribute substantively to the complex trait of sporadic congenital scoliosis.

Generally, the molecular mechanism of human mendelian disorders associated with heterozygous gene deletions or null mutations is haplo-insufficiency, in which half the gene dosage is not sufficient for normal function.<sup>34,35</sup> In contrast, the effect of diminished *TBX6* dosage in congenital scoliosis is complex. We found that an additional *TBX6* hypomorphic allele is required to cause a further decrement in gene-expression dosage beyond haploinsufficiency for penetrance of the congenital scoliosis phenotype (Fig. 3B).

A precedent for the mechanism of compound inheritance was provided by a recent study of human thrombocytopenia with absent radii (TAR) associated with the gene *RBM8A* in 1q21.1.<sup>36</sup> The compound inheritance of *TBX6* in congenital scoliosis, like that of *RBM8A* in TAR and analogous to a recessive trait, requires two mutant alleles at a genetic locus for trait manifestation. This model may also pertain to phenotypic variation and penetrance of particular endophenotypes manifested in other genomic disorders that are associated with large-scale copy-number variants.<sup>37-39</sup>

The phenotypes of sporadic congenital scoliosis can be dramatically different. However, the 23 persons whom we found to have *TBX6*-associated congenital scoliosis were phenotypically consistent with one another, having one or more hemivertebrae. A heterozygous *TBX6* stop-loss mutation (p.\*437Cext\*81) was reported in a pedigree (p.\*437Cext\*81) was reported in a pedigree with dominantly inherited spondylocostal dysostosis, a specific group of vertebral abnormalities characterized by multiple segmentation defects of the vertebrae, malalignment of the ribs (though often with a symmetric thoracic cage), and non-progressive scoliosis.<sup>40,41</sup> Functional studies showed that this mutation caused partial reduction in the ability of *TBX6* to activate transcription.<sup>41</sup> We further analyzed this previously reported pedigree and found that this *TBX6* stop-loss mutation was located within the T-C-A risk haplotype. The combined reduction in *TBX6* expression and transcriptional activation might therefore account for spondylocostal dysostosis in this family. Alternatively, the 81 amino acids added to the C-terminal may affect the ability of *TBX6* to interact with other proteins that are critical in vertebral formation, such as *MESP2* and *RIPPLY2*.<sup>42,43</sup>

Environmental factors could also contribute to the variety of phenotypes of *TBX6*-associated congenital scoliosis. Mouse embryos are susceptible to gestational hypoxia, and



developmental processes such as somitogenesis are affected by cellular hypoxia.<sup>4,44</sup> Moreover, gene–environment interactions can affect the penetrance of vertebral phenotypes in mouse embryos that are genetically predisposed to vertebral defects. Similar mechanisms could affect the phenotypes of persons with variants in and deletions of *TBX6*.<sup>4</sup> Our data suggest that *TBX6* mutations are a common genetic cause of congenital scoliosis in the populations from which we drew our samples. Our finding that compound inheritance of a rare *TBX6* null mutation and a common risk haplotype causes congenital scoliosis will enable the molecular diagnosis of the disorder and genetic counseling for some persons who have or are at risk for congenital scoliosis.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Authors

N. Wu, X. Ming, J. Xiao, Z. Wu, X. Chen, M. Shinawi, Y. Shen, G. Yu, J. Liu, H. Xie, Z.S. Gucev, S. Liu, N. Yang, H. Al-Kateb, J. Chen, Jian Zhang, N. Hauser, T. Zhang, V. Tasic, P. Liu, X. Su, X. Pan, C. Liu, L. Wang, Joseph Shen, Jianxiong Shen, Y. Chen, T. Zhang, Jianguo Zhang, K.W. Choy, Jun Wang, Q. Wang, S. Li, W. Zhou, J. Guo, Y. Wang, C. Zhang, H. Zhao, Y. An, Y. Zhao, Jiucun Wang, Z. Liu, Y. Zuo, Y. Tian, X. Weng, V.R. Sutton, H. Wang, Y. Ming, S. Kulkarni, T.P. Zhong, P.F. Giampietro, S.L. Dunwoodie, S.W. Cheung, X. Zhang, L. Jin, J.R. Lupski, G. Qiu, and F. Zhang

## Affiliations

Department of Orthopedic Surgery, Peking Union Medical College Hospital (N.W., Z.W., J.L., S. Liu, J.C., X.S., Jianxiong Shen, Jianguo Zhang, S. Li, Y.W., H.Z., Y. Zhao, Z.L., Y. Zuo, Y.T., X.W., G.Q.), the PET-CT Center, Cancer Hospital (Y.M.), and McKusick–Zhang Center for Genetic Medicine and State Key Laboratory of Medical Molecular Biology, Institute of Basic Medical Sciences (X.Z.), Peking Union Medical College and Chinese Academy of Medical Sciences, and the Beijing Municipal Key Laboratory of Child Development and Nutriomics (X.C., H.X., T.Z. [M.D.], J.G.), and Department of Neurology, Affiliated Children’s Hospital (L.W., Jun Wang), Capital Institute of Pediatrics, Beijing, the State Key Laboratory of Genetic Engineering and Ministry of Education Key Laboratory of Contemporary Anthropology (X.M., J.X., G.Y., N.Y., Jian Zhang, X.P., Y.C., Q.W., W.Z., C.Z., Y.A., Jiucun Wang, H.W., T.P.Z., L.J., F.Z.), and Collaborative Innovation Center for Genetics and Development (J.X., H.W., T.P.Z., L.J., F.Z.), School of Life Sciences, Fudan University, and the School of Medicine (Y.S.) and the Department of Gastroenterology, Hepatology, and Nutrition, Shanghai Children’s Hospital (T.Z. [M.D., Ph.D.]), Shanghai Jiao Tong University, Shanghai, the Department of Biotechnology, School of Basic Medical Science, Nanjing Medical University, Nanjing (C.L.), the Department of Obstetrics and Gynaecology, Chinese University of Hong Kong, Hong Kong (K.W.C.), and Fudan–Taizhou Institute of Health Sciences, Taizhou (Jiucun Wang, L.J., F.Z.) — all in China; the Department of

Pediatrics, Division of Genetics and Genomic Medicine (M.S.), and the Department of Pathology and Immunology (H.A.-K., S.K.), Washington University School of Medicine, St. Louis; the Department of Pathology, Harvard Medical School, and the Department of Laboratory Medicine, Boston Children's Hospital — both in Boston (Y.S.); the Department of Endocrinology and Genetics, University Children's Hospital, Medical Faculty Skopje, Skopje, Macedonia (Z.S.G., V.T.); the Department of Medical Genetics and Metabolism, Children's Hospital Central California, Madera (N.H., Joseph Shen); the Departments of Molecular and Human Genetics (P.L., V.R.S., S.W.C., J.R.L.) and Pediatrics (J.R.L.), Baylor College of Medicine, and Texas Children's Hospital (V.R.S., J.R.L.) — both in Houston; the Department of Pediatrics, University of Wisconsin–Madison, Madison (P.F.G.); and the Developmental and Stem Cell Biology Division, Victor Chang Cardiac Research Institute, and St. Vincent's Clinical School Faculty of Medicine and the School of Biotechnology and Biomolecular Sciences Faculty of Science, University of New South Wales — all in Sydney (S.L.D.)

## Acknowledgments

Supported in part by grants from the National Basic Research Program of China (2012CB944600 and 2011CBA00401), the National Natural Science Foundation of China (81222014, 81130034, 81100841, 31171210, 81271942, 81370708, 81171673, and 81371903), the 111 Project (B13016), the Beijing Natural Science Foundation (7144235), the Shu Guang Project (12SG08), the Shanghai Pujiang Program (10PJ1400300), the Recruitment Program of Global Experts, U.S. National Institute of Neurological Disorders and Stroke (R01NS058529), and the U.S. National Human Genome Research Institute (U54HG006542) and by an Australian National Health and Medical Research Council Senior Research Fellowship (ID1042002).

(Funded by the National Basic Research Program of China and others.)

We thank Yunzeng Zou for providing the P19CL6 cells, Yanli Zhu for neurodevelopmental evaluation, Bo Yin and Pei Ma for DNA extraction, Lu Chen for microarray design, Renqian Du for analysis of copy-number variants, and Sangang Xu for data analysis.

## Appendix


The authors' full names and academic degrees are as follows: Nan Wu, M.D., Xuan Ming, B.S., Jianqiu Xiao, Ph.D., Zhihong Wu, M.D., Xiaoli Chen, M.D., Ph.D., Marwan Shinawi, M.D., Yiping Shen, Ph.D., Guangju Yu, B.S., Jiaqi Liu, M.D., Hua Xie, M.D., Zoran S. Gucev, M.D., Ph.D., Sen Liu, M.D., Nan Yang, M.Sc., Hussam Al-Kateb, Ph.D., Jun Chen, Ph.D., Jian Zhang, M.Sc., Natalie Hauser, M.D., Ting Zhang, M.D., Ph.D., Velibor Tasic, M.D., Ph.D., Pengfei Liu, Ph.D., Xinlin Su, M.D., Xuedong Pan, B.S., Chunyu Liu, B.S., Liwen Wang, M.D., Joseph Shen, M.D., Jianxiong Shen, M.D., Yulin Chen, B.S., Ting Zhang, M.D., Jianguo Zhang, M.D., Kwong Wai Choy, Ph.D., Jun Wang, M.D., Qiqi Wang, B.S., Shugang Li, M.D., Weichen Zhou, Ph.D., Jin Guo, M.D., Yipeng Wang, M.D., Cheng Zhang, Ph.D., Hong Zhao, M.D., Yu An, Ph.D., Yu Zhao, M.D., Jiucun Wang, Ph.D., Zhenlei Liu, M.D., Yuzhi Zuo, M.D., Ye Tian, M.D., Xisheng Weng, M.D., V. Reid Sutton, M.D., Hongyan Wang, Ph.D., Yue Ming, M.D., Shashikant Kulkarni, Ph.D., Tao P. Zhong, Ph.D., Philip F. Giampietro, M.D., Ph.D., Sally L. Dunwoodie, Ph.D., Sau Wai Cheung, Ph.D., Xue Zhang, M.D., Ph.D., Li Jin, Ph.D., James R. Lupski, M.D., Ph.D., D.Sc., Guixing Qiu, M.D., and Feng Zhang, Ph.D.

## References

1. Wynne-Davies R. Congenital vertebral anomalies: aetiology and relationship to spina bifida cystica. *J Med Genet.* 1975; 12:280–8. [PubMed: 1100836]
2. Giampietro PF, Dunwoodie SL, Kusumi K, et al. Progress in the understanding of the genetic etiology of vertebral segmentation disorders in humans. *Ann N Y Acad Sci.* 2009; 1151:38–67. [PubMed: 19154516]
3. Pourquié O. Vertebrate segmentation: from cyclic gene networks to scoliosis. *Cell.* 2011; 145:650–63. [PubMed: 21620133]
4. Sparrow DB, Chapman G, Smith AJ, et al. A mechanism for gene-environment interaction in the etiology of congenital scoliosis. *Cell.* 2012; 149:295–306. [PubMed: 22484060]
5. Ghebranious N, Blank RD, Raggio CL, et al. A missense T (Brachyury) mutation contributes to vertebral malformations. *J Bone Miner Res.* 2008; 23:1576–83. [PubMed: 18466071]
6. Rosenfeld JA, Coppinger J, Bejjani BA, et al. Speech delays and behavioral problems are the predominant features in individuals with developmental delays and 16p11.2 microdeletions and microduplications. *J Neurodev Disord.* 2010; 2:26–38. [PubMed: 21731881]
7. Weiss LA, Shen Y, Korn JM, et al. Association between microdeletion and microduplication at 16p11.2 and autism. *N Engl J Med.* 2008; 358:667–75. [PubMed: 18184952]
8. Walters RG, Jacquemont S, Valsesia A, et al. A new highly penetrant form of obesity due to deletions on chromosome 16p11.2. *Nature.* 2010; 463:671–5. [PubMed: 20130649]
9. Jacquemont S, Reymond A, Zufferey F, et al. Mirror extreme BMI phenotypes associated with gene dosage at the chromosome 16p11.2 locus. *Nature.* 2011; 478:97–102. [PubMed: 21881559]
10. Shimojima K, Inoue T, Fujii Y, Ohno K, Yamamoto T. A familial 593-kb micro-deletion of 16p11.2 associated with mental retardation and hemivertebrae. *Eur J Med Genet.* 2009; 52:433–5. [PubMed: 19770079]
11. Fernandez BA, Roberts W, Chung B, et al. Phenotypic spectrum associated with de novo and inherited deletions and duplications at 16p11.2 in individuals ascertained for diagnosis of autism spectrum disorder. *J Med Genet.* 2010; 47:195–203. [PubMed: 19755429]
12. Shen Y, Chen X, Wang L, et al. Intra-family phenotypic heterogeneity of 16p11.2 deletion carriers in a three-generation Chinese family. *Am J Med Genet B Neuropsychiatr Genet.* 2011; 156:225–32. [PubMed: 21302351]
13. Al-Kateb H, Khanna G, Filges I, et al. Scoliosis and vertebral anomalies: additional abnormal phenotypes associated with chromosome 16p11.2 rearrangement. *Am J Med Genet A.* 2014; 164A:1118–26. [PubMed: 24458548]
14. Shen Y, Irons M, Miller DT, et al. Development of a focused oligonucleotide-array comparative genomic hybridization chip for clinical diagnosis of genomic imbalance. *Clin Chem.* 2007; 53:2051–9. [PubMed: 17901113]
15. Boone PM, Bacino CA, Shaw CA, et al. Detection of clinically relevant exonic copy-number changes by array CGH. *Hum Mutat.* 2010; 31:1326–42. [PubMed: 20848651]
16. Li L, Gong H, Yu H, et al. Knockdown of nucleosome assembly protein 1-like 1 promotes dimethyl sulfoxide-induced differentiation of P19CL6 cells into cardiomyocytes. *J Cell Biochem.* 2012; 113:3788–96. [PubMed: 22807403]
17. Gavrilov S, Nührenberg TG, Ashton AW, et al. Tbx6 is a determinant of cardiac and neural cell fate decisions in multipotent P19CL6 cells. *Differentiation.* 2012; 84:176–84. [PubMed: 22721678]
18. Okita K, Matsumura Y, Sato Y, et al. A more efficient method to generate integration-free human iPS cells. *Nat Methods.* 2011; 8:409–12. [PubMed: 21460823]
19. Beers J, Gulbranson DR, George N, et al. Passaging and colony expansion of human pluripotent stem cells by enzyme-free dissociation in chemically defined culture conditions. *Nat Protoc.* 2012; 7:2029–40. [PubMed: 23099485]
20. Lian X, Zhang J, Azarin SM, et al. Directed cardiomyocyte differentiation from human pluripotent stem cells by modulating Wnt/β-catenin signaling under fully defined conditions. *Nat Protoc.* 2013; 8:162–75. [PubMed: 23257984]


21. Giampietro PF, Raggio CL, Reynolds CE, et al. An analysis of PAX1 in the development of vertebral malformations. *Clin Genet*. 2005; 68:448–53. [PubMed: 16207213]
22. 1000 Genomes Project Consortium. Abecasis GR, Auton A, et al. An integrated map of genetic variation from 1,092 human genomes. *Nature*. 2012; 491:56–65. [PubMed: 23128226]
23. Rosenfeld JA, Coe BP, Eichler EE, Cuckle H, Shaffer LG. Estimates of penetrance for recurrent pathogenic copy-number variations. *Genet Med*. 2013; 15:478–81. [PubMed: 23258348]
24. Chapman DL, Papaioannou VE. Three neural tubes in mouse embryos with mutations in the T-box gene *Tbx6*. *Nature*. 1998; 391:695–7. [PubMed: 9490412]
25. White PH, Farkas DR, McFadden EE, Chapman DL. Defective somite patterning in mouse embryos with reduced levels of *Tbx6*. *Development*. 2003; 130:1681–90. [PubMed: 12620991]
26. Fei Q, Wu Z, Wang H, et al. The association analysis of *TBX6* polymorphism with susceptibility to congenital scoliosis in a Chinese Han population. *Spine (Phila Pa 1976)*. 2010; 35:983–8. [PubMed: 20228709]
27. MacArthur DG, Tyler-Smith C. Loss-of-function variants in the genomes of healthy humans. *Hum Mol Genet*. 2010; 19:R125–R130. [PubMed: 20805107]
28. Cheng Y, Miura RM, Tian B. Prediction of mRNA polyadenylation sites by support vector machine. *Bioinformatics*. 2006; 22:2320–5. [PubMed: 16870936]
29. Skelly DA, Ronald J, Akey JM. Inherited variation in gene expression. *Annu Rev Genomics Hum Genet*. 2009; 10:313–32. [PubMed: 19630563]
30. ENCODE Project Consortium. An integrated encyclopedia of DNA elements in the human genome. *Nature*. 2012; 489:57–74. [PubMed: 22955616]
31. Johnson DS, Mortazavi A, Myers RM, Wold B. Genome-wide mapping of in vivo protein-DNA interactions. *Science*. 2007; 316:1497–502. [PubMed: 17540862]
32. Fujimi TJ, Aruga J. Upstream stimulatory factors, *USF1* and *USF2* are differentially expressed during *Xenopus* embryonic development. *Gene Expr Patterns*. 2008; 8:376–81. [PubMed: 18585979]
33. Watabe-Rudolph M, Schlautmann N, Papaioannou VE, Gossler A. The mouse rib-vertebrae mutation is a hypomorphic *Tbx6* allele. *Mech Dev*. 2002; 119:251–6. [PubMed: 12464437]
34. Lupski JR, Stankiewicz P. Genomic disorders: molecular mechanisms for rearrangements and conveyed phenotypes. *PLoS Genet*. 2005; 1(6):e49. [PubMed: 16444292]
35. Cirulli ET, Goldstein DB. Uncovering the roles of rare variants in common disease through whole-genome sequencing. *Nat Rev Genet*. 2010; 11:415–25. [PubMed: 20479773]
36. Albers CA, Paul DS, Schulze H, et al. Compound inheritance of a low-frequency regulatory SNP and a rare null mutation in exon-junction complex subunit *RBM8A* causes TAR syndrome. *Nat Genet*. 2012; 44:435–9. [PubMed: 22366785]
37. Kurotaki N, Shen JJ, Touyama M, et al. Phenotypic consequences of genetic variation at hemizygous alleles: Sotos syndrome is a contiguous gene syndrome incorporating coagulation factor twelve (*FXII*) deficiency. *Genet Med*. 2005; 7:479–83. [PubMed: 16170239]
38. Wapner RJ, Martin CL, Levy B, et al. Chromosomal microarray versus karyotyping for prenatal diagnosis. *N Engl J Med*. 2012; 367:2175–84. [PubMed: 23215555]
39. Girirajan S, Rosenfeld JA, Coe BP, et al. Phenotypic heterogeneity of genomic disorders and rare copy-number variants. *N Engl J Med*. 2012; 367:1321–31. Erratum, *N Engl J Med* 2012;367:2362. [PubMed: 22970919]
40. Gucev ZS, Tasic V, Pop-Jordanova N, et al. Autosomal dominant spondylocostal dysostosis in three generations of a Macedonian family: negative mutation analysis of *DLL3*, *MESP2*, *HES7*, and *LFNG*. *Am J Med Genet A*. 2010; 152A:1378–82. [PubMed: 20503311]
41. Sparrow DB, McInerney-Leo A, Gucev ZS, et al. Autosomal dominant spondylocostal dysostosis is caused by mutation in *TBX6*. *Hum Mol Genet*. 2013; 22:1625–31. [PubMed: 23335591]
42. Hitachi K, Danno H, Kondow A, et al. Physical interaction between *Tbx6* and *mespb* is indispensable for the activation of bowline expression during *Xenopus* somitogenesis. *Biochem Biophys Res Commun*. 2008; 372:607–12. [PubMed: 18510946]

43. Kawamura A, Koshida S, Takada S. Activator-to-repressor conversion of T-box transcription factors by the Ripply family of Groucho/TLE-associated mediators. *Mol Cell Biol.* 2008; 28:3236–44. [PubMed: 18332117]
44. O'Reilly VC, Lopes Floro K, Shi H, et al. Gene-environment interaction demonstrates the vulnerability of the embryonic heart. *Dev Biol.* 2014; 391:99–110. [PubMed: 24657234]




**Figure 1. Effect of the 16p11.2 Deletion on *TBX6***

Panel A shows the region affected by the proximal 16p11.2 deletion based on the human genome assembly hg19. A pair of long direct genomic repeats (shown in orange) can mediate 0.6-Mb recurrent deletions. The genes affected by this deletion, including *TBX6* (circled), are shown. Panel B shows three common single-nucleotide polymorphisms (SNPs) in *TBX6*. Two SNPs, rs3809624 and rs3809627, are located in the 5' noncoding region, whereas rs2289292 is a synonymous SNP in the last exon. Panels C and D show two pedigrees with 16p11.2 deletions (SE1 and SE2). Squares denote male pedigree members, circles female pedigree members, solid symbols members with congenital scoliosis, and open symbols unaffected members; the probands are indicated by black arrows. The red bracket represents the deletion allele, and the nonreference alleles of rs2289292, rs3809624, and rs3809627 on the nondeletion chromosomes are shown in blue. The slash denotes a deceased family member. NA denotes not available.



**Figure 2. Spinal Radiographs in Patients with TBX6-Associated Congenital Scoliosis**  
The vertebrae with malformations are indicated by arrows. The patient number is shown above each radiograph.





**Figure 3. Results of In Vitro Functional Assays of Common TBX6 Variants and the Mechanistic Model of TBX6-Associated Congenital Scoliosis**

Panel A shows the results of luciferase reporter assays. Two cell types (P19CL6 cells and human induced pluripotent stem cells [hiPSCs]) with high levels of expression of TBX6 during induced differentiation were used. The nonreference alleles of rs3809624 and rs3809627 are shown in blue, and the risk haplotype is underlined. At least three independent experiments of normalized luciferase activity were quantified, with the graph showing mean values and their standard errors (I bars). Panel B shows a simplified genetic model of TBX6-associated congenital scoliosis. TBX6 expression is critical for normal vertebral formation (the reference allele of TBX6 is indicated by a black line). Heterozygous hypomorphic variants (shown in blue) cause only a moderate reduction in TBX6 expression. Even homozygous hypomorphic variants do not reduce TBX6 expression dramatically. Heterozygous TBX6 null mutations (indicated by red brackets) may reduce gene expression by one half. Independently, these mutations hardly reach the gene-dosage threshold for congenital scoliosis. In combination, however, a null allele and a hypomorphic allele of TBX6 cause further reductions in gene expression that may confer a high risk of congenital scoliosis.

Table 1

Genetic and Clinical Characteristics of Patients with *TBX6*-Associated Congenital Scoliosis.

Patient No.	Sex	Age (yr)	TBX6 Allele		Vertebral Malformation*	Other Congenital Malformation
			Null Mutation at First Allele	Risk Haplotype at Second Allele <sup>†</sup>		
<b>Congenital scoliosis series 1 (discovery)</b>						
XH004	F	4	16p11.2 del	T-C-A	L1 hemivertebra (left)	None
XH025	M	3	16p11.2 del	T-C-A	T11 hemivertebra (left)	Rib abnormality (missing left 11th rib)
XH042	M	6	16p11.2 del	T-C-A	T9 hemivertebra (left)	Patent foramen ovale and rib abnormalities (missing right 9th and left 12th ribs)
XH141	F	18	16p11.2 del	T-C-A	T4 hemivertebra (left)	Syringomyelia, spina bifida, and rib abnormalities (fused left 3rd through 6th ribs)
XH149	F	7	16p11.2 del	T-C-A	T12 hemivertebra (right)	Rib abnormality (missing left 12th rib)
XH186	M	4	16p11.2 del	T-C-A	T11 hemivertebra (left)	Rib abnormality (missing right 11th rib)
XH237	M	13	16p11.2 del	T-C-A	T12 hemivertebra (left) and L3 butterfly vertebra	Rib abnormality (missing right 12th rib)
XH265	M	4	16p11.2 del	T-C-A	L1 hemivertebra (right)	None
XH270	M	14	16p11.2 del	T-C-A	T11 and T12 hemivertebrae (left)	Rib abnormalities (missing right 12th rib and bifurcation of right 2nd rib)
XH292	F	11	16p11.2 del	T-C-A	T6 hemivertebra (left) and T12 wedge vertebra	Rib abnormality (missing right 6th rib)
XH300	M	3	16p11.2 del	T-C-A	T10 hemivertebra (left)	Rib abnormalities (missing right 10th rib and bifurcation of right 9th rib)
XH303	M	5	16p11.2 del	T-C-A	T7 butterfly vertebra and T12 hemivertebra (left)	Rib abnormality (missing left 12th rib)
XH101	F	7	c.1250_1251insT frame shift	T-C-A	L2 hemivertebra (left)	Atrial septal defect and rib abnormalities (missing bilateral 12th ribs)
XH122	M	5	c.266_267insC frame shift	T-C-A	Hemivertebra (left) between T12 and L1	Rib abnormality (missing right 12th rib)
XH170	M	5	c.704_705insG frame shift	T-C-A	T12 hemivertebra (left)	Rib abnormality (missing right 12th rib)
XH286	M	2	c.1169_1170insC frame shift	T-C-A	L2 hemivertebra (left) and T8 butterfly vertebra	None
XH148	F	16	c.844C→T (p.R282X)	T-C-A	T1 and L3 butterfly vertebrae, T4 hemivertebra (right), and T7 hemivertebra (left)	None
<b>Congenital scoliosis series 2 (replication)</b>						
XR345	F	13	16p11.2 del	T-C-A	Hemivertebra (left) between T12 and L1	None

Patient No.	Sex	Age (yr)	TBX6 Allele		Vertebral Malformation*	Other Congenital Malformation
			Null Mutation at First Allele	Risk Haplotype at Second Allele <sup>†</sup>		
XR353	M	11	16p11.2 del	T-C-A	T11 hemivertebra (right)	Rib abnormality (missing left 11th rib)
XR402	M	11	16p11.2 del	T-C-A	C5 hemivertebra (left), T5 hemivertebra (right), and T8 hemivertebra (left)	Rib abnormalities (missing left 5th and right 8th ribs)
XR434	F	16	16p11.2 del	T-C-A	T12 hemivertebra (left) and T9 butterfly vertebra	None
XR439	F	7	16p11.2 del	T-C-A	L3 hemivertebra (right)	None
XR341	M	2	c.1179_1180delAG frame shift	T-C-A	T12 hemivertebra (right)	Rib abnormality (missing left 12th rib)

\* C denotes cervical, L lumbar, and T thoracic.

<sup>†</sup>T-C-A represents the haplotype defined by three common *TBX6* single-nucleotide polymorphisms (reference/nonreference): rs2289292 (C/T), rs3809624 (T/C), and rs3809627 (C/A).

**Table 2**Distribution of Cases of *TBX6*-Associated Congenital Scoliosis.

Series	No. of Patients	Cases of Congenital Scoliosis Explained by <i>TBX6</i> Compound Inheritance	P Value
		<i>percent</i>	
Series 1: patients with congenital scoliosis	161	11	$<3.8 \times 10^{-6}$ *
Series 2: patients with congenital scoliosis	76	8	$<8.4 \times 10^{-4}$ †
Series 3: patients with 16p11.2 deletion	42‡	83	0.004§

\*The P value was calculated on the basis of 17 of 161 patients with congenital scoliosis versus 0 of 166 controls.

†The P value was calculated on the basis of 6 of 76 patients with congenital scoliosis versus 0 of 166 controls.

‡Six of 42 patients with 16p11.2 deletion had a congenital scoliosis phenotype.

§The P value was calculated on the basis of 5 of 6 deletion carriers with congenital scoliosis versus 5 of 30 deletion carriers without scoliosis.