

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

NEUTRAL CURRENT PHENOMENOLOGY OF SUPERSYMMETRIC  $SU(2) \times U(1) \times U(1)$  MODELS

### Permalink

<https://escholarship.org/uc/item/4vg9h7s2>

### Author

Kang, Y.S.

### Publication Date

1982-06-01



# Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Physics, Computer Science &  
Mathematics Division

RECEIVED  
LAWRENCE  
BERKELEY LABORATORY

JUN 2 1982

LIBRARY AND  
DOCUMENTS SECTION

Submitted for publication

NEUTRAL CURRENT PHENOMENOLOGY OF SUPERSYMMETRIC  
 $SU(2) \times U(1) \times \tilde{U}(1)$  MODELS

Young S. Kang

June 1982



LBL-14656  
c.2

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

June 1982

LBL-14656

Neutral Current Phenomenology  
of  
Supersymmetric  $SU(2) \times U(1) \times \tilde{U}(1)$  Models\*

Young S. KANG

Lawrence Berkeley Laboratory  
and  
Department of Physics  
University of California  
Berkeley, California 94720

ABSTRACT

The neutral current sector of a class of supersymmetric  $SU(2) \times U(1) \times \tilde{U}(1)$  models is parametrized. Bounds on the neutral boson masses are obtained from the low energy data, and the implications of future experimental findings for these models are discussed.

---

\* This work was supported by the Director, Office of Energy Research, Office of High Energy and Nuclear Physics, Division of High Energy Physics of the U.S. Department of Energy under Contract DE-AC03-76SF00098.

In some models with spontaneously broken supersymmetry, a new  $\tilde{U}(1)$  gauge symmetry is introduced to give large masses to the unseen scalar partners of leptons and quarks [1, 2]. In addition to the requirements on theoretical consistency such as the correct vacuum structure and anomaly-freedom, model-building of this sort is subject to phenomenological constraints, as the modified neutral current structure may spoil the success of the standard model.

It is the purpose of this paper to perform the same type of phenomenological analysis on a class of supersymmetric  $SU(2) \times U(1) \times \tilde{U}(1)$  models as the variously motivated multi-boson electroweak models of the past [3]. Bounds on the masses of the neutral bosons are obtained from the data for neutrino scattering, electron-nucleon interactions, and electron-positron annihilation.

Any supersymmetric  $\tilde{U}(1)$  model with a realistic mass spectrum should possess the following minimum features:

(I) The axial part of the  $\tilde{U}(1)$  charge of a 4-component fermion is greater in magnitude than its vector part. This allows the fermion to be lighter than both of its scalar partners.

(II) There are at least two Higgs isodoublets acquiring non-zero VEV's to give mass to both charge  $\frac{2}{3}$  and  $-\frac{1}{3}$  quarks through Yukawa terms at the tree level.

A simple supersymmetric generalization of the standard model particle assignment satisfies (I); left-handed matter supermultiplets consist of  $Q_L = \begin{pmatrix} U_L \\ D_L \end{pmatrix}$ ,  $\bar{U}_R$ ,  $\bar{D}_R$ ,  $L_L = \begin{pmatrix} N_L \\ E_L \end{pmatrix}$ ,  $\bar{E}_R$ , all of which have the same  $\tilde{U}(1)$  charge,  $\frac{\tilde{y}}{2}$ . Quarks and charged leptons then have the identical, purely axial  $\tilde{U}(1)$  charge,  $\tilde{y}$ , and  $\nu$  has a V-A coupling,  $\frac{\tilde{y}}{2}$ , to  $\tilde{U}(1)$ . Any extension of this minimal fermionic structure, leading to a proliferation of parameters, finds little theoretical motivation in most

models, and is not considered here [F1].

The simplest Higgs structure satisfying (II) consists of two Higgs isodoublets  $\phi_1$  and  $\phi_2$ , whose U(1) charges are  $\frac{1}{2}$  and  $-\frac{1}{2}$ , and whose neutral components obtain the same VEV. The  $\hat{U}(1)$  sector then remains detached from SU(2) x U(1). It was observed by Fayet [6], however, that given the fermionic  $\hat{U}(1)$  couplings outlined above, this minimal Higgs structure leads to unacceptable values for axial parameters in neutrino scattering.

We seek extension of this minimal model in two ways [F2]. First, we allow mixing between  $\hat{U}(1)$  and SU(2) x U(1) by having  $\langle \phi_1 \rangle = \begin{pmatrix} 0 \\ h_1 \end{pmatrix}$ ,  $\langle \phi_2 \rangle = \begin{pmatrix} h_2 \\ 0 \end{pmatrix}$  with  $h_1 \neq h_2$  in general. This defines the Minimal Mixing Model (MMM) which is probably the most economical supersymmetric  $\hat{U}(1)$  model that can be made consistent with the present data. Secondly, we allow Higgs fields  $\phi_0^i$  ( $i = 1$  to  $N$ ) which are singlets under the standard gauge group and have  $\hat{U}(1)$  charges  $\tilde{y}_0^i$ , to obtain VEV's  $h_0^i$ . This is the situation often found in actual models [2]. We call the class of models with both mixing and  $\phi_0^i$  the Extended Mixing Model (EMM).

Now we parametrize the neutral current sector of EMM, of which MMM is a special case. The neutral gauge boson mass<sup>2</sup> matrix is:

$$M^2 = \frac{1}{4} (h_1^2 + h_2^2) \begin{matrix} & B^\mu & W_3^\mu & \hat{B}^\mu \\ \begin{bmatrix} g^2 & -gg_2 & -gg\bar{\epsilon} \\ -gg_2 & g_2^2 & g_2\bar{g}\bar{\epsilon} \\ -gg\bar{\epsilon} & g_2\bar{g}\bar{\epsilon} & \bar{g}^2 t \end{bmatrix} & & & \end{matrix} \quad (1)$$

where  $\bar{g} = 2\tilde{y}g$ ,  $\bar{\epsilon} = (h_1^2 - h_2^2)/(h_1^2 + h_2^2)$ ,  $t = 1 + \sum_1^N (\tilde{y}_0^i h_0^i)^2 / [\tilde{y}^2 (h_1^2 + h_2^2)]$ , and  $g, g_2, \tilde{g}$  are the couplings and  $B^\mu, W_3^\mu, \hat{B}^\mu$  the boson fields associated with U(1), SU(2),  $\hat{U}(1)$ , respectively.

The mass eigenstates are written as follows:

$$\begin{pmatrix} A^\mu \\ Z^\mu \\ \tilde{Z}^\mu \end{pmatrix} = \begin{pmatrix} \cos\theta & \sin\theta & 0 \\ -\cos\alpha \sin\theta & \cos\alpha \cos\theta & \sin\alpha \\ \sin\alpha \sin\theta & -\sin\alpha \cos\theta & \cos\alpha \end{pmatrix} \begin{pmatrix} B^\mu \\ W_3^\mu \\ \tilde{B}^\mu \end{pmatrix}, \quad (2)$$

where  $\sin^2\theta = g^2 / (g^2 + g_2^2)$  is the standard model mixing parameter, and  $\cos^2\alpha = \frac{1}{2} \{ 1 + [1 + 4A \epsilon^2 / (1 - At)^2]^{-1/2} \}$ ,  $A = g^2 / (g^2 + g_2^2)$ .

One of the eigenvalues of the matrix (1) is of course zero, corresponding to the photon, and the other two are given as follows:

$$\begin{pmatrix} \rho \\ \tilde{\rho} \end{pmatrix} = \frac{1}{2} \{ 1 + At \pm (1 - At) [1 + 4A \epsilon^2 / (1 - At)^2]^{1/2} \}, \quad (3)$$

where  $\rho = \frac{m_Z^2 \cos^2\theta}{m_W^2}$ ,  $\tilde{\rho} = \frac{m_{\tilde{Z}}^2 \cos^2\theta}{m_W^2}$ , i.e., mass squared in units

of the standard model Z mass squared. Notice that  $(1 - \rho)(1 - \tilde{\rho}) \leq 0$ , which satisfies the Georgi-Weinberg theorem [7], although its premise is not met.

There are six independent parameters for EMM:  $g, g_2, \tilde{g}, h_1, h_2, t$ . Apart from  $e = gg_2 / (g^2 + g_2^2)^{1/2}$ ,  $G_F = 1 / [\sqrt{2}(h_1^2 + h_2^2)]$ , we choose to work with the following four free parameters:  $\sin^2\theta$ , as in the standard model, and  $A \geq 0$ ,  $|\epsilon| \leq 1$ ,  $t \geq 1$ , which measure the relative strength of the  $\tilde{U}(1)$  coupling, the degree of mixing between the two massive neutral bosons, and the isosinglet Higgs contribution to boson masses, respectively.

In order to compare the predictions of these models with the data, we make use, as usual, of low energy model independent parameters [8] defined in the following processes.

(i) Neutrino interactions:

$$\mathcal{L}_{\text{eff}} = - \frac{G_F}{\sqrt{2}} \bar{\nu} \gamma^\alpha (1 - \gamma_5) \nu \bar{f} \gamma_\alpha (g_V^f + g_A^f \gamma_5) f,$$

where  $f = e, u, \text{ or } d$ .

(ii) Asymmetry ( $A_{eD}$ ) in electron-deuteron scattering:

$$\frac{A_{eD}}{Q^2} = - \frac{9 G_F}{20\sqrt{2}\pi\alpha_{EM}} \left[ a + \frac{1 - (1-y)^2}{1 + (1-y)^2} b \right].$$

Atomic experiments provide further, if not complete, information on the four parameters describing parity-violating electron-quark reactions, but are not considered here.

The expressions for the following eight combinations of the above model independent parameters have been derived in terms of the model parameters, and listed in Table 1:

$$g_V^e, g_A^e, \alpha \equiv g_V^u - g_V^d, \beta \equiv g_A^u - g_A^d, \gamma \equiv g_V^u + g_V^d, \delta \equiv g_A^u + g_A^d, a, c \equiv a + \frac{b}{4}.$$

Also shown are the corresponding standard model expressions, and the data from various neutrino interactions and SLAC eD experiments.

We notice from Table 1 that  $A$  does not enter into these expressions, since they are, in the zero momentum transfer limit, functions of  $\tilde{g}^2/m_{\tilde{Z}}^2$  which is independent of  $\tilde{g}$ . Information on the strength of the  $\tilde{U}(1)$  coupling will come from  $e^+e^- \rightarrow \mu^+\mu^-$  at PETRA energies.

In fitting the model parameters  $\sin^2\theta, \epsilon, t$  to data, we first consider MMM ( $t = 1$ ). That non-zero mixing is needed in this case as mentioned earlier is seen immediately from the expressions for  $g_A^e$  and  $\delta$  (the isoscalar axial coupling); for  $\epsilon = 0$ , they are more than 4  $\sigma$ 's off. The other neutrino parameters (vector, and isovector axial) and the eD parameters in this non-mixing case are the same as in the standard model, since the  $e, u, d$   $\tilde{U}(1)$  neutral currents are purely axial.

It is not a priori clear that there should exist a region in the parameter space consistent with the data, since for no value of  $(A, \sin^2\theta, \epsilon)$  MMM reduces to the standard model. However, the best fit gives  $\epsilon = -.44, \sin^2\theta = .295$ , for which agreement with the data is good



for all the eight parameters, as shown in the last column of Table 1. The location of the best fit is not surprising, since for  $\epsilon = -\frac{1}{2}$  all  $\nu$  parameters are reduced to their standard model expressions, and the change in eD parameters is compensated by a larger value for  $\sin^2\theta$  than in the standard model, as seen in Fig. 1. This value of  $\sin^2\theta$  may not be a problem in view of the absence of a realistic supersymmetric GUT including  $\tilde{U}(1)$ .

If some isosinglet Higgs obtain VEV's (EMM), for large  $t$  the standard model is asymptotically recovered, as in models where the scale of the  $\tilde{U}(1)$  breaking is made large compared to  $O(100)$  GeV. The best fit for EMM parameters to data gives  $t$  of the order of a few hundred, which corresponds to  $h_0^i \sim h_{1,2} \times O(10)$ , unless  $N$  is large.

This value of  $t$  may seem too large in view of the theoretical prejudice regarding naturalness. However, the value of  $t$  is very sensitive to data, and it turns out that the current data can be accommodated for a wide range of  $t$  (including  $t \sim 5$ ) as well as or better than in the standard model. The present experimental accuracy thus places virtually no useful constraint on the allowed range of  $t$ . It should be noted that for smaller  $t$  the allowed range of  $\epsilon$  becomes more restricted, which has implications for the upper bound on the lower boson mass.

As an illustration of these points, we consider the case where  $\epsilon = -\frac{1}{2}$  for which all the  $\nu$  parameters are the same as in the standard model, whereas eD parameters depend on an additional parameter  $t$ . In this respect, this model resembles the model of Deshpande and Iskandar [9]. Taking the one-parameter fit to the  $\nu$  data from previous analyses [8],  $\sin^2\theta = .239 \pm .010$ , we find  $t = 7.85 \pm_{-3.35}^{12.9}$  from the best fit to SLAC eD data. Fig. 2 shows the sensitivity of  $t$  to data.

Since  $e$  and  $\mu$  have axial  $\hat{U}(1)$  charges, the exchange of a new gauge boson contributes to the forward-backward asymmetry ( $A^{\mu\mu}$ ) in  $e^+e^- \rightarrow \mu^+\mu^-$ . The interference terms between the photon and the massive boson exchanges give, to lowest order,

$$A^{\mu\mu} = - \frac{3\sigma}{32 \sin^2\theta \cos^2\theta} \left\{ \frac{(1+t+2\epsilon) - \sigma}{t(1-\sigma) - \epsilon^2} + \left[ \frac{\sigma(1+\epsilon-\sigma)^2}{t(1-\sigma) - \epsilon^2} \right] \frac{1}{[t(1-\sigma) - \epsilon^2]A - (\sigma - \sigma^2)} \right\}, \quad (4)$$

where 
$$\sigma = \frac{s \cos^2\theta}{m_w^2}$$

The dependence on  $A$  is isolated in the second term in braces, which is always positive for the CM energy ( $\sqrt{s}$ ) less than the smaller boson mass, including where the above approximation is valid. The minimum of  $|A^{\mu\mu}|$  (corresponding to  $A \rightarrow \infty$ ) is larger than  $|A^{\mu\mu}|$  in the standard model for the same value of  $\sin^2\theta$ . For the values of the MMM parameters determined earlier, and for  $\sqrt{s} = 34$  GeV, (4) becomes

$$A^{\mu\mu} = -(.117 + \frac{.005}{A - .226}) \quad (5)$$

While some experimental groups (CELLO [10], MARK J [11]) reported data consistent with the standard model prediction, other groups (TASSO [12], JADE [13]) find the asymmetry considerably more negative. In the former case, the errors are not small enough to rule out MMM, and  $A$  will have a lower bound. In the latter, if we take the TASSO data ( $A^{\mu\mu} = -.16 \pm .03$ ), for example, the bounds on  $A$  are:

$.29 \leq A \leq .61$ , which corresponds to  $43 \text{ GeV} \leq m_z \leq 53 \text{ GeV}$ , and  $85 \text{ GeV} \leq m_z \leq 90 \text{ GeV}$ . Fig. 3 shows the minimum  $|A^{\mu\mu}|$  for MMM, along with the data from various groups.

For EMM, any appreciable deviation ( $\Delta$ ) from the standard model pre-

diction for  $A^{\mu\mu}$  favors small  $t$  ( $< 5$ ). This is because for large  $t$ ,  $A \sim \frac{\text{const.}}{t^2 \Delta} + \frac{\sigma}{t}$ , so that unless  $\Delta$  is very small,  $A \sim \frac{\sigma}{t}$ , which implies that the current CM energy is very close to the lower boson mass. Thus the version of EMM given before ( $\epsilon = -\frac{1}{2}$ ), for example, would be ruled out by the TASSO or JADE data. If small  $t$  is indeed preferred, large mixing ( $\epsilon$ ) could put a nontrivial upper bound on the lower boson mass, since  $\tilde{\rho} \leq 1 - \frac{\epsilon^2}{t}$ .

The contribution to  $(g - 2)/2$  of the muon from the two neutral bosons in EMM is given by:

$$\frac{m_\mu^2 G_F}{24\sqrt{2} \pi^2} \frac{1}{t - \epsilon^2} [(1 - 4 \sin^2 \theta) t - 5(1 + t - 2\epsilon)] \quad (6)$$

For EMM no useful bound on  $t$  is obtained, again because of the sensitivity of  $t$  to experimental errors. For MMM, in a wide range of  $\epsilon$  including the best fit from low energy data quoted earlier, (6) is more negative and cancels more of the charged weak current contribution (which has the opposite sign) than in the standard model. Thus no useful constraints on the model parameters are obtained from the current data for the muon anomalous magnetic moment.

In conclusion, the large mixing (both  $\epsilon$  and  $\sin^2 \theta$ ) required in MMM places the upper bound of 74 GeV on the mass of the lighter boson. Depending on which  $e^+e^- \rightarrow \mu^+\mu^-$  asymmetry data we take, the prediction for the lower boson mass can be as low as 43 GeV. If a neutral boson does not turn up in the upcoming  $p\bar{p}$  experiments until around 90 GeV, MMM is probably ruled out.

As for EMM, if the asymmetry is indeed considerably larger than predicted in the standard model, the isosinglet contribution to the boson mass does not dominate. This further implies that the mixing ( $\epsilon$ )

would have to be small if a low-mass boson is not found. On the other hand, if the asymmetry is not large, EMM survives for a wide range of  $t$ , regardless of the location of the lightest boson.

#### ACKNOWLEDGEMENTS

We thank Ian Hinchliffe for the suggestion of this problem and discussions, and Bob Cahn for encouragement and valuable comments. This work was supported by the Director, Office of Energy Research, Office of High Energy and Nuclear Physics, Division of High Energy Physics of the U.S. Department of Energy under Contract DE-AC03-76SF00098.

## REFERENCES

- [1] P. Fayet, Phys. Lett. 69B (1977) 489; 70B (1977) 461.
- [2] L. J. Hall and I. Hinchliffe, Phys. Lett. 112B (1982) 351.
- [3] See, for example, E. H. de Groot, G. J. Gounaris and D. Schildknecht, Phys. Lett. 85B (1979) 399; N. G. Deshpande and R. J. Johnson, Oregon preprint, OITS-188 (1982).
- [4] M. Suzuki, Berkeley preprint, UCB - PTH 82/7 (1982).
- [5] J. Ellis and D. V. Nanopoulos, Phys. Lett. 110B (1982) 44.
- [6] P. Fayet, Phys. Lett. 95B (1980) 285; 96B (1980) 83.
- [7] H. Georgi and S. Weinberg, Phys. Rev. D17 (1978) 275.
- [8] For a review, see P. Q. Hung and J. J. Sakurai, Ann. Rev. Nucl. Part. Sci. 31 (1981) 375; J. E. Kim, P. Langacker, M. Levine, H. H. Williams, Rev. Mod. Phys. 53 (1981) 211.  
Note our axial parameters have the opposite sign from theirs.
- [9] N. G. Deshpande and D. Iskandar, Phys. Lett. 87B (1979) 383.
- [10] CELLO Collab., DESY 82-019.
- [11] MARK J Collab., MIT Technical Report #124 (1982).
- [12] TASSO Collab., DESY 82-002.
- [13] JADE Collab., DESY 81-072.

## FOOTNOTES

F1: In fact, there are arguments suggesting that the scalar partners of  $u$  and  $d$  (pairwise) [4], and also of different generation copies [5] are almost degenerate in mass. They support in part the minimal fermionic structure considered here.

F2: Fayet [6] considered isosinglets and vector fermionic  $\hat{U}(1)$  couplings, and made observations relevant to the case with no mixing, based on the neutrino and  $eD$  data.

Table 1.

Model Independent Parameters	SU(2) x U(1) x $\tilde{U}(1)$	Standard Model	Data [8]	MMM Best Fit
$g_V^e$	$-\frac{1}{2}(1-4x)\frac{t+\epsilon/2}{t-\epsilon^2}$	$-\frac{1}{2}(1-4x)$	$.06 \pm .08$	.09
$g_A^e$	$\frac{1}{2}\frac{t-(1+\epsilon)/2}{t-\epsilon^2}$	$\frac{1}{2}$	$.52 \pm .06$	.45
$\alpha$	$(1-2x)\frac{t+\epsilon/2}{t-\epsilon^2}$	$1-2x$	$.589 \pm .067$	.396
$\beta$	$-\frac{t+\epsilon/2}{t-\epsilon^2}$	-1	$-.937 \pm .062$	-.967
$\gamma$	$-\frac{2}{3}x\frac{t+\epsilon/2}{t-\epsilon^2}$	$-\frac{2}{3}x$	$-.273 \pm .081$	-.190
$\delta$	$-\frac{1/2+\epsilon}{t-\epsilon^2}$	0	$-.101 \pm .093$	-.074
a	$(1-\frac{20}{9}x)\frac{t-\epsilon}{t-\epsilon^2}$	$1-\frac{20}{9}x$	$.60 \pm .16$	.62
c	$(1-\frac{20}{9}x)\frac{t-\epsilon}{t-\epsilon^2}$ $+ (\frac{1}{4}-x)\frac{t+\epsilon/3}{t-\epsilon^2}$	$\frac{5}{4}-\frac{29}{9}x$	$.53 \pm .05$	.57

Note:  $x \equiv \sin^2\theta$

## FIGURE CAPTIONS

Fig. 1: Regions in the  $\epsilon - \sin^2\theta$  plane consistent with data for  $\nu$  scattering (I) and SLAC eD experiments (II). The best fit to MMM is also shown.

Fig. 2: The shaded region in the  $\sin^2\theta - t$  plane is allowed by eD data to within one standard deviation for EMM with  $\epsilon = -\frac{1}{2}$ . The best fit for  $t$  as a function of  $\sin^2\theta$  is also shown.

Fig. 3: Predictions for the asymmetry in the standard model and in MMM. The shaded region is allowed by MMM, and the solid line corresponds to the minimum  $|A^{\mu\mu}|$ . Also shown are the data from various groups.






Fig. 1




Fig. 2




Fig. 3

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

7      210  
7      210

TECHNICAL INFORMATION DEPARTMENT  
LAWRENCE BERKELEY LABORATORY  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720