

Lawrence Berkeley National Laboratory

Recent Work

Title

RESOLUTION OF THE E--MASS ANOMALY

Permalink

<https://escholarship.org/uc/item/4vh2j80v>

Authors

Barkas, Walter H.

Dyer, John N.

Heckman, Harry H.

Publication Date

1963-05-24

University of California
Ernest O. Lawrence
Radiation Laboratory

RESOLUTION OF THE Σ^- -MASS ANOMALY

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

10837
ay 2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA
Lawrence Radiation Laboratory
Berkeley, California
Contract No. W-7405-eng-48

RESOLUTION OF THE Σ^- -MASS ANOMALY

Walter H. Barkas, John N. Dyer, and Harry H. Heckman

May 24, 1963

RESOLUTION OF THE Σ^- -MASS ANOMALY

Walter H. Barkas, John N. Dyer, and Harry H. Heckman

Lawrence Radiation Laboratory
University of California
Berkeley, California

May 24, 1963

The masses of the charged Σ -hyperons and the K^- meson are determined most accurately from range measurements in emulsion on the products of the following processes: 1) $\Sigma^+ \rightarrow p + \pi^0$, 2) $\Sigma^+ \rightarrow n + \pi^+$, 3) $K^- + p \rightarrow \Sigma^+ + \pi^-$, and 4) $K^- + p \rightarrow \Sigma^- + \pi^+$. Prior to 1960 the masses were determined, but not overdetermined, from the ranges of the charged baryons alone. The Σ^+ mass was found from the proton range in reaction (1). The K^- mass was determined from the Σ^+ range in reaction (3), and the $\Sigma^- - \Sigma^+$ mass difference was deduced from the $\Sigma^+ - \Sigma^-$ range difference in reactions (3) and (4). For such measurements, events are used in which the Σ^+ decays at rest and in which the K^- is captured by a free proton after it comes to rest.

In 1959-1960 we began measurements of the pion ranges with a dual purpose: to confirm the measurements by overdetermining the masses, and to study further the high-velocity range-energy relation in emulsion. We soon found¹ that there was an apparent lack of momentum balance in reaction (4), although the other reactions seemed to balance satisfactorily.

We have recently carried out new measurements in a large emulsion stack. The results are practically identical with those reported previously, and we cannot doubt now that a real effect exists.^{1,2,3} In this letter we present our data, and propose that a new physical effect is the reason for the measurement contradictions. We suggest that slow negative hyperons lose energy at a lower rate than do positive particles of the same velocity.

This means that the existing range-energy (RE) relation,^{4,5} which is based on positive particle ranges, will not give correct values for quantities derived from the Σ^+ range in (4). We are led to this conclusion after rejecting several alternative possibilities, none of which could be made at all compatible with our understanding of the measurement errors.

Our ranges in emulsion of specific gravity 3.815 are given in Table I.

Table I. Observed Ranges

Process	Measured	Mean Range	Number of emulsion stacks used
1. $\Sigma^+ \rightarrow p + \pi^0$	144 p	1677 $\pm 2\mu$	5
2. $\Sigma^+ \rightarrow n + \pi^+$	48 π^+	92.74 ± 0.34 mm	2
3. $K+p \rightarrow \Sigma^+ + \pi^-$	40 Σ^+	818.8 $\pm 1.7\mu$	6
	24 π^-	88.58 ± 0.51 mm	3
4. $K+p \rightarrow \Sigma^- + \pi^+$	94 Σ^-	708.9 $\pm 1.5\mu$	6
	63 π^+	78.45 ± 0.25 mm	3

The standard deviation of each range is in close agreement with Bohr's straggling theory except in the case of the Σ^- , where theory predicts 10μ and we observe 15μ .

The pion ranges were measured independently by two observers with an automated coordinate readout microscope. Some ranges were also measured graphically by plotting the grid coordinates (a 1mm grid is photographed on each pellicle) of the pion track. The two methods of measurement agreed within 1% for each track. The baryon ranges were

measured in accord with our previously established⁶ standards. The data from all stacks are consistent. By careful measurements, the collinearity of tracks from processes (3) and (4) also was tested. There is, in any case, no possibility of mistaking K^- -capture by a bound proton for one by a free proton because of the large difference in Q -values.

The low velocity ($\beta < 0.28$) portion of the proton RE relation has been much studied, and in connection with the present program, was rechecked. There is less empirical data on which to base the high velocity range table needed for the pions. The original tables^{4,6} were calculated from theory using a mean excitation potential, I , of 331 eV and the shell corrections calculated by Walske⁷. Later high velocity data by Barkas and von Friesen⁸ gave a range of I values from 304 to 328 eV, and other information cited by them indicated that 331 eV might be somewhat high. In the meantime range tables with more elaborate shell corrections at a series of I -values were prepared.⁹ These tables and the measurements of Table I now enable us to improve our estimate of the emulsion I -value.

Reaction (1) uses only the well established portion of the range-energy curve. From it we obtain for the mass of the Σ^+ hyperon: 1189.36 ± 0.17 MeV. In order to obtain the same Σ^+ mass from reaction (2) as from reaction (1) we assume that I is an adjustable parameter. To obtain agreement between reactions (1) and (2) we must take $I = 319 \pm 9$ eV. We also require momentum balance between the hyperon and pion in reaction (3). To bring the pion momentum into agreement with that deduced from the Σ^+ range and mass, we must choose $I = 329 \pm 20$ eV. The best estimate

is therefore $I = 321 \pm 8$ eV. It is notable that raising the I-value raises the calculated mass from reaction (2) and lowers it in reaction (3).

In what follows we used the existing^{4,5} RE tables for $\beta < 0.28$, and for high velocities use the new table of reference 9 with $I = 320$ eV. They agree at $\beta = 0.28$. We estimate the uncertainty of the tabulated ranges at high velocities to be $\pm 1/2\%$.

Derived Masses

We combine the Σ^+ mass estimate from reaction (2) (which is partially dependent on reaction (1)) with the value from reaction (1) to obtain a best estimate of the Σ^+ mass. It is 1189.35 ± 0.15 MeV. The K^- mass is best obtained from the Σ^+ range in process (3). For it we find 493.7 ± 0.3 MeV.

The Σ^- hyperon mass is now determined from the pion ranges of reactions (3) and (4) and the mass of the Σ^+ . We find

$$M_{\Sigma^-} = \left[2(M_P + M_K)(T_{\pi^-} - T_{\pi^+}) + M_{\Sigma^+}^2 \right]^{1/2} = 1197.6 \pm 0.5 \text{ MeV.}$$

The error includes the uncertainties in the Σ^+ and K^- masses, and the uncertainty in the pion energy difference $(T_{\pi^-} - T_{\pi^+})$. The Σ^- mass is insensitive to the RE relation and to the K^- mass.

This result changes substantially our mass estimate for the Σ^- hyperon and also that for the Σ^0 , which is tied to it. (The new value for Σ^0 is 1193.2 ± 0.7 MeV.) In a recent tabulation¹⁰ of particle masses a footnote reference to the anomaly was made, but the Σ^- mass was derived using its own range in reaction (4).

The Energy-Loss Defect of the Negative Hyperon

The momentum of the pion in reaction (4) is 172.74 ± 0.38 Mev/c. Were the stopping cross sections of positive and negative particles equal, a Σ^- hyperon with this momentum and the above mass would have a range of $684 \pm 5\mu$. We observe $703.9 \pm 1.5\mu$.

It is well known that stopping theory based on the first Born approximation fails when the particle velocity becomes comparable to the velocities of many of the electrons in the stopping material. It is perhaps not surprising that there should appear a small difference between the stopping cross sections of positive and negative particles in this region, because their predicted equality is also a first Born approximation result, while a difference exists in the second approximation. At high velocities this effect is not expected. At intermediate velocities it is interesting to note that this effect probably is the explanation for an apparent difference between the positive and negative pion masses, which has remained unexplained for many years^{11,12}. We know of no other measurements of sufficient accuracy for the effect to have been observed before. A quantitative estimation of the effect is being attempted.

REFERENCES

1. J. N. Dyer, W. H. Barkas, H. H. Heckman, C. J. Mason, N. A. Nickols, and F. M. Smith, Bull. Am. Phys. Soc. 5, 4 (1960).
2. W. H. Barkas, J. N. Dyer, and H. H. Heckman, Bull. Am. Phys. Soc. 7, 469 (1962).
3. J. N. Dyer, Ph.D. Thesis, University of California Lawrence Radiation Laboratory Report UCRL-9450, 1960 (unpublished).

