Lawrence Berkeley National Laboratory

Recent Work

Title

Microbial community structure with trends in methylation gene diversity and abundance in mercury-contaminated rice paddy soils in Guizhou, China.

Permalink

<https://escholarship.org/uc/item/4xh9c1q5>

Journal

Environmental science. Processes & impacts, 20(4)

ISSN

2050-7887

Authors

Vishnivetskaya, Tatiana A
Hu, Haiyan
Van Nostrand, Joy D
et al.

Publication Date

2018-04-01

DOI

10.1039/c7em00558j

Peer reviewed

Microbial community structure with trends in methylation gene diversity and abundance in mercury-contaminated rice paddy soils in Guizhou, China

[Tatiana A. Vishnivetskaya](#),^a [Haiyan Hu](#),^{bc} [Joy D. Van Nostrand](#),^d [Ann M. Wymore](#),^a [Xiaohang Xu](#),^c [Guangle Qiu](#),^c [Xinbin Feng](#),^c [Jizhong Zhou](#),^d [Steven D. Brown](#),^a [Craig C. Brandt](#),^a [Mircea Podar](#),^a [Baohua Gu](#)^{*b} and [Dwayne A. Elias](#)^{*a}
[Author affiliations](#)

Abstract

Paddy soils from mercury (Hg)-contaminated rice fields in Guizhou, China were studied with respect to total mercury (THg) and methylmercury (MeHg) concentrations as well as Bacterial and Archaeal community composition. Total Hg (0.25–990 $\mu\text{g g}^{-1}$) and MeHg (1.3–30.5 ng g^{-1}) varied between samples. Pyrosequencing (454 FLX) of the hypervariable v1–v3 regions of the 16S rRNA genes showed that *Proteobacteria*, *Actinobacteria*, *Chloroflexi*, *Acidobacteria*, *Euryarchaeota*, and *Crenarchaeota* were dominant in all samples. The Bacterial α -diversity was higher in samples with relatively Low THg and MeHg and decreased with increasing THg and MeHg concentrations. In contrast, Archaeal α -diversity increased with increasing of MeHg concentrations but did not correlate with changes in THg concentrations. Overall, the methylation gene *hgcAB* copy number increased with both increasing THg and MeHg concentrations. The microbial communities at High THg and High MeHg appear to be adapted by species that are both Hg resistant and carry *hgcAB* genes for MeHg production. The relatively high abundance of both sulfate-reducing δ -*Proteobacteria* and methanogenic Archaea, as well as their positive correlations with increasing THg and MeHg concentrations, suggests that

these microorganisms are the primary Hg-methylators in the rice paddy soils in Guizhou, China.

- This article is part of the themed collection: [Mercury Biogeochemistry, Exposure, and Impacts](#)

[About](#)

[Cited by](#)

[Related](#)

[Back to tab navigation](#)

[Download this article](#) [PDF format](#) [Article HTML](#)