

UCSF

UC San Francisco Previously Published Works

Title

Patterns of Opacification in Coronary CT Angiography: Contrast Differences and Gradients

Permalink

<https://escholarship.org/uc/item/4xk198qz>

Journal

Current Cardiovascular Imaging Reports, 7(10)

ISSN

1941-9066

Authors

Rybicki, Frank J
Juan, Yu-Hsiang
Saboo, Sachin S
[et al.](#)

Publication Date

2014-10-01

DOI

10.1007/s12410-014-9291-z

Peer reviewed



Published in final edited form as:

Curr Cardiovasc Imaging Rep. 2014 October 1; 7(10): 9291–. doi:10.1007/s12410-014-9291-z.

Patterns of Opacification in Coronary CT Angiography: Contrast Differences and Gradients

Frank J. Rybicki, MD, PhD^{1,*}, Yu-Hsiang Juan, MD¹, Sachin S. Saboo, MD¹, Elizabeth George, MD¹, Rani Bhivasankar, MD¹, and Dimitrios Mitsouras, PhD¹

¹Applied Imaging Science Laboratory, Department of Radiology, Brigham and Women's Hospital & Harvard Medical School, 75 Francis Street, Boston, MA 02115

Abstract

Iodinated contrast delivery is a key component of coronary CT angiography. However, the purpose of contrast delivery has been limited to morphology alone. Specifically, iodine opacification of the coronary lumen has been used to separate it from the coronary artery wall and lesions within the coronary arteries. Because contrast is delivered to the coronary arteries according to the coronary blood flow, there is flow information encoded within the contrast opacification which, depending on CT hardware and acquisition protocol, can be recognized in coronary CT angiography. In addition, metrics related to flow have been identified and studied. They include coronary contrast opacification differences and contrast opacification gradients.

Keywords

coronary CT angiography; coronary artery disease; contrast opacification gradients; contrast opacification differences

Introduction

Coronary Computed Tomography Angiography (CTA) is an important non-invasive tool for the evaluation of suspected coronary artery disease (CAD) in stable patients. Its largest benefit to date has been the high negative predictive value when invasive coronary angiography (ICA) is used as a reference standard. However, CT has lower specificity for predicting the functional significance of coronary lesions, a current role for other cardiac imaging modalities including nuclear or magnetic resonance methods. For CT, the reported positive predictive values vary widely between roughly 50–80%, particularly in arteries with moderate (50–70% diameter) stenosis [1]. Thus, a commonly used algorithm is to reserve CT for patients in whom a normal, or near-normal, study is likely. One example is a patient with atypical chest pain and a low pre-text likelihood of coronary artery stenoses. In such a

*Corresponding author: Frank J. Rybicki, MD, PhD, FAHA, FACR, Applied Imaging Science Laboratory, Department of Radiology, Brigham and Women's Hospital & Harvard Medical School, 75 Francis Street, Boston, MA 02115, Phone: 617-732-7206, Fax: 617-264-5245, frybicki@partners.org.

Conflict of Interest

Frank J. Rybicki receives research support from Toshiba Medical Systems Corporation.

Yu-Hsiang Juan, Sachin S. Saboo, Elizabeth George, Rani Bhivasankar, and Dimitrios Mitsouras declare that they have no conflict of interest.

patient, a normal CT scan provides reassurance. However, if the CT scan is abnormal, the patient will in fact have CAD, and the further workup will generally include cardiac imaging at stress. If ischemia is suspected the patient would then proceed to ICA with or without fractional flow reserve (FFR), potentially followed by intervention. While it is well validated and accepted, this strategy is overall a less efficient utilization of resources than the development of a single imaging study that provides both anatomical and functional coronary assessment [2, 3].

Improvements in CT technology have led to significant progress toward achieving this goal. One method detailed elsewhere is the combination of CTA and CT Perfusion (CTP) performed using adenosine stress. To date, the most comprehensive assessment of this methodology [4] assessed the combination of CTA plus CTP with respect to reference standard ICA plus Single Photon Emission Computed Tomography (SPECT) [5]. CTP was shown to significantly enhance the accuracy of CTA alone, based on independent core laboratory reads from all four imaging modalities [2]. This study used only the morphology of the coronary arteries from the perspective of anatomical evaluation of coronary stenosis.

Newer generation CT hardware has provided improved spatial and temporal resolution and larger volume coverage [6, 7]. Specifically, the larger volume coverage enables the extraction of flow-dependent information from contrast enhancement patterns in first pass CTA images. In this way, current technologies are capable of characterizing the attenuation of contrast medium in coronary arteries [8], and these data can in turn be used to evaluate both the degree and direction of contrast flow, as it is carried along by the blood [9, 10].

Based on this concept, there is a new family of parameters that have been developed to quantify the contrast density variations in routine coronary CTA DICOM images [11]. All of these parameters are based upon one of the author's (FR) clinical observation and clinical follow-up regarding patients who have CT lesions that ultimately undergo an intervention for CAD at Brigham and Women's Hospital in Boston, MA, USA. The observation is that CT attenuation as measured in HU differs from the proximal to the distal coronary artery, and those coronary arteries that are associated with events in follow-up have a greater HU drop off from proximal to distal artery than those vessels that are not associated with events. While this observation is simple and clinically reliable for a technically adequate scan, the physics of blood flow are very complex, and there are many potential methods to determine those scans that should be considered adequate. For cardiovascular imagers who acquire, and interpret coronary CTA images for patient management decisions, there has been a discussion regarding the ability to see substantial HU drop offs with the naked eye versus making ROI measurements. It is one of the author's (FR) opinion that many coronary CTA interpretations are performed with too much ambient room lighting, and simply adjusting the reading room to one tailored for clinical interpretation of images can make several aspects of the interpretation more favorable. Moreover, there are now multiple software packages that will compute these parameters along with the coronary vessel segmentation. When used, these software package results should be correlated with the appearance of the images, similar to using software that automatically judges the degree of stenosis from an individual coronary lesion.

This paper reviews and interprets the several publications that test this family of new coronary CTA parameters for the hemodynamic significance of coronary stenosis. There are two basic parameters, and both are based on HU measurements, i.e. Regions of Interest (ROI) measurements, in the coronary lumen. The first parameter reflects the difference between two points in the coronary artery such as the proximal and distal Left Anterior Descending (LAD). Another way to place the two points more closely would be just proximal and just distal to a coronary lesion. This measurement is called a contrast opacification difference, or for simplicity a “contrast difference”. For the second parameter, many, as opposed to two, points are placed within a coronary artery. These points then map the opacification pattern along the vessel. The contrast opacification can then be plotted with respect to a standard denominator such as the distance along the coronary artery. This is termed the contrast opacification gradient for the coronary distance, and it is defined as the linear regression coefficient of the Hounsfield Units (HU) versus distance from the coronary ostium (Figure 1). There is no theoretical difference between the contrast opacification gradient with respect to distance and the transluminal attenuation gradient, or “TAG”, except for the fact that the concept was hypothesized, tested, and then presented [12] and published [13]** using initial CT images acquired from 320-detector row CT hardware. The “TAG” was later applied first to 64-detector row CT acquisitions, and this abbreviation was subsequently applied to 320 CT acquisitions. As noted below, the sampling rate along the coronary artery has varied in the reported data. In summary, we recommend that all these data together be termed “contrast gradients” or “gradients” to reflect the distinction from contrast differences.

In addition to the definition and historical description above, the purpose of this article is to instill among cardiac imagers the fact that CT contrast enhancement in the coronary lumen provides more information than the simple separation of the lumen and the surrounding structures. Simply put, the data in the contrast opacification is related to blood flow, for it is that flow that delivered the contrast at the point in time when the CT data was acquired. The text below is intended to illustrate these techniques, summarize the results to date, and review limitations that require future studies. It is likely that in the near future such novel yet simple approaches to quantify the flow of contrast will allow enhanced prediction of functionally significant stenoses from coronary CTA by enhancing specificity, particularly in cases where evaluation is otherwise difficult [14–17].

Contrast Opacification in Coronary CT Images

The initial evaluation of 320-detector row coronary CTA reported consistently lower contrast density in distal normal coronary arteries compared to proximal arteries [8]. Further analysis of this phenomenon led to the discovery of the contrast gradient, namely, that this drop-off in contrast enhancement along the length of a coronary artery is both continuous and highly linear, as well as smaller in normal coronaries than in those with significant diameter stenosis $\geq 75\%$ [13].

The physical principles describing the time-varying concentration of a contrast bolus injection at the entry and exit of a circulatory sub-system, under the assumption of no recirculation, were described in the mid-1950's and early 1960's [18, 19]. These basic

principles can be applied to the first pass iodinated contrast passage in all CTA data (including the coronary arteries) because the Hounsfield unit (HU) of blood linearly relates to the concentration of the contrast agent [20]. Based on this theory, both brain [21] and myocardial [22] flow have been measured by CT, although widespread clinical use is limited by technical and radiation dose considerations because accurate data requires continuous imaging throughout the bolus passage. Nonetheless, since the proximal and distal segments of a single artery without any intervening bifurcations also form a subsystem with a single entry and exit, contrast bolus passage curves at any two such arterial locations can similarly be used to quantify blood flow. The discovery of the contrast gradient in normal and obstructed coronary arteries using CTA [13] first suggested that CT is sufficiently sensitive to detect contrast concentration differences that reflect the time delay in the contrast bolus passage between two coronary arterial locations.

Our initial findings were further validated *in vitro* by Lackner et al who thoroughly analyzed the temporal patterns of CT attenuation in conduits with caliber similar to coronary arteries, and with varying stenosis severities [9]. Coronary arteries were simulated using 3 mm inner-diameter silicon tubes with 15 mm-long stenoses between 50 and 90% diameter reduction. Dynamic imaging of these phantoms was performed using both 16- and 64-detector row CT during the injection of contrast in saline pumped with a pulsatile flow profile. For 16-detector CT at a temporal resolution of 210 ms, stenoses 70%, 80% and 90% could be excluded with 99% confidence using cut-offs for the slope of peak density increase versus time. With 64-detector row CT at 270 ms temporal resolution, cut-off values of the slopes could exclude stenosis 80% or 90%, but stenosis 70% could not be confidently excluded [9]. The results of this study validated that current CT technology can detect the level of variations in contrast concentration that are induced by blood flow.

Contrast Opacification Gradients

The introduction of gradients evaluated both normal coronary arteries and LAD arteries with stenosis using 320-detector row CT. The gradient values were computed from 1 mm equidistant intervals along the vessel using automated post-processing software [13]. The “distance gradient” was described as a linear regression between intraluminal contrast opacification (mean HU) and distance from the ostium. In addition to distance gradient, lumen cross-sectional area (G_a) and lumen short axis diameter (G_s) gradients were assessed (Figure 1); all three metrics revealed high reproducibility with insignificant variations between cardiac phases, heart rates, body mass index, and between readers. This study reported the linear relationship of HU reduction against vessel distance, with higher rate of HU reduction with significant artery stenosis. The G_a and G_s assessments revealed significant non-zero slope or gradient ($p < 0.0001$), with excellent or good goodness-of-fit in all 3 gradients and no difference between the three coronary arteries for G_a and G_s . The comparison of the gradients revealed a significant mean value difference between patients with or without significant stenosis ($p = 0.021$), with G_s providing the most consistent difference in the mean value [13]. For assessment of distance gradient across a single lesion in a coronary artery, differences in the attenuation values across the lesion was divided by the length of the lesion and studies have proven larger gradients to exist across lesions as compared to normal coronary arteries.

One way to measure the gradient is to make intraluminal ROI measurements of the average attenuation in cross-sections of the artery orthogonal to the centerline at fixed intervals along its course, starting at the coronary ostium and extending to the distal artery. These manual tracings can be performed at a 5–10 mm interval, which has been assumed sufficient. Using semi-automated software, tighter spacing of 0.5–1 mm intervals have been used to obtain more data points [8, 11, 13, 14, 16, 17, 23]. Another variable is the most distal spot in the coronary artery to make the HU measurement. To date this has been chosen using a diameter cutoff of 2 – 2.5 mm or an area cutoff of 2 mm² [8, 11, 13–17, 23]. Similarly, the ROI used to perform the intraluminal contrast attenuation measurement has been operator-dependent, and has been either the entire luminal contour, or a small ROI of fixed 2mm² area placed manually within the lumen.

As noted above, when the number of HU measurement is reduced to two, a proximal and distal point along a coronary artery, the gradient is subsequently reduced to an opacification difference. Although the difference alone does not include distance traveled by contrast between measurement sites, one of the requirements to calculate flow velocity, its usefulness has been established for coronary artery lesions [14, 23, 24]. The first comparison of contrast opacification changes in a vessel with respect to ICA was performed by Chow et al who developed and compared opacification differences against both anatomic severity by ICA and functional severity in terms of blood flow as defined by the Thrombolysis in Myocardial Infarction (TIMI) grade in 52 patients [24]. This study used 64-detector row sub-volume CT, and because there is temporal non-uniformity in the coronary arteries, the proximal and distal coronary HU values were normalized by dividing the coronary ROI mean HU by the attenuation in the aorta on the same axial slice. The contrast difference in normal arteries was found to be 0.100 ± 0.042 . The difference was significantly different than that found both in arteries with obstructive CAD as defined by ICA ($> 50\%$ diameter stenosis; difference = 0.191 ± 0.214), as well as in arteries with TIMI flow grade < 3 (difference = 0.406 ± 0.226), although it was not changed (difference = 0.078 ± 0.078) when compared to arteries with TIMI flow grade of 3 [24]. That is, while the opacification difference was not changed from normal for arteries with non-flow obstructive lesions, it was somewhat higher than normal for anatomically obstructive lesions ($p = 0.004$) and much higher ($p < 0.001$) for vessels with abnormal resting coronary flow [24].

Based on a thorough analysis of the variability of the opacification difference in normal arteries, a cut-off for the difference of > 0.184 was used to detect abnormal coronary flow (TIMI flow grade < 3) with an overall accuracy of 89% [24]. In addition, when comparing vessels with stenoses $> 50\%$ against normal vessels, the minimum normalized contrast opacification found proximal to a stenosis was slightly higher than normal vessels ($p = 0.006$), while the minimum normalized contrast opacification post-stenosis was significantly lower ($p < 0.001$). This finding correlated to decreased contrast flow through the stenotic lesion, resulting in more contrast medium accumulation in the pre-stenosis region as compared to post-stenosis segments. The study noted that measurement of the opacification difference may be particularly beneficial in arteries otherwise difficult to evaluate, such as those with severe coronary calcification or stents [24]. It additionally stressed the importance of correcting for the temporal variation in coronary contrast opacification at different heart beats as occurs with 64-detector CT, using the opacification in the descending

aorta acting as the normalizing factor, since assessment of CAD severity was not possible by contrast opacification measurements without this correction [24].

Choi et al first assessed gradients for the detection of significant stenosis in comparison to both ICA and TIMI flow measurements in 126 patients [14]. Although they used 64-detector row hardware, there was no correction to account for the sub-volume imaging. Nonetheless, their results mirrored those from our group [13], namely that the gradient became more abnormal according to increasing stenosis severity by CT, as well as in accordance with quartiles of segmental stenosis and segment-at-risk scores. A consistent and significantly lower gradient value with increasing maximum stenosis severity on a per artery basis was also observed against invasive assessment by quantitative ICA, with gradient values ranging from -2.37 ± -4.67 HU/10 mm for lesions with less than 50% stenosis to -13.46 ± -9.59 HU/10mm for diameter stenosis of 100% ($p < 0.0001$) [14]. Importantly, in keeping with our study [13], gradient values were different among the three coronary territories, with marked differences between left and right coronaries. For example, among left anterior descending (LAD) and left circumflex (LCX) arteries with <50% stenosis by ICA, gradient values were -2.93 ± 5.14 and -3.62 ± 5.16 HU/10 mm respectively, while in right coronary arteries (RCA) with <50% stenosis, they were -0.16 ± 2.08 HU/10 mm. This difference persisted across stenosis severity levels; for example in 70–99% stenosis, the values for the LAD, LCX, and RCA were -11.6 ± 5.91 , -13.99 ± 9.18 and -9.24 ± 6.3 HU/10 mm, respectively.

In terms of invasively measured resting flow, these results mirrored those of Chow et al [24], with gradients correlating to both TIMI grade as well as corrected TIMI frame count. Additionally, in arteries with 100% stenosis or TIMI grade 0, the gradient correlated ($p < 0.0001$ for all comparisons) to Rentrop grade classification of collateral flow [14]. Finally, the addition of the gradient to the CT based stenosis classification significantly enhanced accuracy compared to CT alone for the detection of obstructive CAD $\geq 50\%$ stenosis by ICA (area under the curve [AUC] of the receiver operating characteristic [ROC] curve 0.951 versus 0.932). This was also true for either calcified or non-calcified plaques independently (ROC AUCs of 0.880 versus 0.825 and 0.978 versus 0.960, respectively). However, results were less encouraging for the reclassification of patients, with the net reclassification improvement index compared to CT alone becoming significant for only calcified ($> 30\%$ calcification by volume) plaques [14].

Contrast Opacification Gradients and the Hemodynamically Significant Coronary Lesion

The first study of the physiological significance of gradients and opacification differences was also performed by Choi et al in 97 arteries with luminal diameter stenosis $\geq 50\%$, and invasive FFR less than or equal to 0.8 as the reference standard for myocardial ischemia [23]. Forty one per-cent of vessels had a lesion with significant FFR. For the prediction of significant FFR, gradients had low sensitivity (47.5%) but high specificity (91.2%), in contrast to CT that had high sensitivity (92.5%) but low specificity (52.6%). The contrast differences alone had moderate sensitivity and specificity (65 and 61.4%, respectively). The addition of gradients to the CTA read of morphology alone significantly increased the ROC AUC (0.809 ± 0.044 versus 0.726 ± 0.056), but addition of the opacification differences to

CT did not (AUC 0.784 ± 0.048). However, addition of the gradients to the CT did not lead to significant improvement in the net reclassification index, while addition of differences to the CT reads led to its impairment [23]. The main explanation noted in these results was that both contrast metrics were obtained from a rest coronary CTA, while FFR is assessed under pharmacologically-induced stress [11, 23]. Rest flow does not adequately correlate to coronary hemodynamics for a lesion with $<85\%$ stenosis [25].

A second study of 64-detector row coronary CTA to predict invasive FFR was published by Yoon et al [17]. This study focused on the comparison of the diagnostic value of gradients and coronary CTA-derived FFR. This is typically denoted FFR_{CT} , and results are detailed [26–28] in other articles. Among 82 vessels, 39% of which had lesions with $FFR \leq 0.8$ [17], both the gradient and FFR_{CT} were significantly lower in vessels with lesion-specific ischemia than in those without. The ROC AUC of the gradient alone was 0.63 (gradient sensitivity and specificity of 37.5% and 88%, respectively) and not significantly different than that of coronary CTA, while that of FFR_{CT} was 0.94 (sensitivity and specificity of 81.3% and 94%, respectively), significantly higher than that of coronary CTA. However, in the subset of calcified lesions, none of the three tests had significantly different AUCs. Interestingly, the gradient results in this study were equivalent to those found by Choi et al, including sensitivity, specificity and AUC measures. Unfortunately, however, this study opted for a “head-to-head” comparison of FFR_{CT} against gradients as stand-alone diagnostic tests, rather than using the combined test of gradients plus coronary CTA, for which prior studies had established that the high specificity of the gradients could be combined with the high sensitivity of CT.

Wong et al first evaluated the diagnostic accuracy of gradients and differences using single heart beat 320-detector row coronary CTA images against invasive FFR in 78 vessels [16]*. These images have temporal uniformity since the 16 cm craniocaudal FOV covers the entire heart [29]. Of note, at the time of writing, gradients have not been analyzed using high-pitch dual source CT; such acquisitions occur at different time points in diastole over a single R-R interval [30]. In the Wong et al study, the gradients were significantly lower in vessels with lesions with $FFR \leq 0.8$ than in those with $FFR > 0.8$ (median -21 vs. -11 HU/10 mm). A gradient cutoff value of -15.1 HU/10mm had sensitivity and specificity to detect functionally significant ischemia of 77% and 74%, and PPV and NPV of 67% and 83%, respectively. The gradient provided incremental predictive value to CT, significantly increasing the ROC AUC (0.89 for TAG plus coronary CTA, versus 0.814 for the gradient alone, versus 0.79 for coronary CTA alone), and with evidence that incorporation of the gradient enhances both the sensitivity and specificity of coronary CTA to detect lesion-specific ischemia [16]. Finally, no difference in diagnostic accuracy was found based on plaque composition. All contrast opacification differences were significantly higher in arteries with $FFR \leq 0.8$ as compared to arteries with no functionally significant lesions. However, the opacification difference was not independently predictive of $FFR \leq 0.8$ when the CT reads alone plus the gradients were already included in the multivariate model, and no further analysis of the difference was warranted, since overall accuracy of using the gradient (i.e. more data points) was higher than the HU difference between the proximal and distal coronary artery [16].

Another contrast enhancement pattern that can be used clinically has been termed the reverse attenuation gradient sign, or a “reverse gradient”. This term was coined by Li et al [31], although Choi et al noted in the literature that gradients correlate with Rentrop collateralization grade [14]. The reverse gradient is very common and also can be seen with the naked eye; it refers to a drop off in attenuation from proximal to distal along a coronary artery, after which there is a paradoxical increase in attenuation. In coronary CT image interpretation, there is typically a sharp drop in HU followed by a dramatic rise in enhancement. The reason is collateral flow. Accordingly, Li et al described the published the finding as a highly specific predictor of chronic total coronary occlusion [31]. Chronic total occlusion is reported to be found in more than 20% of patients with suspected coronary artery disease and as many as one-third of patients with at least one vessel with > 50% stenosis at ICA [32–35]. At our institution, cardiac CT has been used for a decade to identify the features that render candidates suitable for successful intervention [36], and aid the operator in percutaneous procedure planning [32]. Incidentally, one of the authors (FR) has used the same finding in the abdomen to determine total versus subtotal occlusion in the mesentery, where collateral flow in patients with an occluded celiac axis or superior mesenteric artery is very common.

Mathematically, patients with a reverse gradient have a positive (increasing) HU trend in segments distal to the lesion in question. In distinction, patients with a subtotal occlusion, for example a high-grade stenosis with retained antegrade flow, maintain a negative (decreasing) HU trend distal to the stenosis. It is reported that when the reverse gradient sign is absent, there may be some collateral flow, and similarly, bridging collaterals can still lead to the absence of the reverse gradient sign in patients with chronic total occlusion. Nonetheless, in a series of 94 occlusive lesions, Li et al demonstrated a gradient of $5.1 \text{ HU}/10\text{mm} \pm 13.4$ for lesions with chronic total occlusion versus $-13.4 \text{ HU}/10 \text{ mm} \pm 8.7$ for subtotal lesions. These data were acquired using single source 128-detector row scanner. The reverse gradient was present in 65% of chronic occlusions and 7% of high-grade but patent lesions.

Conclusion

Before the advent of hardware and techniques that allowed the analysis of first pass CTA contrast opacification within the course of the coronary arteries, iodine enhancement was used only to identify the vessel lumen and differentiate it from the wall and intervening lesions. In our opinion, and supported by the data reviewed in this article, this represents an underutilization of CTA data in general, and specifically an underutilization of coronary CTA where flow dependent information can be critical for patient management. The second take-home message is that contrast differences and gradients, i.e. the metrics themselves, are no more than a mathematical representation of what one can observe when reading a coronary CTA study. The fluid mechanics of the flow and how it is imperfectly captured by a CT scanner are orders of magnitude more complex, despite the fact that the clinical observation is simple. This is the reason why gradients may not “work” as cleanly as one might expect if a simple equation is applied to a coronary lesion. However, there are large amounts of visual data in the enhancement patterns, and these data should be considered in image interpretation. Finally, we believe that contrast differences and gradients have the

favorable quality of being simple to understand and simple to see and compute. Thus, visual assessment and quantitative information can be easily incorporated into cardiac imaging workflow.

Acknowledgments

Dimitrios Mitsouras is supported by NIBIB 1-K01-EB015868 (Mentor Frank J. Rybicki).

References

1. Mark DB, Berman DS, Budoff MJ, et al. ACCF/ACR/AHA/NASCI/SAIP/SCAI/SCCT 2010 expert consensus document on coronary computed tomographic angiography: a report of the American College of Cardiology Foundation Task Force on Expert Consensus Documents. *Journal of the American College of Cardiology*. 2010; 55(23):2663–2699. [PubMed: 20513611]
2. Rochitte CE, George RT, Chen MY, et al. Computed tomography angiography and perfusion to assess coronary artery stenosis causing perfusion defects by single photon emission computed tomography: the CORE320 study. *European heart journal*. 2013
3. Yoon YE, Koo BK. Non-invasive functional assessment using computed tomography: when will they be ready for clinical use? *Cardiovascular diagnosis and therapy*. 2012; 2(2):106–112. [PubMed: 24282704]
4. Vavere AL, Simon GG, George RT, et al. Diagnostic performance of combined noninvasive coronary angiography and myocardial perfusion imaging using 320 row detector computed tomography: design and implementation of the CORE320 multicenter, multinational diagnostic study. *Journal of cardiovascular computed tomography*. 2011; 5(6):370–381. [PubMed: 22146496]
5. George RT, Arbab-Zadeh A, Cerci RJ, et al. Diagnostic performance of combined noninvasive coronary angiography and myocardial perfusion imaging using 320-MDCT: the CT angiography and perfusion methods of the CORE320 multicenter multinational diagnostic study. *AJR American journal of roentgenology*. 2011; 197(4):829–837. [PubMed: 21940569]
6. Hsiao EM, Rybicki FJ, Steigner M. CT coronary angiography: 256-slice and 320-detector row scanners. *Current cardiology reports*. 2010; 12(1):68–75. [PubMed: 20425186]
7. Kumamaru KK, Hoppel BE, Mather RT, Rybicki FJ. CT angiography: current technology and clinical use. *Radiologic clinics of North America*. 2010; 48(2):213–235. vii. [PubMed: 20609871]
8. Rybicki FJ, Otero HJ, Steigner ML, et al. Initial evaluation of coronary images from 320-detector row computed tomography. *The international journal of cardiovascular imaging*. 2008; 24(5):535–546. [PubMed: 18368512]
9. Lackner K, Bovenschulte H, Stutzer H, Just T, Al-Hassani H, Krug B. In vitro measurements of flow using multislice computed tomography (MSCT). *The international journal of cardiovascular imaging*. 2011; 27(6):795–804. [PubMed: 20972834]
10. Nagao M, Kido T, Watanabe K, et al. Functional assessment of coronary artery flow using adenosine stress dual-energy CT: a preliminary study. *The international journal of cardiovascular imaging*. 2011; 27(3):471–481. [PubMed: 20686853]
11. de Feyter PJ. CT functional imaging using intracoronary gradient analysis: an indispensable boost for CT coronary angiography. *European heart journal cardiovascular Imaging*. 2012; 13(12):971–972. [PubMed: 22909793]
12. Mitsouras D, Otero HJ, Steigner M, Nallamshetty L, Levit NA, Rybicki FJ. Contrast Opacification Gradients between Normal and Stenotic Coronary Arteries Obtained from 320-Detector Row CT Coronary Angiography. *Proceedings of the Radiological Society of North America [serial online]*. 2008 vol SSJ03-06.
13. Steigner ML, Mitsouras D, Whitmore AG, et al. Iodinated contrast opacification gradients in normal coronary arteries imaged with prospectively ECG-gated single heart beat 320-detector row computed tomography. *Circulation Cardiovascular imaging*. 2010; 3(2):179–186. [PubMed: 20044512]

14. Choi JH, Min JK, Labounty TM, et al. Intracoronary transluminal attenuation gradient in coronary CT angiography for determining coronary artery stenosis. *JACC Cardiovascular imaging*. 2011; 4(11):1149–1157. [PubMed: 22093264]
15. Einstein AJ. TAG-is it it?: improving coronary computed tomography angiography with the isotemporal transluminal contrast attenuation gradient. *Journal of the American College of Cardiology*. 2013; 61(12):1280–1282. [PubMed: 23414793]
16. Wong DT, Ko BS, Cameron JD, et al. Transluminal attenuation gradient in coronary computed tomography angiography is a novel noninvasive approach to the identification of functionally significant coronary artery stenosis: a comparison with fractional flow reserve. *Journal of the American College of Cardiology*. 2013; 61(12):1271–1279. [PubMed: 23414792] : This is the most recent validation of 320 detector row CT contrast opacification gradients. This paper shows correlation between gradient values and fractional flow reserve
17. Yoon YE, Choi JH, Kim JH, et al. Noninvasive diagnosis of ischemia-causing coronary stenosis using CT angiography: diagnostic value of transluminal attenuation gradient and fractional flow reserve computed from coronary CT angiography compared to invasively measured fractional flow reserve. *JACC Cardiovascular imaging*. 2012; 5(11):1088–1096. [PubMed: 23153908]
18. Meier P, Zierler KL. On the theory of the indicator-dilution method for measurement of blood flow and volume. *Journal of applied physiology*. 1954; 6(12):731–744. [PubMed: 13174454]
19. Sherman H. On the theory of indicator-dilution methods under varying blood-flow conditions. *The bulletin of mathematical biophysics*. 1960; 22(4):417–424.
20. Bae KT. Intravenous contrast medium administration and scan timing at CT: considerations and approaches. *Radiology*. 2010; 256(1):32–61. [PubMed: 20574084]
21. Axel L. Cerebral blood flow determination by rapid-sequence computed tomography: theoretical analysis. *Radiology*. 1980; 137(3):679–686. [PubMed: 7003648]
22. Wang T, Wu X, Chung N, Ritman EL. Myocardial blood flow estimated by synchronous, multislice, high-speed computed tomography. *IEEE transactions on medical imaging*. 1989; 8(1): 70–77. [PubMed: 18230502]
23. Choi JH, Koo BK, Yoon YE, et al. Diagnostic performance of intracoronary gradient-based methods by coronary computed tomography angiography for the evaluation of physiologically significant coronary artery stenoses: a validation study with fractional flow reserve. *European heart journal cardiovascular Imaging*. 2012; 13(12):1001–1007. [PubMed: 22802430]
24. Chow BJ, Kass M, Gagne O, et al. Can differences in corrected coronary opacification measured with computed tomography predict resting coronary artery flow? *Journal of the American College of Cardiology*. 2011; 57(11):1280–1288. [PubMed: 21392642]
25. Gould KL, Lipscomb K. Effects of coronary stenoses on coronary flow reserve and resistance. *The American journal of cardiology*. 1974; 34(1):48–55. [PubMed: 4835753]
26. Min JK, Leipsic J, Pencina MJ, et al. Diagnostic accuracy of fractional flow reserve from anatomic CT angiography. *JAMA : the journal of the American Medical Association*. 2012; 308(12):1237–1245.
27. Nakazato R, Park HB, Berman DS, et al. Noninvasive fractional flow reserve derived from computed tomography angiography for coronary lesions of intermediate stenosis severity: results from the DeFACTO study. *Circulation Cardiovascular imaging*. 2013; 6(6):881–889. [PubMed: 24081777]
28. Norgaard BL, Leipsic J, Gaur S, et al. Diagnostic performance of non-invasive fractional flow reserve derived from coronary CT angiography in suspected coronary artery disease: The NXT trial. *Journal of the American College of Cardiology*. 2014
29. Steigner ML, Otero HJ, Cai T, et al. Narrowing the phase window width in prospectively ECG-gated single heart beat 320-detector row coronary CT angiography. *The international journal of cardiovascular imaging*. 2009; 25(1):85–90. [PubMed: 18663599]
30. Achenbach S, Marwan M, Schepis T, et al. High-pitch spiral acquisition: a new scan mode for coronary CT angiography. *J Cardiovasc Comput Tomogr*. 2009; 3(2):117–121. [PubMed: 19332343]

31. Li M, Zhang J, Pan J, Lu Z. Obstructive coronary artery disease: reverse attenuation gradient sign at CT indicates distal retrograde flow--a useful sign for differentiating chronic total occlusion from subtotal occlusion. *Radiology*. 2013; 266(3):766–772. [PubMed: 23220898]
32. Hoe J. CT coronary angiography of chronic total occlusions of the coronary arteries: how to recognize and evaluate and usefulness for planning percutaneous coronary interventions. *The international journal of cardiovascular imaging*. 2009; 25(Suppl 1):43–54. [PubMed: 19165621]
33. Prasad A, Rihal CS, Lennon RJ, Wiste HJ, Singh M, Holmes DR Jr. Trends in outcomes after percutaneous coronary intervention for chronic total occlusions: a 25-year experience from the Mayo Clinic. *Journal of the American College of Cardiology*. 2007; 49(15):1611–1618. [PubMed: 17433951]
34. Rolf A, Werner GS, Schuhback A, et al. Preprocedural coronary CT angiography significantly improves success rates of PCI for chronic total occlusion. *The international journal of cardiovascular imaging*. 2013; 29(8):1819–1827. [PubMed: 23793727]
35. Stone GW, Kandzari DE, Mehran R, et al. Percutaneous recanalization of chronically occluded coronary arteries: a consensus document: part I. *Circulation*. 2005; 112(15):2364–2372. [PubMed: 16216980]
36. Gasparovic H, Rybicki FJ, Millstine J, et al. Three dimensional computed tomographic imaging in planning the surgical approach for redo cardiac surgery after coronary revascularization. *European journal of cardio-thoracic surgery : official journal of the European Association for Cardio-thoracic Surgery*. 2005; 28(2):244–249. [PubMed: 15993621]




Figure 1.

Illustration of contrast opacification gradients. The top left image illustrates the course of the LAD with respect to the myocardium. The straightened reformatted image with corresponding region of interest Hounsfield Unit measurements appear on the top right. The lower panels show the regression lines that define the gradients. The distance gradient (bottom left) is most commonly reported.