

# Lawrence Berkeley National Laboratory

## Lawrence Berkeley National Laboratory

### Title

Neutrinoless Double Beta Decay in Light of SNO Salt Data

### Permalink

<https://escholarship.org/uc/item/4xr2058f>

### Author

Murayama, Hitoshi

### Publication Date

2009-09-21

# Neutrinoless Double Beta Decay in Light of SNO Salt Data

Hitoshi Murayama and Carlos Peña-Garay

*School of Natural Sciences, Institute for Advanced Study, Einstein Dr, Princeton, NJ 08540, USA*

---

This work was supported in part by the Director, Office of Science, Office of High Energy Physics, of the U.S. Department of Energy under Contract No. DE-AC03-76SF00098.

#### DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor The Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or The Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or The Regents of the University of California.

# Neutrinoless Double Beta Decay in Light of SNO Salt Data

Hitoshi Murayama\* and Carlos Peña-Garay

*School of Natural Sciences, Institute for Advanced Study, Einstein Dr, Princeton, NJ 08540, USA*

(Dated: February 1, 2008)

In the SNO data from its salt run, probably the most significant result is the consistency with the previous results without assuming the  $^8\text{B}$  energy spectrum. In addition, they have excluded the maximal mixing at a very high confidence level. This has an important implication on the double beta decay experiments. For the inverted or degenerate mass spectrum, we find  $|\langle m_\nu \rangle_{ee}| > 0.013$  eV at 95% CL, and the next generation experiments can discriminate Majorana and Dirac neutrinos if the inverted or degenerate mass spectrum will be confirmed by the improvements in cosmology, tritium data beta decay, or long-baseline oscillation experiments.

In the past five years, there had been an amazing progress in neutrino physics. The atmospheric neutrinos showed a large up-down asymmetry in the SuperKamiokande (SK) experiment which came as the first significant evidence for a finite neutrino mass [1] and hence the incompleteness of the Standard Model of particle physics. SuperKamiokande also improved the accuracy in solar neutrino studies greatly using the elastic scattering (ES) process. The Sudbury Neutrino Observatory (SNO) experiment has studied the charged-current (CC) and neutral-current (NC) process in addition to the ES process, and has shown that the solar neutrinos change their flavors from the electron type to other active types (muon and tau neutrinos) [2]. Finally, the KamLAND reactor anti-neutrino oscillation experiments reported a significant deficit in reactor anti-neutrino flux over approximately 180 km of propagation [3]. Further combined with the pioneering Homestake experiment [4] and Gallium-based experiments [5], the decades-long solar neutrino problem [6] appears solved. The so-called Large Mixing Angle (LMA) solution [7], where the electron neutrinos produced at Sun's core propagate adiabatically to a heavier mass eigenstate due to the matter effect [8], is the only viable explanation of the data.

On September 7, 2003, SNO published the result from their salt run with an enhanced sensitivity to the NC process [9]. Most importantly, the new result agrees well with previous results, confirming the LMA solution to the solar neutrino problem. In addition, they have reported a much better determination of the mixing angle  $\theta_{12}$ , which excludes the maximal mixing  $\theta_{12} = \pi/4$  at a very high significance: 5.4 sigma.

The exclusion of the maximal mixing has an important impact on another crucial question in neutrino physics: Is neutrino its own anti-particle? If yes, neutrinos are Majorana fermions; if not, Dirac. This question is even deeper than it sounds. For instance, if neutrinos and anti-neutrinos are identical, there could have been a process in Early Universe that affected the balance between particles and anti-particles, leading to the matter anti-matter asymmetry we need to exist. In fact, so-called leptogenesis models directly link the Majorana nature of neutrinos to the observed baryon asymmetry [10] [37].




FIG. 1: Three possible mass spectra of neutrinos. The wider splitting is  $\Delta m_{\text{atm}}^2 \simeq 2 \times 10^{-3} \text{ eV}^2$ , while the smaller one is  $\Delta m_{\text{solar}}^2 \simeq 7 \times 10^{-5} \text{ eV}^2$ .

This question can in principle be resolved if a neutrinoless double beta decay is observed. Because such a phenomenon will violate the lepton number by two units, it cannot be caused if the neutrino is different from the anti-neutrino [38]. Many experimental proposals exist that will increase the sensitivity to such a phenomenon dramatically over the next ten years [13]. The crucial question is if a negative result from such experiments can lead to a definitive statement about the nature of neutrinos. In particular, the matrix element of neutrinoless double beta decay is proportional to the effective electron-neutrino mass [14]

$$\langle m_\nu \rangle_{ee} = m_1 U_{e1}^2 + m_2 U_{e2}^2 + m_3 U_{e3}^2, \quad (1)$$

which may have cancellation among three terms that makes it difficult to assess the result of a negative search. However, the exclusion of the maximal mixing in  $\theta_{12}$  actually helps to eliminate such an unfortunate situation. Note that the proposed experiments are aiming at the sensitivity reaching  $|\langle m_\nu \rangle_{ee}| \sim 0.01$  eV [13].

Within three generations of neutrinos and given all neutrino oscillation data [39], there are three possible mass spectra: degenerate, normal hierarchy and inverted hierarchy (see Fig. 1) [40]. Given that the third mixing angle  $\theta_{13} = \arcsin |U_{e3}|$  is known to be small from the CHOOZ limit [18], one can obtain a lower bound on the effective electron-neutrino mass. For the

\*On leave of absence from Department of Physics, University of California, Berkeley, California 94720, USA




FIG. 2: Left:  $\Delta\chi^2$  vs  $\cos 2\theta_{12}$  with all solar neutrino data with and without KamLAND, marginalized on  $\Delta m_{12}^2$ . Right:  $\Delta\chi^2$  vs  $\sqrt{\Delta m_{\text{atm}}^2}$  with SuperKamiokande without the update [15] or with the update [16], marginalized on  $\sin^2 2\theta_{23}$  and combined with K2K.

degenerate spectrum of the nearly common mass  $m$ , we can ignore  $m_3 U_{e3}^2$  relative to two other terms, and find

$$\begin{aligned} |\langle m_\nu \rangle_{ee}| &\simeq |mU_{e1}^2 + mU_{e2}^2| \\ &\geq m(|U_{e1}|^2 - |U_{e2}|^2) = m \cos 2\theta_{12}. \end{aligned} \quad (2)$$

For the inverted hierarchy,  $m_1 \simeq m_2 \geq \sqrt{\Delta m_{\text{atm}}^2}$ , and we can again ignore  $m_3 U_{e3}^2$  relative to two other terms. Therefore,

$$\begin{aligned} |\langle m_\nu \rangle_{ee}| &\geq \sqrt{\Delta m_{\text{atm}}^2} |U_{e1}^2 + U_{e2}^2| \\ &\geq \sqrt{\Delta m_{\text{atm}}^2} (|U_{e1}|^2 - |U_{e2}|^2) = \sqrt{\Delta m_{\text{atm}}^2} \cos 2\theta_{12} \end{aligned} \quad (3)$$

Note that the bound for the inverted hierarchy is weaker than that for the degenerate spectrum by definition, because the degeneracy requires  $m \gtrsim \sqrt{\Delta m_{\text{atm}}^2}$ . Therefore, Eq. (3) is our master equation for most of our discussions.

Unfortunately, for the normal hierarchy, one cannot obtain a similar rigorous lower limit. On the other hand, the improvement in the cosmological data [19] and the KATRIN experiment on the end point of the tritium beta decay [20] may positively establish the degenerate spectrum, or the long baseline neutrino oscillation experiments may positively establish the inverted hierarchy [21]. If either of them happens, and if the neutrinoless double beta won't be seen within these bounds, the neutrinos will be found to be Dirac particles [41].

$\cos 2\theta_{12}$  now has a robust lower bound given the new SNO result. To best of our knowledge, it was pointed out first in [23] that the less than maximal mixing leads to a lower bound on  $|\langle m_\nu \rangle_{ee}|$  for the degenerate and inverted spectra. More recent papers [24] studied the bound quantitatively before the

recent SNO result when the lower bound was not quite robust, because the exclusion of the maximal mixing was reported at different confidence levels among different analyses and depended crucially on Homestake data [4].

There are obviously two main ingredients in the lower bound. One is  $\Delta m_{\text{atm}}^2$  from SuperKamiokande experiment which had recently been updated [25], and the other is  $\theta_{12}$  from the solar neutrino data which includes the recent SNO result. The last ingredient is  $\theta_{13}$  which we assume to be zero throughout our discussions. We will come back to the little effect of non-vanishing  $\theta_{13}$  at the end of the letter.

First on  $\Delta m_{\text{atm}}^2$ . The analysis of the atmospheric (SK) and accelerator (K2K) data was done in the general case of  $3\nu$  oscillations in [15], and we show the marginalized  $\Delta\chi^2$  as a function of  $\sqrt{\Delta m_{\text{atm}}^2}$  in the right pane of Fig. 2. The constraint  $\theta_{13} = 0$  does not modify the shape of these functions. This analysis uses the data available before updates this summer [25]. The SK preliminary analysis of atmospheric data show a shift of the allowed region to lower  $\Delta m_{\text{atm}}^2$ , due to several improvements in their analysis: new neutrino flux with updated primary cosmic ray flux, hadron interaction model and calculation methods (3D), and improved neutrino interactions, detector simulation and event reconstruction. We included the SK update [16].

Second on  $\theta_{12}$ . The analysis of solar and reactor data is done as described in [15], except that  $\theta_{13}$  is set to zero, the Gallium rate is updated [5] and the latest SNO data (NC, CC and ES measured in phase-II [9]) is included [26]. The  $\Delta\chi^2$  is shown as a function of  $\cos 2\theta_{12}$  in the right pane of Fig. 2.

Combining  $\Delta m_{\text{atm}}^2$  and  $\theta_{12}$  discussed above, we obtain the final result on the effective electron-neutrino mass. The lower




FIG. 3: Lower bound on  $|\langle m_\nu \rangle_{ee}|$  vs confidence levels for the inverted hierarchy spectrum. Note that what is shown here is  $\sqrt{\Delta m_{\text{atm}}^2} \cos 2\theta_{12}$ , which is the minimum value of  $|\langle m_\nu \rangle_{ee}|$  allowing the maximum cancellation between  $m_1$  and  $m_2$ . The solid line is based on the atmospheric neutrino data before the update, while the dashed line with the update.

bounds are shown at different confidence levels in Fig. 3. One can see that

$$|\langle m_\nu \rangle_{ee}| > 0.013 \text{ (011) eV} \quad 95\% \text{ CL (99\% CL)}. \quad (4)$$

The result is quite robust in the sense that one has to require an extremely high confidence level (99.7%) to bring  $|\langle m_\nu \rangle_{ee}|$  below 0.01 eV. Recall that the proposed experiments are aiming at the sensitivity reaching  $|\langle m_\nu \rangle_{ee}| \sim 0.01$  eV [13].

In all of the above discussions, we ignored  $\theta_{13}$ . First of all,  $\theta_{13}$  is small due to the limit from CHOOZ reactor experiment [18]. Even setting the CHOOZ limit aside, it is well-known, however, that  $\theta_{13}$  has very little effect on the determination of  $\Delta m_{\text{atm}}^2$  [15], and also can only decrease the preferred values of  $\theta_{12}$  [27, 36]. Therefore, the impact of a non-vanishing  $\theta_{13}$  on  $\Delta m_{\text{atm}}^2$  and  $\theta_{12}$  can only strengthen our result.

One may also worry about corrections to the approximate formula Eq. (3) due to  $\Delta m_{\text{solar}}^2$  and  $\theta_{13}$ . To minimize  $|\langle m_\nu \rangle_{ee}|$ , we can study the case where both  $U_{e2}^2$  and  $U_{e3}^2$  have the opposite sign from  $U_{e1}^2$ , giving

$$|\langle m_\nu \rangle_{ee}| = |(m_1 - m_3)c_{13}^2 \cos 2\theta_{12} - (m_2 - m_1)c_{13}^2 s_{12}^2 + m_3(c_{13}^2 \cos 2\theta_{12} - s_{13}^2)|. \quad (5)$$

In the limit  $\Delta m_{\text{solar}}^2 = 0$  ( $m_2 = m_1$ ) and  $\theta_{13} = 0$ , it reduces to Eq. (3) due to the first term above. The suppression factor due to  $c_{13}^2$  is at most 4.4% (95% CL) thanks to the CHOOZ limit. The second term does not vanish due to  $\Delta m_{\text{solar}}^2 \neq 0$ , and gives a correction at most of

$$\frac{\Delta m_{\text{solar}}^2}{2\Delta m_{\text{atm}}^2} \frac{s_{12}^2}{\cos 2\theta_{12}} \lesssim 3\% \quad (95\% \text{ CL}). \quad (6)$$

Finally, the last term cannot be negative given the CHOOZ limit and only strengthens our limit. Overall, our lower bound can change at most by 8%.

The bound on  $|\langle m_\nu \rangle_{ee}|$  is expected to improve further as more data will become available. As for long-baseline (LBL) accelerator-based neutrino oscillation experiments, K2K will double the data set, while MINOS, ICARUS, and OPERA are expected to come online around 2005. If approved, the neutrino beam from J-PARC will be available around 2007. They will improve the accuracy on  $\Delta m_{\text{atm}}^2$  dramatically [28]. SNO will install dedicated Neutral Current Detector (NCD) this fall, which will allow event-by-event separation of CC/ES and NC events and lead to a more accurate measurement of  $\theta_{12}$  [29]. Later, measurements of low-energy solar neutrino fluxes ( ${}^7\text{Be}$  and  $pp$ ) will allow even better determination of  $\theta_{12}$  [30]. The corrections due to  $\theta_{13}$  will also be constrained better by LBL experiments as well as new multiple-baseline reactor anti-neutrino oscillation experiments [31].

It is useful to recall the cosmological bound. The combination of WMAP, 2dFGRS, and Lyman  $\alpha$  data leads to an upper bound [32] (see [33] for a slightly weaker bound)

$$\sum_i m_{\nu_i} < 0.70 \text{ eV}, \quad (7)$$

which translates to [34]

$$|\langle m_\nu \rangle_{ee}| < 0.23 \text{ eV}, \quad (8)$$

allowing the maximum constructive interference between three mass eigenstates. This follows from the fact that neutrinos are degenerate in this mass range and the inequality

$$|U_{e1}^2 + U_{e2}^2 + U_{e3}^2| \leq |U_{e1}^2| + |U_{e2}^2| + |U_{e3}^2| = 1. \quad (9)$$

For a comparison [34], the reported evidence for the neutrinoless double beta decay suggest  $|\langle m_\nu \rangle_{ee}| = (0.11\text{--}0.56)$  eV [35], while the reanalysis in [14] gives 0.4–1.3 eV using a different set of nuclear matrix elements.

To summarize, we have obtained a robust lower bound on the effective electron-neutrino mass relevant to the neutrinoless double beta decay. For the degenerate and inverted mass spectra, the next generation experiments that have sensitivity on  $|\langle m_\nu \rangle_{ee}|$  down to 0.01 eV can determine if neutrino is its own anti-particle. For the normal hierarchy, the effective electron-neutrino mass may even vanish. However, if the large-scale structure cosmological data, improved data on the tritium beta decay, or the long-baseline neutrino oscillation experiments establish the degenerate or inverted mass spectrum, the null result from such double-beta decay experiments will lead to a definitive result.

### Acknowledgments

This work was supported by Institute for Advanced Study, funds for Natural Sciences. CPG is grateful to John Bahcall and Concha Gonzalez-Garcia for using partial results of the ongoing work on the analyses of solar and reactor data after

SNO phase II data. CPG was supported in part with the aid from the W.M. Keck Foundation and in part by the NSF grant PHY-0070928. HM was also supported in part by the DOE

Contract DE-AC03-76SF00098 and in part by the NSF grant PHY-0098840.

- 
- [1] Y. Fukuda *et al.* [Super-Kamiokande Collaboration], Phys. Rev. Lett. **81**, 1562 (1998) [hep-ex/9807003].
- [2] Q. R. Ahmad *et al.* [SNO Collaboration], Phys. Rev. Lett. **89**, 011301 (2002) [nucl-ex/0204008].
- [3] K. Eguchi *et al.* [KamLAND Collaboration], Phys. Rev. Lett. **90**, 021802 (2003) [hep-ex/0212021].
- [4] B. T. Cleveland *et al.*, Astrophys. J. **496**, 505 (1998).
- [5] E. Bellotti, Talk at VIIIth International Conference on Topics in Astroparticle and Underground Physics (TAUP03), Seattle, (Sep 5-9, 2003); V. Gavrin, *ibid.*
- [6] J. N. Bahcall, Phys. Rev. Lett. **12**, 300 (1964); R. Davis, Phys. Rev. Lett. **12**, 303 (1964).
- [7] For a recent review, see M. C. Gonzalez-Garcia and Y. Nir, Rev. Mod. Phys. **75**, 345 (2003) [hep-ph/0202058].
- [8] L. Wolfenstein, Phys. Rev. D **17**, 2369 (1978); S. P. Mikheev and A. Y. Smirnov, Sov. J. Nucl. Phys. **42**, 913 (1985) [Yad. Fiz. **42**, 1441 (1985)].
- [9] S.N. Ahmed *et al.* [SNO Collaboration], [nucl-ex/0309004].
- [10] M. Fukugita and T. Yanagida, Phys. Lett. B **174**, 45 (1986).
- [11] E. K. Akhmedov, V. A. Rubakov and A. Y. Smirnov, Phys. Rev. Lett. **81**, 1359 (1998) [arXiv:hep-ph/9803255]; K. Dick, M. Lindner, M. Ratz and D. Wright, Phys. Rev. Lett. **84**, 4039 (2000) [hep-ph/9907562]; H. Murayama and A. Pierce, Phys. Rev. Lett. **89**, 271601 (2002) [hep-ph/0206177].
- [12] M. Fukugita and T. Yanagida, "Physics Of Neutrinos And Applications To Astrophysics," Springer Verlag (2003), Berlin, Germany.
- [13] Talk by G. Gratta at XXI International Symposium on Lepton and Photon Interactions at High Energies, 11-16 August 2003, Fermi National Accelerator Laboratory, Batavia, Illinois USA.
- [14] P. Vogel, "Limits From Neutrinoless Double-Beta Decay (Rev.)," in Particle Data Group (K. Hagiwara *et al.*), Phys. Rev. D **66**, 010001 (2002).
- [15] M. C. Gonzalez-Garcia and C. Peña-Garay, hep-ph/0306001.
- [16] We thank the SuperKamiokande collaboration, in particular T. Kajita and C. Yanagisawa, for providing us a plot of  $\Delta\chi^2$  vs  $\Delta m^2$  which we used to produce our result.
- [17] A. Aguilar *et al.* [LSND Collaboration], Phys. Rev. D **64**, 112007 (2001) [hep-ex/0104049].
- [18] M. Apollonio *et al.*, Eur. Phys. J. C **27**, 331 (2003) [hep-ex/0301017].
- [19] See, *e.g.*, W. Hu, D. J. Eisenstein and M. Tegmark, Phys. Rev. Lett. **80**, 5255 (1998) [astro-ph/9712057]; . Hannestad, Phys. Rev. D **67**, 085017 (2003) [arXiv:astro-ph/0211106]; M. Kaplinghat, L. Knox and Y. S. Song, arXiv:astro-ph/0303344.
- [20] A. Osipowicz *et al.* [KATRIN Collaboration], hep-ex/0109033.
- [21] See, *e.g.*, C. Albright *et al.*, hep-ex/0008064.
- [22] J. F. Beacom, N. F. Bell, D. Hooper, J. G. Learned, S. Pakvasa and T. J. Weiler, arXiv:hep-ph/0307151.
- [23] S. M. Bilenky, C. Giunti, C. W. Kim and S. T. Petcov, Phys. Rev. D **54**, 4432 (1996) [hep-ph/9604364].
- [24] F. Feruglio, A. Strumia and F. Vissani, Nucl. Phys. B **637**, 345 (2002) [Addendum-*ibid.* B **659**, 359 (2003)] [arXiv:hep-ph/0201291]; S. Pascoli and S. T. Petcov, Phys. Lett. B **544**, 239 (2002) [arXiv:hep-ph/0205022]; S. M. Bilenky, C. Giunti, J. A. Grifols and E. Masso, Phys. Rept. **379** (2003) 69 [hep-ph/0211462]; S. Pascoli, S. T. Petcov and W. Rodejohann, Phys. Lett. B **558**, 141 (2003) [hep-ph/0212113], and references therein; F. R. Joaquim, Phys. Rev. D **68**, 033019 (2003) [arXiv:hep-ph/0304276].
- [25] Y. Hayato, talk presented at International Europhysics Conference On High-Energy Physics (HEP 2003), 17-23 Jul 2003, Aachen, Germany.
- [26] J. N. Bahcall, M. C. Gonzalez-Garcia, and C. Peña-Garay, in preparation.
- [27] G. L. Fogli, E. Lisi, D. Montanino and A. Palazzo, Phys. Rev. D **62**, 013002 (2000) [hep-ph/9912231].
- [28] Talk by K. Nishikawa at XXI International Symposium on Lepton and Photon Interactions at High Energies, 11-16 August 2003, Fermi National Accelerator Laboratory, Batavia, Illinois USA.
- [29] Talk by A. Bellerive at XXI International Symposium on Lepton and Photon Interactions at High Energies, 11-16 August 2003, Fermi National Accelerator Laboratory, Batavia, Illinois USA.
- [30] J. N. Bahcall and C. Peña-Garay, hep-ph/0305159.
- [31] Talk by K. Inoue at XXI International Symposium on Lepton and Photon Interactions at High Energies, 11-16 August 2003, Fermi National Accelerator Laboratory, Batavia, Illinois USA.
- [32] D. N. Spergel *et al.*, astro-ph/0302209.
- [33] S. Hannestad, JCAP **0305**, 004 (2003) [arXiv:astro-ph/0303076].
- [34] A. Pierce and H. Murayama, hep-ph/0302131.
- [35] H. V. Klapdor-Kleingrothaus, A. Dietz, H. L. Harney and I. V. Krivosheina, Mod. Phys. Lett. A **16**, 2409 (2001) [hep-ph/0201231].
- [36] M. C. Gonzalez-Garcia, M. Maltoni, C. Peña-Garay and J. W. Valle, Phys. Rev. D **63**, 033005 (2001) [hep-ph/0009350].
- [37] It is possible, however, that the leptogenesis occurs even with Dirac neutrinos [11].
- [38] Other new physics beyond the Standard Model, *e.g.*  $R$ -parity violating supersymmetry, Majoron, doubly charged Higgs, can also cause neutrinoless double-beta decay (see *e.g.*, [12] and references therein). However, such models induce Majorana mass for neutrinos from radiative corrections as well.
- [39] The positive evidence for neutrino oscillation from the LSND experiment [17] does not fit into the standard three-generation framework. We ignore this evidence in this letter.
- [40] The degenerate spectrum can further be either of normal or inverted hierarchy.
- [41] This statement assumes that there are only three light neutrinos mass eigenstates. HM thanks Xerxes Tata on this point. See [22] for a related discussion.