UC San Diego

UC San Diego Previously Published Works

Title

Overestimate of committed warming.

Permalink

<https://escholarship.org/uc/item/4xz8q3cw>

Journal

Nature, 547(7662)

ISSN

0028-0836

Authors

Schmidt, Gavin A
Severinghaus, Jeff
Abe-Ouchi, Ayako
et al.

Publication Date

2017-07-01

DOI

10.1038/nature22803

Peer reviewed

Overestimate of committed warming

ARISING FROM C. W. Snyder *Nature* **538**, 226–228 (2016); doi:10.1038/nature19798

Palaeoclimate variations are an essential component in constraining future projections of climate change as a function of increasing abundances of anthropogenic greenhouse gases¹. The Earth system sensitivity (ESS) describes the multi-millennial response of Earth (in terms of the change in global-mean temperature) to a doubling of atmospheric CO₂ concentrations. A recent study² used a correlation of inferred temperatures and radiative forcing from greenhouse gases over the past 800,000 years³ to estimate that the ESS from present-day CO₂ concentrations is about 9 °C and to imply a long-term commitment of 3–7 °C even if greenhouse gas concentrations remain at present-day levels. However, we demonstrate that the methodology of ref. 2 does not reliably estimate the ESS in the presence of orbital forcing of ice-age cycles and therefore conclude that the inferred² present-day committed warming is considerably overestimated. There is a Reply to this Comment by Snyder, C. W. *Nature* **547**, <http://dx.doi.org/10.1038/nature22804> (2017).

The previous analysis² was based on the assumption that greenhouse gases were solely responsible for long-term global-mean glacial–interglacial temperature changes. This is not correct^{4–7}. Although it is clear that greenhouse gases have a large role, quantifying that role is difficult because of simultaneous changes in many factors that also influence the energy balance of Earth (such as the extent of the ice sheets, snow cover, vegetation, dust load and cloud cover)⁴. However, it is widely accepted that orbital forcing is the ultimate trigger for glacial–interglacial cycles^{4–7}, enhanced by fast and slow feedbacks that involve the ice albedo, clouds, the carbon cycle, vegetation, and so on¹, sometimes resulting in hysteresis behaviour⁶. Therefore, the strong correlation that is seen in the datasets analysed in ref. 2 is a conflation of the sensitivity of the climate to CO₂ and the response of the carbon cycle to variations in temperature and ice-sheet extent. The Charney climate sensitivity (which includes fast atmospheric feedbacks, but not long-term changes in ice-sheet extent or in vegetation) can be constrained by these data by treating the long-term factors as forcings⁸. However, estimating the long-term sensitivity to greenhouse gas forcing alone requires constraints from periods that are not affected by the

interaction between orbital forcing and ice sheets, or that include a model-based assessment of the response to other forcings^{1,9–11}.

To illustrate the lack of connection between the ESS and the scaled regression of temperature and greenhouse gas forcing, we use a simple, coupled, three-component model¹² for land ice, temperature and carbon that incorporates the effects of orbital forcing on ice sheets, short- and long-term feedbacks to changes in atmospheric concentrations of CO₂, and two-way coupling between temperature and ice. On the basis of approximate differences in sea level, temperature and CO₂ concentrations between the pre-industrial era and the Last Glacial Maximum⁸, we fix the response of CO₂ to temperature (20 p.p.m. K^{−1}) and the radiative forcing related to ice (0.025 W m^{−2} per metre of sea-level equivalent), and vary the non-Planck climate feedback and the ice-sheet response to temperature to span a wide but plausible range of Charney and Earth system sensitivities. The system is driven by an external 80-kyr periodic signal that is applied directly to the ice-sheet component (Fig. 1a). We calculate the model ESS and the scaled regression over the glacial cycles (the linear slope in K W^{−1} m² multiplied by the canonical estimate of 2 × CO₂ forcing, 3.7 W m^{−2}; equivalent to the ESS presented in ref. 2) and plot their ratio (Fig. 1b). If the latter were a good estimate of the response to CO₂ forcing alone, then the ratio would be close to unity everywhere. This is clearly not the case—biases are very large and pervasive. For the chosen ranges of the model parameters, the scaled regression is a considerable overestimate of the actual model ESS.

This result shows that applying an aggregate regression from glacial periods, in which orbital forcing as well as greenhouse gases cause temperature variations, to the committed warming from current radiative forcing will probably overestimate future warming. In addition, given the current estimate of the radiative imbalance¹², the future changes to vegetation and ice sheets that would be required in response to current and committed short-term warming in order to produce long-term warming of 3–7 °C necessitate at least a doubling of the original forcing of 3.7 W m^{−2}. But such a doubling seems implausible, owing to the limited extent to which the areal coverage of current ice sheets can

Figure 1 | Glacial–interglacial cycles in a simple model with varying Earth system sensitivity. **a**, Two examples (with parameters as indicated in **b**) of synthetic temperature (black) and CO₂ (red) cycles over one million years, driven by an 80-kyr cycle in ice-sheet forcing. The cycles shown are obtained from a simple, three-component model for temperature, CO₂ and glacial ice (see Supplementary Information). A temperature anomaly

of 0 °C corresponds to a nominal average temperature over the glacial–interglacial cycles (see Supplementary Information). **b**, The ratio of the scaled regression (the linear regression coefficient in K W^{−1} m² multiplied by 3.7 W m^{−2}; equivalent to the ESS presented in ref. 2) to the model ESS over a plausible range of parameter space.

change and the expected range of effects on vegetation. Furthermore, the response to global forcing probably depends on the state of the climate^{1,13}. Any palaeo-derived ESS, if it is to be applied to the present-day radiative imbalance, must be defined in such a way that it estimates the effect of external radiative forcing only, and should be drawn from evidence from non-glacial base climate states.

In summary, we demonstrate that the ESS of about 9 °C defined in ref. 2 cannot be used to project future warming and that there is no reason to alter the most recent assessment of the present-day committed warming¹⁴.

Data Availability The R code to generate the data in Fig. 1 is available as Supplementary Information.

Gavin A. Schmidt¹, Jeff Severinghaus², Ayako Abe-Ouchi^{3,4}, Richard B. Alley⁵, Wallace Broecker⁶, Ed Brook⁷, David Etheridge⁸, Kenji Kawamura^{9,10,11}, Ralph F. Keeling², Margaret Leinen², Kate Marvel^{1,12} & Thomas F. Stocker¹³

¹NASA Goddard Institute for Space Studies, New York, New York 10025, USA. email: gavin.a.schmidt@nasa.gov

²Scripps Institution of Oceanography, University of California, San Diego, La Jolla, California 92093, USA.

³Atmosphere and Ocean Research Institute, University of Tokyo, Kashiwa, Japan.

⁴Japan Agency for Marine-Earth Science and Technology, Yokohama 236-0001, Japan.

⁵Department of Geosciences, Earth and Environmental Systems Institute, Pennsylvania State University, University Park, Pennsylvania 16802, USA.

⁶Lamont-Doherty Earth Observatory, Columbia University, Palisades, New York, New York 10964, USA.

⁷College of Earth, Ocean, and Atmospheric Sciences, Oregon State University, Corvallis, Oregon 97331, USA.

⁸CSIRO Climate Science Centre, Aspendale, Victoria, Australia.

⁹National Institute for Polar Research, Tachikawa, Tokyo 190-8518, Japan.

¹⁰Department of Polar Science, SOKENDAI (The Graduate University for Advanced Studies), Tachikawa, Tokyo 190-8518, Japan.

¹¹Institute of Biogeosciences, Japan Agency for Marine-Earth Science and Technology, Yokosuka 237-0061, Japan.

Snyder replies

REPLYING TO G. A. Schmidt *et al.* *Nature* **547**, <http://dx.doi.org/10.1038/nature22803> (2017)

The Earth system sensitivity (ESS) summarizes the feedback behaviour of Earth's climate system and includes ice sheets, vegetation and dust as internal feedbacks. This definition of the ESS is in contrast to that of the equilibrium climate sensitivity (ECS), for which those feedbacks are considered external forcings^{1,2}. As previously quantified from palaeoclimate records, the ESS^{3–7} and palaeoclimate sensitivity parameter $S_{\text{[GHG]}}$ (ref. 2) do not test causation, but summarize the past aggregate, correlational relationships among those feedbacks. In the accompanying Comment⁸, Schmidt *et al.* contend that I miscalculated the ESS by defining it to include all changes to ice sheets as internal feedbacks⁹. However, previous research had applied the same definition of the ESS^{3–7} or $S_{\text{[GHG]}}$ (ref. 2), yielding estimates of the ESS of 8.6 ± 2.8 °C (1σ)², 7.4 °C (6.2–9.1 °C, 95% confidence interval)³ and 6 °C (refs 4, 5) from the late Pleistocene and exceeding 6 °C from Cenozoic periods with large ice sheets⁶. These values are comparable to my estimate of 9 °C (7–13 °C, 95% credible interval)⁹. Moreover,

¹²Department of Applied Physics and Applied Mathematics, Columbia University, New York, New York 10025, USA.

¹³Climate and Environmental Physics, Physics Institute, University of Bern, Sidlerstrasse 5, CH-3012 Bern, Switzerland.

Received 17 November 2016; accepted 3 April 2017.

1. PALAEOSSENS Project Members. Making sense of palaeoclimate sensitivity. *Nature* **491**, 683–691 (2012).
2. Snyder, C. W. Evolution of global temperature over the past two million years. *Nature* **538**, 226–228 (2016).
3. Bereiter, B. *et al.* Revision of the EPICA Dome C CO₂ record from 800 to 600 kyr before present. *Geophys. Res. Lett.* **42**, 542–549 (2015).
4. Hays, J. D., Imbrie, J. & Shackleton, N. J. Variations in the Earth's orbit: pacemaker of the ice ages. *Science* **194**, 1121–1132 (1976).
5. Raymo, M. E., Lisiecki, L. E. & Nisancioglu, K. H. Plio-Pleistocene ice volume, Antarctic climate, and the global $\delta^{18}\text{O}$ record. *Science* **313**, 492–495 (2006).
6. Abe-Ouchi, A. *et al.* Insolation-driven 100,000-year glacial cycles and hysteresis of ice-sheet volume. *Nature* **500**, 190–193 (2013).
7. Paillard, D. Quaternary glaciations: from observations to theories. *Quat. Sci. Rev.* **107**, 11–24 (2015).
8. Köhler, P. *et al.* What caused Earth's temperature variations during the last 800,000 years? Data-based evidence on radiative forcing and constraints on climate sensitivity. *Quat. Sci. Rev.* **29**, 129–145 (2010).
9. Hansen, J. *et al.* Target atmospheric CO₂: where should humanity aim? *Open Atmos. Sci. J.* **2**, 217–231 (2008).
10. Lunt, D. J. *et al.* Earth system sensitivity inferred from Pliocene modelling and data. *Nat. Geosci.* **3**, 60–64 (2010).
11. Pagani, M., Liu, Z., LaRiviere, J. & Ravelo, A. C. High Earth-system climate sensitivity determined from Pliocene carbon dioxide concentrations. *Nat. Geosci.* **3**, 27–30 (2010).
12. Johnson, G. C. *et al.* in *State of the Climate in 2015* (eds Blunden, J. & Arndt, D. S.) *Bull. Am. Meteorol. Soc.* **97** (Suppl.), S66–S70 (2016).
13. von der Heydt, A. S. *et al.* Lessons on climate sensitivity from past climate changes. *Curr. Clim. Change Rep.* **2**, 148–158 (2016).
14. Collins, M. *et al.* in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (eds Stocker, T. F. *et al.*) Ch. 12 (Cambridge Univ. Press, 2013).

Supplementary Information accompanies this Comment.

Author Contributions G.A.S. and J.S. jointly conceived this Comment, G.A.S. and K.M. developed the simple model, and all authors contributed equally to the writing.

Competing Financial Interests Declared none.

doi:10.1038/nature22803

my estimates of the ESS⁹ are consistent with recent IPCC estimates of the ECS¹.

The primary debate regards the utility of the concept of ESS for providing insights relevant to future warming. I estimated the ESS from the late Pleistocene and applied this estimate to current greenhouse gas concentrations to quantify the implied future warming over millennial timescales⁹. The purpose of this was to provide a general perspective on potential warming under the assumption that the correlational relationship between greenhouse gas concentrations and global temperature from mid-glacial and interglacial conditions over the past 800 kyr will be similar in the future. This approach has two main sources of uncertainty.

First, it assumes that, at equilibrium, the relationships between the internal climate feedbacks will be the same for different causal triggers—that is, regardless of the initial source of change to the global energy balance (such as changes in orbital forcing, ice-sheet extent, temperature or

greenhouse gas concentrations), the resultant internal feedbacks iterate until they converge at equilibrium to the same aggregate relationship. In complexity theory, such a phenomenon is referred to as emergence. If instead the ESS depends on the source of the initial trigger, then the ESS calculated from the past 800 kyr would not be applicable to future warming triggered by anthropogenic emissions.

Second, although I found that the ESS was constant across mid-glacial and interglacial conditions over the past 800 kyr, and was higher under those conditions than under deep glacial conditions⁹, states warmer than the present day may have different ESS.

How much Earth will warm over millennial timescales in response to anthropogenic emissions remains uncertain. For example, in ref. 2 (see figure 4) it is assumed that no longer-timescale feedbacks occur during future warming. Yet we already observe that some longer-timescale feedbacks that are not included in the ECS, such as changes in ice-sheet extent, sea level and vegetation, are occurring today¹.

The modelling presented in Schmidt *et al.*⁸ does not add further insight to this discussion. They assert that a certain amount of change in ice-sheet extent is caused by orbital forcing that is entirely independent of interactive changes with temperature or greenhouse gas concentrations, and therefore that a certain amount of changes to ice sheets should not be included as internal feedbacks in calculating the ESS. This conflicts with previous work on the ESS^{3–7}. Although the causes of the glacial–interglacial quasi-cycles of the late Pleistocene continue to be debated^{10–18}, most theories include iterative feedbacks between temperature, ice sheets, greenhouse gases and other key climate features regardless of the initial trigger^{10–18}.

The simple model of Schmidt *et al.*⁸ does not test their assertions about the ESS. Rather, the construction of their model (via the term F_1 for orbitally driven changes in ice-sheet extent) and the equation that they use to quantify the ESS (in which $F_1 = 0$) predetermine their new estimate of the ESS. These factors also predetermine that the ratio that Schmidt *et al.* present between the two different calculations of the ESS is variable, not equal to one. Moreover, Schmidt *et al.*⁸ do not provide justification from published research or palaeoclimate data for their exclusion from the calculation of the ESS of specific changes to ice sheets.

The views expressed here are those of the author and do not necessarily reflect the views or policies of the US Environmental Protection Agency.

Carolyn W. Snyder¹

¹Interdisciplinary Program in Environment and Resources, Stanford University, Stanford, California 94305, USA.
email: carolyn.snyder@gmail.com

1. Stocker, T. F. *et al.* (eds) *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (Cambridge Univ. Press, 2013).
2. PALAESENS Project Members. Making sense of palaeoclimate sensitivity. *Nature* **491**, 683–691 (2012).
3. Martínez-Botí, M. A. *et al.* Plio-Pleistocene climate sensitivity evaluated using high-resolution CO₂ records. *Nature* **518**, 49–54 (2015).
4. Hansen, J. *et al.* Climate change and trace gases. *Phil. Trans. R. Soc. Lond. A* **365**, 1925–1954 (2007).
5. Hansen, J. *et al.* Target atmospheric CO₂: where should humanity aim? *Open Atmos. Sci. J.* **2**, 217–231 (2008).
6. Royer, D. L., Pagani, M. & Beerling, D. J. Geobiological constraints on Earth system sensitivity to CO₂ during the Cretaceous and Cenozoic. *Geobiology* **10**, 298–310 (2012).
7. Pagani, M., Liu, Z., LaRiviere, J. & Ravelo, A. C. High Earth-system climate sensitivity determined from Pliocene carbon dioxide concentrations. *Nat. Geosci.* **3**, 27–30 (2010).
8. Schmidt, G. A. *et al.* Overestimate of committed warming. *Nature* **547**, <http://dx.doi.org/10.1038/nature22803> (2017).
9. Snyder, C. W. Evolution of global temperature over the past two million years. *Nature* **538**, 226–228 (2016).
10. Raymo, M. E. The timing of major climate terminations. *Paleoceanography* **12**, 577–585 (1997).
11. Tziperman, E. & Gildor, H. On the mid-Pleistocene transition to 100-kyr glacial cycles and the asymmetry between glaciation and deglaciation times. *Paleoceanography* **18**, 1–1–8 (2003).
12. Huybers, P. & Wunsch, C. A depth-derived Pleistocene age model: uncertainty estimates, sedimentation variability, and nonlinear climate change. *Paleoceanography* **19**, PA1028 (2004).
13. Huybers, P. Early Pleistocene glacial cycles and the integrated summer insolation forcing. *Science* **313**, 508–511 (2006).
14. Clark, P. U. *et al.* The middle Pleistocene transition: characteristics, mechanisms, and implications for long-term changes in atmospheric pCO₂. *Quat. Sci. Rev.* **25**, 3150–3184 (2006).
15. Imbrie, J. Z., Imbrie-Moore, A. & Lisiecki, L. E. A phase-space model for Pleistocene ice volume. *Earth Planet. Sci. Lett.* **307**, 94–102 (2011).
16. Ganopolski, A. & Calov, R. The role of orbital forcing, carbon dioxide and regolith in 100 kyr glacial cycles. *Clim. Past* **7**, 1415–1425 (2011).
17. McClymont, E. L., Sosdian, S. M., Rosell-Melé, A. & Rosenthal, Y. Pleistocene sea-surface temperature evolution: early cooling, delayed glacial intensification, and implications for the mid-Pleistocene climate transition. *Earth Sci. Rev.* **123**, 173–193 (2013).
18. Abe-Ouchi, A. *et al.* Insolation-driven 100,000-year glacial cycles and hysteresis of ice-sheet volume. *Nature* **500**, 190–193 (2013).

doi:10.1038/nature22804