

UC Berkeley

Faculty Publications

Title

Compatibility and accessibility: lattice representations for semantics of non-classical and modal logics

Permalink

<https://escholarship.org/uc/item/4z83s9z9>

Author

Holliday, Wesley Halcrow

Publication Date

2022-07-07

Data Availability

The data associated with this publication are available at:

<https://github.com/wesholliday/compat-frames>

Copyright Information

This work is made available under the terms of a Creative Commons Attribution-NonCommercial-NoDerivatives License, available at

<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Peer reviewed

Compatibility and accessibility: lattice representations for semantics of non-classical and modal logics

Wesley H. Holliday

University of California, Berkeley

Forthcoming in *Advances in Modal Logic*, Vol. 14.

Abstract

In this paper, we study three representations of lattices by means of a set with a binary relation of compatibility in the tradition of Ploščica. The standard representations of complete ortholattices and complete perfect Heyting algebras drop out as special cases of the first representation, while the second covers arbitrary complete lattices, as well as complete lattices equipped with a negation we call a proto-complementation. The third topological representation is a variant of that of Craig, Haver, and Priestley. We then extend each of the three representations to lattices with a multiplicative unary modality; the representing structures, like so-called graph-based frames, add a second relation of accessibility interacting with compatibility. The three representations generalize possibility semantics for classical modal logics to non-classical modal logics, motivated by a recent application of modal orthologic to natural language semantics.

Keywords: lattices, representation theorems, ortholattices, orthologic, Heyting algebras, intuitionistic logic, Boolean algebras, modal logic, negation, graph-based frames, possibility semantics

1 Introduction

Semantics for non-classical and modal logics may be seen as arising from more basic algebraic representation theorems. For example, traditional semantics for intuitionistic logic, orthologic, and classical modal logic may be understood in terms of the following well-known representations:

- Any Heyting algebra H embeds into the lattice of downsets of a poset, and the embedding is an isomorphism if H is complete and its completely join-irreducible elements are join-dense (see, e.g., [22], [17, Prop. 1.1]).
- Any ortholattice L embeds into the lattice of \perp -closed sets of an orthoframe (X, \perp) equipped with an orthocomplementation \neg induced by the relation \perp , and the embedding is an isomorphism if L is complete (see [29] for the orthoframe description and [6, §§ 32-4], [19] for other descriptions).
- Any Boolean algebra B equipped with a multiplicative unary operation \square

embeds into the powerset of a set W equipped with an operation \Box_R induced by a binary relation R on W , and the embedding is an isomorphism if B is complete and atomic and \Box completely multiplicative (see [46,38,47]).

In each case, adding topology to the relevant relational structures allows one to characterize topologically the image of the relevant embedding [22,30,5,27,44].

Here we study representations that subsume and go beyond all of those mentioned above. In § 2, we explain how to go from a set together with a binary relation of “compatibility” to a complete lattice. In § 3, we study three ways of going back: one economical representation of certain complete lattices, including but not limited to Heyting and ortholattice cases; one less economical but fully general representation of complete lattices, including complete lattices equipped with a type of negation that we call a protocomplementation; and one representation of arbitrary lattices. In § 4, we extend the three representations to lattices with a multiplicative unary modality \Box , by adding a second relation of accessibility interacting with compatibility. We conclude in § 5.

After writing this paper, I discovered Ploščica’s [43] representation of bounded lattices using certain compatibility frames¹ as in Definition 2.1 with a topology, as well as Craig et al.’s [16] modification of Ploščica’s approach. In § 3.3, we briefly cover a variant of this representation with a different topology. In addition, a referee informed me that the addition of modal accessibility interacting with compatibility (Definition 4.2) appears in the *graph-based frames* of Conradie et al. [10]. We will return to this connection in § 4.

A Jupyter notebook with code to verify examples and investigate conjectures and questions is available at github.com/wesholliday/compat-frames.

2 From compatibility frames to lattices

2.1 Basic concepts

Our starting point is a certain way of going from a set with a binary relation to a complete lattice. For a comparison with other ways of realizing complete lattices using doubly ordered structures and polarities, see [34].

Definition 2.1 A *relational frame* is a pair $\mathcal{F} = (X, \triangleleft)$ where X is a nonempty set and \triangleleft is a binary relation on X . A *compatibility frame* is a relational frame in which \triangleleft is reflexive.

We read $x \triangleleft y$ as “ x is compatible with y ,” also written $y \triangleright x$.²

Convention 2.2 In diagrams, such as Fig. 1, an arrow with a triangle arrowhead from y to x indicates $y \triangleright x$. Thus, we draw the directed graph (X, \triangleright) to represent the compatibility frame (X, \triangleleft) . Reflexive loops are not shown.

Recall that a unary operation on a lattice is a *closure operator* if c is inflationary ($x \leq c(x)$), idempotent ($c(c(x)) = c(x)$), and monotone ($x \leq y$ implies

¹ Also see [15] for *TiRS graphs*, which are compatibility frames with extra properties, which we do not require here (as is crucial for a number of our results and for Conjecture 3.13).

² In [34], we wrote $x \check{<} y$ and $y \check{>}^{-1} x$ instead of $x \triangleleft y$ and $y \triangleright x$, respectively.

$c(x) \leq c(y)$). We will use the compatibility relation \triangleleft to define a closure operator on $\wp(X)$, whose fixpoints give us a complete lattice as in the following classic result (see, e.g., [8, Thm. 5.2]).

Proposition 2.3 *Let X be a nonempty set and c a closure operator on $\wp(X)$. Then the fixpoints of c , i.e., those $A \subseteq X$ with $c(A) = A$, ordered by \subseteq form a complete lattice with*

$$\bigwedge_{i \in I} A_i = \bigcap_{i \in I} A_i \text{ and } \bigvee_{i \in I} A_i = c\left(\bigcup_{i \in I} A_i\right).$$

Definition 2.4 Given a relational frame (X, \triangleleft) , define $c_{\triangleleft} : \wp(X) \rightarrow \wp(X)$ by

$$c_{\triangleleft}(A) = \{x \in X \mid \forall x' \triangleleft x \exists x'' \triangleright x' : x'' \in A\}.$$

Thus, x is in $c_{\triangleleft}(A)$ iff every state compatible with x is compatible with some state in A . Given a compatibility frame, we are interested in the c_{\triangleleft} -fixpoints, i.e., those $A \subseteq X$ such that $c_{\triangleleft}(A) = A$. Looking at a diagram of a compatibility frame, one can check that $c_{\triangleleft}(A) = A$ by checking that the following holds:

- from any $x \in X \setminus A$, you can step forward along an arrow to a state x' that cannot step backward along an arrow into A .

Informally, “from x you can see a state that cannot be seen from A .”

Example 2.5 Consider the cycle on three elements on the left of Fig. 1, regarded as a compatibility frame according to Convention 2.2: $\{y\}$ is a c_{\triangleleft} -fixpoint because z and x can both see x , which cannot be seen from $\{y\}$. Yet $\{y, z\}$ is not a c_{\triangleleft} -fixpoint, because x cannot see a state that cannot be seen from $\{y, z\}$, since both x and y can be seen from $\{y, z\}$.

We get the reverse verdicts on $\{y\}$ and $\{y, z\}$ in the acyclic (ignoring loops) but non-transitive frame on the right of Fig. 1: $\{y\}$ is *not* a c_{\triangleleft} -fixpoint, because now z cannot see a state that cannot be seen from $\{y\}$; but $\{y, z\}$ *is* a c_{\triangleleft} -fixpoint, because x can see a state, namely x , that cannot be seen from $\{y, z\}$.




Fig. 1. Two compatibility frames drawn according to Convention 2.2 with their c_{\triangleleft} -fixpoints (except X and \emptyset) outlined, followed by their associated lattices.

Proposition 2.6 *For any relational frame, c_{\triangleleft} is a closure operator on $\wp(X)$.*

Proof. That $Y \subseteq c_{\triangleleft}(Y)$ and that $Y \subseteq Z$ implies $c_{\triangleleft}(Y) \subseteq c_{\triangleleft}(Z)$ are obvious. To see $c_{\triangleleft}(c_{\triangleleft}(Y)) \subseteq c_{\triangleleft}(Y)$, suppose $x \in c_{\triangleleft}(c_{\triangleleft}(Y))$ and $x' \triangleleft x$. Hence there is an $x'' \triangleright x'$ with $x'' \in c_{\triangleleft}(Y)$. This implies there is an $x''' \triangleright x'$ with $x''' \in Y$. Thus, for any $x' \triangleleft x$ there is an $x''' \triangleright x'$ with $x''' \in Y$. Therefore, $x \in c_{\triangleleft}(Y)$. \square

Given Propositions 2.3 and 2.6, we have the following immediate corollary.

Corollary 2.7 *For any relational frame (X, \triangleleft) , the c_{\triangleleft} -fixpoints ordered by \subseteq form a complete lattice $\mathfrak{L}(X, \triangleleft)$ with meet and join as in Proposition 2.3.*

Example 2.8 We see in Fig. 1 that the \mathbf{M}_3 lattice (ignoring the dashed arrows for now) arises from the cycle on three elements, while the \mathbf{N}_5 lattice arises from the acyclic but non-transitive frame on three elements.

We can relate Corollary 2.7 to possible world semantics for classical and intuitionistic logic as follows. Let $=_W$ be the identity relation on the set W .

Proposition 2.9

- (i) *Given any set W , the pair $(W, =_W)$ is a compatibility frame, and $\mathfrak{L}(W, =_W)$ is the Boolean algebra of all subsets of W .*
- (ii) *Given any preorder \leq on a set P , the pair (P, \leq) is a compatibility frame, and $\mathfrak{L}(P, \leq)$ is the Heyting algebra of all downsets of (P, \leq) .*

Part (ii) appears in [13, Prop. 4.1.1] but we include a proof for convenience.

Proof. Part (i) is obvious. For (ii), let A be a downset and $x \in P \setminus A$. Setting $x' = x$, we have $x' \triangleleft x$, and for all $x'' \triangleright x'$, i.e., all $x'' \geq x'$, $x'' \notin A$, since A is a downset. Thus, $c_{\triangleleft}(A) = A$. Conversely, suppose $c_{\triangleleft}(A) = A$, $x \in A$, and $y \leq x$. Let $y' \triangleleft y$, so $y' \leq y$ and hence $y' \leq x$. Then setting $y'' = x$, we have $y' \triangleleft y'' \in A$. Since $c_{\triangleleft}(A) = A$, it follows that $y \in A$. Thus, A is a downset. \square

Example 2.10 To illustrate part (ii), if we add to the non-transitive frame in Fig. 1 the arrow from x to z required by transitivity, then instead of realizing \mathbf{N}_5 , we realize the four-element chain in Fig. 2, which is a Heyting algebra.




Fig. 2. A compatibility frame realizing a Heyting algebra.

Some appealing aspects of working with downsets of a preorder also apply to our fixpoints; let ∂ , \oplus , $\overline{\oplus}$, $\dot{\cup}$, and \times be the dual, linear sum, vertical sum, disjoint union, and product operations [18, § 1.31, § 1.24, Ex. 1.18, Prop. 1.32].

Proposition 2.11 *For any relational frames \mathcal{F} and \mathcal{G} , (i) $\mathfrak{L}(\mathcal{F}^\partial) \cong \mathfrak{L}(\mathcal{F})^\partial$; (ii) $\mathfrak{L}(\mathcal{F} \oplus \mathcal{G}) \cong \mathfrak{L}(\mathcal{F}) \overline{\oplus} \mathfrak{L}(\mathcal{G})$; and (iii) $\mathfrak{L}(\mathcal{F} \dot{\cup} \mathcal{G}) \cong \mathfrak{L}(\mathcal{F}) \times \mathfrak{L}(\mathcal{G})$.*

Proof. Let $\mathcal{F} = (X, \triangleleft)$, so $\mathcal{F}^\partial = (X, \triangleright)$. For an isomorphism φ from $\mathfrak{L}(\mathcal{F}^\partial)$ to $\mathfrak{L}(\mathcal{F})^\partial$, let $\varphi(A) = \{x \in X \mid \forall y \triangleleft x \ y \notin A\}$. From $\mathfrak{L}(\mathcal{F}) \overline{\oplus} \mathfrak{L}(\mathcal{G})$ to $\mathfrak{L}(\mathcal{F} \oplus \mathcal{G})$, $\varphi(A) = A$ if $A \in \mathfrak{L}(\mathcal{F})$, and $\varphi(A) = A \cup X$ if $A \in \mathfrak{L}(\mathcal{G})$. Part (iii) is also easy. \square

We can also relate our approach to that of realizing complete Boolean algebras as in forcing [45] or possibility semantics [33] for classical logic as follows.

Proposition 2.12 *Given a preordered set (P, \leq) , define \triangleleft on P by: $x \triangleleft y$ if $\exists z \in P: z \leq x$ and $z \leq y$. Then (P, \triangleleft) is a compatibility frame, and $\mathfrak{L}(P, \triangleleft)$ is the Boolean algebra $\mathcal{RO}(P, \leq)$ of all regular open downsets of (P, \leq) .*

Proof. Observe that any c_{\triangleleft} -fixpoint is a \leq -downset, and for any \leq -downset A , (i) $\forall x' \leq x \exists x'' \leq x' : x'' \in A$ and (ii) $\forall x' \triangleleft x \exists x'' \triangleright x' : x'' \in A$ are equivalent. It follows that A is a regular open downset iff A is a c_{\triangleleft} -fixpoint. \square

Now (X, \triangleleft) gives us not only $\mathfrak{L}(X, \triangleleft)$ but also an operation \neg_{\triangleleft} on $\mathfrak{L}(X, \triangleleft)$.

Proposition 2.13 *For any relational frame (X, \triangleleft) and $A \subseteq X$, the set $\neg_{\triangleleft}A = \{x \in X \mid \forall y \triangleleft x \ y \notin A\}$ is a c_{\triangleleft} -fixpoint.*

Proof. If $x \in X \setminus \neg_{\triangleleft}A$, then $\exists x' \triangleleft x$ with $x' \in A$, so $\forall x'' \triangleright x', x'' \notin \neg_{\triangleleft}A$. \square

To characterize the \neg_{\triangleleft} operation, let us recall some terminology.

Definition 2.14 Let L be a bounded lattice and $a \in L$. An $x \in L$ is a *semicomplement* of a if $a \wedge x = 0$, a *complement* of a if $a \wedge x = 0$ and $a \vee x = 1$, and a *pseudocomplement* of a if x is the maximum in L of $\{y \in L \mid a \wedge y = 0\}$.

A unary operation \neg on L is a *semicomplementation* (resp. *complementation*, *pseudocomplementation*) if for all $a \in L$, $\neg a$ is a semicomplement (resp. complement, pseudocomplement) of a . It is *antitone* if for all $a, b \in L$, $a \leq b$ implies $\neg b \leq \neg a$, *involutive* if $\neg \neg a = a$ for all $a \in L$, and *anti-inflationary* if $a \not\leq \neg a$ for all nonzero $a \in L$. An *ortholattice* is a bounded lattice equipped with an involutive antitone complementation, called an *orthocomplementation*. A *p-algebra* is a bounded lattice equipped with a pseudocomplementation.

Finally, for a non-standard piece of terminology, we say \neg is a *protocomplementation* if \neg is an antitone semicomplementation such that $\neg 0 = 1$.

An antitone \neg is anti-inflationary iff it is a semicomplementation. Also recall that the operation in a Heyting algebra H defined by $\neg a = a \rightarrow 0$ is a pseudocomplementation, so H may also be regarded as a p-algebra; and the complementation in a Boolean algebra B is an orthocomplementation, so B is an ortholattice. As for the operation \neg_{\triangleleft} , the following is easy to check.

Proposition 2.15 *For any compatibility (resp. relational) frame (X, \triangleleft) , \neg_{\triangleleft} is a protocomplementation (resp. antitone and such that $\neg_{\triangleleft}1 = 0$) on $\mathfrak{L}(X, \triangleleft)$.*

In our diagrams of lattices arising from compatibility frames, the dashed arrows represent the operation \neg_{\triangleleft} . We omit arrows representing $\neg 0 = 1$ and $\neg 1 = 0$.

2.2 Frames for ortholattices

If we assume that \triangleleft is *symmetric*, then we get a standard representation (as in [29] via “proximity frames”) of ortholattices. The proof is straightforward.

Proposition 2.16 *For any compatibility frame (X, \triangleleft) , if \triangleleft is symmetric, then \neg_{\triangleleft} is an orthocomplementation on $\mathfrak{L}(X, \triangleleft)$.*

Example 2.17 Fig. 3 shows two symmetric compatibility frames and their associated ortholattices, \mathbf{MO}_2 or \mathbf{M}_4 (left) and the Benzene ring \mathbf{O}_6 (right).




Fig. 3. Compatibility frames realizing ortholattices.

2.3 Frames for Heyting and Boolean algebras

We conclude this section with sufficient conditions on \triangleleft for realizing Heyting and Boolean algebras, weaker than those in Proposition 2.9 (where \triangleleft was a preorder or the identity relation). First, we define some auxiliary notions.

Definition 2.18 Given a compatibility frame (X, \triangleleft) and $x, y \in X$:

- (i) x *pre-refines* y , written $x \sqsubseteq_{pr} y$, if for all $z \in X$, $z \triangleleft x$ implies $z \triangleleft y$;
- (ii) x *post-refines* y , written $x \sqsubseteq_{po} y$, if for all $z \in X$, $x \triangleleft z$ implies $y \triangleleft z$;
- (iii) x *refines* y , written $x \sqsubseteq y$, if x pre-refines and post-refines y ;
- (iv) x is *compossible with* y if there is a $w \in X$ that refines x and pre-refines y .

Note that if \triangleleft is symmetric, then pre-refinement and post-refinement are equivalent, and x is compossible with y just in case they have a common refinement.

Lemma 2.19 For any compatibility frame (X, \triangleleft) , \sqsubseteq_{pr} and \sqsubseteq_{po} are preorders on X . Moreover, each c_{\triangleleft} -fixpoint is a \sqsubseteq_{pr} -downset.

Proof. The preorder part is obvious. Next suppose A is a c_{\triangleleft} -fixpoint, $x \in A$, and $y \sqsubseteq_{pr} x$. Toward showing that $y \in A$, consider any $y' \triangleleft y$. Then since $y \sqsubseteq_{pr} x$, we have $y' \triangleleft x$, so taking $y'' = x$, we have shown that for every $y' \triangleleft y$ there is a $y'' \triangleright y'$ with $y'' \in A$. Since A is a c_{\triangleleft} -fixpoint, it follows that $y \in A$. \square

Definition 2.20 A *compossible compatibility frame* is a compatibility frame (X, \triangleleft) in which for any $x, y \in X$, if $x \triangleleft y$, then x is compossible with y .

Theorem 2.21

- (i) If (X, \triangleleft) is a compossible compatibility frame, then $\mathfrak{L}(X, \triangleleft)$ is a Heyting algebra with \rightarrow defined by $A \rightarrow B = \{x \in X \mid \forall y \sqsubseteq_{pr} x (y \in A \Rightarrow y \in B)\}$, and \neg_{\triangleleft} is the Heyting pseudocomplementation.
- (ii) If (X, \triangleleft) is a compossible symmetric compatibility frame, then $\mathfrak{L}(X, \triangleleft)$ is the Boolean algebra $\mathcal{RC}(X, \sqsubseteq)$, and \neg_{\triangleleft} is the Boolean complementation.

Proof. For part (i), recall that a *nucleus* on a Heyting algebra H is a closure operator that is also multiplicative, i.e., for all $x, y \in H$, we have $c(x) \wedge c(y) \leq c(x \wedge y)$. This follows from c being a closure operator such that for all $a, b \in H$, $a \wedge c(b) \leq c(a \wedge b)$. For then setting $a = c(x)$ and $b = y$, $c(x) \wedge c(y) \leq c(c(x) \wedge y) = c(y \wedge c(x))$. Then setting $a = y$ and $b = x$, we have $y \wedge c(x) \leq c(y \wedge x)$, in which

case monotonicity and idempotence yield $c(y \wedge c(x)) \leq c(c(y \wedge x)) = c(y \wedge x) = c(x \wedge y)$. Thus, combining the two long strings of equations, $c(x) \wedge c(y) \leq c(x \wedge y)$.

Now it is well known that the fixpoints of a nucleus j on a (complete) Heyting algebra H form a (complete) Heyting algebra H_j under the restricted lattice order (see, e.g., [21, p. 71]), where for $a, b \in H_j$, $a \rightarrow_{H_j} b = a \rightarrow_H b$.

Let H be the Heyting algebra of all \sqsubseteq_{pr} -downsets. By the results above, to prove the first part of (i), it suffices to show that c_{\triangleleft} restricted to H is a nucleus, for which it suffices to show that for all $A, B \in H$, $A \cap c_{\triangleleft}(B) \subseteq c_{\triangleleft}(A \cap B)$. Suppose $x \in A \cap c_{\triangleleft}(B)$ but $x \notin c_{\triangleleft}(A \cap B)$. Since $x \notin c_{\triangleleft}(A \cap B)$, there is a $y \triangleleft x$ such that (\star) for all $y' \triangleright y$, we have $y' \notin A \cap B$. Since $y \triangleleft x$, by compossibility there is a z that refines y and pre-refines x . Since z pre-refines x , $z \triangleleft x$. Then since $x \in c_{\triangleleft}(B)$, there is a $z' \triangleright z$ with $z' \in B$. Since $z \triangleleft z'$, there is a w that refines z and pre-refines z' . Since w pre-refines z' , from $z' \in B$ we have $w \in B$ by Lemma 2.19. Since w pre-refines z and z pre-refines x , w pre-refines x , so $x \in A$ implies $w \in A$. Thus, $w \in A \cap B$. Moreover, since w post-refines z and z post-refines y , w post-refines y . Hence $y \triangleleft w$. But then by (\star) , $w \notin A \cap B$. This is a contradiction. Finally, for the claim about \neg_{\triangleleft} , observe that for any \sqsubseteq_{pr} -downset A , we have $A \rightarrow \emptyset = \{x \in X \mid \forall y \sqsubseteq_{pr} x \ y \notin A\} = \neg_{\triangleleft} A$; for the second equality, the right-to-left inclusion uses that $y \sqsubseteq_{pr} x$ implies $y \triangleleft x$, while the left-to-right inclusion uses the assumption of compossibility.

For part (ii), to see that $\mathfrak{L}(X, \triangleleft)$ is isomorphic to $\mathcal{RO}(X, \sqsubseteq)$, by Proposition 2.12 it suffices to show that for all $x, y \in X$, we have $x \triangleleft y$ iff there is a $z \in X$ such that $z \sqsubseteq x$ and $z \sqsubseteq y$. From left to right, if $x \triangleleft y$, then since (X, \triangleleft) is a compossible compatibility frame, there is a z that refines x and pre-refines y , which implies that z refines y by the symmetry of \triangleleft . From right to left, $z \sqsubseteq x$ implies $x \triangleleft z$, which with $z \sqsubseteq y$ implies $x \triangleleft y$. Finally, for the claim about \neg_{\triangleleft} , recall that $\neg A$ in $\mathcal{RO}(X, \sqsubseteq)$ is $\{x \in X \mid \forall y \sqsubseteq x \ y \notin A\}$, which is equal to $\neg_{\triangleleft} A$ by reasoning analogous to that at the end of the previous paragraph. \square

3 From lattices to compatibility frames

3.1 Representation of special complete lattices via join-dense sets

In this section, we give an economical representation of certain complete lattices L using compatibility frames based on a join-dense set V of nonzero elements of L , so the frame representing L is smaller than L . In all our figures, the frame can be seen as obtained from the lattice via this representation.

Let $CJI(L)$ be the set of completely join-irreducible elements of L . Recall that a complete Heyting algebra is *perfect* if $CJI(L)$ is join-dense in L . The standard representations of complete perfect Heyting algebras and complete ortholattices drop out of the representation in this section as special cases.

3.1.1 Suitable compatibility relations

Before giving the crucial definition of the compatibility relation used in our representation (Definition 3.5(i) below), it is instructive to distill conditions on a compatibility relation sufficient for a successful representation.

Definition 3.1 Let L be a lattice and V a set of elements of L . A binary

relation \triangleleft on V is *suitable for L* if \triangleleft is reflexive and:

- (i) for $a \in V$ and $b \in L$, if $a \not\leq b$, then $\exists a' \triangleleft a \forall a'' \triangleright a' a'' \not\leq b$;
- (ii) if $a \in V$, B is a c_{\triangleleft} -fixpoint of (V, \triangleleft) with a join b in L , and $a \leq b$, then $a \in B$.

We call (i) and (ii) the first and second *suitability conditions*, respectively.

Proposition 3.2 *Let L be a lattice, V a join-dense set of elements of L , and \triangleleft a reflexive relation on V . For $b \in L$, define $\varphi(b) = \{x \in V \mid x \leq b\}$.*

- (i) *If \triangleleft satisfies the first suitability condition, then φ embeds L into $\mathfrak{L}(V, \triangleleft)$.*
- (ii) *If L is complete and \triangleleft suitable, φ is an isomorphism from L to $\mathfrak{L}(V, \triangleleft)$.*

Proof. For part (i), clearly φ is order-preserving: if $a \leq b$, then $\varphi(a) \subseteq \varphi(b)$. Moreover, φ is order-reflecting: since V is join-dense in L , we have $a = \bigvee A$ for some $A \subseteq V$, so $a \not\leq b$ implies that for some $a_0 \in A$, we have $a_0 \not\leq b$, so $a_0 \in \varphi(a)$ but $a_0 \notin \varphi(b)$, and hence $\varphi(a) \not\subseteq \varphi(b)$.

Next we claim $\varphi(b)$ is a c_{\triangleleft} -fixpoint. Suppose for $a \in V$ that $a \notin \varphi(b)$, so $a \not\leq b$. Then by the first suitability condition, there is an $a' \triangleleft a$ such that for all $a'' \triangleright a'$, $a'' \not\leq b$ and hence $a'' \notin \varphi(b)$. Thus, $\varphi(b)$ is a c_{\triangleleft} -fixpoint.

For (ii), φ is surjective. Let B be a c_{\triangleleft} -fixpoint and $b = \bigvee B$. We claim $B = \varphi(b)$. For $B \subseteq \varphi(b)$, if $b_0 \in B$, then $b_0 \leq b$, so $b_0 \in \varphi(b)$. For $B \supseteq \varphi(b)$, suppose $a \in \varphi(B)$, so $a \leq b$. Then by the second suitability condition, $a \in B$. \square

Our strategy for defining suitable compatibility relations on a join-dense set V of elements of L will be to assume that L comes equipped with an anti-inflationary operation \neg , e.g., as in the case of an ortholattice with orthocomplementation \neg or a p-algebra or Heyting algebra with pseudocomplementation \neg or a lattice to which we have added an anti-inflationary operation \neg as in Fig. 1. We will then use \neg to define a compatibility relation \triangleleft_{\neg} on V .

3.1.2 The first suitability condition

In the case of an ortholattice, our defined compatibility relation will be equivalent to $x \triangleleft y$ if $y \not\leq \neg x$ (Lemma 3.7(i)). To see why this compatibility relation satisfies the first suitability condition, the following concept is useful.

Definition 3.3 Let L be a lattice equipped with a unary operation \neg and V a set of elements of L . Given $a, b \in L$, we say that a *escapes b in V with \neg* if there is some $c \in V$ such that $a \not\leq \neg c$ but $b \leq \neg c$.

Lemma 3.4 *Let L be an ortholattice, V a join-dense set of elements of L , and $a, b \in L$. If $a \not\leq b$, then a escapes b in V with the orthocomplementation \neg .*

Proof. Suppose $a \not\leq b$, so $\neg b \not\leq \neg a$. Since V is join-dense, we have $\neg b = \bigvee C$ for some $C \subseteq V$. Then $\neg b \not\leq \neg a$ implies $c \not\leq \neg a$ for some $c \in C$. From $c \not\leq \neg a$ we have $a \not\leq \neg c$, and from $c \in C$ we have $c \leq \neg b$ and hence $b \leq \neg c$. \square

Using Lemma 3.4, it is easy to see that the relation \triangleleft on ortholattices defined by $x \triangleleft y$ if $y \not\leq \neg x$ satisfies the first suitability condition.

However, in lattices with \neg that are not ortholattices, we can have $a \not\leq b$ while a cannot escape b , as shown with two examples in Fig. 4.




Fig. 4. Lattices with \neg (dashed arrows) in which $a \not\leq b$ but a cannot escape b .

In order to deal with lattices in which $a \not\leq b$ does not imply that a can escape b , we introduce the key definition of this section.

Definition 3.5 Let L be a lattice, \neg an anti-inflationary operation on L , and V a set of nonzero elements of L .

- (i) Define \triangleleft_V on V by: $x \triangleleft_V y$ if both $y \not\leq \neg x$ and for all $z \in L$, if $y \leq z$ but $x \not\leq z$, then x escapes z in V with \neg .
- (ii) Given $a \in V$ and $b \in L$, we say that a *compatibly escapes* b in V with \neg if there is some $c \in V$ such that $c \triangleleft_V a$ and $b \leq \neg c$.
- (iii) L has *compatible escape with \neg in V* if for all $a \in V$ and $b \in L$, if a escapes b in V with \neg , then a compatibly escapes b in V with \neg .

The following is easy to check.

Lemma 3.6 Under the assumptions of Definition 3.5, (i) \triangleleft_V is reflexive, (ii) $x \leq y$ implies $x \triangleleft_V y$, and (iii) $x \triangleleft_V y \leq y'$ implies $x \triangleleft_V y'$.

It will turn out (Proposition 3.8) that L having compatible escape ensures that \triangleleft_V satisfies the first suitability condition. First, we show that ortholattices and Heyting algebras are alike in having compatible escape, and \triangleleft_V reduces to familiar relations in ortholattices and complete perfect Heyting algebras.

Proposition 3.7 Let L be a lattice and V a join-dense set of nonzero elements.

- (i) If L is an ortholattice, then $x \triangleleft_V y$ iff $y \not\leq \neg x$ for $x, y \in V$; hence L has compatible escape in V with \neg .
- (ii) If L is a p -algebra, then L has compatible escape in V with \neg .
- (iii) If L is a complete Heyting algebra and $V = CJI(L)$, then $x \triangleleft_V y$ iff $x \leq y$ for $x, y \in V$.

Proof. For part (i), by definition, $x \triangleleft_V y$ implies $y \not\leq \neg x$. Conversely, suppose $y \not\leq \neg x$. To show $x \triangleleft_V y$, suppose $y \leq z$ but $x \not\leq z$. From $x \not\leq z$, it follows by Lemma 3.4 that x escapes z in V with \neg . This shows $x \triangleleft_V y$.

For (ii), suppose a escapes b in V , so for some $c \in V$, $a \not\leq \neg c$ but $b \leq \neg c$. Since $a \not\leq \neg c$ and \neg is pseudocomplementation, we have $a \wedge c \neq 0$. From $b \leq \neg c$, we have $b \leq \neg(a \wedge c)$. Since V is join-dense in L , there is a $d \in V$ such that $d \leq a \wedge c$, so $\neg(a \wedge c) \leq \neg d$. Then since $b \leq \neg(a \wedge c)$, we have $b \leq \neg d$. Moreover, by Lemma 3.6(ii), $d \leq a$ implies $d \triangleleft_V a$. Hence we have found a $d \triangleleft_V a$ such that $b \leq \neg d$. Thus, a compatibly escapes b in V with \neg .

For (iii), given Lemma 3.6(ii), we need only show that $x \triangleleft_{\nabla} y$ implies $x \leq y$. Suppose $x \triangleleft_{\nabla} y$ but $x \not\leq y$. Let $w = \bigvee \{z \in L \mid y \leq z, x \not\leq z\}$. Since $x \in CJI(L)$ and L is a complete Heyting algebra, x is completely join-prime, which implies $x \not\leq w$. Then since $x \triangleleft_{\nabla} y$ and $y \leq w$, x escapes w , so for some $v \in V$, $x \not\leq \neg v$ but $w \leq \neg v$ and thus $y \leq \neg v$. Hence $x \not\leq v$, for otherwise $\neg v \leq \neg x$ and so $y \leq \neg x$, contradicting $x \triangleleft_{\nabla} y$. From $x \not\leq w$ and $x \not\leq v$, we have $x \not\leq w \vee v$, while $y \leq w \vee v$. Since $w \leq \neg v$, we also have $w < w \vee v$. But together $y \leq w \vee v$, $x \not\leq w \vee v$, and $w < w \vee v$ contradict the definition of w . \square

Crucially, there are other lattices with compatible escape besides ortholattices and p-algebras, as in \mathbf{N}_5 equipped with the \neg operation in Fig. 4 (left).

Proposition 3.8 *Let L be a lattice, \neg an anti-inflationary operation on L , and V a set of nonzero elements of L . If L has compatible escape in V with \neg , then \triangleleft_{∇} satisfies the first suitability condition.*

Proof. For $a \in V$ and $b \in L$, suppose $a \not\leq b$.

Case 1: a escapes b . Then since L has compatible escape in V with \neg , a compatibly escapes b using some $c \in V$, so $c \triangleleft_{\nabla} a$ and $b \leq \neg c$. Let $a' = c$, so $a' \triangleleft_{\nabla} a$. Suppose $a'' \triangleright_{\nabla} a'$, so $a'' \not\leq \neg c$. Then since $b \leq \neg c$, we have $a'' \not\leq b$.

Case 2: a does not escape b . Then let $a' = a$, so $a' \triangleleft_{\nabla} a$. Suppose $a'' \triangleright_{\nabla} a'$. If $a'' \leq b$, then since $a \not\leq b$, from $a \triangleleft_{\nabla} a''$ it follows that a escapes b , contradicting the assumption of the case. Hence $a'' \not\leq b$. \square

Corollary 3.9 *Let L be a lattice and V a join-dense set of nonzero elements.*

- (i) *If L is an ortholattice, then \triangleleft_{∇} satisfies the first suitability condition.*
- (ii) *If L is a p-algebra, then \triangleleft_{∇} satisfies the first suitability condition.*

3.1.3 The second suitability condition

We can treat the second suitability condition more quickly.

Proposition 3.10 (i) *If L is a complete ortholattice and V a join-dense set of nonzero elements of L , then \triangleleft_{∇} satisfies the second suitability condition.*

(ii) *If L is a complete Heyting algebra and $V = CJI(L)$ is join-dense in L , then \triangleleft_{∇} satisfies the second suitability condition.*

Proof. Suppose L is an ortholattice, $a \in V$, B is a $c_{\triangleleft_{\nabla}}$ -fixpoint, and $a \leq b = \bigvee B$. Suppose $a' \triangleleft_{\nabla} a$, so $a \not\leq \neg a'$ and hence $b \not\leq \neg a'$. Then for some $b_0 \in B$, we have $b_0 \not\leq \neg a'$. Now let $a'' = b_0$, so $a'' \in B$. Since $a'' \not\leq \neg a'$, by Proposition 3.7(i) we have $a' \triangleleft_{\nabla} a''$. Given that B is a $c_{\triangleleft_{\nabla}}$ -fixpoint, this shows $a \in B$.

Suppose L is a complete Heyting algebra, $a \in V$, B is a $c_{\triangleleft_{\nabla}}$ -fixpoint, and $a \leq \bigvee B$. Since L is a complete Heyting algebra and $V = CJI(L)$, $a \in V$ implies that a is completely join-prime. Hence $a \leq b_0$ for some $b_0 \in B$. Now consider any $a' \triangleleft_{\nabla} a$. Let $a'' = b_0$, so $a'' \in B$. Since $a' \triangleleft_{\nabla} a \leq a''$, by Lemma 3.6(iii) we have $a' \triangleleft_{\nabla} a''$. Given that B is a $c_{\triangleleft_{\nabla}}$ -fixpoint, this shows $a \in B$. \square

3.1.4 Representation theorem

Combining Propositions 3.2-3.10, we have the following representation theorem.

Theorem 3.11 *Let L be a complete lattice and V a join-dense set of nonzero elements of L .*

- (i) *If L is an ortholattice, then L is isomorphic to $\mathfrak{L}(V, \triangleleft_{\neg})$.³*
- (ii) *If L is a Heyting algebra and $V = CJI(L)$, L is isomorphic to $\mathfrak{L}(V, \triangleleft_{\neg})$.*
- (iii) *If \neg is an anti-inflationary operation on L such that L has compatible escape in V with \neg , and \triangleleft_{\neg} satisfies the second suitability condition, then L is isomorphic to $\mathfrak{L}(V, \triangleleft_{\neg})$.*

Part (iii) applies to many non-ortholattice and non-Heyting examples, as in Fig. 1, but its precise scope is an open question.

Question 3.12 *For which complete lattices L is there a set V of nonzero elements and an anti-inflationary \neg on L such that L is isomorphic to $\mathfrak{L}(V, \triangleleft_{\neg})$?*

The Jupyter notebook cited in § 1 verifies that every lattice L up to size 8 is such an L . It also verifies the following for every lattice up to size 16 (cf. Footnote 4).

Conjecture 3.13 *For any nondegenerate finite lattice L , there is a compatibility frame (X, \triangleleft) with $|X| < |L|$ such that L is isomorphic to $\mathfrak{L}(X, \triangleleft)$.*

Finally, Appendix A.1 modifies (iii) to represent not only L but also (L, \neg) .

3.2 Representation of arbitrary complete lattices

In this section, we turn to the representation of arbitrary complete lattices. Instead of representing a lattice L using special elements of L as in § 3.1, here we use a potentially less economical representation in terms of pairs of elements, as in the birelational representations of complete lattices in [2,3,34,41].

Definition 3.14 Let L be a lattice and P a set of pairs of elements of L . Define a binary relation \triangleleft on P by $(a, b) \triangleleft (c, d)$ if $c \not\leq b$. Then we say P is *separating* if for all $a, b \in L$:

- (i) if $a \not\leq b$, then there is a $(c, d) \in P$ with $c \leq a$ and $c \not\leq b$;
- (ii) for all $(c, d) \in P$, if $c \not\leq b$, then there is a $(c', d') \triangleleft (c, d)$ such that for all $(c'', d'') \triangleright (c', d')$, we have $c'' \not\leq b$.

Proposition 3.15 *Let L be a lattice and P a separating set of pairs of elements of L . For $a \in L$, define $\varphi(a) = \{(x, y) \in P \mid x \leq a\}$. Then:*

- (i) *φ is a lattice embedding of L into $\mathfrak{L}(P, \triangleleft)$;*
- (ii) *if L is complete, then φ is an isomorphism from L to $\mathfrak{L}(P, \triangleleft)$.*

Proof. For part (i), condition (ii) of Definition 3.14 implies that $\varphi(b)$ is a c_{\triangleleft} -fixpoint for each $b \in L$. Clearly φ preserves meet. For join, if $(x, y) \in \varphi(a \vee b)$ and $(x', y') \triangleleft (x, y)$, then $a \not\leq y'$ or $b \not\leq y'$, so (i) of Definition 3.14 yields an $(x'', y'') \in \varphi(a) \cup \varphi(b)$ with $(x', y') \triangleleft (x'', y'')$. It also ensures that φ is injective.

For part (ii), we claim φ is surjective. Given a c_{\triangleleft} -fixpoint A , define $a = \bigvee \{a_i \mid \exists b_i : (a_i, b_i) \in A\}$. We claim $A = \varphi(a)$. For $A \subseteq \varphi(a)$, suppose

³ The orthocomplementation of L is represented as $\neg_{\triangleleft_{\neg}}$ by Propositions A.1 and A.2.

$(a_i, b_i) \in A$. Then by definition of a , $a_i \leq a$, so $(a_i, b_i) \in \varphi(a)$. For $A \supseteq \varphi(a)$, suppose $(c, d) \in \varphi(a)$, so $c \leq a$. Since A is a c_{\triangleleft} -fixpoint, to show $(c, d) \in A$, it suffices to show that for every $(c', d') \triangleleft (c, d)$ there is a $(c'', d'') \triangleright (c', d')$ with $(c'', d'') \in A$. Suppose $(c', d') \triangleleft (c, d)$, so $c \not\leq d'$, which with $c \leq a$ implies $a \not\leq d'$. Then for some $(a_i, b_i) \in A$, we have $a_i \not\leq d'$. Setting $(c'', d'') = (a_i, b_i)$, from $a_i \not\leq d'$ we have $(c', d') \triangleleft (c'', d'')$, and $(c'', d'') \in A$, so we are done. \square

Every lattice has a separating set of pairs, and for ortholattices and Heyting algebras we can cut down the sets of pairs, as in Proposition 3.16. Compare part (i) to the representation of a complete lattice by join-dense and meet-dense sets in [39,40] or in the fundamental theorem of concept lattices [18, Thm. 3.9].

Proposition 3.16 *Let L be a lattice, V a join-dense set of elements of L , and Λ a meet-dense set of elements of L . Then:*

- (i) *if P_1 is a subset of $P_0 = \{(a, b) \mid a \in V, b \in \Lambda, a \not\leq b\}$ such that for each $a \in V$ there is a $b \in \Lambda$ with $(a, b) \in P_1$, and for each $b \in \Lambda$ there is an $a \in V$ with $(a, b) \in P_1$, then P_1 is separating;⁴*
- (ii) *if L is an ortholattice, then $P_2 = \{(a, \neg a) \mid a \in V, a \neq 0\}$ is separating;*
- (iii) *if L is Heyting, then $P_3 = \{(a, a \rightarrow b) \mid a \in V, b \in \Lambda, a \not\leq b\}$ is separating.*

Proof. In each case, that condition (i) of Definition 3.14 is satisfied is obvious, so we focus on condition (ii).

For P_1 , suppose $(c, d) \in P_1$ and $c \not\leq b$. Then where $b = \bigwedge \{b_i \in \Lambda \mid i \in I\}$, there is some $b_i \in \Lambda$ such that $c \not\leq b_i$, but $b \leq b_i$, and some $a \in V$ with $(a, b_i) \in P_1$. Let $(c', d') = (a, b_i)$, so $(c', d') \triangleleft (c, d)$. Now consider any $(c'', d'') \in P_1$ with $(c', d') \triangleleft (c'', d'')$. Then $c'' \not\leq d' = b_i$, so $c'' \not\leq b$.

For the ortholattice case with P_2 , suppose $(c, d) = (c, \neg c) \in P_2$ and $c \not\leq b$. Hence $\neg b \not\leq \neg c$. Then where $\neg b = \bigvee \{x_i \in V \mid i \in I\}$, for some $i \in I$ we have $x_i \not\leq \neg c$, but $x_i \leq \neg b$, so $c \not\leq \neg x_i$ and $b \leq \neg x_i$. Let $(c', d') = (x_i, \neg x_i)$. Since $c \not\leq \neg x_i$, we have $(c', d') \triangleleft (c, d)$. Now consider any $(c'', d'') \in P_2$ with $(c', d') \triangleleft (c'', d'')$. Then $c'' \not\leq d' = \neg x_i$, which with $b \leq \neg x_i$ implies $c'' \not\leq b$.

For the Heyting case with P_3 , suppose $(c, d) = (c, c \rightarrow e) \in P_3$ and $c \not\leq b$. Then where $b = \bigwedge \{b_i \in \Lambda \mid i \in I\}$, there is some $b_i \in \Lambda$ such that $c \not\leq b_i$, but $b \leq b_i$. From $c \not\leq b_i$, we have $c \not\leq c \rightarrow b_i$. Then where $(c', d') = (c, c \rightarrow b_i)$, we have $(c', d') \in P_3$ and $(c', d') \triangleleft (c, d)$. Now consider any $(c'', d'') \in P_3$ such that $(c', d') \triangleleft (c'', d'')$, so $c'' \not\leq d' = c \rightarrow b_i$. Then $c'' \wedge c \not\leq b_i$, which with $b \leq b_i$ implies $c'' \wedge c \not\leq b$ and hence $c'' \not\leq b$. \square

By Propositions 3.15 and 3.16, L embeds into $\mathfrak{L}(P_1, \triangleleft)$. As the image of the embedding is join-dense and meet-dense in $\mathfrak{L}(P_1, \triangleleft)$, it follows that $\mathfrak{L}(P_1, \triangleleft)$ is (up to isomorphism) the MacNeille completion of L (see [26, Thm. 2.2]), and similarly for $\mathfrak{L}(P_2, \triangleleft)$ and $\mathfrak{L}(P_3, \triangleleft)$ in the ortholattice⁵ and Heyting cases.

⁴ Here P_1 is an edge cover for the bipartite graph (V, Λ, P_0) . If for every finite lattice L , the smallest edge cover for the bipartite graph of its join- and meet-irreducibles has cardinality less than $|L|$ (as we verified for all L with $|L| \leq 16$ using [23]), then Conjecture 3.13 is true.

⁵ It is also easy to check that φ respects the orthocomplementation: $\varphi(\neg a) = \neg_{\triangleleft} \varphi(a)$.

Recall from Propositions 2.16 and 2.21 how symmetric, compossible, and symmetric compossible compatibility frames give rise to ortholattices, Heyting algebras, and Boolean algebras, respectively. We now prove a converse.

Proposition 3.17 *Let L be a lattice, V a join-dense set of elements of L , and Λ a meet-dense set of elements of L . Let P_1 , P_2 , and P_3 be defined from V and Λ as in Proposition 3.16. Then:*

- (i) (P_1, \triangleleft) is a compatibility frame;
- (ii) if L is an ortholattice, then (P_2, \triangleleft) is a symmetric frame;
- (iii) if L is a Heyting algebra, then (P_3, \triangleleft) is a compossible frame;
- (iv) if L is a Boolean algebra, then (P_2, \triangleleft) is a symmetric compossible frame.

Proof. For each part, $(a, b) \in P_i$ implies $a \not\leq b$, so $(a, b) \triangleleft (a, b)$. Hence \triangleleft is reflexive. For part (ii), if $(a, \neg a) \triangleleft (b, \neg b)$, so $b \not\leq \neg a$, then $a \not\leq \neg b$, so $(b, \neg b) \triangleleft (a, \neg a)$. Hence \triangleleft is symmetric.

For (iii), toward showing that (P_3, \triangleleft) is a compossible compatibility frame, suppose $(a, a \rightarrow b) \triangleleft (c, c \rightarrow d)$. We must show there is an $(x, x \rightarrow y) \in P_3$ that refines $(a, a \rightarrow b)$ and pre-refines $(c, c \rightarrow d)$. Since $(a, a \rightarrow b) \triangleleft (c, c \rightarrow d)$, we have $c \not\leq a \rightarrow b$, so $c \wedge a \not\leq b$. Then where $c \wedge a = \bigvee \{x_i \in V \mid i \in I\}$ and $b = \bigwedge \{b_k \in \Lambda \mid k \in K\}$, there are $i \in I$ and $k \in K$ such that $x_i \not\leq b_k$ and hence $x_i \not\leq x_i \rightarrow b_k$. We claim that $(x_i, x_i \rightarrow b_k)$ refines $(a, a \rightarrow b)$ and pre-refines $(c, c \rightarrow d)$. To see that $(x_i, x_i \rightarrow b_k)$ pre-refines $(a, a \rightarrow b)$ and $(c, c \rightarrow d)$, suppose $(w, w \rightarrow v) \triangleleft (x_i, x_i \rightarrow b_k)$, so $x_i \not\leq w \rightarrow v$. Then since $x_i \leq a$, we have $a \not\leq w \rightarrow v$, so $(w, w \rightarrow v) \triangleleft (a, a \rightarrow b)$; and since $x_i \leq c$, we have $c \not\leq w \rightarrow v$, so $(w, w \rightarrow v) \triangleleft (c, c \rightarrow d)$. To see that $(x_i, x_i \rightarrow b_k)$ post-refines $(a, a \rightarrow b)$, suppose $(x_i, x_i \rightarrow b_k) \triangleleft (w, w \rightarrow v)$, so $w \not\leq x_i \rightarrow b_k$. It follows that $w \wedge x_i \not\leq b_k$ and hence $w \wedge x_i \not\leq b$, which with $x_i \leq a$ implies $w \wedge a \not\leq b$, so $w \not\leq a \rightarrow b$ and hence $(a, a \rightarrow b) \triangleleft (w, w \rightarrow v)$.

For (iv), symmetry follows from part (ii). As for compossibility, if $(a, \neg a) \triangleleft (b, \neg b)$, so $b \not\leq \neg a$, then $a \wedge b \neq 0$. Hence there is some nonzero $c \in V$ with $c \leq a \wedge b$, which implies that $(c, \neg c) \in P_2$ refines both $(a, \neg a)$ and $(b, \neg b)$. \square

Combining Propositions 3.15-3.17, we have the following.

Theorem 3.18 *If L is a lattice (resp. ortholattice, Heyting algebra, Boolean algebra), then L embeds into the lattice of c_{\triangleleft} -fixpoints of a compatibility frame (resp. symmetric, compossible, symmetric compossible compatibility frame), and if L is complete, the embedding is an isomorphism.*

Thus, compossible compatibility frames yield a semantics for intermediate logics as general as complete Heyting algebras in the “semantic hierarchy” of [3].

Finally, we prove that we can also represent any complete lattice expanded with a protocomplementation using a compatibility frame.

Theorem 3.19 *For any bounded lattice L equipped with a protocomplementation \neg , the expansion (L, \neg) embeds into the lattice of c_{\triangleleft} -fixpoints of a compatibility frame equipped with \neg_{\triangleleft} , and if L is complete, the embedding is an isomorphism.*

Proof. First, we claim $P = \{(a, b) \mid a, b \in L, a \not\leq b, \neg a \leq b\}$ is separating. For part (i) of Definition 3.14, take $(c, d) = (a, \neg a)$. For (ii), suppose $(c, d) \in P$ and $c \not\leq b$. Let $(c', d') = (1, b)$. Since $b \neq 1$ and $\neg 1 = 0 \leq b$, $(1, b) \in P$, and since $c \not\leq b$, $(c', d') \triangleleft (c, d)$. Now consider any $(c'', d'') \in P$ with $(c', d') \triangleleft (c'', d'')$. Then $c'' \not\leq d' = b$, so (ii) holds. Thus, by Proposition 3.15, L embeds into $\mathfrak{L}(P, \triangleleft)$ via the map φ , which is an isomorphism if L is complete. Also observe that \triangleleft is reflexive on P . It only remains to show $\varphi(\neg a) = \neg_{\triangleleft} \varphi(a)$. Suppose $(x, y) \in \varphi(\neg a)$, so $x \leq \neg a$, and $(x', y') \triangleleft (x, y)$. If $x' \leq a$, then $\neg a \leq \neg x'$, which with $x \leq \neg a$ implies $x \leq \neg x'$, which with $\neg x' \leq y'$ implies $x \leq y'$, contradicting $(x', y') \triangleleft (x, y)$. Thus, $x' \not\leq a$, so $(x', y') \notin \varphi(a)$. Hence $(x, y) \in \neg_{\triangleleft} \varphi(a)$. Conversely, let $(x, y) \in P \setminus \varphi(\neg a)$, so $x \not\leq \neg a$. Since $\neg 0 = 1$, it follows that $a \neq 0$, so $(a, \neg a) \in P$, and $(a, \neg a) \triangleleft (x, y)$, so $(x, y) \notin \neg_{\triangleleft} \varphi(a)$. \square

Note that any bounded lattice can be equipped with the protocomplementation such that $\neg 0 = 1$ and $\neg a = 0$ for $a \neq 0$ (in which case the set P in the proof of Theorem 3.19 coincides with P_0 from Proposition 3.16 where \vee and \wedge are the non-minimum and non-maximum elements of L , respectively), so Theorem 3.19 generalizes the part of Theorem 3.18 concerning bounded lattices.

3.3 Representation of arbitrary lattices

There is another way of representing any lattice L as a sublattice of the lattice of c_{\triangleleft} -fixpoints of a compatibility frame, which is now the canonical extension of L (see [25,14]) rather than its MacNeille completion.⁶ The sublattice can then be characterized in a simple way topologically. This approach uses disjoint filter-ideal pairs and appears already in [16], building on [48,1], though we use a different topology in order to generalize the choice-free Stone duality of [4]. Given a lattice L , define $\text{FI}(L) = (X, \triangleleft)$ as follows: X is the set of all pairs (F, I) such that F is a filter in L , I is an ideal in L , and $F \cap I = \emptyset$; and $(F, I) \triangleleft (F', I')$ iff $I \cap F' = \emptyset$. Given $a \in L$, let $\hat{a} = \{(F, I) \in X \mid a \in F\}$. Let $\mathbf{S}(L)$ be $\text{FI}(L)$ endowed with the topology generated by $\{\hat{a} \mid a \in L\}$.

In Appendix A.2, we prove the following (without choice).

Theorem 3.20 *For any lattice (resp. bounded lattice) L , the map $a \mapsto \hat{a}$ is (i) a lattice (resp. bounded lattice) embedding of L into $\mathfrak{L}(\text{FI}(L))$ and (ii) an isomorphism from L to the sublattice of $\mathfrak{L}(\text{FI}(L))$ consisting of c_{\triangleleft} -fixpoints that are compact open in the space $\mathbf{S}(L)$.*

In Appendix A.2 we also prove the following characterization of spaces equipped with a relation \triangleleft that are isomorphic to $\mathbf{S}(L)$ for some L in a manner analogous to the characterization of UV-spaces in [4]. Let X be a topological space and \triangleleft a reflexive relation on X . Let $\text{COFix}(X, \triangleleft)$ be the set of all compact open sets of X that are also c_{\triangleleft} -fixpoints. Given $U, V \in \text{COFix}(X, \triangleleft)$, $U \vee V = c_{\triangleleft}(U \cup V)$. Given $x \in X$, let

$$\begin{aligned} \mathbf{F}(x) &= \{U \in \text{COFix}(X, \triangleleft) \mid x \in U\} \\ \mathbf{I}(x) &= \{U \in \text{COFix}(X, \triangleleft) \mid \forall y \triangleright x \ y \notin U\}. \end{aligned}$$

⁶ We do not have space to discuss (L, \neg) under this approach, so we save this for the future.

Proposition 3.21 *For any space X and reflexive binary relation \triangleleft on X , there is a lattice L such that (X, \triangleleft) and $\mathbf{S}(L)$ are homeomorphic as spaces and isomorphic as relational frames iff the following conditions hold for all $x, y \in X$: (i) $x = y$ iff $(F(x), I(x)) = (F(y), I(y))$; (ii) $\text{COFix}(X, \triangleleft)$ is closed under \cap and \vee and forms a basis for X ; (iii) each disjoint filter-ideal pair from $\text{COFix}(X, \triangleleft)$ is $(F(x), I(x))$ for some $x \in X$; (iv) $x \triangleleft y$ iff $I(x) \cap F(y) = \emptyset$.*

4 Compatibility and accessibility frames

In this section, we extend the three representations from §§ 3.1, 3.2, and 3.3 to lattices equipped with a modal operation \Box (we defer other modalities not definable from \Box to future work). First, we add an accessibility relation R to compatibility frames and require that the standard modal operation \Box_R sends c_{\triangleleft} -fixpoints to c_{\triangleleft} -fixpoints. These frames are similar to the *graph-based frames* of [10, Definition 2], which have been applied in [9,13,12]. Conradie et al. [10, Theorem 1] use the filter-ideal frame $\mathbf{FI}(L)$ equipped with accessibility relations to prove completeness of the minimal non-distributive modal logic with respect to graph-based frames (compare our Theorem 4.10); in addition, they treat Sahlqvist correspondence theory for graph-based frames.

There are many related approaches to representing lattices with modalities in the literature (see, e.g., [42,24,11,31,32,28,20] and references therein). The approach below using just two binary relations on a single set is of special interest to us as a non-classical generalization of classical “possibility semantics” for modal logic [37,35,33,49,50]. Our motivation for such a non-classical generalization comes from a recent application of the approach below to modal ortholattices for natural language semantics [36].

All proofs in this section are deferred to Appendix A.3.

Definition 4.1 A *necessity lattice* is a pair (L, \Box) where L is a lattice and \Box is a unary operation on L that is multiplicative, i.e., $\Box(a \wedge b) = \Box a \wedge \Box b$ for all $a, b \in L$, and $\Box 1 = 1$ if L contains a maximum element 1. We say \Box is *completely multiplicative* if for any $A \subseteq L$, if $\bigwedge A$ exists in L , then $\Box \bigwedge A = \bigwedge \{\Box a \mid a \in A\}$.

Definition 4.2 A *compatibility and accessibility (CA) frame* is a triple (X, \triangleleft, R) such that (X, \triangleleft) is a compatibility frame and R is a binary relation on X such that for any $A \subseteq X$, if A is a c_{\triangleleft} -fixpoint, then so is

$$\Box_R A = \{x \in X \mid R(x) \subseteq A\},$$

where $R(x) = \{y \in X \mid xRy\}$.

Stronger conditions on the interplay of \triangleleft and R could be imposed (see [10, Def. 2] and Proposition 4.5 below) but Definition 4.2 suffices for the following.

Proposition 4.3 *For any CA frame (X, \triangleleft, R) , the pair $(\mathfrak{L}(X, \triangleleft), \Box_R)$ is a complete necessity lattice with \Box_R completely multiplicative.*

Proof. That $\mathfrak{L}(X, \triangleleft)$ is a complete lattice is Corollary 2.7. Recall that meet is intersection. Then the complete multiplicativity of \Box_R is obvious. \square

Example 4.4 Fig. 5 shows a CA frame (left) where a dotted line from w to v means wRv . Observe that $\Box_R(\{x\}) = \{x\}$, $\Box_R(\{y, z\}) = \{z\}$, and $\Box_R(\{z\}) = \{z\}$. Thus, \Box_R sends c_{\triangleleft} -fixpoints to c_{\triangleleft} -fixpoints. The \Box_R operation on the lattice of c_{\triangleleft} -fixpoints (right) is represented by the double-shafted arrows.




Fig. 5. CA frame and associated necessity lattice.

Under certain assumptions about \triangleleft , the condition that \Box_R sends c_{\triangleleft} -fixpoints to c_{\triangleleft} -fixpoints corresponds to a first-order condition on R and \triangleleft . For example, when \triangleleft is a preorder, in light of Proposition 2.9(ii), the condition that \Box_R sends c_{\triangleleft} -fixpoints to c_{\triangleleft} -fixpoints corresponds to \Box_R sending downsets to downsets and hence to the standard interaction condition for intuitionistic modal frames [7]: if $y \in R(x)$ and $x \triangleleft x'$, then there is a $y' \in R(x')$ with $y \triangleleft y'$. When \triangleleft is symmetric, we get the first-order condition in the following proposition. For useful notation, define $z \triangleleft_R x \Leftrightarrow \exists y : z \triangleleft y \in R(x)$.

Proposition 4.5 *If (X, \triangleleft) is a compatibility frame and R a binary relation on X , then (X, \triangleleft, R) is a CA frame if the following condition holds: for all $x, z \in X$, if $z \triangleleft_R x$, then $\exists x' \triangleleft x \forall x'' \triangleright x' z \triangleleft_R x''$.*

Moreover, if (X, \triangleleft) is a symmetric compatibility frame, then (X, \triangleleft, R) is a CA frame if and only if the stated condition holds.

In fact, the proof of Proposition 4.10 below shows that any necessity lattice can be represented using a CA frame satisfying the condition in Proposition 4.5, so in that sense we can work with such CA frames without loss of generality.

Next we turn to extending the representation theorems of § 3.

Proposition 4.6 *Let L be a complete lattice satisfying the hypotheses of Proposition 3.2, so L is isomorphic to $\mathfrak{L}(V, \triangleleft)$ via $b \mapsto \varphi(b) = \{x \in V \mid x \leq b\}$. Given a completely multiplicative operation \Box on L , define R on V by xRy iff $y \leq \bigwedge \{a \in L \mid x \leq \Box a\}$. Then (V, \triangleleft, R) is a CA frame, and φ is an isomorphism from (L, \Box) to $(\mathfrak{L}(V, \triangleleft), \Box_R)$.*

Example 4.7 Fig. 6 shows a necessity ortholattice (L, \Box) (right) along with the CA frame (left) that comes from the representation of (L, \Box) via join-irreducible elements given by Theorem 3.11 and Proposition 4.6. It is argued in [36] that this necessity ortholattice captures some important logical entailments involving the epistemic modals ‘must’ (formalized as \Box) and ‘might’ (formalized as $\diamond = \neg\Box\neg$) in natural language, including the phenomenon of “epistemic contradiction” whereby sentences of the form “ p , but it might be that $\neg p$ ” are judged contradictory, even though “it might be that $\neg p$ ” does not entail $\neg p$.




Fig. 6. CA frame realizing a necessity ortholattice (with \diamond defined by $\neg \square \neg$).

We can extend our other representation results to the modal setting as well. Given (L, \square) and P a separating set of pairs of elements of L as in § 3.2, define a relation R on P by $(x, x')R(y, y')$ iff xRy as defined in Proposition 4.6.

Proposition 4.8 *If (L, \square) is a complete necessity lattice with \square completely multiplicative and P a separating set of pairs of elements of L , then (P, \triangleleft, R) is a CA frame and (L, \square) is isomorphic to $(\mathfrak{L}(P, \triangleleft), \square_R)$.*

Using Theorem 3.19 we can similarly represent complete necessity lattices equipped with a protocomplementation.

Proposition 4.9 *If (L, \square) is a complete necessity lattice with \square completely multiplicative and \neg is a protocomplementation on L , then there is a CA frame (P, \triangleleft, R) such that (L, \square, \neg) is isomorphic to $(\mathfrak{L}(P, \triangleleft), \square_R, \neg_{\triangleleft})$.*

Finally, define $\text{FI}(L, \square)$ just like $\text{FI}(L)$ in § 3.3 but with the addition of a relation R with $(F, I)R(F', I')$ if for all $a \in L$, $\square a \in F$ implies $a \in F'$.

Proposition 4.10 *For any necessity lattice (L, \square) , $\text{FI}(L, \square)$ is a CA frame, and the map $a \mapsto \hat{a}$ is (i) an embedding of (L, \square) into $(\mathfrak{L}(\text{FI}(L)), \square_R)$ and (ii) an isomorphism from (L, \square) to the subalgebra of $(\mathfrak{L}(\text{FI}(L)), \square_R)$ consisting of c_{\triangleleft} -fixpoints that are compact open in the space $\mathfrak{S}(L)$ (recall § 3.3).*

5 Conclusion

We have investigated three representations of complete lattices by means of compatibility frames, as well as modal analogues thereof. For future work, we hope to make progress on Question 3.12 and Conjecture 3.13, as well as applications of the representations studied here to lattice-based logics. For modal logic in particular, for reasons in [36] we would like to understand the lattice of modal orthologies, for which we hope that CA frames will be useful.

Acknowledgements

I thank Peter Jipsen, Guillaume Massas, and the referees for *Advances in Modal Logic* for helpful feedback.

A Appendix

A.1 Representation of (L, \neg)

In Theorem 3.11(iii), the conclusion is that L is isomorphic to $\mathfrak{L}(V, \triangleleft_{\neg})$, not that (L, \neg) is isomorphic to $(\mathfrak{L}(V, \triangleleft_{\neg}), \neg_{\triangleleft_{\neg}})$. To represent (L, \neg) , we ask for a *third suitability condition* on \triangleleft_{\neg} , namely that for $x \in V$ and $y \in L$:

if $x \not\leq \neg y$, then there is a $y_0 \in V$ such that $y_0 \leq y$ and $y_0 \triangleleft_{\neg} x$.

Proposition A.1 *If L is a complete lattice, V is a join-dense set of nonzero elements, \neg is a protocomplementation on L such that L has compatible escape in V with \neg , and \triangleleft_{\neg} satisfies the second and third suitability conditions, then (L, \neg) is isomorphic to $(\mathfrak{L}(V, \triangleleft_{\neg}), \neg_{\triangleleft_{\neg}})$.*

Proof. We need only add to the proof of Theorem 3.11(iii) that $\varphi(\neg a) = \neg_{\triangleleft_{\neg}} \varphi(a)$. Suppose $x \in \varphi(\neg a)$, so $x \leq \neg a$, and $x' \triangleleft_{\neg} x$, so $x \not\leq \neg x'$. If $x' \leq a$, then $\neg a \leq \neg x'$, which with $x \leq \neg a$ implies $x \leq \neg x'$, contradicting the previous sentence. Thus, $x' \not\leq a$, so $x' \notin \varphi(a)$. Hence $x \in \neg_{\triangleleft_{\neg}} \varphi(a)$. Conversely, suppose $x \in V \setminus \varphi(\neg a)$, so $x \not\leq \neg a$. Then by the third suitability condition, there is an $a_0 \in V$ such that $a_0 \leq a$, so $a_0 \in \varphi(a)$, and $a_0 \triangleleft_{\neg} x$, so $x \notin \neg_{\triangleleft_{\neg}} \varphi(a)$. \square

It remains to be seen how broadly Proposition A.1 applies to lattices expanded with a protocomplementation. In the Jupyter notebook cited in § 1, we show there are such expansions (L, \neg) that cannot be represented by any compatibility frame with $|X| \leq |L|$. Of course, Proposition A.1 applies to all ortholattices.

Proposition A.2 *If (L, \neg) is a complete ortholattice, and V is a join-dense set of nonzero elements, then \triangleleft_{\neg} satisfies the third suitability condition.*

Proof. If $x \not\leq \neg y$, then $x \not\leq \neg \bigvee \{z \in V \mid z \leq y\} = \bigwedge \{\neg z \mid z \in V, z \leq y\}$, so there is $y_0 \in V$ with $y_0 \leq y$ and $x \not\leq \neg y_0$, so $y_0 \triangleleft_{\neg} x$ by Proposition 3.7(i). \square

Recall that using the less economical representation of § 3.2, any complete lattice expanded with a protocomplementation is representable (Theorem 3.19).

A.2 Proofs for § 3.3

Theorem 3.20 *For any lattice (resp. bounded lattice) L , the map $a \mapsto \widehat{a}$ is (i) a lattice (resp. bounded lattice) embedding of L into $\mathfrak{L}(\text{Fl}(L))$ and (ii) an isomorphism from L to the sublattice of $\mathfrak{L}(\text{Fl}(L))$ consisting of c_{\triangleleft} -fixpoints that are compact open in the space $\mathfrak{S}(L)$.*

Proof. First observe that for any $a \in L$, \widehat{a} is a c_{\triangleleft} -fixpoint. It suffices to show that if $(F, I) \notin \widehat{a}$, then there is an $(F', I') \triangleleft (F, I)$ such that for all $(F'', I'') \triangleright (F', I')$, we have $(F'', I'') \notin \widehat{a}$. Suppose $(F, I) \notin \widehat{a}$, so $a \notin F$. Let $F' = F$ and $I' = \downarrow a$. Then $a \notin F$ implies $F' \cap I' = \emptyset$. Thus, $(F', I') \in \text{Fl}(L)$. Now consider any (F'', I'') such that $(F', I') \triangleleft (F'', I'')$, so $I' \cap F'' = \emptyset$. Then since $a \in I'$, we have $a \notin F''$, so $(F'', I'') \notin \widehat{a}$, as desired.

Next, the map $a \mapsto \widehat{a}$ is clearly injective: if $a \not\leq b$, then $\uparrow a \cap \downarrow b = \emptyset$, so $(\uparrow a, \downarrow b) \in \text{Fl}(L)$, $(\uparrow a, \downarrow b) \in \widehat{a}$, and $(\uparrow a, \downarrow b) \notin \widehat{b}$. If L has bounds, then $\widehat{1} = X$ and $\widehat{0} = \emptyset$. The map also preserves \wedge : $\widehat{a \wedge b} = \{(F, I) \in X \mid a \wedge b \in F\} = \{(F, I) \in X \mid a, b \in F\} = \{(F, I) \in X \mid a \in F\} \cap \{(F, I) \in X \mid b \in F\} = \widehat{a} \cap \widehat{b} = \widehat{a \wedge b}$.

To complete part (i), we show $\widehat{a \vee b} \subseteq \widehat{a} \vee \widehat{b}$, as the converse inclusion follows from meet preservation. Recall from Proposition 2.3 that $\widehat{a} \vee \widehat{b} = c_{\triangleleft}(\widehat{a} \cup \widehat{b})$. Suppose $(F, I) \in \widehat{a \vee b}$, so $a \vee b \in F$. Consider any $(F', I') \triangleleft (F, I)$, so $I' \cap F = \emptyset$ and hence $a \vee b \notin I'$. Then since I' is an ideal, $a \notin I'$ or $b \notin I'$. Without loss of generality, suppose $a \notin I'$. Then setting $F'' = \uparrow a$ and $I'' = \downarrow c$ for any $c \in I'$, we have $(F'', I'') \in \text{FI}(L)$ and $I' \cap F'' = \emptyset$, so $(F', I') \triangleleft (F'', I'')$, and $(F'', I'') \in \widehat{a}$. Thus, we have shown that for any $(F', I') \triangleleft (F, I)$ there is an $(F'', I'') \triangleright (F', I')$ with $(F'', I'') \in \widehat{a} \cup \widehat{b}$. Hence $(F, I) \in \widehat{a} \vee \widehat{b}$.

For part (ii), we first show that \widehat{a} is compact open. Since \widehat{b} 's form a basis, we need only show that if $\widehat{a} \subseteq \bigcup \{\widehat{b}_i \mid i \in I\}$, then there is a finite subcover. Suppose $a \not\leq b_i$ for some $i \in I$. Then since $(\uparrow a, \downarrow b_i) \in \widehat{a}$, we have $(\uparrow a, \downarrow b_i) \in \widehat{b}_j$ for some $j \in I$, which implies $a \leq b_j$. Thus, $a \leq b_k$ for some $k \in I$, so $\widehat{a} \subseteq \widehat{b}_k$.

Finally, we show that the map $a \mapsto \widehat{a}$ is onto the set of compact open c_{\triangleleft} -fixpoints. Suppose U is compact open, so $U = \widehat{a}_1 \cup \dots \cup \widehat{a}_n$ for some $a_1, \dots, a_n \in L$. Further suppose U is a c_{\triangleleft} -fixpoint, so $c_{\triangleleft}(U) = U$. Where $d = a_1 \vee \dots \vee a_n$, an obvious induction using part (i) and the fact that $c_{\triangleleft}(c_{\triangleleft}(A) \cup B) = c_{\triangleleft}(A \cup B)$ for any $A, B \subseteq X$ yields $\widehat{d} = c_{\triangleleft}(\widehat{a}_1 \cup \dots \cup \widehat{a}_n)$, so $\widehat{d} = c_{\triangleleft}(U) = U$. \square

Proposition 3.21 *For any space X and reflexive binary relation \triangleleft on X , there is a lattice L such that (X, \triangleleft) and $\mathbf{S}(L)$ are homeomorphic as spaces and isomorphic as relational frames iff the following conditions hold for all $x, y \in X$: (i) $x = y$ iff $(\mathbf{F}(x), \mathbf{l}(x)) = (\mathbf{F}(y), \mathbf{l}(y))$; (ii) $\text{COFix}(X, \triangleleft)$ is closed under \cap and \vee and forms a basis for X ; (iii) each disjoint filter-ideal pair from $\text{COFix}(X, \triangleleft)$ is $(\mathbf{F}(x), \mathbf{l}(x))$ for some $x \in X$; (iv) $x \triangleleft y$ iff $\mathbf{l}(x) \cap \mathbf{F}(y) = \emptyset$.*

Proof. Suppose there is such an L . It suffices to show $\mathbf{S}(L)$ satisfies (i)–(iv) in place of (X, \triangleleft) . That (ii) holds for $\text{COFix}(\mathbf{S}(L))$ and $\mathbf{S}(L)$ follows from the proof of Theorem 3.20. Let φ be the isomorphism $a \mapsto \widehat{a}$ from L to $\text{COFix}(\mathbf{S}(L))$ in Theorem 3.20, which induces a bijection $(F, I) \mapsto (\varphi[F], \varphi[I])$ between disjoint filter-ideal pairs of L and of $\text{COFix}(\mathbf{S}(L))$. Parts (i), (iii), and (iv) follow from the fact that for any $x = (F, I) \in \mathbf{S}(L)$, $(\varphi[F], \varphi[I]) = (\mathbf{F}(x), \mathbf{l}(x))$. First, $\widehat{a} \in \varphi[F]$ iff $a \in F$ iff $x \in \widehat{a}$ iff $\widehat{a} \in \mathbf{F}(x)$. Second, $\widehat{a} \in \varphi[I]$ iff $a \in I$, and we claim that $a \in I$ iff $\widehat{a} \in \mathbf{l}(x)$, i.e., for all $(F', I') \triangleright (F, I)$, $(F', I') \notin \widehat{a}$, i.e., $a \notin F'$. If $a \in I$ and $(F, I) \triangleleft (F', I')$, then $a \notin F'$ by definition of \triangleleft . Conversely, if $a \notin I$, let $F' = \uparrow a$ and $I' = I$; then $(F, I) \triangleleft (F', I')$ and $a \in F'$. Now for (i), given $x, y \in \mathbf{S}(L)$ with $x = (F, I)$ and $y = (F', I')$, we have $(F, I) = (F', I')$ iff $(\varphi[F], \varphi[I]) = (\varphi[F'], \varphi[I'])$ iff $(\mathbf{F}(x), \mathbf{l}(x)) = (\mathbf{F}(y), \mathbf{l}(y))$; similarly, for (iv), $(F, I) \triangleleft (F', I')$ iff $I \cap F' = \emptyset$ iff $\varphi[I] \cap \varphi[F'] = \emptyset$ iff $\mathbf{l}(x) \cap \mathbf{F}(y) = \emptyset$. Finally, for (iii), if $(\mathcal{F}, \mathcal{I})$ is a disjoint filter-ideal pair from $\text{COFix}(\mathbf{S}(L))$, then setting $x = (\varphi^{-1}[\mathcal{F}], \varphi^{-1}[\mathcal{I}])$, we have $x \in \mathbf{S}(L)$ and $(\mathcal{F}, \mathcal{I}) = (\mathbf{F}(x), \mathbf{l}(x))$.

Assuming X satisfies the conditions, $\text{COFix}(X, \triangleleft)$ is a lattice, and we define a map ϵ from (X, \triangleleft) to $\mathbf{S}(\text{COFix}(X, \triangleleft))$ by $\epsilon(x) = (\mathbf{F}(x), \mathbf{l}(x))$. The proof that ϵ is a homeomorphism using (i)–(iii) is analogous to the proof of Thm. 5.4(2) in [4]. That ϵ preserves and reflects \triangleleft follows from (iv). \square

A.3 Proofs for § 4

Proposition 4.5 *If (X, \triangleleft) is a compatibility frame and R a binary relation on X , then (X, \triangleleft, R) is a CA frame if the following condition holds: for all $x, z \in X$, if $z \triangleleft_R x$, then $\exists x' \triangleleft x \forall x'' \triangleright x' z \triangleleft_R x''$.*

Moreover, if (X, \triangleleft) is a symmetric compatibility frame, then (X, \triangleleft, R) is a CA frame if and only if the stated condition holds.

Proof. For the first part, we must show that $\Box_R A$ is a c_{\triangleleft} -fixpoint for any c_{\triangleleft} -fixpoint A . That is, we must show that $x \in X \setminus \Box_R A \Rightarrow \exists x' \triangleleft x \forall x'' \triangleright x' x'' \notin \Box_R A$. Suppose $x \notin \Box_R A$, so there is some $y \in R(x)$ with $y \notin A$. Then since A is a c_{\triangleleft} -fixpoint, there is a $z \triangleleft y$ such that (\star) for all $z' \triangleright z$, we have $z' \notin A$. Since $z \triangleleft y \in R(x)$, by the condition we have $\exists x' \triangleleft x \forall x'' \triangleright x' \exists y' : z \triangleleft y' \in R(x'')$. Now $z \triangleleft y'$ implies $y' \notin A$ by (\star) , which with $y' \in R(x'')$ implies $x'' \notin \Box_R A$.

For the second part, assume \triangleleft is symmetric and \Box_R sends c_{\triangleleft} -fixpoints to c_{\triangleleft} -fixpoints. Toward proving the condition, suppose $z \triangleleft y \in R(x)$. Hence $y \notin \neg_{\triangleleft} c_{\triangleleft}(\{z\})$, so $x \notin \Box_{R \neg_{\triangleleft} c_{\triangleleft}}(\{z\})$. Since by assumption $\Box_{R \neg_{\triangleleft} c_{\triangleleft}}(\{z\})$ is a c_{\triangleleft} -fixpoint, it follows that there is an $x' \triangleleft x$ such that for all $x'' \triangleright x'$, we have that $x'' \notin \Box_{R \neg_{\triangleleft} c_{\triangleleft}}(\{z\})$. Thus, there is a $w \in R(x'')$ such that $w \notin \neg_{\triangleleft} c_{\triangleleft}(\{z\})$, so for some $w' \triangleleft w$, we have $w' \in c_{\triangleleft}(\{z\})$, which means that for all $w'' \triangleleft w'$, there is a $w''' \triangleright w''$ such that $w''' \in \{z\}$, i.e., for all $w'' \triangleleft w'$, $w'' \triangleleft z$. Since \triangleleft is symmetric, from $w' \triangleleft w$, we have $w \triangleleft w'$, so setting $w'' = w$, we conclude $w \triangleleft z$, so $z \triangleleft w$. Thus, $z \triangleleft w \in R(x'')$, i.e., $z \triangleleft_R x''$. \square

Proposition 4.6 *Let L be a complete lattice satisfying the hypotheses of Proposition 3.2, so L is isomorphic to $\mathfrak{L}(\mathbb{V}, \triangleleft)$ via $b \mapsto \varphi(b) = \{x \in \mathbb{V} \mid x \leq b\}$. Given a completely multiplicative operation \Box on L , define R on \mathbb{V} by xRy iff $y \leq \bigwedge \{a \in L \mid x \leq \Box a\}$. Then $(\mathbb{V}, \triangleleft, R)$ is a CA frame, and φ is an isomorphism from (L, \Box) to $(\mathfrak{L}(\mathbb{V}, \triangleleft), \Box_R)$.*

Proof. First, recall the key fact provided by complete multiplicativity of \Box : if $x \not\leq \Box b$, then $\bigwedge \{a \in L \mid x \leq \Box a\} \not\leq b$. For if $\bigwedge \{a \in L \mid x \leq \Box a\} \leq b$, then $\Box \bigwedge \{a \in L \mid x \leq \Box a\} \leq \Box b$ and hence $\bigwedge \{\Box a \in L \mid x \leq \Box a\} \leq \Box b$, so $x \leq \Box b$.

Now we show that for all $b \in L$, $\varphi(\Box b) = \Box_R \varphi(b)$. Suppose $x \in \varphi(\Box b)$, so $x \leq \Box b$. Then for all $y \in R(x)$, we have $y \leq b$ and hence $y \in \varphi(b)$. Thus, $x \in \Box_R \varphi(b)$. Now suppose $x \notin \varphi(\Box b)$, so $x \not\leq \Box b$. Hence $\bigwedge \{a \in L \mid x \leq \Box a\} \not\leq b$ as above. Then since \mathbb{V} is join-dense, there is a $y \in \mathbb{V}$ such that $y \leq \bigwedge \{a \in L \mid x \leq \Box a\}$ but $y \not\leq b$. Hence xRy and $y \notin \varphi(b)$, so $x \notin \Box_R \varphi(b)$.

Finally, we prove that $(\mathbb{V}, \triangleleft, R)$ is indeed a CA frame: if B is a c_{\triangleleft} -fixpoint of $(\mathbb{V}, \triangleleft)$, so is $\Box_R B$. By the surjectivity of φ , $B = \varphi(b)$ for some $b \in B$. Then $\Box_R B = \Box_R \varphi(b) = \varphi(\Box b)$, and $\varphi(\Box b)$ is a c_{\triangleleft} -fixpoint, so we are done. \square

Proposition 4.8 *If (L, \Box) is a complete necessity lattice with \Box completely multiplicative and P a separating set of pairs of elements of L , then (P, \triangleleft, R) is a CA frame and (L, \Box) is isomorphic to $(\mathfrak{L}(P, \triangleleft), \Box_R)$.*

Proof. We showed in the proof of Proposition 3.15 that $a \mapsto \varphi(a) = \{(x, y) \in P \mid x \leq a\}$ is an isomorphism from L to $\mathfrak{L}(P, \triangleleft)$. It only remains to show that $\varphi(\Box b) = \Box_R \varphi(b)$. Suppose $(x, y) \in \varphi(\Box b)$, so $x \leq \Box b$. Then $(x, y)R(x', y')$

implies $x' \leq b$ and hence $(x', y') \in \varphi(b)$. Thus, $(x, y) \in \square_R \varphi(b)$. Conversely, suppose $(x, y) \notin (\square b)$, so $x \not\leq \square b$. Since \square is completely multiplicative, it follows that $\bigwedge \{a \in L \mid x \leq \square a\} \not\leq b$. Then since P is separating, there is some $(c, d) \in P$ with $c \leq \bigwedge \{a \in L \mid x \leq \square a\}$ but $c \not\leq b$. Hence $(x, y)R(c, d)$ but $c \notin \varphi(b)$, so $(x, y) \notin \square_R \varphi(b)$. Now the proof that (P, \triangleleft, R) is a CA frame is analogous to the last paragraph of the previous proof. \square

Proposition 4.9 *If (L, \square) is a complete necessity lattice with \square completely multiplicative and \neg is a protocomplementation on L , then there is a CA frame (P, \triangleleft, R) such that (L, \square, \neg) is isomorphic to $(\mathfrak{L}(P, \triangleleft), \square_R, \neg_{\triangleleft})$.*

Proof. Where $P = \{(a, b) \mid a, b \in L, a \neq 0, a \not\leq b, \neg a \leq b\}$, we showed in the proof of Theorem 3.19 that P is separating. Hence by the proof of Proposition 4.8, (L, \square) is isomorphic to $(\mathfrak{L}(P, \triangleleft), \square_R)$ via the map $a \mapsto \varphi(a) = \{(x, y) \in P \mid x \leq a\}$. We also showed in the proof of Theorem 3.19 that φ preserves the protocomplementation. Hence (L, \square, \neg) is isomorphic to $(\mathfrak{L}(P, \triangleleft), \square_R, \neg_{\triangleleft})$. \square

Proposition 4.10 *For any necessity lattice (L, \square) , $\text{FI}(L, \square)$ is a CA frame, and the map $a \mapsto \widehat{a}$ is (i) an embedding of (L, \square) into $(\mathfrak{L}(\text{FI}(L)), \square_R)$ and (ii) an isomorphism from (L, \square) to the subalgebra of $(\mathfrak{L}(\text{FI}(L)), \square_R)$ consisting of c_{\triangleleft} -fixpoints that are compact open in the space $\mathfrak{S}(L)$.*

Proof. In the proof of Proposition 4.5, we showed that for any (X, \triangleleft) and binary relation R on X , if the first-order condition in Proposition 4.5 holds, then (X, \triangleleft, R) is a CA frame. We claim $\text{FI}(L, \square)$ satisfies the condition. Suppose $(G, H) \triangleleft (G', H') \in R((F, I))$, which implies $H \cap \{a \in L \mid \square a \in F\} = \emptyset$. Then where $F' = F$ and I' is the ideal generated by $\{\square a \mid a \in H\}$, we claim $F' \cap I' = \emptyset$, so $(F', I') \triangleleft (F, I)$. For if $b \in F' \cap I'$, then for some $a_1, \dots, a_n \in H$, $b \leq \square a_1 \vee \dots \vee \square a_n$, which implies $b \leq \square(a_1 \vee \dots \vee a_n)$, so $\square(a_1 \vee \dots \vee a_n) \in F$, whence $a_1 \vee \dots \vee a_n \notin H$, contradicting $a_1, \dots, a_n \in H$. Now suppose $(F', I') \triangleleft (F'', I'')$, so $I' \cap F'' = \emptyset$. Let J be the filter generated by $\{b \in L \mid \square b \in F''\}$. We claim $J \cap H = \emptyset$. For if $a \in J$, then $b_1 \wedge \dots \wedge b_n \leq a$ for $\square b_i \in F''$, which implies $\square a \in F''$, so $\square a \notin I'$, whence $a \notin H$. Thus, $(G, H) \triangleleft (J, H) \in R((F'', I''))$, which establishes the desired condition.

Now for (i)–(ii), we need only add to Theorem 3.20 that $\widehat{\square a} = \square_R \widehat{a}$. Suppose $(F, I) \in \widehat{\square a}$, so $\square a \in F$. Then if $(F, I)R(F', I')$, we have $a \in F'$ and hence $(F', I') \in \widehat{a}$. Thus, $(F, I) \in \square_R \widehat{a}$. Conversely, suppose $(F, I) \notin \widehat{\square a}$, so $\square a \notin F$. Let F' be the filter generated by $\{b \in L \mid \square b \in F\}$ and $I' = \downarrow a$. Since $\square a \notin F$ and \square is multiplicative, it follows that $F' \cap I' = \emptyset$. Hence $(F', I') \in \text{FI}(L, \square)$, $(F', I') \notin \widehat{a}$, and by construction of F' , $(F, I)R(F', I')$. Thus $(F, I) \notin \square_R \widehat{a}$. \square

References

- [1] Allwein, G. and C. Hartonas, *Duality for bounded lattices* (1993), Indiana University Logic Group, Preprint Series, IULG-93-25 (1993).
- [2] Allwein, G. and W. MacCaull, *A Kripke semantics for the logic of Gelfand quantales*, *Studia Logica* **68** (2001), pp. 173–228.
- [3] Bezhanishvili, G. and W. H. Holliday, *A semantic hierarchy for intuitionistic logic*, *Indagationes Mathematicae* **30** (2019), pp. 403–469.
- [4] Bezhanishvili, N. and W. H. Holliday, *Choice-free Stone duality*, *The Journal of Symbolic Logic* **85** (2020), pp. 109–148.
- [5] Bimbó, K., *Functorial duality for ortholattices and De Morgan lattices*, *Logica Universalis* **1** (2007), pp. 311–333.
- [6] Birkhoff, G., “Lattice Theory,” American Mathematical Society, New York, 1940.
- [7] Božic, M. and K. Došen, *Models for normal intuitionistic modal logics*, *Studia Logica* **43** (1984), pp. 217–245.
- [8] Burris, S. N. and H. P. Sankappanavar, “A Course in Universal Algebra,” Springer-Verlag, New York, 1981.
- [9] Conradie, W., A. Craig, A. Palmigiano and N. Wijnberg, *Modelling competing theories*, in: *Proceedings of the 11th Conference of the European Society for Fuzzy Logic and Technology (EUSFLAT 2019)*, Atlantis Studies in Uncertainty Modelling **1**, 2019, pp. 721–739.
- [10] Conradie, W., A. Craig, A. Palmigiano and N. M. Wijnberg, *Modelling informational entropy*, in: R. Iemhoff, M. Moortgat and R. Queiroz, editors, *Logic, Language, Information, and Computation. WoLLIC 2019*, Lectures Notes in Computer Science **11541**, 2019, pp. 140–160.
- [11] Conradie, W., S. Frittella, A. Palmigiano, M. Piazzai, A. Tzimoulis and N. M. Wijnberg, *Categories: How I learned to stop worrying and love two sorts*, in: J. Väänänen, A. Hirvonen and R. de Queiroz, editors, *Logic, Language, Information, and Computation. WoLLIC 2016*, Lectures Notes in Computer Science **9803**, 2016, pp. 145–164.
- [12] Conradie, W., A. Palmigiano, C. Robinson, A. Tzimoulis and N. Wijnberg, *Modelling socio-political competition*, *Fuzzy Sets and Systems* **407** (2021), pp. 115–141.
- [13] Conradie, W., A. Palmigiano, C. Robinson and N. Wijnberg, *Non-distributive logics: from semantics to meaning*, in: A. Rezus, editor, *Contemporary Logic and Computing*, Landscapes in Logic, College Publications, 2020 pp. 38–86.
- [14] Craig, A. and M. Haviar, *Reconciliation of approaches to the construction of canonical extensions of bounded lattices*, *Mathematica Slovaca* **64** (2014), pp. 1335–1356.
- [15] Craig, A. P. K., M. J. Gouveia and M. Haviar, *TiRS graphs and TiRS frames: a new setting for duals of canonical extensions*, *Algebra Universalis* **74** (2015), pp. 123–138.
- [16] Craig, A. P. K., M. Haviar and H. A. Priestley, *A fresh perspective on canonical extensions for bounded lattices*, *Applied Categorical Structures* **21** (2013), pp. 725–749.
- [17] Davey, B. A., *On the lattice of subvarieties*, *Houston Journal of Mathematics* **5** (1979), pp. 183–192.
- [18] Davey, B. A. and H. A. Priestley, “Introduction to Lattices and Order,” Cambridge University Press, New York, 2002, 2nd edition.
- [19] Dishkant, H., *Semantics of the minimal logic of quantum mechanics*, *Studia Logica* **30** (1972), pp. 23–30.
- [20] Dmitrieva, A., “Positive modal logic beyond distributivity: duality, preservation and completeness,” Master’s thesis, University of Amsterdam (2021).
- [21] Dragalin, A. G., “Mathematical Intuitionism: Introduction to Proof Theory,” Translations of Mathematical Monographs, American Mathematical Society, Providence, Rhode Island, 1988.
- [22] Esakia, L., “Heyting Algebras. Duality Theory,” Springer, Cham, 2019, english translation by A. Evseev, eds. G. Bezhanishvili and W. H. Holliday.
- [23] Gebhardt, V. and S. Tawn, *Constructing unlabelled lattices*, *Journal of Algebra* **545** (2020), pp. 213–236.

- [24] Gehrke, M., *Generalized Kripke frames*, *Studia Logica* **84** (2006), pp. 241–275.
- [25] Gehrke, M. and J. Harding, *Bounded lattice expansions*, *Journal of Algebra* **238** (2001), pp. 345–371.
- [26] Gehrke, M., J. Harding and Y. Venema, *MacNeille completions and canonical extensions*, *Transactions of the American Mathematical Society* **358** (2005), pp. 573–590.
- [27] Goldblatt, R., “Metamathematics of Modal Logic,” Ph.D. thesis, Victoria University, Wellington (1974).
- [28] Goldblatt, R., *Morphisms and duality for polarities and lattices with operators* (2019), arXiv:1902.09783 [math.LO].
- [29] Goldblatt, R. I., *Semantic analysis of orthologic*, *Journal of Philosophical Logic* **3** (1974), pp. 19–35.
- [30] Goldblatt, R. I., *The Stone space of an ortholattice*, *Bulletin of the London Mathematical Society* **7** (1975), pp. 45–48.
- [31] Hartonas, C., *Discrete duality for lattices with modal operators*, *Journal of Logic and Computation* **29** (2018), pp. 71–89.
- [32] Hartonas, C. and E. Orłowska, *Representation of lattices with modal operators in two-sorted frames*, *Fundamenta Informaticae* **166** (2019), pp. 29–56.
- [33] Holliday, W. H., *Possibility semantics*, in: M. Fitting, editor, *Selected Topics from Contemporary Logics*, *Landscapes in Logic*, College Publications, 2021 pp. 363–476.
- [34] Holliday, W. H., *Three roads to complete lattices: Orders, compatibility, polarity*, *Algebra Universalis* **82** (2021).
- [35] Holliday, W. H., *Possibility frames and forcing for modal logic*, *The Australasian Journal of Logic* (Forthcoming).
URL <https://escholarship.org/uc/item/0tm6b30q>
- [36] Holliday, W. H. and M. Mandelkern, *The orthologic of epistemic modals* (2021), arXiv:2203.02872 [cs.LO].
- [37] Humberstone, I. L., *From worlds to possibilities*, *Journal of Philosophical Logic* **10** (1981), pp. 313–339.
- [38] Jónsson, B. and A. Tarski, *Boolean algebras with operators. Part I.*, *American Journal of Mathematics* **73** (1951), pp. 891–939.
- [39] Markowsky, G., *Some combinatorial aspects of lattice theory*, in: *Proc. Univ. of Houston Lattice Theory Conf.*, 1973, pp. 36–68.
- [40] Markowsky, G., *The factorization and representation of lattices*, *Transactions of the American Mathematical Society* **203** (1975), pp. 185–200.
- [41] Massas, G., *B-frame duality for complete lattices* (2020), manuscript.
- [42] Orłowska, E. and D. Vakarelov, *Lattice-based modal algebras and modal logics*, in: *Logic, methodology and philosophy of science. Proceedings of the 12th international congress*, 2005, pp. 147–170.
- [43] Ploščica, M., *A natural representation of bounded lattices*, *Tatra Mountains Mathematical Publication* **5** (1995), pp. 75–88.
- [44] Sambin, G. and V. Vaccaro, *Topology and duality in modal logic*, *Annals of Pure and Applied Logic* **37** (1988), pp. 249–296.
- [45] Takeuti, G. and W. M. Zaring, “Axiomatic Set Theory,” Springer-Verlag, New York, 1973.
- [46] Tarski, A., *Zur Grundlegung der Bool’schen Algebra. I.*, *Fundamenta Mathematicae* **24** (1935), pp. 177–198.
- [47] Thomason, S. K., *Categories of frames for modal logic*, *The Journal of Symbolic Logic* **40** (1975), pp. 439–442.
- [48] Urquhart, A., *A topological representation theory for lattices*, *Algebra Universalis* **8** (1978), pp. 45–58.
- [49] Yamamoto, K., *Results in modal correspondence theory for possibility semantics*, *Journal of Logic and Computation* **27** (2017), pp. 2411–2430.
- [50] Zhao, Z., *Algorithmic correspondence and canonicity for possibility semantics*, *Journal of Logic and Computation* **31** (2021), pp. 523–572.