# UC Irvine UC Irvine Previously Published Works

## Title

Neutrophil P2X7 receptors mediate NLRP3 inflammasome-dependent IL-1 $\beta$  secretion in response to ATP

Permalink https://escholarship.org/uc/item/4zh671p0

**Journal** Nature Communications, 7(1)

ISSN

2041-1723

## Authors

Karmakar, Mausita Katsnelson, Michael A Dubyak, George R <u>et al.</u>

## **Publication Date**

2016

## DOI

10.1038/ncomms10555

## **Copyright Information**

This work is made available under the terms of a Creative Commons Attribution License, available at <u>https://creativecommons.org/licenses/by/4.0/</u>

Peer reviewed



## ARTICLE

Received 26 Oct 2015 | Accepted 28 Dec 2015 | Published 15 Feb 2016

DOI: 10.1038/ncomms10555

**OPEN** 

# Neutrophil P2X<sub>7</sub> receptors mediate NLRP3 inflammasome-dependent IL-1 $\beta$ secretion in response to ATP

Mausita Karmakar<sup>1,2</sup>, Michael A. Katsnelson<sup>1</sup>, George R. Dubyak<sup>1</sup> & Eric Pearlman<sup>2,3,†</sup>

Although extracellular ATP is abundant at sites of inflammation, its role in activating inflammasome signalling in neutrophils is not well characterized. In the current study, we demonstrate that human and murine neutrophils express functional cell-surface P2X<sub>7</sub>R, which leads to ATP-induced loss of intracellular K<sup>+</sup>, NLRP3 inflammasome activation and IL-1 $\beta$  secretion. ATP-induced P2X<sub>7</sub>R activation caused a sustained increase in intracellular [Ca<sup>2+</sup>], which is indicative of P2X<sub>7</sub>R channel opening. Although there are multiple polymorphic variants of P2X<sub>7</sub>R, we found that neutrophils from multiple donors express P2X<sub>7</sub>R, but with differential efficacies in ATP-induced increase in cytosolic [Ca<sup>2+</sup>]. Neutrophils were also the predominant P2X<sub>7</sub>R-expressing cells during *Streptococcus pneumoniae* corneal infection, and P2X<sub>7</sub>R was required for bacterial clearance. Given the ubiquitous presence of neutrophils and extracellular ATP in multiple inflammatory conditions, ATP-induced P2X<sub>7</sub>R activation and IL-1 $\beta$  secretion by neutrophils likely has a significant, wide ranging clinical impact.

<sup>&</sup>lt;sup>1</sup> Department of Physiology and Biophysics, Case Western Reserve University, Cleveland, Ohio 44106, USA. <sup>2</sup> Department of Ophthalmology and Visual Sciences, Case Western Reserve University, Cleveland, Ohio 44106, USA. <sup>3</sup> Department of Physiology and Biophysics and Department of Ophthalmology, University of California, Irvine, California 92697, USA. <sup>†</sup> Present address: University of California, 843 Health Sciences Road, Irvine, CA, USA. Correspondence and requests for materials should be addressed to E.P. (email: eric.pearlman@uci.edu).

he inflammatory process results in tissue damage that can impair organ function, which can result in clinical manifestations. Almost all mammalian cells including myeloid cells, platelets, leukocytes, epithelial and endothelial cells can release ATP<sup>1</sup>, which can then lead to paracrine or autocrine activation of downstream purinergic signalling and exacerbation of the inflammatory response<sup>2</sup>. ATP activates plasma membrane purinergic receptors of the P2X and P2Y families that are expressed on many cell types, including myeloid and lymphoid cells<sup>3</sup>. Extracellular ATP has been implicated in multiple *in vivo* inflammatory responses, including lung inflammation and fibrosis, systemic inflammation and tissue damage during endotoxemia<sup>4-6</sup>.

The ionotropic P2X<sub>7</sub> receptor (P2X<sub>7</sub>R) is expressed on macrophages and dendritic cells, which is activated by extracellular ATP to induce NLRP3 inflammasome assembly and caspase-1-dependent processing and release of the proinflammatory cytokines interleukin (IL)-1 $\beta$  and IL-18 (ref. 7). In addition to macrophages and dendritic cells, expression of functional P2X<sub>7</sub>R has been described in other human and murine hematopoietic lineage cells, including mast cells, B and T lymphocytes, monocytes, microglial cells and osteoclasts<sup>8</sup>; however, reports of P2X<sub>7</sub>R expression and its function on human and murine neutrophils are conflicting, and therefore the role of P2X<sub>7</sub>R in these cells remains uncertain.

P2X<sub>7</sub>R messenger RNA expression was reported in both human peripheral blood neutrophils and in the HL-60 human promyelocytic leukemia cells that were differentiated into a granulocyte lineage<sup>9,10</sup>. Another report showed the presence of P2X<sub>7</sub>R protein in the cytosol, but not on the cell surface of human neutrophils<sup>11</sup>, whereas other investigators did not detect P2X7R messenger RNA or protein in human neutrophils<sup>11-14</sup>. We are aware of only one report showing ATP-induced activation of inflammasome signalling in murine neutrophils<sup>15</sup>, although those investigators did not identify the ATP receptor. Notably, no systemic studies on P2X<sub>7</sub>R RNA or protein expression in murine neutrophils have been performed. Given that neutrophils are the predominant cell type in multiple causes of acute infection and inflammation, and that ATP is released at sites of inflammation, we addressed these apparent contradictions in the literature by using multiple approaches to characterize P2X7R expression and function in human peripheral blood neutrophils and in murine bone marrow neutrophils.

We now report that extracellular ATP triggers rapid increase in cytosolic Ca<sup>2+</sup> and K<sup>+</sup> efflux and robust IL-1 $\hat{\beta}$  secretion in human and murine neutrophils via a P2X7R-regulated inflammasome platform that requires NLRP3, ASC and caspase-1. We further extended our analysis of human peripheral blood neutrophils by testing multiple healthy donors (n = 11) from different ethnic backgrounds and confirmed cell-surface P2X7R expression as well as P2X<sub>7</sub>R-mediated increases in cytosolic Ca<sup>2+</sup> and IL-1 $\beta$  secretion in response to ATP. Although the magnitude of the responses was variable, there was no association with donor ethnicity. Lastly, using a murine model of corneal infection, we demonstrated the in vivo relevance of these findings by showing that P2X7R-expressing neutrophils are recruited into Streptococcus pneumoniae infected corneas, and that P2X7R expression on neutrophils regulates bacterial survival in the tissues. Given the abundance of extracellular ATP generated during inflammatory processes, P2X7R activation on neutrophils may be a potential target to regulate tissue damage.

#### Results

ATP induces IL-1 $\beta$  secretion by C57BL/6 neutrophils. To examine IL-1 $\beta$  secretion by neutrophil in response to extracellular ATP, bone marrow neutrophils from C57BL/6 mice were primed

with lipopolysaccharide (LPS), and stimulated with ATP in the presence or absence of apyrase, which catalyses rapid hydrolysis of ATP to ADP and AMP.

As shown in Fig. 1a, ATP stimulated IL-1 $\beta$  secretion by LPS-primed neutrophils in a dose-dependent manner. Further, apyrase ablated ATP-induced IL-1 $\beta$  secretion, but had no effect on IL-1 $\beta$  secretion induced by the K<sup>+</sup> ionophore nigericin (Fig. 1b). Similarly, neutrophils that were primed with heat killed S. pneumoniae (hkSP) secreted IL-1B following ATP stimulation, which was ablated in the presence of apyrase (Supplementary Fig. 1B,C). Notably, millimolar concentrations of ATP (effector concentration for half-maximum response (EC<sub>50</sub>) ~ 1.3 mM) were required for robust IL-1 $\beta$  release, which is consistent with the low affinity of P2X<sub>7</sub>R for ATP-a defining hallmark of P2X<sub>7</sub>R pharmacology<sup>16</sup>. In contrast, IL-1β production by LPS-primed neutrophils stimulated with 3-O-benzoylbenzoic acid-derivatized ATP analogue (BzATP) was maximal at 300 µM (Fig. 1c), consistent with the well-characterized higher affinity of P2X7R for BzATP<sup>17</sup>. IL-1 $\beta$  was not produced when neutrophils were stimulated with UTP, which targets the G-protein-coupled P2Y<sub>2</sub> receptors that are highly expressed in murine and human neutrophils (Supplementary Fig. 1D).

To examine whether ATP-induced IL-1 $\beta$  secretion is dependent on the NLRP3/ASC inflammasome, bone marrow neutrophils from C57BL/6, *Nlrp3<sup>-/-</sup>*, *Asc<sup>-/-</sup>* and *Caspase1/11<sup>-/-</sup>* mice were primed with LPS, and stimulated with ATP or nigericin. As shown in Fig. 1d, IL-1 $\beta$  secretion induced by either ATP or nigericin was significantly lower in *Nlrp3<sup>-/-</sup>*, *Asc<sup>-/-</sup>* and Caspase-1/11<sup>-/-</sup> neutrophils compared with C57BL/6 cells, indicating dependence on this pathway. Further, neutrophils stimulated with ATP or nigericin rapidly secreted precursor and mature forms of caspase-1 and IL-1 $\beta$ , which was detected in the extracellular supernatants (Fig. 1e,f).

As caspase-1 mediates pyroptosis in macrophages and dendritic cells<sup>18</sup>, we examined whether ATP-stimulated neutrophils undergo pyroptotic cell death by measuring release of lactate dehydrogenase (LDH) and uptake of propidium iodide (PI). As shown in Fig. 1g, there was no increased LDH release in ATP-stimulated neutrophils compared with unstimulated neutrophils. Similarly, <2% of neutrophils stimulated with LPS plus ATP had intracellular PI, whereas 48.5% of bone-marrow-derived macrophages (BMDMs) were PI + after 45 min (Fig. 1h).

Thus, the release of precursor and mature forms of both caspase-1 and IL-1 $\beta$  by ATP-stimulated neutrophils is mediated by the NLRP3 inflammasome and is regulated by a non-lytic mechanism rather than neutrophil pyroptosis.

**Murine neutrophils express functional P2X**<sub>7</sub>R. Although P2X<sub>7</sub>R mediates ATP-induced IL-1 $\beta$  secretion by macrophages and dendritic cells<sup>16,19</sup>, the role of this receptor on murine neutrophils has yet to be characterized. The high concentration of extracellular ATP required for IL-1 $\beta$  secretion by murine neutrophils (Fig. 1) is consistent with P2X<sub>7</sub>R characteristics. To examine this hypothesis, neutrophils and BMDMs from C57BL/6 and  $P2X_7^{-/-}$  mice were primed with LPS, and analysed for P2X<sub>7</sub>R expression using antibodies that recognize epitopes in either the intracellular C terminus domain (C-term antibody) or the extracellular domain (ecto antibody) of the receptor.

BMDMs from C57BL/6 mice exhibited robust constitutive expression P2X<sub>7</sub>R (Fig. 2a), as reported previously<sup>16,19</sup>. Further, C57BL/6 neutrophils constitutively expressed P2X<sub>7</sub>R as detected by western blot analysis (Fig. 2a). In contrast,  $P2X_7^{-/-}$  neutrophil cell lysate did not react with the C terminus antibody, but showed low-level immunoreactivity with the ectodomain antibody indicating the presence of truncated C-terminal receptor variants described for the Pfizer  $P2X_7^{-/-}$  mice<sup>20</sup>.



**Figure 1 | ATP—induced IL-1** $\beta$  secretion by murine bone marrow neutrophils. (a,b). IL-1 $\beta$  in the cell supernatants of C57BL/6 bone-marrow-derived neutrophils primed 3 h with LPS (500 ng ml<sup>-1</sup>) and stimulated 45 min with ATP (0.5-3.0 mM) or 10  $\mu$ M nigericin. Apyrase (10U ml<sup>-1</sup>) was added 30 min before stimulation with ATP or nigericin, and IL-1 $\beta$  secretion was quantified by ELISA. (c). IL-1 $\beta$  production by LPS-primed C57BL/6 murine neutrophils stimulated 45 min with BzATP. (d). IL-1 $\beta$  in supernatants of LPS-primed bone-marrow-derived neutrophils from C57BL/6, *Nlrp3<sup>-/-</sup>*, *Asc<sup>-/-</sup>* and *Caspase1,11<sup>-/-</sup>* mice after 45 min stimulation with 3 mM ATP or 10  $\mu$ M nigericin. (e,f). Pro- and cleaved forms of caspase-1 and IL-1 $\beta$  in total cell lysates and TCA precipitated supernatant of LPS-primed C57BL/6 neutrophils after 45 min stimulation with 3 mM ATP or 10  $\mu$ M nigericin. (b). Propidium iodide (PI) uptake by bone marrow neutrophils and bone-marrow-derived macrophages following LPS priming and 45 min ATP (3 mM) stimulation. Data were generated by flow cytometry. Data points are mean ± s.d. of at least four replicates per treatment and are representative of three independent experiments.

To discriminate between cell surface and intracellular  $P2X_7R$  protein, LPS-primed C57BL/6 neutrophils were permeabilized before incubating with the  $P2X_7$  ecto-domain antibody, and  $P2X_7R$  expression was examined by flow cytometry and fluorescence microscopy. We found similar levels of total  $P2X_7R$  protein in permeabilized neutrophils that were either unstimulated or LPS-primed (Fig. 2b,c).  $P2X_7R$  was also detected on the cell surface of non-permeabilized cells (Fig. 2d,e), indicating that  $P2X_7R$  is constitutively expressed on the cell surface of murine neutrophils.

To examine the ionotropic function of cell-surface  $P2X_7R$ , cytosolic  $[Ca^{2+}]$  bone marrow neutrophils from C57BL/6 and  $P2X_7^{-/-}$  mice were primed with LPS and loaded with Fluo-4-AM Ca<sup>2+</sup> indicator dye. Stimulation with 3 mM ATP elicited a rapid and sustained rise in intracellular  $[Ca^{2+}]$ 

concentration (Fig. 2f). In contrast, ATP-stimulated  $P2X_7^{-/-}$  neutrophils (Fig. 2f), and C57BL/6 neutrophils that were incubated with the highly specific P2X<sub>7</sub> antagonists AZ10606120 or A438079 (Fig. 2g,h) exhibited only a transient increase in cytosolic [Ca<sup>2+</sup>], which rapidly decayed to baseline. This transient increase likely indicates rapid mobilization of intracellular Ca<sup>2+</sup> stores by G-protein-coupled P2Y<sub>2</sub> receptors, which are also activated by ATP, and is consistent with an earlier study showing that ATP elicits transient Ca<sup>2+</sup> mobilization in  $P2X_7^{-/-}$  macrophages<sup>21</sup>.

Quantification of the areas under the curve (AUC) for each Ca<sup>2+</sup> transient showed significant reduction in the integrated increase in intracellular  $[Ca^{2+}]$  in  $P2X_7^{-/-}$  neutrophils and in C57BL/6 neutrophils treated with P2X<sub>7</sub> antagonists (Fig. 2i-k).

## ARTICLE



**Figure 2** | **P2X**<sub>7</sub> receptor expression and function in murine bone marrow neutrophils. (a). Western blot analysis of P2X<sub>7</sub> receptors on unstimulated and LPS-primed bone marrow neutrophils and macrophages from C57BL/6 and  $P2X_7^{-/-}$  mice. (**b**-e). Flow cytometry and fluorescence microscopy of P2X<sub>7</sub>R ecto domain (using the Hano-3 antibody) in LPS-primed bone marrow neutrophils that were either permeabilized with 0.1% TX-100 to detect total P2X<sub>7</sub>R (**b**,**c**) or were non-permeabilized to detect only cell surface P2X<sub>7</sub>R (**d**,**e**). (**c**,**e**) Cells were also stained with DAPI to detect the nucleus; Scale bar, 5 µm). (**f**-h). Representative profiles of ATP-induced Ca<sup>2+</sup> influx in bone marrow neutrophils from C57BL/6 and  $P2X_7^{-/-}$  mice (**f**), and in C57BL/6 neutrophils after stimulation with ATP (3 mM) in the presence of P2X<sub>7</sub>R antagonists AZ10606120 (10 µM) (**g**), or with different concentrations of A438079 (**h**). (**i**-**k**). Mean ± s.e.m. pf Area under the curve (AUC) of intracellular Ca<sup>2+</sup> from three independent experiments. A *P* value ≤ 0.05 was considered significant using an unpaired Student's *t*-test.

As additional controls, neutrophils from C57BL/6 and  $P2X_7^{-/-}$  mice were stimulated with fMLP or opsonized zymosan, and IL-1 $\beta$  secretion was examined. Although fMLP induced increased cytosolic Ca<sup>2+</sup>, there was no detectable IL-1 $\beta$  secretion (Supplementary Fig. 2A,B). In contrast, opsonized zymosan did induce IL-1 $\beta$  secretion from murine neutrophils; however, this was independent of P2X<sub>7</sub>R (Supplementary Fig. 2C).

Taken together, these data clearly demonstrate that murine bone marrow neutrophils express functional cell-surface  $P2X_7R$  that respond to extracellular ATP by facilitating sustained increase in cytosolic  $[Ca^{2+}]$ , which is indicative of non-selective cation channels. These findings also indicate that in the absence of  $P2X_7R$  expression or in the presence of highly selective  $P2X_7$  antagonists, activation of other ATP receptors, such as  $P2Y_2$ , elicits only a transient increase in cytosolic  $[Ca^{2+}]$ . **P2X<sub>7</sub>R-mediated IL-1β secretion by murine neutrophils.** To examine the role of P2X<sub>7</sub>R in ATP-stimulated IL-1β secretion, LPS-primed C57BL/6 and P2X<sub>7</sub><sup>-/-</sup> neutrophils were stimulated with extracellular ATP or nigericin, and IL-1β secretion was quantified by enzyme-linked immunosorbent assay (ELISA). We found significantly less ATP-stimulated IL-1β in  $P2X_7^{-/-}$  compared with C57BL/6 neutrophils (Fig. 3a), whereas there was no difference in nigericin-induced release of IL-1β (Fig. 3b). Consistent with this observation, ATP-induced IL-1β secretion by C57BL/6 neutrophils was completely inhibited in the presence of P2X<sub>7</sub> antagonists AZ10606120 or A438079 (Fig. 3c,d). AZ10606120 and A438079 also inhibited IL-1β secretion induced by BzATP (Supplementary Fig. 2D).

To ascertain whether the secreted IL-1 $\beta$  as detected in the ELISA is bioactive, we incubated cells supernatants from ATP-stimulated neutrophils with the HEK-Blue-IL-1R reporter



**Figure 3** | **P2X**<sub>7</sub>**R-dependent IL-1** $\beta$  secretion by **ATP-stimulated murine neutrophils.** Total IL-1 $\beta$  secreted by bone marrow neutrophils from C57BL/6 and  $P2X_7^{-/-}$  mice (**a**,**b**), or from C57BL/6 neutrophils incubated with the P2X<sub>7</sub> antagonists AZ10606120 (10  $\mu$ M) (**c**) or A438079 (25  $\mu$ M) (**d**) following LPS priming and stimulation with ATP or nigericin measured by ELISA. (**e-g**). Bioactive IL-1 from the same samples detected using HEK-Blue-IL-1R reporter cells. (**h**). IL-1 $\alpha$  and IL-1 $\beta$  from the C57BL/6 neutrophil supernatants stimulated 45 min with 3 mM ATP and 10  $\mu$ M nigericin were quantified by ELISA. Data points are mean ± s.d. of at least three replicates per treatment and are representative of three independent experiments. Using an ANOVA with Tukey *post hoc* analysis, a *P* value ≤ 0.05 was considered significant.

cell line. As shown in Fig. 3e, bioactive IL-1 was detected in LPS-primed C57BL/6 neutrophils stimulated with ATP or nigericin; however, production of biologically active IL-1 was significantly reduced in ATP-stimulated  $P2X_7^{-/-}$  neutrophils and in C57BL/6 neutrophils incubated with P2X<sub>7</sub>R antagonists AZ10606120 and A438079 (Fig. 3e-g). Further, neither gene knockout nor pharmacological antagonism of P2X<sub>7</sub>R suppressed production of biologically active IL-1 by nigericin-stimulated neutrophils. Given the small amount of IL-1 $\alpha$  production compared with IL-1 $\beta$  (Fig. 3h), the biologically active IL-1 released from the ATP-activated neutrophils appears to be primarily due to IL-1 $\beta$ .

Taken together, these data demonstrate that the  $P2X_7R$  on murine neutrophils mediates production of bioactive IL-1 $\beta$  in response to extracellular ATP.

Functional P2X7R expression on human neutrophils. To examine whether ATP also elicits IL-1 $\beta$  secretion from human neutrophils, peripheral blood neutrophils were isolated from healthy donors, primed with LPS for 3 h, and stimulated with ATP or nigericin. As shown in Fig. 4a, IL-1 $\beta$  produced by ATP-stimulated human neutrophils was completely abrogated in the presence of apyrase, whereas there was no effect of apyrase on nigericin-induced IL-1β. As with murine neutrophils, millimolar concentrations of ATP were required for IL-1ß release, and ATPinduced IL-1ß secretion was significantly inhibited in the presence of the P2X7 antagonists AZ10606120 and AZ11645373 (Fig. 4b,c), indicating that ATP induces IL-1ß production by human neutrophils through the P2X7 receptor. Human peripheral blood neutrophils constitutively express NLRP3 and ASC protein, although NLRP3 can be enhanced following TLR stimulation (Supplementary Fig. 2F).

 $P2X_7$  receptor expression and function in human neutrophils is controversial, with one study reporting intracellular, but not cell-surface  $P2X_7$  receptor expression<sup>11</sup>, and polymorphic  $P2X_7$ receptor variants reported in this gene, which have been linked to quantitative or qualitative differences in  $P2X_7$  receptor function in mice<sup>22–25</sup> and in humans (Refs. 57-60). Given this heterogeneity, we assayed  $P2X_7$  receptor expression and function in human peripheral blood neutrophils from 11 healthy donors from multiple ethnicities. Neutrophils were primed for 3 h with LPS, and  $P2X_7$ protein expression was assayed by western blot and flow cytometry using antibodies against the C terminus and the extracellular domain of the receptor.

As shown for two donors, full length  $P2X_7R$  was detected in unstimulated and LPS-primed neutrophils indicating constitutive  $P2X_7R$  expression (Fig. 4d). Consistent with this observation,  $P2X_7R$  surface expression was detected in non-permeabilized, unstimulated and LPS-stimulated neutrophils using the ecto-domain antibody (Fig. 4e), thereby demonstrating constitutive cell surface expression of  $P2X_7$ receptors on human neutrophils.  $P2X_7$  receptors were also constitutively expressed on neutrophils from different ethnic groups (Table 1).

To characterize ionotropic  $P2X_7R$  function, neutrophils were incubated with LPS for 10 min to establish a stable baseline  $[Ca^{2+}]$ , then stimulated with 4 mM ATP in the presence of the  $P2X_7$  receptor antagonist AZ10606120. As shown in Fig. 4f, ATP induced a rapid increase in cytosolic  $[Ca^{2+}]$ , followed by a gradual decay over 30 min; however, the response was markedly attenuated in the presence of AZ10606120, indicating  $P2X_7R$  activation. Increases in cytosolic  $[Ca^{2+}]$  were detected within 5 seconds of adding ATP (Supplementary Fig. 2E).



**Figure 4 | P2X<sub>7</sub>R expression and function on human peripheral blood neutrophils. (a-c)**. IL-1 $\beta$  measured by ELISA in the supernatant of human peripheral blood neutrophils primed with LPS (500 ng ml<sup>-1</sup>) for 3 h and stimulated with 4 mM ATP or 10  $\mu$ M nigericin in the presence or absence of 10U ml<sup>-1</sup> apyrase (**a**), or with increasing concentrations of ATP in the presence of P2X<sub>7</sub> antagonists AZ10606120 (10  $\mu$ M) or AZ11645373 (5  $\mu$ M) (**b,c**). (**d**) Western blot analysis of P2X<sub>7</sub>R of unstimulated and LPS-primed human neutrophils from two donors that had been LPS primed for 3 h.  $\beta$  actin was used as a loading control. (**e**). Flow cytometry detection of P2X<sub>7</sub>R cell-surface expression on human neutrophils using antibody against the extracellular domain of the receptor. (**f**). Cytosolic [Ca<sup>2+</sup>] in LPS-primed human neutrophils from all donors (*n* = 11) in the presence of AZ10606120 (10  $\mu$ M). (**g**). Area under the curve (AUC) for ATP induced cytosolic Ca<sup>2+</sup> in neutrophils from all donors (*n* = 11) in the presence of P2X<sub>7</sub>R inhibitor (*P* values were obtained by paired *t*-tests, and a *P* value ≤0.05 was considered significant). Individual donors were for (**a**), Donor 8, (**b**), Donor 9, (**c**), Donor 7, (**d**), Donors 1 and 12, (**e**)—Donor 1, (**f**), Donor 2. Histograms or data points are mean ± s.d. of at least five replicates per group and data shown are representative of three independent experiments with three different donor neutrophils (for **a-c**).

Although neutrophils from all the donors tested expressed cellsurface  $P2X_7$  receptors as quantified by flow cytometry, we noted donor-specific differences in  $P2X_7$  receptor expression and function as indicated by both the magnitude of the ATP-induced cytosolic Ca<sup>2+</sup> increases and the relative sensitivity of these responses to attenuation by the selective  $P2X_7$  antagonists AZ10606120 and AZ11645373 (Table 1).

ATP-induced cytosolic  $[Ca^{2+}]$  transients for each individual are shown in Supplementary Fig. 3, and combined data showing the AUC for ATP-induced cytosolic  $[Ca^{2+}] \pm P2X_7R$ inhibitor over a 30-min period are shown in Fig. 4g. Neutrophils from 10 of the 11 individuals responded to ATP by increased cytosilic  $[Ca^{2+}]$ ; however, the AUC values were highly variable, ranging from 26 to 56%. Further, whereas the two P2X<sub>7</sub>R antagonists completely suppressed ATP-induced IL-1 $\beta$  release from human neutrophils, these inhibitors only partially attenuated the ATP-stimulated increases in cytosolic  $[Ca^{2+}]$ . Although we did not examine this directly, partial inhibition ATP-induced increases in cytosolic  $[Ca^{2+}]$  is consistent with the well-defined role of G-protein-coupled P2Y<sub>2</sub> receptors in mediating ATPtriggered mobilization of IP<sub>3</sub>-sensitive Ca<sup>2+</sup> stores in these leukocytes.

**P2X<sub>7</sub>R-induced K<sup>+</sup> efflux.** As non-selective cation channels, ATP-activated P2X<sub>7</sub>R mediates K<sup>+</sup> efflux, and we and others reported that NLRP3 inflammasome activation in macrophages and dendritic cells is dependent on rapid K<sup>+</sup> efflux, and can be dissociated from changes in cytosolic  $[Ca^{2+}]^{26,27}$ . As we also recently demonstrated that NLRP3 inflammasome and IL-1 $\beta$  secretion by pneumolysin-stimulated neutrophils requires a decrease in intracellular K<sup>+</sup> concentration<sup>28</sup>, we examined whether there is a requirement for K<sup>+</sup> efflux in ATP-induced IL-1 $\beta$  secretion from neutrophils. LPS-primed C57BL/6 neutrophils were stimulated with 3 mM ATP, and the intracellular K<sup>+</sup> content was quantified by atomic absorbance spectrometry. Figure 5a shows a ~30% decrease in intracellular K<sup>+</sup> within 5 min of ATP stimulation, which was abrogated in presence of the P2X<sub>7</sub>R antagonist AZ10606120.

To determine whether this decrease in intracellular K<sup>+</sup> was sufficient to drive inflammasome activation and IL-1 $\beta$  secretion, primed neutrophils were incubated in medium with elevated extracellular [K<sup>+</sup>] saline (130 mM KCl), which eliminates the outwardly directed K<sup>+</sup> gradient that normally favors K<sup>+</sup> efflux via gated cation channels. As shown in Fig. 5b, elimination of the K<sup>+</sup> gradient in 130 mM KCl resulted in significantly lower ATPinduced IL-1 $\beta$  secretion compared with 5 mM isotonic KCl, indicating that P2X<sub>7</sub>R-driven IL-1 $\beta$  release from neutrophils is dependent on K<sup>+</sup> efflux.

To ascertain whether the rapid decrease in intracellular K<sup>+</sup> during the initial several minutes of ATP stimulation was sufficient to trigger NLRP3 inflammasome assembly and IL-1 $\beta$  secretion, LPS-primed neutrophils were stimulated with 3 mM ATP in the presence of apyrase, and IL-1 $\beta$  secretion was quantified after 45 min. ATP—induced IL-1 $\beta$  secretion was completely abrogated when cells were incubated with apyrase before stimulation with ATP (Fig. 5c). However, addition of apyrase 10 min after stimulation with ATP did not inhibit IL-1 $\beta$ secretion, indicating that only a transient activation of P2X<sub>7</sub>R

#### Table 1 | P2X<sub>7</sub>R expression and function in peripheral blood neutrophils.

Donor	Ethnicity	Surface P2X7R (% positive cells [MFI])	Ca <sup>2+</sup> influx (AUC)		% Inhibition (AUC) of Ca <sup>2+</sup> influx in presence of P2X7 inhibitor (AZ106 or AZ116)
#1	Caucasian	Unstim—79.1% (8961)	LPS	2,591	44.2%
		LPS—68.2% (8417)	LPS + ATP	7,554	
			LPS + ATP + AZ106	4,220	
#2	Caucasian	Unstim—87.5% (8637)	LPS	3,998	43.6%
		LPS—83.8% (11,355)	LPS + ATP	13,331	
			LPS + ATP + AZ106	7,527	
#3	Caucasian	Unstim—53.5% (14.580.60)	LPS	2,323	48.6%
		LPS—52.32% (15,845.83)	LPS + ATP	10,903	
			LPS + ATP + AZ106	5,606	
#4	Caucasian	Unstim—91.9% (43,739.63)	LPS	838	28.4%
		LPS-86.9% (35,508.18)	LPS + ATP	6,640	
			LPS + ATP + AZ116	4,827	
#5	African American-Caucasian	Unstim—99.6% (30,889.34)	LPS	858	45.5%
		LPS-99.4% (24,296.22	LPS + ATP	4,460	
			LPS + ATP + AZ116	2,432	
#6	Asian	Unstim—85.6% (12,772.51)	LPS	2,307	55.9%
		LPS-86.7% (14,020.63)	LPS + ATP	7,558	
			LPS + ATP + AZ106	3,334	
#7	South east Asian	Unstim—99.2% (14,653.43)	LPS	888	33.4%
		LPS-92.9% (10,036.8)	LPS + ATP	2,782	
			LPS + ATP + AZ116	1,854	
#8	Hispanic	Unstim—72.3% (16,183.67)	LPS	4,079	26.2%
		LPS-62.9% (14,452.63)	LPS + ATP	6,120	
			LPS + ATP + AZ116	4,522	
#9	Hispanic	Unstim—95.5% (15,629.02)	LPS	1,980	33.7%
		LPS—94.3% (16,336)	LPS + ATP	4,143	
			LPS + ATP + AZ116	2,749	
#10	Caucasian	Unstim—98.1% (70,563.26)	LPS	1,353	0%
		LPS—95.9% (45,937.79)	LPS + ATP	5,325	
			LPS + ATP + AZ116	5,543	
#11	African	Unstim—93.7% (15,332.15)	LPS	1,017	42.6%
	American	LPS—75.9% (14,587.90)	LPS + ATP	2,128	
			LPS + ATP + AZ116	1,216	

AUC, area under the curve; FSC, forward scatter; SSC, side scatter.

Granulocytes from healthy donors were gated based on SSC (side scatter) and FSC (forward scatter), and 30,000 cells (unstimulated or primed with LPS for 3 h) were analyzed for surface P2X<sub>7</sub> expression (extracellular antibody) within the gated population.

ATP-induced cytosolic  $Ca^{2+}$  increases and their relative sensitivity to attenuation by the selective P2X<sub>7</sub> antagonists were also demonstrated.  $Ca^{2+}$  transients for each donor were quantified by measuring AUC. Attenuation of cytosolic  $[Ca^{2+}]$  in presence of P2X<sub>7</sub> antagonists is represented as % inhibition compared to LPS + ATP-treated cells. Individual curves for cytosolic  $[Ca^{2+}]$  are shown in Supplementary Fig. 3.



**Figure 5** | **Role of K**<sup>+</sup> **efflux in P2X<sub>7</sub>R-mediated IL-1** $\beta$  **secretion by murine neutrophils. (a)**. Intracellular K<sup>+</sup> content in LPS-primed C57BL/6 neutrophils stimulated with ATP (3 mM) alone and in the presence of AZ10606120 (10  $\mu$ M) measured by atomic absorbance spectroscopy. (b). IL-1 $\beta$  production by LPS primed and ATP (3 mM) or nigericin (10  $\mu$ M) stimulated neutrophils in the presence of 130 mM KCl or 5 mM KCl as quantified by ELISA. (c). IL-1 $\beta$  from LPS-primed C57BL/6 neutrophils in the presence of apyrase (10U ml<sup>-1</sup>) added either 30 min before or 10 min after stimulation with ATP or nigericin. Data points are mean ± s.d. of at least four replicates per treatment and are representative of three independent experiments. A *P* value ≤0.05 was considered significant using ANOVA with Tukey *post hoc* analysis.

channels is required to initiate the cascade of NLRP3 inflammasome assembly that results in proteolytic maturation and IL-1 $\beta$ secretion by neutrophils.

**P2X<sub>7</sub>R expression in** *S. pneumoniae* **corneal infection**. Bacterial and fungal infections of the cornea are leading causes of blindness worldwide, and murine models reveal neutrophils to be prominent infiltrating cells, especially early after infection<sup>28–32</sup>. To determine whether neutrophils express P2X<sub>7</sub>R *in vivo*, corneas of C57BL/6 mice were infected with *S. pneumoniae*, and after 24 h, corneas were digested with collagenase, and Ly6G<sup>+</sup> neutrophils and P2X<sub>7</sub>R<sup>+</sup> cells were detected by flow cytometry. Figure 6a shows a distinct population of P2X<sub>7</sub>R-expressing neutrophils in the cornea, which comprise >40% of the total corneal cells. When gated on the total corneal P2X<sub>7</sub><sup>+</sup> population, >90% were Ly6G<sup>+</sup> (Fig. 6b), indicating that neutrophils are the predominant P2X<sub>7</sub>R-expressing cells in the *S. pneumoniae* infected corneas.

As a second approach, neutrophils were depleted following intraperitoneal injection with NIMP-R14 antibody (anti-Ly6G) or control IgG. Corneas were then infected with *S. pneumoniae*, and P2X<sub>7</sub>R-expressing cells were detected by flow cytometry. As shown

in Fig. 6c, IgG-treated C57BL/6 mice showed a distinct clear peak of infiltrating neutrophils (~21%) and P2X<sub>7</sub>R-expressing cells (~17%) in infected corneas. In contrast, NIMP-R14-treated mice had no infiltrating corneal neutrophils indicative of systemic depletion of neutrophils. Moreover, with depletion of neutrophils, corneal P2X<sub>7</sub>R-expressing cells were also significantly reduced (Fig. 6d). Figure 6e shows significantly less P2X<sub>7</sub>R-expressing cells in infected corneas from NIMP-R14-treated compared with control mice. Further, neither P2X<sub>7</sub>R nor mature IL-1 $\beta$  (p17) were detected in NIMP-R14-treated mice (Fig. 6f).

Taken together, these data indicate that neutrophils are the primary source of  $P2X_7R$  and mature IL-1 $\beta$  *in vivo* during *S. pneumoniae* corneal infection.

**Neutrophil P2X<sub>7</sub>R-mediated bacterial killing** *in vivo*. To assess the role of P2X<sub>7</sub>R in *S. pneumoniae* infection, corneas of C57BL/6 and  $P2X_7^{-/-}$  mice were infected, and numbers of infiltrating corneal neutrophils and macrophages and bacterial colony forming units (CFU) in the eyes were quantified 24 h post infection. As shown in Fig. 7a, there was no significant difference between C57BL/6 and  $P2X_7^{-/-}$  mice in the total number of infiltrating neutrophils, and there were no differences in the



**Figure 6 | Neutrophil P2X<sub>7</sub> receptor expression in** *S. pneumoniae* **corneal infection.** P2X<sub>7</sub> receptor expression on total cells (**a**), and on Ly6G + (1A8) neutrophils (**b**) 24 h after *S. pneumoniae* corneal infection. Cells were obtained following collagenase digestion, incubated with 1A8 and antibodies against the extracellular domain of the P2X<sub>7</sub> receptor and analysed by flow cytometry. (**c**-**f**) Neutrophil depletion following intraperitoneal injection of NIMP-R14 (Ly6G) or control rat IgG. (**c**,**d**) 1A8 and P2X<sub>7</sub>-expressing cells in the corneas 24 h post *S. pneumoniae* infection. One representative cornea is shown from each group. (**e**) Percentage and standard error of P2X<sub>7</sub>-receptor-expressing cells in IgG and NIMP-R14-treated mice (n = 8 corneas). (**f**) P2X<sub>7</sub> receptor, and pro- and mature IL-1 $\beta$  in infected corneas. Data are representative of two independent experiments with at least 5 mice per group. A *P* value ≤0.05 was considered significant using a Student's *t*-test.

relatively few macrophages in the cornea at this time point. However, there were significantly more CPU in infected corneas from  $P2X_7^{-/-}$  compared with C57BL/6 mice (Fig. 7b), indicating impaired bacterial clearance in the absence of P2X\_7R. The cleaved forms of caspase-1 and IL-1 $\beta$  were also not detected in corneas from infected  $P2X_7^{-/-}$  mice (Fig. 7c), thereby demonstrating that inflammasome activation *in vivo* is driven by P2X\_7R.

To examine whether P2X7R expression on neutrophils is required for host defense against S. pneumoniae, we used a neutrophil adoptive transfer experiment as described previously<sup>31,32</sup>.  $Cd18^{-/-}$  mice exhibit impaired neutrophil extravasation as they are unable to bind to ICAM-1 on vascular endothelial cells<sup>33</sup>. Adoptive transfer of wild type or  $P2X_7^{-/-}$ bone-marrow-derived neutrophils into the  $Cd18^{-/-}$  mice will therefore ascertain the specific role of neutrophil P2X7R expression during S. pneumoniae corneal infection. Cd18-/mice were infected with  $1 \times 10^5$  S. pneumoniae and 3 h post infection,  $8 \times 10^6$  C57BL/6 and  $P2X_7^{-/-}$  bone-marrow-derived neutrophils were transferred by tail vein injection and 24 h later infected corneas were isolated for flow cytometry and quantification of CFU. The purity of donor bone marrow neutrophil preparation was estimated by FACS analysis using Ly6G antibody (Supplementary Fig. 4A). As shown in Fig. 7d, flow cytometry analysis showed no difference in the number of infiltrating neutrophils in  $Cd18^{-/-}$  mice with either C57BL/6 or  $P2X_7^{-/-}$  donor cells. Figure 7e shows significantly higher bacterial CFU in the corneas of infected  $\text{CD18}^{-/-}$  mice compared with C57BL/6 mice indicative of impaired bacterial killing in absence of neutrophil infiltration in  $Cd18^{-/-}$  mice. Interestingly,  $Cd18^{-/-}$  mice that received  $P2X_7^{-/-}$  donor neutrophils had significantly higher bacterial CFU compared with mice that received C57BL/6 donor neutrophils (Fig. 7f).

Taken together, these data indicate that the  $P2X_7R$  expression on neutrophils is required for NLRP3 inflammaomse activation and bacterial killing in *S. pneumoniae* corneal infection.

### Discussion

Although most studies on inflammasomes and IL-1 $\beta$  processing have examined macrophages and dendritic cells, several reports indicate that neutrophils are also a major source of IL-1 $\beta$  in infectious and inflammatory diseases. Inflammasome driven caspase-1 activation by neutrophils is the major driver of IL-1 $\beta$ cleavage in *Salmonella typhimurium*, *Staphylococcus aureus* and *S. pneumonia* infections, and in nigericin-stimulated cells<sup>15,28,31,34</sup>. Also, inflammasome- and caspase-1-independent IL-1 $\beta$  secretion by neutrophils has been reported in murine models of *Pseudomonas aeruginosa* infection and rheumatoid arthritis<sup>30,35</sup>. We recently demonstrated that human and murine neutrophils express NLRP3, and that secondary activation with bacterial pneumolysin induces K<sup>+</sup> efflux and formation of multiple NLRP3/ASC specks together with active caspase-1, resulting in secretion of mature IL-1 $\beta$  (ref. 28). The presence of multiple specks



**Figure 7 | The role of neutrophil P2X<sub>7</sub> receptor expression on** *S. pneumoniae* **corneal infection.** (**a**-c). *S. pneumoniae* infected corneas of C57BL/6 and  $P2X_7^{-/-}$  mice. Total neutrophils and macrophages quantified by flow cytometry (**a**), and bacterial CFU (**b**) 24 h post infection. (**c**). Western blot analysis of pro- and cleaved caspase-1 and IL-1 $\beta$  from 24 h infected corneas. (**d**-**f**).  $CD18^{-/-}$  mice were infected with *S. pneumoniae* and 3 h later injected intravenously with donor bone marrow neutrophils from C57BL/6 or  $P2X_7^{-/-}$  mice. Total neutrophils in infected corneas of recipient  $CD18^{-/-}$  mice given C57BL/6 or  $P2X_7^{-/-}$  mice (**b**); and in infected  $CD18^{-/-}$  mice given donor bone marrow neutrophils (**d**); CFU in corneas of infected C57BL/6 and  $CD18^{-/-}$  mice (**e**); and in infected  $CD18^{-/-}$  mice given donor bone marrow neutrophils from C57BL/6 or  $P2X_7^{-/-}$  mice (**f**). These experiments were repeated twice with similar results. 24 h post infection mice were sacrificed and and bacterial quantified. Each data point represents one cornea and histograms are mean ± s.d. of at least three corneas per group. A *P* value  $\leq 0.05$  was considered significant using a Student's t-test.

is distinct from macrophages, which except under conditions of NLRC4 gain of function mutations<sup>36</sup>, form single ASC specks<sup>37</sup>. Also, in contrast to macrophages, activation of caspase-1 in neutrophils does not lead to pyroptosis<sup>28,34</sup>.

Although neutrophils do not undergo pyroptosis, the precursor and mature forms of caspase-1 and IL-1 $\beta$  are secreted after ATP or nigericin stimulation. A similar mechanism occurs in human monocytes, where pro IL-1 $\beta$  and caspase-1 are packaged together in late endosomes or in secretory lysosomes, resulting in release of pro- and mature forms simultaneously<sup>38,39</sup>. Similarly, precursor caspase-1 and IL-1 $\beta$  are compartmentalized in microvesicles in ATP-stimulated human monocytes and dendritic cells, resulting in IL-1 $\beta$  processing before release of pro- and mature forms of the proteins<sup>21,40,41</sup>. It has yet to be determined if neutrophils utilize a similar non-lytic mechanism to concomitantly release pro- and mature forms of caspase-1 and IL-1 $\beta$ .

We reported that the exogenous pore-forming bacterial toxin pneumolysin drives  $K^+$  efflux-dependent NLRP3 inflammasomes activation in neutrophils<sup>28</sup>, and in that report and in the current study, we show that IL-1 $\beta$  secretion by murine neutrophils requires priming with either LPS or with heat killed *S. pneumoniae* before stimulation with pneumolysin or ATP. However, during *S. pneumoniae* corneal infection, live bacteria appear to provide both the priming and inflammasome activation signals required to generate the cleaved form of IL-1 $\beta$ .

Results from the current study also demonstrate that endogenously expressed ion channels trigger this pathway in murine and human neutrophils.  $P2X_7R$  is a non-selective cation channel that mediates rapid influx of  $Ca^{2+}$  and  $Na^+$ , and efflux of K<sup>+</sup> in response to high concentrations of ATP, which results in NLRP3 activation in macrophages and dendritic cells<sup>16,42,43</sup>. However, several reports indicate that  $P2X_7R$  is either not expressed on the cell surface of human neutrophils, or is present but non-functional<sup>11-14</sup>.

In the current study, we demonstrate constitutive surface expression of P2X7R on murine and human neutrophils, and show ATP-driven NLRP3/caspase-1 activation and cleavage and secretion of IL-1 $\beta$  by these cells. Our findings also provide a mechanistic context for earlier reports showing P2X7R transcripts in HL-60 promyelocytes, differentiated HL-60 granulocytes and human blood neutrophils<sup>9,10</sup>, and support studies showing that LPS-primed human neutrophils secrete IL-1ß in response to ATP via a caspase-1-dependent pathway<sup>44,45</sup>, although these investigators did not identify the ATP-sensing receptor. Similarly, there is one report of ATP-induced IL-1 $\beta$  secretion by murine neutrophils<sup>15</sup>; however, as with human neutrophils, the ATP-sensing receptor was not defined. In addition to P2X<sub>7</sub>R, ATP also activates the G-protein-coupled P2Y receptors, which in neutrophils potentiates chemotaxis, superoxide production and exocytosis of primary granules<sup>46,47</sup>. However, our current findings used specific antagonists and  $P2X_7^{-/-}$  mice to demonstrate that P2X<sub>7</sub>R is the critical ATP sensing receptor for NLRP3 inflammasome signalling and IL-1ß secretion in murine and human neutrophils.

High concentrations of extracellular ATP generated during inflammation engage low-affinity  $P2X_7R$  in mononuclear cells, whereas low concentrations of extracellular ATP promote either immunostimulatory or immunosuppressive effects that are mediated by activating high ATP-affinity G-protein-coupled P2Y receptors<sup>2,7</sup>. Specifically, high concentrations of ATP stimulate macrophage production of IL-1 $\alpha$ , IL-6 and tumour necrosis factor- $\alpha$ , resulting in enhanced neutrophil chemotactic activity and increased adhesion to endothelial cells<sup>48–50</sup>.

Because local accumulation of extracellular ATP is critical for  $P2X_7R$  activation, several questions regarding  $P2X_7R$  function in neutrophils need to be considered. The requirement for

millimolar levels of extracellular ATP to activate P2X<sub>7</sub>R is in contrast to the six other P2XR subtypes, which are gated by micromolar ATP levels. We suggest that the low ATP-affinity status of P2X<sub>7</sub>R minimizes inadvertent activation of maladaptive inflammatory signalling in neutrophils until they encounter damaged host cells or bacteria at sites of local tissue infection. However, given the high ATP concentration required for P2X<sub>7</sub>R gating, and that active ecto-ATPases are present in infected tissues, the physiological relevance has to be considered.

Junger and colleagues presented compelling evidence that neutrophils can release ATP at their leading edge when migrating in response to chemoattractant gradients<sup>51</sup>. In those experiments, the locally released ATP was shown to trigger G-protein-coupled  $P2Y_2$  receptor signalling that amplifies the rapid cytoskeletal remodelling required for efficient motility. It is therefore likely that similar autocrine and paracrine pathways exist in infected and injured tissues that may transiently and locally increase interstitial ATP levels to the millimolar levels required for activation of  $P2X_7R$  signaling. In support of this concept, Kubes and colleagues reported activation of *in situ*  $P2X_7R$  signaling responses by live-tissue imaging in their models of highly localized sterile injury to the liver<sup>52</sup>.

In the current study, we used both genetic and pharmacological approaches to demonstrate that ATP-induced IL-1 $\beta$  secretion is dependent on the P2X<sub>7</sub>R in murine and human neutrophils. First, we showed that ATP-induced IL-1 $\beta$  secretion is impaired in  $P2X_7^{-/-}$  neutrophils and by highly selective P2X<sub>7</sub>R antagonists. Moreover, ATP stimulation of neutrophils caused rapid efflux of K<sup>+</sup>, which was inhibited by P2X<sub>7</sub>R antagonists, and was upstream of NLRP3 inflammasome activity. In addition, we showed constitutive surface expression of the P2X<sub>7</sub>R in murine and human neutrophils, and that in contrast to macrophages<sup>53</sup>, receptor expression was not increased following LPS priming. However, we found that neutrophils also constitutively expressed an intracellular pool of P2X<sub>7</sub>R, which raises the possibility that these receptors may be mobilized to the cell surface during inflammation.

Examination of functional P2X<sub>7</sub>R is complicated by the presence of both C- and N-terminal splice variants of the receptor in mice<sup>20,54</sup>. P2X<sub>7</sub> knockout mice have been generated either by inserting a lacZ and a neomycin cassette into exon1 (Glaxo), or by inserting a neomycin cassette into exon 13 (Pfizer). Macrophages and T cells from Glaxo  $P2X_7^{-/-}$  mice show enhanced IL-6 production and P2X<sub>7</sub>R-mediated responses, respectively<sup>22,23</sup>. This was later explained by the presence of a novel transcript variant-P2X<sub>7</sub>K, in the Glaxo mice, which had higher sensitivity to P2X<sub>7</sub>R agonist<sup>24</sup>. In the current study, we used bone-marrow-derived neutrophils from the Pfizer  $P2X_7^{-/-}$  mice, and showed the presence of low levels of C-terminal transcript variants in neutrophils as detected by western blot using antibody against the extracellular domain of the receptor. Moreover,  $P2X_7^{-/-}$ neutrophils exhibited an incomplete inhibition of IL-1ß secretion, which is in agreement with reduced function of  $\Delta C$  splice variants in the Pfizer  $P2X_7^{-/-}$  mice<sup>20</sup>.

C57BL/6 mice harbour an allelic mutation (P451L) in the intracellular C-terminal domain of the P2X<sub>7</sub>R, whereas the non-mutated form is expressed in BALB/c mice. C57BL/6T cells expressing this Leu-451 variant exhibit a reduced rate of ATP-induced Ca<sup>2+</sup> influx and attenuated apoptotic cell surface phosphatidylserine relative to BALB/c T cells expressing the Pro-451 P2X<sub>7</sub>R (ref. 25). Similarly, HEK cells transfected with P2X<sub>7</sub>-P451L complementary DNA showed reduced calcium influx compared with cells transfected with non-mutated P2X<sub>7</sub> complementary DNA<sup>25</sup>.

However, BMDMs from C57BL/6 mice exhibit robust  $P2X_7R$ mediated  $Ca^{2+}$  influx responses to millimolar ATP levels that are absent in macrophages from the Pfizer  $P2X_7^{-/-}$  mice<sup>21</sup>. We also compared ATP-induced processing and release of IL-1 $\beta$  in BMDM from C57BL/6 and BALB/c mice, and found that although BALB/c macrophages produced more IL-1 $\beta$  in response to lower concentrations of ATP compared with C57BL/6 macrophages, IL-1 $\beta$  release was equivalent at >3 mM ATP<sup>55</sup>. These findings are consistent with data presented in the current study, which used 3 mM ATP to stimulate C57BL/6 neutrophils to produce a robust P2X<sub>7</sub>-mediated Ca<sup>2+</sup> influx, K<sup>+</sup> efflux and processing and secretion of IL-1 $\beta$ .

Heterogeneity in human P2X7R function has been reported in healthy individuals, with marked variation in macrophage responses to ATP<sup>56</sup>. For example, although one report showed a correlation between ATP response and surface expression of human P2X<sub>7</sub>R (ref. 57), others have shown that a singlenucleotide polymorphism in the exon 13 region of human P2X<sub>7</sub>R results in loss of receptor function despite no loss of surface expression<sup>58</sup>, and other P2X<sub>7</sub>R polymorphisms have been identified that alter the function or the localization of the receptor<sup>59</sup>. Further, a single-nucleotide polymorphism (496  $E \rightarrow$ A) is present at low frequency in the Caucasian population, which results in a non-functional P2X7 protein when expressed as a homozygous mutation compared with partial function in heterozygous, although surface expression was normal in both groups<sup>58</sup>. Although P2X<sub>7</sub>R had no role in a murine model of Mycobacterium tuberculosis infection [REF 60 Myers], the 496  $E \rightarrow A$  loss-of-function polymorphism was found to be associated with increased susceptibility to tuberculosis<sup>60,61</sup>. Conversely, the  $348A \rightarrow T$  gain-of-function mutation is associated with increased resistance to congenital and ocular toxoplasmosis<sup>61,62</sup>.

Because extracellular ATP can stimulate increases in cytosolic [Ca<sup>2+</sup>] via engagement of multiple P2Y and P2X receptor subtypes, we used highly selective  $P2X_7$  antagonists to assess the contribution of  $P2X_7R$ -mediated  $Ca^{2+}$  influx to the overall ATP-induced  $Ca^{2+}$  mobilization in human neutrophils. In the current study, we tested peripheral blood neutrophils from 11 healthy donors of different ethnicities and found that: (1) all donors responded to ATP; (2) 10 of 11 donors (90.1%) exhibited a  $P2X_7R$ -dependent increase in cytosolic  $[Ca^{2+}]$ , with only one donor who was completely insensitive to P2X7 antagonism, indicating that this donor expressed a non-functional P2X7 variant; (3) there was considerable variation in the P2X<sub>7</sub>R antagonist-sensitive responses among donors; and (4) there was no apparent relationship between P2X7R function and ethnic origin (Table 1), although a much larger number of donors would need to be examined to determine whether there is a role for ethnicity in P2X7R-dependent responses with ATP. However, variation in the relative expression levels in functional P2X7R receptor in the neutrophils of different human subjects may in part explain discrepancies in previous studies of P2X7R expression in human neutrophils.

Finally, we investigated the relevance of our *in vitro* findings using a mouse model of acute *S. pneumoniae* corneal infection in which neutrophils are the major source of precursor and cleaved IL-1 $\beta$  (ref. 28). We demonstrate here that neutrophils are also the major P2X<sub>7</sub>R-expressing cells in the cornea, and showed by adoptive transfer studies that neutrophil P2X<sub>7</sub>R is required for IL-1 $\beta$  processing *in vivo*, and for optimal bacterial clearance during *S. pneumoniae* corneal infection.

In conclusion, we provide evidence of functional  $P2X_7R$  expression in murine and human neutrophils, which mediates ATP induced NLRP3 inflammasome activation and IL-1 $\beta$  secretion. Since neutrophils are abundant in early stages of microbial infections, sterile inflammation, and auto-inflammatory diseases where extracellular ATP is generated, activation of P2X<sub>7</sub>R on these cells exacerbates inflammatory

responses by mediating NLRP3 inflammasome assembly and IL-1 $\beta$  secretion.

#### Methods

**Source of mice**. C57BL/6 mice (6- to 10-week old) were from The Jackson Laboratory (Bar Harbor, ME). *Caspase1/11<sup>-/-</sup>* mice were generated by Richard Flavell (Yale University, CT) as *Caspase1<sup>-/-</sup>* mice and subsequently found to be deficient in caspase-11 (ref. 62). *Nlrp3<sup>-/-</sup>* and *Asc<sup>-/-</sup>* mice were generated by Millennium Pharmaceuticals (Cambridge, MA). *Cd18<sup>-/-</sup>* mice were kindly provided by Claire Doerschuk (University of North Carolina, Chapel Hill, NC). A *P2X<sub>7</sub><sup>-/-</sup>* mouse strain was originally provided by Pfizer Global Research and Development. All the gene knockout mice were on a C57BL/6 background. Animals were housed in pathogen free conditions in microisolator cages and were treated according to institutional guidelines after approval by the Case Western Reserve University IACUC.

**Source of reagents.** P2X<sub>7</sub>R inhibitors AZ10606120 dihydrochloride and AZ11645373, A438079 hydrochloride (Tocris Bioscience) were dissolved in DMSO and used at concentrations noted in the results. BzATP triethylammonium salt (Tocris Bioscience) was dissolved in sterile water and used at millimolar concentrations. Nigericin (Calbiochem) was used at 10  $\mu$ M concentration for all *in vitro* assays. Apyrase (Sigma Aldrich) was used at 10 U ml<sup>-1</sup>. ATP disodium salt hydrate (Sigma Aldrich) was dissolved in water and used at micromolar concentrations. Ultrapure LPS (Adipogen) was dissolved in endotoxin free water and used at 500 ng ml<sup>-1</sup> for *in vitro* assays.

Isolation of human and mouse bone marrow neutrophils. Human neutrophils were isolated from the peripheral blood of healthy volunteers following informed consent as approved by the Institutional Review Board of University Hospitals of Cleveland. Heparinized blood was mixed with 3% dextran (Sigma Aldrich) in PBS for 20 min at room temperature (RT). The top clear layer containing leukocytes was transferred to a fresh tube and the cells were under laid with 10 ml of Ficoll Paque Plus (GE Healthcare), and centrifuged at 300g for 20 min. The overlying plasma was aspirated and PBMC layer was collected. The underlying neutrophil/RBC pellet was suspended in  $1 \times RBC$  lysis buffer (eBiosciences) and neutrophils were washed in sterile PBS and suspended in RPMI + 1-glutamine media (Hyclone) supplemented with 2% fetal bovine serum (FBS; Mediatech). This protocol routinely yielded > 97% cell purity as assessed by flow cytometry and Wrights–Giemsa stain (Sigma Aldrich).

For mouse neutrophils, total bone marrow cells were collected from tibias and femurs, and neutrophils were isolated from the total bone marrow cells by negative selection using magnetic bead-based EasySep Mouse Neutrophil Enrichment Kit (Stem Cell Technologies). Neutrophils were then washed with PBS and suspended in RPMI + L-glutamine with 1% FBS. This procedure routinely yielded >94% pure neutrophils as quantified by flow cytometry and Wright-Giemsa stain (Supplementary Fig. 1A).

Western blot analysis. Neutrophil lysates  $(20-30 \ \mu g \ protein)$  were fractionated in 12% SDS-PAGE, transferred onto nitrocellulose membranes, and incubated with primary antibodies—mouse IL-1 $\beta$  (R&D Systems, AF-401-NA at 1:1,300), mouse Caspase-1 p10 (Santa Cruz, SC-514 at 1:200), or with antibodies targeted to the intracellular C terminus or extracellular region of P2X<sub>7</sub>R (Alomone Labs, catalog# APR 004 for the C terminal, and APR 008 for the ecto-domain, both at 1:200 dilution). These antibodies recognize mouse and human receptors. Loading controls were shown using antibodies to  $\beta$  actin (Sigma Aldrich A3854, 1:50 000). Reactivity was determined using HRP-conjugated secondary antibodies (Santa Cruz) and developed using Supersignal West Femto Maximum Sensitivity Substrate (Pierce). Images were cropped for presentation; full size images are presented in Supplementary Fig. 5.

**Flow cytometry analysis.** Neutrophils were washed twice in PBS after *in vitro* stimulation. Fc receptors were blocked for 20 min at RT with anti-mouse or anti-human CD16/32 antibody (eBiosciences 16-0161-81) followed by incubation with FITC tagged anti-Ly6G (clone 1A8, Biolegend, 127613, 0.5 µg added to  $1 \times 10^6$  murine neutrophils in 100 µl) and anti-P2X<sub>7</sub> receptor antibodies as described above. F(ab')2 donkey anti-rabbit IgG-PE (eBiosciences) was used as secondary antibody for P2X<sub>7</sub>. Following two washes in FACS Buffer (1% FBS in PBS), cells were fixed in 0.5% PFA for analysis by flow cytometry using an Accuri C6 Flow cytometre (Becton Dickinson).

**Intracellular Ca<sup>2+</sup> assay.** Human and mouse neutrophils were plated at  $1 \times 10^6$  per well in a 24-well plate, and incubated with LPS for 3 h. Neutrophils were then washed with PBS and transferred to basal salt solution media (130 mM NaCl, 4 mM KCl, 1.5 mM CaCl<sub>2</sub>, 1 mM MgCl<sub>2</sub>, 25 mM HEPES, 5 mM p-glucose, 1% BSA, pH 7.4) containing 1  $\mu$ M Fluo-4-AM (Life Technologies) and 2.5 mM probenecid (Sigma Aldrich) for 30 min at 37 °C. Fluo-4-AM was mixed with 20% pluronic acid F-127 in a 1:1 ratio before additing to BSS media. After incubation, Fluo-4-AM

## ARTICLE

containing media was aspirated, cells were washed in BSS media and then resuspended in BSS media containing 2.5 mM probenecid. Baseline readings were taken at 37 °C using a fluorescent plate reader (Synergy HT, BioTek) at 485/528 nm for 10 min, after which ATP was added to the cells and kinetic readings were obtained over 30 min. Cells were permeabilized with 1% Triton-X 100 to obtain the maximum Ca<sup>2+</sup>-dependent fluorescence, followed by Tris/EGTA addition to obtain Ca<sup>2+</sup>-independent background fluorescence.

**Fluorescence microscopy**. Neutrophils were fixed with 4% PFA (Fisher) at RT for 15 min. Cells were then blocked in 10% goat serum (Vector Laboratories) in PBS for 1 h at RT, followed by staining with P2X<sub>7</sub> antibody (clone Hano3, Enzo Life Science ALX-802-027, 1:5) for 1 h at 37°C. Neutrophils were then washed three times in PBS and incubated 40 min at RT with Alexa-488 goat anti-rat IgG (Jackson Immuno Research 112-007-003, 1:300), washed three times in PBS and counter stained with DAPI before microscopy. Images were collected on a Leica DMI 6000 B inverted microscope using a × 63 objective connected to a Retiga EXi Aqua Blue camera (Q-imaging). Resulting image stacks were subjected to deconvolution using Autoquant software (Mediacybernetics). Further analysis was performed using Metamorph Imaging Software (Molecular Devices).

Atomic absorbance spectroscopy. Neutrophils were washed in potassium-free isotonic buffer (135 mM sodium gluconate, 1.5 mM CaCl<sub>2</sub>, 1 mM MgCl<sub>2</sub>, and 25 mM HEPES). Cell pellets were then extracted in 1 ml of 10% HNO<sub>3</sub>, and K<sup>+</sup> content in the nitric acid extracts was quantified by atomic absorbance spectrometry (Agilent 55B AA)<sup>28</sup>. Triplicate samples were run for all test conditions in each experiment.

**LDH release assay.** Neutrophil supernatants were collected and LDH release was quantified using CytoTox 96 Non-Radioactive Cytotoxicity Assay (Promega) according to the manufacturer's instructions. Percentage cytotoxicity was calculated based on LDH release in total cell lysates.

**ELISA**. Half-well cytokine assays were performed using Duoset ELISA assay kits for murine and human IL-1 $\beta$  and mouse IL-1 $\alpha$  according to manufacturer's protocol (R&D Systems).

**IL-1 Bioassay.** HEK-Blue-IL-1R reporter cell line (Invivogen) was used to measure production of biologically active IL-1 (mature form of the cytokine). This reporter cell line stably expresses murine IL-1 receptor and a secreted alkaline phosphatase (SEAP) reporter gene under the control of a minimal IFN-β promoter fused to NFκB and AP-1 binding sites. The cells were cultured in 24-well plates at 2.5 × 10<sup>5</sup> cells in 500 µl. They were then incubated with 5 µl supernatant from stimulated bone marrow neutrophils or with recombinant murine IL-1β standards. After 15–18 h, 20 µl aliquots of conditioned medium (either undiluted or diluted 1:5 or 1:10) from the stimulated HEK- Blue-IL-1R cells were transferred to 96-well plates containing 180 µl of QUANTI-Blue SEAP assay reagent per well. Production of blue SEAP product was measured by absorbance (620 nm) in BioTek Synergy HT plate reader, and the concentration was derived from the standard curve (known IL-1β standards) and expressed as pg per ml. As this IL-1R-based bioassay also detects IL-1α, the same samples of neutrophil-conditioned medium were assayed for murine IL-1β by ELISA (R&D).

**Murine model of Streptococcus pneumoniae corneal infection.** S. pneumoniae TIGR4 (serotype IV) were grown at 5%  $CO_2$  in Todd-Hewitt Broth (Neogen) supplemented with 0.5% yeast extract. Bacteria were grown to mid-exponential phase ( $1x10^8$  CFU per ml), and  $1 \times 10^5$  bacteria in 2 µl sterile PBS were injected into the corneal stroma. After 24 h, corneas were dissected, and either digested in 82 U of type- I collagenase (Sigma) to recover cells for flow cytometry, homogenized in lysis buffer for western blot analysis, or whole eyes were homogenized and viable bacteria were quantified by CFU.

For flow cytometry, cell suspensions from infected corneas were passed through a 30-µm filter to remove undigested tissue, and cells were identified and quantified by flow cytometry. For CFU, whole eyes were homogenized in 1 ml sterile PBS using a Mixer Mill MM300 (Retsch). Serial log dilutions were plated on blood agar plates, incubated in a CO<sub>2</sub> incubator at 37 °C for 18 h, and CFU were counted manually.

For *in vivo* neutrophil depletion, mice were injected intraperitoneally with 250 µg of the Ly6G antibody NIMP-R14 24 h before corneal infection (NIMP-R14 was generated in-house from the hybridoma). In neutrophil adoptive transfer model experiments, corneas of recipient  $Cd18^{-/-}$  mice were infected with *S. pneumoniae* as described above, and after 3 h,  $8 \times 10^6$  purified bone marrow neutrophils from donor C57BL/6 or  $P2X_7^{-/-}$  mice were injected intravenously to the infected  $Cd18^{-/-}$  mice. After 24 h, corneas were collagenase digested and cells were examined by flow cytometry.

**Statistical analysis**. Student's *t*-test or analysis of variance with Tukey *post hoc* analysis (Prism, Graphpad Software) were used as indicated in the figure legends. A *P* value of < 0.05 was considered significant.

#### References

- Fitz, J. G. Regulation of cellular ATP release. Trans. Am. Clin. Climatol. Assoc. 118, 199–208 (2007).
- Junger, W. G. Immune cell regulation by autocrine purinergic signalling. *Nat. Rev. Immunol.* 11, 201–212 (2011).
- 3. Trautmann, A. Extracellular ATP in the immune system: more than just a "danger signal". *Sci. Signal.* **2**, pe6 (2009).
- Cauwels, A., Rogge, E., Vandendriessche, B., Shiva, S. & Brouckaert, P. Extracellular ATP drives systemic inflammation, tissue damage and mortality. *Cell Death Dis.* 5, e1102 (2014).
- Riteau, N. et al. Extracellular ATP is a danger signal activating P2X7 receptor in lung inflammation and fibrosis. Am. J. Respir. Crit. Care Med. 182, 774–783 (2010).
- Shah, D., Romero, F., Stafstrom, W., Duong, M. & Summer, R. Extracellular ATP mediates the late phase of neutrophil recruitment to the lung in murine models of acute lung injury. *Am. J. Physiol. Lung Cell Mol. Physiol.* 306, L152–L161 (2014).
- Gombault, A., Baron, L. & Couillin, I. ATP release and purinergic signaling in NLRP3 inflammasome activation. *Front. Immunol.* 3, 414 (2012).
- Burnstock, G. & Knight, G. E. Cellular distribution and functions of P2 receptor subtypes in different systems. *Int. Rev. Cytol.* 240, 31–304 (2004).
- Suh, B. C., Kim, J. S., Namgung, U., Ha, H. & Kim, K. T. P2X7 nucleotide receptor mediation of membrane pore formation and superoxide generation in human promyelocytes and neutrophils. *J. Immunol.* 166, 6754–6763 (2001).
- Chen, Y., Shukla, A., Namiki, S., Insel, P. A. & Junger, W. G. A putative osmoreceptor system that controls neutrophil function through the release of ATP, its conversion to adenosine, and activation of A2 adenosine and P2 receptors. J. Leukoc. Biol. 76, 245–253 (2004).
- Gu, B. J. et al. Expression of P2X(7) purinoceptors on human lymphocytes and monocytes: evidence for nonfunctional P2X(7) receptors. Am. J. Physiol. Cell Physiol. 279, C1189–C1197 (2000).
- Martel-Gallegos, G. et al. Human neutrophils do not express purinergic P2X7 receptors. Purinergic Signal. 6, 297–306 (2010).
- Vaughan, K. R. et al. Inhibition of neutrophil apoptosis by ATP is mediated by the P2Y11 receptor. J. Immunol. 179, 8544–8553 (2007).
- 14. Mohanty, J. G., Raible, D. G., McDermott, L. J., Pelleg, A. & Schulman, E. S. Effects of purine and pyrimidine nucleotides on intracellular Ca2 + in human eosinophils: activation of purinergic P2Y receptors. J. Allergy Clin. Immunol. 107, 849–855 (2001).
- Mankan, A. K., Dau, T., Jenne, D. & Hornung, V. The NLRP3/ASC/Caspase-1 axis regulates IL-1beta processing in neutrophils. *Eur. J. Immunol.* 42, 710–715 (2012).
- Surprenant, A. & North, R. A. Signaling at purinergic P2X receptors. Annu. Rev. Physiol. 71, 333–359 (2009).
- Donnelly-Roberts, D. L., Namovic, M. T., Han, P. & Jarvis, M. F. Mammalian P2X7 receptor pharmacology: comparison of recombinant mouse, rat and human P2X7 receptors. *Br. J. Pharmacol.* 157, 1203–1214 (2009).
- Miao, E. A., Rajan, J. V. & Aderem, A. Caspase-1-induced pyroptotic cell death. Immunol. Rev. 243, 206–214 (2011).
- Surprenant, A., Rassendren, F., Kawashima, E., North, R. A. & Buell, G. The cytolytic P2Z receptor for extracellular ATP identified as a P2X receptor (P2X7). Science 272, 735–738 (1996).
- Masin, M. et al. Expression, assembly and function of novel C-terminal truncated variants of the mouse P2X7 receptor: re-evaluation of P2X7 knockouts. Br. J. Pharmacol. 165, 978–993 (2012).
- Qu, Y., Franchi, L., Nunez, G. & Dubyak, G. R. Nonclassical IL-1 beta secretion stimulated by P2X7 receptors is dependent on inflammasome activation and correlated with exosome release in murine macrophages. *J. Immunol.* 179, 1913–1925 (2007).
- Chessell, I. P. et al. Disruption of the P2X7 purinoceptor gene abolishes chronic inflammatory and neuropathic pain. Pain 114, 386–396 (2005).
- 23. Taylor, S. R. *et al.* Lymphocytes from P2X7-deficient mice exhibit enhanced P2X7 responses. *J. Leukoc. Biol.* **85**, 978–986 (2009).
- 24. Nicke, A. *et al.* A functional P2X7 splice variant with an alternative transmembrane domain 1 escapes gene inactivation in P2X7 knock-out mice. *J. Biol. Chem.* 284, 25813–25822 (2009).
- Adriouch, S. *et al.* Cutting edge: a natural P451L mutation in the cytoplasmic domain impairs the function of the mouse P2X7 receptor. *J. Immunol.* 169, 4108–4112 (2002).
- Munoz-Planillo, R. *et al.* K(+) efflux is the common trigger of NLRP3 inflammasome activation by bacterial toxins and particulate matter. *Immunity* 38, 1142–1153 (2013).

- Katsnelson, M. A., Rucker, L. G., Russo, H. M. & Dubyak, G. R. K + efflux agonists induce NLRP3 inflammasome activation independently of Ca2 + signaling. J. Immunol. 194, 3937–3952 (2015).
- Karmakar, M. *et al.* Neutrophil IL-1beta processing induced by pneumolysin is mediated by the NLRP3/ASC inflammasome and caspase-1 activation and is dependent on K + efflux. *J. Immunol.* **194**, 1763–1775 (2015).
- 29. Hazlett, L. D. Bacterial infections of the cornea (*Pseudomonas aeruginosa*). Chem. Immunol. Allergy. **92**, 185–194 (2007).
- 30. Karmakar, M., Sun, Y., Hise, A. G., Rietsch, A. & Pearlman, E. Cutting edge: IL-1beta processing during Pseudomonas aeruginosa infection is mediated by neutrophil serine proteases and is independent of NLRC4 and caspase-1. *J. Immunol.* 189, 4231–4235 (2012).
- 31. Leal, Jr. S. M. *et al.* Fungal antioxidant pathways promote survival against neutrophils during infection. *J. Clin. Invest.* **122**, 2482–2498 (2012).
- Taylor, P. R. et al. Activation of neutrophils by autocrine IL-17 A-IL-17RC interactions during fungal infection is regulated by IL-6, IL-23, RORgammat and dectin-2. Nat. Immunol. 15, 143–151 (2014).
- Wilson, R. W. et al. Gene targeting yields a CD18-mutant mouse for study of inflammation. J. Immunol. 151, 1571–1578 (1993).
- Chen, K. W. *et al.* The neutrophil NLRC4 inflammasome selectively promotes IL-1beta maturation without pyroptosis during acute Salmonella challenge. *Cell Rep.* 8, 570–582 (2014).
- Guma, M. *et al.* Caspase 1-independent activation of interleukin-1beta in neutrophil-predominant inflammation. *Arthritis Rheum.* 60, 3642–3650 (2009).
- Romberg, N. et al. Mutation of NLRC4 causes a syndrome of enterocolitis and autoinflammation. Nat. Genet. 46, 1135–1139 (2014).
- Latz, E., Xiao, T. S. & Stutz, A. Activation and regulation of the inflammasomes. *Nat. Rev. Immunol.* 13, 397–411 (2013).
- Andrei, C. *et al.* The secretory route of the leaderless protein interleukin 1beta involves exocytosis of endolysosome-related vesicles. *Mol. Biol. Cell* 10, 1463–1475 (1999).
- Andrei, C. *et al.* Phospholipases C and A2 control lysosome-mediated IL-1 beta secretion: Implications for inflammatory processes. *Proc. Natl Acad. Sci. USA* 101, 9745–9750 (2004).
- 40. MacKenzie, A. *et al.* Rapid secretion of interleukin-1beta by microvesicle shedding. *Immunity* **15**, 825–835 (2001).
- Pizzirani, C. *et al.* Stimulation of P2 receptors causes release of IL-1 beta-loaded microvesicles from human dendritic cells. *Blood* 109, 3856–3864 (2007).
- 42. Di Virgilio, F. Liaisons dangereuses: P2X(7) and the inflammasome. *Trends Pharmacol. Sci.* 28, 465–472 (2007).
- Sanz, J. M. & Di Virgilio, F. Kinetics and mechanism of ATP-dependent IL-1 beta release from microglial cells. J. Immunol. 164, 4893–4898 (2000).
- 44. Bakele, M. et al. Localization and functionality of the inflammasome in neutrophils. J. Biol. Chem. 289, 5320–5329 (2014).
- 45. Gabelloni, M. L. *et al.* NADPH oxidase derived reactive oxygen species are involved in human neutrophil IL-1beta secretion but not in inflammasome activation. *Eur. J. Immunol.* **43**, 3324–3335 (2013).
- Coutinho-Silva, R., Correa, G., Sater, A. A. & Ojcius, D. M. The P2X(7) receptor and intracellular pathogens: a continuing struggle. *Purinergic Signal.* 5, 197–204 (2009).
- Freyer, D. R., Boxer, L. A., Axtell, R. A. & Todd, 3rd R. F. Stimulation of human neutrophil adhesive properties by adenine nucleotides. *J. Immunol.* 141, 580–586 (1988).
- Mehta, V. B., Hart, J. & Wewers, M. D. ATP-stimulated release of interleukin (IL)-1beta and IL-18 requires priming by lipopolysaccharide and is independent of caspase-1 cleavage. J. Biol. Chem. 276, 3820–3826 (2001).
- Solle, M. et al. Altered cytokine production in mice lacking P2X(7) receptors. J. Biol. Chem. 276, 125–132 (2001).
- Tonetti, M., Sturla, L., Giovine, M., Benatti, U. & De Flora, A. Extracellular ATP enhances mRNA levels of nitric oxide synthase and TNF-alpha in lipopolysaccharide-treated RAW 264.7 murine macrophages. *Biochem. Biophys. Res. Commun.* 214, 125–130 (1995).
- 51. Chen, Y. *et al.* ATP release guides neutrophil chemotaxis via P2Y2 and A3 receptors. *Science* **314**, 1792–1795 (2006).

- 52. McDonald, B. *et al.* Intravascular danger signals guide neutrophils to sites of sterile inflammation. *Science* **330**, 362–366 (2010).
- Humphreys, B. D. & Dubyak, G. R. Induction of the P2z/P2X7 nucleotide receptor and associated phospholipase D activity by lipopolysaccharide and IFN-gamma in the human THP-1 monocytic cell line. *J. Immunol.* 157, 5627–5637 (1996).
- 54. Xu, X. J. *et al.* Splice variants of the P2X7 receptor reveal differential agonist dependence and functional coupling with pannexin-1. *J. Cell Sci.* **125**, 3776–3789 (2012).
- Verhoef, P. A., Kertesy, S. B., Lundberg, K., Kahlenberg, J. M. & Dubyak, G. R. Inhibitory effects of chloride on the activation of caspase-1, IL-1beta secretion, and cytolysis by the P2X7 receptor. J. Immunol. 175, 7623–7634 (2005).
- Lammas, D. A. *et al.* ATP-induced killing of mycobacteria by human macrophages is mediated by purinergic P2Z(P2X7) receptors. *Immunity* 7, 433–444 (1997).
- 57. Li, C. M., Campbell, S. J., Kumararatne, D. S., Hill, A. V. & Lammas, D. A. Response heterogeneity of human macrophages to ATP is associated with P2X7 receptor expression but not to polymorphisms in the P2RX7 promoter. *FEBS Lett.* 531, 127–131 (2002).
- Gu, B. J. et al. A Glu-496 to Ala polymorphism leads to loss of function of the human P2X7 receptor. J. Biol. Chem. 276, 11135–11142 (2001).
- 59. Wiley, J. S., Sluyter, R., Gu, B. J., Stokes, L. & Fuller, S. J. The human P2X7 receptor and its role in innate immunity. *Tissue Antigens* **78**, 321–332 (2011).
- Myers, A. J., Eilertson, B., Fulton, S. A., Flynn, J. L. & Canaday, D. H. The purinergic P2X7 receptor is not required for control of pulmonary Mycobacterium tuberculosis infection. *Infect. Immunol.* 73, 3192–3195 (2005).
- Xiao, J. et al. Metaanalysis of P2X7 gene polymorphisms and tuberculosis susceptibility. FEMS. Immunol. Med. Microbiol. 60, 165–170 (2010).
- Jamieson, S. E. et al. Evidence for associations between the purinergic receptor P2X(7) (P2RX7) and toxoplasmosis. *Genes. Immunol.* 11, 374–383 (2010).

#### Acknowledgements

We thank Drs Scott Howell and Yan Sun for outstanding technical assistance with confocal microscopy and *in vivo* corneal infections. This work was supported by National Institutes of Health Grants R01 EY14362 (E.P.), R01 GM36387 (G.R.D.), and the Visual Sciences Research Center Core grant (P30 EY11373). Additional support for this work was provided by the Research to Prevent Blindness Foundation.

#### Authorcontributions

M.K. and M.A.K. performed the experiments and analysed the data. M.K., G.R.D. and E.P. conceived the project, analysed the data and prepared the manuscript.

#### Additional information

Supplementary Information accompanies this paper at http://www.nature.com/ naturecommunications

Competing financial interests: The authors declare no competing financial interests.

**Reprints and permission** information is available online at http://npg.nature.com/ reprintsandpermissions.

How to cite this article: Karmakar, M. *et al.*, Neutrophil P2X7 receptors mediate NLRP3 inflammasome-dependent IL-1ß secretion in response to ATP. 7:10555 doi: 10.1038/ncomms10555 (2016).

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/