UC Irvine

UC Irvine Previously Published Works

Title

Mechanisms by which metals promote events connected to neurodegenerative diseases.

Permalink

<https://escholarship.org/uc/item/4zs8z7zh>

Journal

Brain research bulletin, 55(2)

ISSN

0361-9230

Authors

Campbell, A
Smith, MA
Sayre, LM
et al.

Publication Date

2001-05-01

DOI

10.1016/s0361-9230(01)00455-5

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Mechanisms by which metals promote events connected to neurodegenerative diseases

Arezoo Campbell,^{1*} Mark A. Smith,² Lawrence M. Sayre,³ Stephen C. Bondy¹ and George Perry²

¹Department of Community and Environmental Medicine, University of California, Irvine, Irvine, CA, USA;
Departments of ²Pathology and ³Chemistry, Case Western Reserve University, Cleveland, OH, USA

ABSTRACT: Although the exact causative phenomenon responsible for the onset and progression of neurodegenerative disorders is at present unresolved, there are some clues as to the mechanisms underlying these chronic diseases. This review addresses mechanisms by which endogenous or environmental factors, through interaction with redox active metals, may initiate a common cascade of events terminating in neurodegeneration. © 2001 Elsevier Science Inc.

KEY WORDS: Copper, Iron, Prion-related protein (PrP), β -amyloid, Aluminum.

INTRODUCTION

As the brain ages, distinct alterations in oxidative and inflammatory events occur that are associated with pathological lesions characteristic of neurodegenerative diseases. Copper, manganese, iron, and other trace redox-active transition metals may be involved in mediating these processes and thus may be involved in the neuropathology of disorders such as Parkinson disease (PD), Alzheimer disease (AD), and amyotrophic lateral sclerosis (ALS). Recently it has been discovered that transition metals bind to proteins involved in neurodegeneration and this association appears to preserve metal redox activity in a manner that is consistent with a pro-oxidant, free radical generating action. The binding of multivalent metals to colloidal aluminum (Al) may parallel this phenomenon. Alterations in the levels of copper- and iron-containing metalloenzymes, involved in processing partially reduced oxygen intermediates, as well as the antioxidant status of cells, may also contribute to altered redox homeostasis in neurodegenerative diseases. Nonetheless, even in familial forms of ALS, linked to mutations in superoxide dismutase, it is not clear whether an altered enzyme activity or, indirectly, a disturbance in transition metal homeostasis is involved in the disease pathogenesis. The purpose of this review is to evaluate recent publications implicating redox-active transition metal ions in promoting oxidative events. The connection between reactive oxygen moieties in promoting glial activation and the resulting pro-inflammatory cascade will be discussed. Both processes are present in the aged and diseased brain and metal-induced exacerbation of these events may contribute to the pathological lesions characteristic of neurodegeneration.

ROLE OF REACTIVE OXYGEN SPECIES IN BIOLOGICAL SYSTEMS

The side-production of superoxide during reduction of molecular oxygen by mitochondria is inevitable in all aerobic organisms. In addition, hydrogen peroxide, produced by oxidases such as monoamine oxidase, increases susceptibility to reactive oxygen of tissues enriched in these enzymes. The principal damaging intermediate of oxidative stress is the hydroxyl radical, which alters biomacromolecules present at its site of generation. In recent years, the explosion of research on the role of nitric oxide (NO) in neurotransmission has led to the recognition of “nitrosative stress”. This is a form of oxidative stress that is directly linked to the reaction of superoxide with NO to give peroxynitrite or to antioxidant activity of NO. Although peroxynitrite also appears capable of hydroxyl radical-like activity, most of these highly reactive intermediates are formed by the Fenton reaction between reduced transition metals (usually iron(II) or copper(I)) and H₂O₂.

Of the various hypotheses that have been suggested for the etiology of idiopathic AD, the one receiving most attention is a role for oxidative stress. Disruption of homeostasis of redox-active metals is particularly significant since substantial evidence for increases in oxidative stress parameters exist [67,76,88,97–99,101, 113]. Whether the presence of enhanced oxidative events is the cause or the end result of the disease is not certain. Antibodies against 4-hydroxynonenal showed that neurons lacking neurofibrillary tangles also displayed hydroxynonenal (HNE)-pyrrole immunoreactivity while the age-matched controls only had background immunoreactivity [88]. Thus, in AD, oxidative stress is a preexisting condition that subsequently leads to neuronal changes associated with the disease process. Furthermore, in Down’s syndrome, which leads to early onset of the pathological changes of AD, evidence of oxidative stress is found long before neuropathological changes, even at the fetal stage [22,77–79].

INFLAMMATORY EVENTS IN THE AGED AND DISEASED BRAIN: RELEVANCE TO METALS

Microglial cells become progressively active with normal aging [103] and the aged hippocampal formation contains increased reactive glia [40]. The amount of activated astrocytes are also increased in both AD and normal age-matched control brains. These reactive astrocytes are associated with both senile plaques and cerebral microvessels [39]. Furthermore, the activation of the pro-inflammatory transcription factor, nuclear factor (NF)- κ B, is

* Address for correspondence: Arezoo Campbell, Department of Community and Environmental Medicine, University of California, Irvine, Irvine, CA 92697-1825, USA. Fax: +1-(949)-824-2070; E-mail: aghadimi@uci.edu

significantly correlated to an elevation in a key inflammatory enzyme COX-2, in both aging and AD temporal lobe neocortex [65].

Cytokines such as interleukin (IL)-1, IL-6, and IL-8 are primarily synthesized by activated microglia and macrophages in response to pathogens and trauma. Chronic production of these factors can result in cytotoxicity because they recruit and activate macrophages that produce high concentrations of reactive oxygen species (ROS) [43]. IL-1 and IL-6 are both elevated in the brain in AD and postischemia [23]. In an animal model of chronic inflammation, induced by infusion of lipopolysaccharide, there was astrogliosis as well as an increase in the levels of amyloid- β protein precursor (A β PP), IL-1, and tumor necrosis factor (TNF) α mRNA levels. This was subsequently followed by hippocampal cell loss and impairment of spatial memory, all of which mirror changes seen in the AD brain [106].

REDOX ACTIVE TRANSITION METALS IN NEURODEGENERATION

Multivalent transition metals such as iron, copper, and manganese are essential in most biological reactions such as the synthesis of DNA, RNA, and proteins. They are also cofactors of numerous enzymes, particularly those involved in respiration. However, abnormal tissue accumulation of redox-active transition metals can be cytotoxic because perturbations in metal homeostasis results in an array of cellular disturbances characterized by increased free radical production. Oxidative stress, defined as the imbalance between biochemical processes leading to production of ROS and the cellular antioxidant cascade, causes molecular damage that can then lead to critical failure of biological functions and ultimately cell death. The levels of copper and iron are increased in the rims of senile plaques (SP) [64] and the presence of these redox reactive metals in both SP and neurofibrillary tangles has been shown to induce hydrogen peroxide dependent oxidation [89]. This indicates that the metals may be playing a role in modulating oxidative events, which may eventually result in neurodegeneration.

COPPER IN NEURODEGENERATION

Copper has a functional role in many enzymes that require oxidation-reduction reactions. For example, the metal is found in the catalytic site of cytochrome c oxidase of the mitochondrial electron transport chain and Cu-Zn superoxide dismutase (SOD). The entry of Cu to the brain is mainly mediated by ceruloplasmin (CP) [61]. This copper binding protein is also involved in the acute phase of inflammation and plays a role in protecting cells against oxidative stress [6]. CP is considered the major Fe(II)-oxidizing enzyme in the central nervous system (CNS), and an inherited metabolic disorder called aceruloplasminemia is associated with impairment in iron homeostasis and consequent neurodegeneration [50]. While CP is increased in brain tissue and cerebrospinal fluid in AD, PD, and Huntington's disease [63], neuronal levels of CP remain unchanged [28]. Thus, while increased CP may indicate a compensatory response to increased oxidative stress in AD, its failure to do so in neurons may play an important role in metal-catalyzed damage. In fact, studies directed at clarifying the relationship between oxidative stress and tissue metal ion levels indicate that the ratio of copper to zinc and the levels of CP are significantly higher with increasing age, and higher yet in cases with neurodegeneration [73]. Redox-inert zinc may serve as an antioxidant by preventing binding of pro-oxidant copper at tissue sites [38].

Iron in Neurodegeneration

Free iron, more than any other transition metal, has been implicated in redox transitions and consequential generation of oxygen free radicals. The potential of iron to cause an increase in oxidative events leading to cellular damage is controlled by a series of iron-binding proteins. Lactoferrin, which has been shown to protect against severe inflammation, is increased in the brain of patients with neurodegenerative diseases [44]. The iron regulatory proteins (IRP-1 and IRP-2), which control other Fe-binding proteins and thus the regulation of cellular iron metabolism, also play a role in decreasing the ROS generating capacity of Fe [60]. IRP-1, but not IRP-2, is rapidly activated by extracellular H₂O₂, establishing a regulatory connection between the control of iron metabolism and response to oxidative stress. Whereas IRP-1 was found to be present in similar levels in both AD and control brain tissue, IRP-2 co-localized with redox-active iron in neurofibrillary tangles (NFT), SP neurites, and neuropil threads [102]. These results suggest that alterations in IRP-2 may be directly linked to impaired iron homeostasis in AD. We have found a marked association of redox-active iron with both NFT and SP in AD [100].

Microglia are the major sites of ferritin bound iron and are thought to be partly responsible for oxidative damage in PD and other neurodegenerative disorders. Abnormally high levels of iron as well as oxidative stress have been demonstrated in a number of neurodegenerative disorders including AD and those characterized by nigral degeneration such as PD, multiple system atrophy, and progressive supranuclear palsy. Microglia stimulated *in vivo* with phorbol ester show increased lipid peroxidation resulting from a superoxide-dependent release of iron from ferritin [112]. In addition, the interplay between oxidative stress and control of iron metabolism in the brain is exemplified by the finding of abnormal iron deposition associated with lipid peroxidation in a transgenic mouse model (expressing IL-6 in astrocytes) of a blood-brain barrier defect associated with progressive neurodegeneration [29].

Iron-Mediated Oxidative Stress and PD

Although a mutation in the α -synuclein gene in certain cases of familial Parkinsonism has been identified, the etiology of the vast majority of cases remains unknown. Nonetheless, several biochemical abnormalities in PD have been identified, including a mitochondrial complex I deficiency, oxidative stress, and excess iron. Fe accumulates in substantia nigra astrocytes of old rats [91,92] and correlating to this is an increase in the Fe(III)/Fe(II) ratio and a decrease in glutathione (GSH) [82]. One explanation is that mitochondrial sequestration of redox-active iron in aging nigral astroglia may be a factor predisposing the senescent nervous system to Parkinsonism and other neurodegenerative disorders. In fact, a persistent condition of oxidative stress, associated with perturbed intracellular redox equilibrium, is recognized as a pathogenic factor underlying neurodegeneration. Moreover, there is circumstantial evidence that the intracellular redox imbalance results in aberrant oxidation of dopamine to 6-hydroxydopamine, which in turn can undergo autooxidation to the corresponding quinone concomitant with generation of superoxide. This reaction, either by itself, or as amplified by redox cycling of this quinone may lead to further generation of ROS at the expense of cellular reductants. This can explain the ultimate demise of these neurons. Studies designed to clarify the mechanism of dopamine oxidation *in vitro* have demonstrated 6-hydroxydopamine conversion in the presence of Fe(II) and either H₂O₂ or alkyl peroxides [80].

THE ROLE OF PROTEIN-METAL BINDING IN NEURODEGENERATION

The misfolding of a normal constitutively expressed protein may lead to formation of protease-resistant neurofibrillary tangles. Aggregated prion-related protein (PrP) and the amyloid- β peptides are two such cases in which the irreversible formation of beta sheeted fibrils lead to microglial activation [7,94].

Prion Protein

PrP is a transmembrane glycoprotein that is constitutively expressed in both glial and neuronal cells. Although its function is unknown, the protein incorporates copper and has SOD activity [14–16,19]. The prion protein (PrP^C) binds copper at its N-terminal region and PrP^C-deficient mice are more prone to copper toxicity [18]. Thus, it may play a role in protecting cells from oxidative damage. The misfolding and consequent aggregation of the protein is generally accepted to be responsible for spongiform encephalopathies (prion diseases), such as the human Gerstmann-Sträussler-Scheinker's syndrome and Creutzfeldt-Jakob disease. These are characterized by the extracellular deposition of the pathological form of the PrP protein and in some cases, the co-existence of amyloid plaques, gliosis, vacuole formation, and neuronal cell death [7,20,59]. These changes parallel the pathological alterations in the brain of AD patients.

Oxidative stress is a common factor in both the aged brain and neurodegeneration and this partially may be due to the increased reactive gliosis that follows both conditions [49,85,86,103]. The prion glycoprotein, expressed on microglial cells, augments activation of these cells [17]. Microglia in turn produce ROS in response to PrP peptide (106–126) and this fragment is toxic to neuronal cell cultures only in the presence of activated glial cells [14]. *In vivo*, the brain of mice infected with scrapie showed astrogliosis and microglial activation early in the course of the disease, followed by neuronal loss [7]. This sequence of events suggests that microglia associated inflammatory events may be contributing to the cell loss.

This view is further strengthened by a recent study demonstrating that scrapie-infected mice brains show an increase in mitochondrial ROS production as well as an enhancement of NF- κ B activation followed by elevated levels of IL-1 α , IL-1 β , TNF- α , IL-6, and inducible NO synthase [59]. Neuronal vacuolation was present in the cortex, hippocampus and especially the thalamus and this was co-localized with NF- κ B activity. Thus it appears that glial activation and consequent cytokine and ROS production may be the factors responsible for neuronal loss.

Amyloid- β Protein

The A β PP is postulated to have neurotropic properties [58,68,72,96]. The role of the protein in protecting neurons is supported by studies that show that A β PP is upregulated during stressful conditions. The synthesis of A β PP in astrocytes increases after treatment with IL-1 [86] and transient ischemic changes modulate the expression of A β PP mRNA in rat cerebral cortex [58]. Amyloid- β is cleaved from A β PP and cultured cells derived from the CNS constitutively produce A β . The soluble forms of the protein are found in both normal and AD cerebrospinal fluid [32].

In many circumstances A β appears to act in a cytokine-like manner and promote inflammatory events. For example, A β and interferon- γ , as well as IL-1 β , activate microglia [27,47,70,90]. These reactive glial cells not only increase NO production [1,54], but they also activate complement [37,84,104]. A β has also been shown to exacerbates oxidative events in cells [8,93,101]. In a recent study, it was demonstrated that A β generated hydroxyl

radical formation was mediated by the reduction of Fe and Cu [52]. Parallel to the function of interleukines, the subsequent secretion of A β may be an integral part of an attempted innate immune response of the CNS against pathological conditions. If the disturbance is not resolved, the chronic production of these potentially harmful substances can then lead to neurodegeneration.

Copper has been shown to bind to A β PP which then reduces the metal from Cu (II) to the ROS generating Cu (I) [75]. This is analogous to the binding of copper to the Prp protein [26]. The addition of antioxidants such as SOD and catalase did not inhibit this ability of the A β PP protein to bind and reduce copper [74]. Copper has been shown to cause aggregation of A β and this increases the toxicity of the protein in neuronal cell cultures [53]. Different fragments of the amyloid- β peptide also reduce copper and generate hydrogen peroxide and it was observed that A β 1–42 is more toxic than A β 1–40. This was associated with the copper reduction capability of the peptides [52]. The generation of hydrogen peroxide occurred at catalytic amounts of Cu and thus these reactions may be occurring *in vivo*. NF- κ B is activated by hydrogen peroxide and copper present as a particulate in air pollution can activate this pro-inflammatory transcription factor [57].

Recent studies suggest that the aggregatory effect of metals depends critically on pH. Cu(II) induced aggregation of A β _{1–40} when the pH was lowered from 7.4 to 6.8. This phenomenon was not common to other metals tested [3]. A mildly acidic environment, together with increased Zn(II) and Cu(II), are common features of inflammation and this could explain the rapid deposition of A β following head injury as well as inflammation associated with increased oxidative damage due to microglial-derived peroxynitrite.

Both A β and aluminum sulfate are able to stabilize ferrous iron, which is the active form of the metal involved in the promotion of hydroxyl radical formation by the Fenton reaction [110]. In isolated systems Al potentiates iron promoted oxidative stress [10] and in the presence of the metal, iron can enhance ROS formation in protein-free liposomes [11]. Aluminum seems able to promote the pro-oxidant potential of iron rather than possessing any intrinsic ROS-generating capacity. It is possible that parallel to PrP and A β , colloidal Al may bind Fe and thus modulate the generation of ROS.

The Potential Role of Colloidal Aluminum in Neurodegeneration

Aluminum exposure and its relevance to neurodegeneration has been the subject of much controversy in the last few decades. In 1976, Alfrey implicated Al as the agent responsible for the outbreak of a dialysis encephalopathy syndrome [4] and in a later study it was demonstrated that the symptoms completely disappear after cessation of oral aluminum intake [87]. This Al-induced encephalopathy has also been reported in patients who have undergone bladder irrigation with 1% alum [81]. Residing in districts with high Al concentrations in the municipal drinking water has been linked to an increased risk of developing AD [69]. However, although it has been demonstrated that Al levels are increased in the brains of AD patients some studies repudiate this and thus the issue remains controversial [9,36,109].

Aluminum has been shown to exacerbate ROS formation *in vivo* and *in vitro*. Intraperitoneal injection of aluminum gluconate increased the rate of ROS formation in cortical tissue [12] and the brain of rats treated with aluminum lactate for 4 weeks showed an increase in lipid peroxidation and a significant decrease in antioxidants [55]. Al increased ROS formation, accompanied with elevated mitochondrial activity and glutathione depletion, in glial but not neuronal cell lines [24]. Because Al salts, complexed to strong acids such as sulfate and chloride, have a strong tendency to

FIG. 1. Transition metal/particle surface complexes may initiate a common chronic inflammatory pathway culminating in neurodegeneration.

form colloidal particles as they age [33], it is possible that the aluminum sulfate-induced increase in ROS generation is due to a glial response to the extracellular aluminum particles.

Several studies demonstrate that aluminum can cause a systemic inflammatory response. Low doses of aluminum, present in parenteral nutrition formula, produced marked portal inflammation [42] and rats exposed to oral doses of Al show an increase in plasma α_1 globulins [30]. Aluminum sensitization can develop in some children vaccinated for diphtheria, tetanus, and pertussis and Al-induced inflammatory nodules are formed in adults revaccinated for hepatitis B because the vaccines contain aluminum hydroxide as an adjuvant [35]. Aluminum, upon entry, may also initiate an inflammatory cascade in the brain. Chronic Al lactate treatment of rabbits increases glial fibrillary acidic protein concentrations in the frontal cortex [111] and the metal also increases NF- κ B activation in human glioblastoma cells [25]. This transcription factor is known to bind to and promote the production of nitric oxide synthase [5] and aluminum can induce the production of NO by microglial cells when it is in an insoluble form [45].

ANTIOXIDANT ENZYMES AND PROTEINS INVOLVED IN TRANSITION METAL HOMEOSTASIS

The development of transgenic mouse models has permitted a range of studies that critically evaluate the extent to which selected biological processes affect cell viability. MnSOD "knock-out" mice, suffering a 50% drop in mitochondrial SOD activity but no reduction of CuZnSOD or GSH peroxidase activity, were found to exhibit increased oxidative damage to mitochondria [108]. In contrast, no damage to cytosolic proteins or to nuclear DNA was

observed. Analysis of homozygote "knock-outs" showed mitochondrial degeneration that could only be followed for the 2 weeks that these mice were alive [71]. These results suggest that decreases in MnSOD activity can explain increased oxidative damage in mitochondria and alterations in the essential function of this organelle.

Overexpression of human CuZnSOD in mice, resulting in a 10-fold higher level in both myocytes and endothelial cells, was able to quench a burst of superoxide (electron probe resonance [EPR] detection) and reduce functional damage following 30 min global ischemia [105]. These results suggest that superoxide is an important factor in protecting against postischemic injury. However, it is surprising that CuZnSOD "knock-out" mice show little if any neurodegenerative phenotype [21]. Nonetheless, it is apparent that decreases in CuZnSOD activity can lead to a perturbation of cellular antioxidant defense mechanisms and promote a pro-oxidant situation.

The CuZnSOD gene is associated with AD neuropathology, and levels of both MnSOD mRNA and CuZnSOD were found to increase in AD, whereas the total antioxidant status was decreased [66]. Similarly, the activity of CuZnSOD in erythrocytes and the lymphocyte mRNA level of MnSOD were significantly increased in AD patients while total plasma antioxidant capacity was diminished [41]. In the brain tissue of AD patients, the activity of catalase and SOD is decreased, followed by an increase in lipid peroxidation [67]. These changes in antioxidant defense mechanisms may be responsible for the increased levels of oxidatively damaged RNA and proteins in AD brains [76].

In the scrapie-infected hamster brain, there is an increase in the

activity of glutathione peroxidase and reductase while the activity of mitochondrial manganese SOD is decreased [31]. This change in mitochondrial antioxidant defense may be responsible for the low level of ATPase activity and the high level of lipid peroxidation found in the cerebral mitochondria of infected animals. In cell culture, prion protein (106–126) induces heme oxygenase-1 (HO-1) in astroglial cells but not neuronal cells [83]. Thus, prion proteins modulate antioxidant status of cells and this may, in part, play a role in the eventual neurodegeneration observed in prion-infected brains.

CuZnSOD Mutations and Familial ALS

An important breakthrough in our understanding of ALS comes from the finding that many of the familial cases (FALS) are associated with mutations in the CuZnSOD gene. The protein products of these mutations retain nearly identical SOD activity, but take on altered properties linked to oxidative stress, possibly involving a gain-of-function peroxidase activity [107]. Transgenic mice overproducing a human FALS CuZnSOD mutant display increases in protein carbonyls suggestive of increased hydroxyl radical production or lipoxidation-derived radicals [2]. Using *in vivo* microdialysis, increased hydroxyl radical production in the striatum was seen for mice overexpressing the mutant enzyme relative to mice overexpressing the wild type human enzyme [13]. The hypothesis that mutant SOD-induced neurodegeneration is associated with disturbances in neuronal free radical homeostasis is further supported by observations made on several neuronal cell cultures expressing the mutant SOD [46]. However, the mechanism underlying the link between the SOD mutations and oxidative stress indicators appears not to be simply increased production of hydroxyl radicals, since no increases in HO• adducts are seen *in vitro* for the Gly93Ala and Ala4Val mutants relative to wild-type enzyme [95].

Recent studies have shown that the mutant and wild-type SODs differ neither in the rates of superoxide dismutation, nor in another enzyme diagnostic, namely H₂O₂-mediated inactivation [48,62]. Structural analysis of the mutant enzymes revealed that in some cases there were alterations such as subunit asymmetry. This suggested aberrant copper-mediated redox chemistry stemming from decreases in the tight folding and thus a more open “active site” [34,51]. Thus, the gain-of-function activity may reflect a “rechanneling” of the enzyme to oxidize biomolecules that normally would never gain access to the SOD oxidative half-reaction. Alternatively, the mutant SOD may possess weakened affinities for zinc or copper leading to a fraction of enzyme with abnormal activity [95] or oxidative reactions associated with leakage of copper [48]. For the predominant, sporadic form of ALS, an imbalance in trace metal ions, possibly tied to increased oxidative stress, has been considered for some time. Recent studies provide evidence for decreases in copper in the cerebrospinal fluid and serum as well as increases in manganese in serum of diseased individuals compared to age-matched control [56].

CONCLUSION

Over the past few years, there has been an increasing awareness of the seminal role that redox-active transition metals play in a variety of neurodegenerative diseases. The physico-chemical properties of aggregated prion proteins, fibrillar β A, and colloidal aluminum are very compatible with the promotion of oxidative stress by providing a complexing surface for redox-active metals and thus allowing activation of pro-oxidant events. These properties are also appropriate for inducing ineffectual glial responses. Failure to disperse these xenobiotic inclusions within the CNS can lead to chronic inflammatory responses and increased ROS pro-

duction that ultimately involve neurons and impair their function (Fig. 1). Presently, the stage is set to critically examine the importance of therapeutic modalities such as antioxidants, chelating agents and non-steroidal anti-inflammatory drugs in the treatment of neurodegenerative disorders.

ACKNOWLEDGEMENTS

The work conducted in the laboratory of the authors was supported by grants from the National Institute of Health and the Alzheimer's Association.

REFERENCES

1. Akama, K. T.; Albanese, C.; Pestell, R. G.; Van Eldik, L. J. Amyloid β peptide stimulates nitric oxide production in astrocytes through an NF- κ B-dependent mechanism. *Proc. Natl. Acad. Sci. USA* 95:5795–5800; 1998.
2. Andrus, P. K.; Fleck, T. J.; Gurney, M. E.; Hall, E. D. Protein oxidative damage in a transgenic mouse model of familial amyotrophic lateral sclerosis. *J. Neurochem.* 71:2041–2048; 1998.
3. Atwood, C. S.; Moir, R. D.; Huang, X.; Scarpa, R. C.; Bacarra, N. M. E.; Romano, D. M.; Hartshorn, M. A.; Tanzi, R. E.; Bush, A. I. Dramatic aggregation of Alzheimer A _{β} by Cu(II) is induced by conditions representing physiological acidosis. *J. Biol. Chem.* 273:12817–12826; 1998.
4. Alfrey, A. C.; leGendre, G. R.; Kaehny, W. D. The dialysis encephalopathy syndrome. Possible aluminum intoxication. *N. Engl. J. Med.* 294:184–188; 1976.
5. Baeuerle, P. A.; Henkel, T. Function and activation of NF- κ B in the immune system. *Annu. Rev. Immunol.* 12:141–179; 1994.
6. Beshgetoor, D.; Hambidge, M. Clinical conditions altering copper metabolism in humans. *Am. J. Clin. Nutr.* 67:1017S–1021S; 1998.
7. Betmouni, S.; Perry, V. H.; Gordon, J. L. Evidence for an early inflammatory response in the central nervous system of mice with scrapie. *Lett. Neurosci.* 74:1–5; 1996.
8. Bianca, V. D.; Dusi, S.; Bianchini, E.; Pra, I. D.; Rossi, F. β -Amyloid activates O₂^{•−} forming NADPH oxidase in microglia, monocytes, and neutrophils. *J. Biol. Chem.* 274:15493–15499; 1999.
9. Bjertness, E.; Candy, J. M.; Torvik, A.; Ince, P.; McArthur, F.; Taylor, G. A.; Johansen, S. W.; Alexander, J.; Gronnesby, J. K.; Bakkevig, L. S.; Edvardson, J. A. Content of brain aluminum is not elevated in Alzheimer disease. *Alzheimer Dis. Assoc. Disord.* 10:171–174; 1996.
10. Bondy, S. C.; Kirshtein, S. The promotion of iron-induced generation of reactive oxygen species in nerve tissue by aluminum. *Mol. Chem. Neuropathol.* 27:185–194; 1996.
11. Bondy, S. C.; Guo-Ross, S. X.; Pien, J. Mechanisms underlying the aluminum-induced potentiation of the pro-oxidant properties of transition metals. *Neurotoxicology* 19:65–72; 1998.
12. Bondy, S. C.; Ali, S. F.; Guo-Ross, S. Aluminum but not iron treatment induces pro-oxidant events in the rat brain. *Mol. Chem. Neuropathol.* 34:219–232; 1998.
13. Bogdanov, M. B.; Ramos, L. E.; Xu, Z.; Beal, M. F. Elevated “hydroxyl radical” generation *in vivo* in an animal model of amyotrophic lateral sclerosis. *J. Neurochem.* 71:1321–1324; 1998.
14. Brown, D. R.; Schmidt, B.; Kretzschmar, H. A. Role of microglia and host prion protein in neurotoxicity of a prion protein fragment. *Nature* 380:345–347; 1996.
15. Brown, D. R.; Schulz-Schaeffer, W. J.; Schmidt, B.; Kretzschmar, H. A. Prion protein-deficient cells show altered response to oxidative stress due to decreased SOD-1 activity. *Exp. Neurol.* 146:104–112; 1997.
16. Brown, D. R.; Besinger, A. Prion protein expression and superoxide dismutase activity. *Biochem. J.* 334:423–429; 1998.
17. Brown, D. R.; Besinger, A.; Herms, J. W.; Kretzschmar, H. A. Microglial expression of the prion protein. *Neuroreport* 9:1425–1429; 1998.
18. Brown, D. R.; Schmidt, B.; Kretzschmar, H. A. Effects of copper on survival of prion protein knockout neurons and glia. *J. Neurochem.* 70:1686–1693; 1998.

19. Brown, D. R.; Wong, B. S.; Hafiz, F.; Clive, C.; Haswell, S. J.; Jones, I. M. Normal prion protein has an activity like that of superoxide dismutase. *Biochem. J.* 344:1–5; 1999.
20. Brown, D. R. Prion protein peptide neurotoxicity can be mediated by astrocytes. *J. Neurochem.* 73:1105–1113; 1999.
21. Bruijn, L. I.; Houseweart, M. K.; Kato, S.; Anderson, K. L.; Anderson, S. D.; Ohama, E.; Reaume, A. G.; Scott, R. W.; Cleveland, D. W. Aggregation and motor neuron toxicity of an ALS-linked SOD1 mutant independent from wild-type SOD1. *Science* 281:1851–1854; 1998.
22. Busciglio, J.; Yankner, B. A. Apoptosis and increased generation of reactive oxygen species in Down's syndrome neurons in vitro. *Nature* 378:776–779; 1995.
23. Cadman, E. D.; Witte, D. G.; Lee, C. M. Regulation of the release of interleukin-6 from human astrocytoma cells. *J. Neurochem.* 63:980–987; 1994.
24. Campbell, A.; Prasad, K. N.; Bondy, S. C. Aluminum induced oxidative events in cell lines: Glioma are more responsive than neuroblastoma. *Free Radic. Biol. Med.* 26:1166–1171; 1999.
25. Campbell, A.; Yang, E.; Hamai, D.; Bondy, S. C. Aluminum promotes reactive oxygen species formation and NF- κ B activation in human glioblastoma cells. (Submitted)
26. Capellari, S.; Zaidi, S. I. A.; Urig, C. B.; Perry, G.; Smith, M. A.; Petersen, R. B. Prion protein glycosylation is sensitive to redox changes. *J. Biol. Chem.* 274:34846–34850; 1999.
27. Casamenti, F.; Prosperi, C.; Scali, C.; Giovannelli, L.; Colivicchi, M. A.; Faussone-Pellegrini, M. S.; Pepeu, G. Interleukin-1 β activates forebrain glial cells and increases nitric oxide production and cortical glutamate and GABA release in vivo: Implications for Alzheimer's disease. *Neuroscience* 91:831–842; 1999.
28. Castellani, R. J.; Smith, M. A.; Nunomura, A.; Harris, P. L. R.; Perry, G. Is increased redox-active iron in Alzheimer disease a failure of the copper-binding protein ceruloplasmin? *Free Radic. Biol. Med.* 26:1508–1512; 1999.
29. Castelnau, P. A.; Garrett, R. S.; Palinski, W.; Witztum, J. L.; Campbell, I. L.; Powell, H. C. Abnormal iron deposition associated with lipid peroxidation in transgenic mice expressing interleukin-6 in the brain. *J. Neuropathol. Exp. Neurol.* 57:268–282; 1998.
30. Cherroret, G.; Capolaghi, B.; Hutin, M. F.; Burnel, D.; Desor, D.; Lehr, P. R. Effects of postnatal aluminum exposure on biological parameters in the plasma. *Toxicol. Lett.* 78:119–125; 1995.
31. Choi, S. I.; Ju, W. K.; Choi, E. K.; Kim, J.; Lea, H. Z.; Carp, R. I.; Wisniewski, H. M.; Kim, Y. S. Mitochondrial dysfunction induced by oxidative stress in the brains of hamsters infected with the 263 K scrapie agent. *Acta Neuropathol.* 96:279–286; 1998.
32. Chung, H.; Brazil, M. I.; Soe, T.; Maxfield, F. R. Uptake, degradation, and release of fibrillar and soluble forms of Alzheimer's amyloid β -peptide by microglial cells. *J. Biol. Chem.* 274:32301–32308; 1999.
33. Corain, B.; Bombi, G. G.; Tapparo, A.; Perazzolo, M.; Zatta, P. Aluminum toxicity and metal speciation: Established data and open questions. *Coord. Chem. Rev.* 149:11–22; 1996.
34. Corson, L. B.; Strain, J. J.; Culotta, V. C.; Cleveland, D. W. Chaperone-facilitated copper binding is a property common to several classes of familial amyotrophic lateral sclerosis-linked superoxide dismutase mutants. *Proc. Natl. Acad. Sci. USA* 95:6361–6366; 1998.
35. Cosnes, A.; Flechet, M. L.; Revuz, J. Inflammatory nodular reactions after hepatitis B vaccination due to aluminum sensitization. *Contact Dermatitis* 23:65–67; 1990.
36. Crapper, D. R.; Krishnan, S. S.; Dalton, A. J. Brain aluminum distribution in Alzheimer's disease and experimental neurofibrillary degeneration. *Science* 180:511–513; 1973.
37. Cribbs, D. H.; Velazquez, P.; Soreghan, B.; Glabe, C. G.; Tenner, A. J. Complement activation by cross-linked truncated and chimeric full-length β -amyloid. *Neuroreport* 8:3457–3462; 1997.
38. Cuajungco, M. P.; Goldstein, L. E.; Nunomura, A.; Smith, M. A.; Lim, J. T.; Atwood, C. S.; Huang, X.; Farrag, Y. W.; Perry, G.; Bush, A. I. Evidence that the β -amyloid plaques of Alzheimer's disease represent the redox-silencing and entombment of A β by zinc. *J. Biol. Chem.* 275:19439–19442; 2000.
39. Cullen, K. M. Perivascular astrocytes within Alzheimer's disease plaques. *Neuroreport* 8:1961–1966; 1997.
40. David, J. P.; Ghazali, F.; Fallet-Bianco, C.; Watzet, A.; Delaire, S.; Boniface, B.; Di Menza, C.; Delacourte, A. Glial reaction in the hippocampal formation is highly concentrated with aging in human brain. *Neurosci. Lett.* 235:53–56; 1997.
41. De Leo, M. E.; Borrello, S.; Passantino, M.; Palazzotti, B.; Mordente, A.; Daniele, A.; Flippini, V.; Galeotti, T.; Masullo, C. Oxidative stress and overexpression of manganese superoxide dismutase in patients with Alzheimer's disease. *Neurosci. Lett.* 250:173–176; 1998.
42. Demircan, M.; Ergun, O.; Coker, C.; Yilmaz, F.; Avanoğlu, S.; Ozok, G. Aluminum in total parenteral nutrition solutions produces portal inflammation in rats. *J. Pediatr. Gastroenterol. Nutr.* 26:274–278; 1998.
43. Dunn, C. J. Cytokines as mediators of chronic inflammatory disease. In: Kimball, E. S., ed. *Cytokines and inflammation*. Boca Raton: CRC Press; 1991:1–35.
44. Fillebeen, C.; Dehouck, B.; Benaissa, M.; Dhennin-Duthille, I.; Cecchelli, R.; Pierce, A. Tumor necrosis factor- α increases lactoferrin transcytosis through the blood-brain barrier. *J. Neurochem.* 73:2491–2500; 1999.
45. Garrel, C.; Lafond, J. L.; Guiraud, P.; Faure, P.; Favier, A. Induction of production of nitric oxide in microglial cells by insoluble form of aluminum. *Ann. NY Acad. Sci.* 738:455–461; 1994.
46. Ghadge, G. D.; Lee, J. P.; Bindokas, V. P.; Jordan, J.; Ma, L.; Miller, R. J.; Roos, R. P. Mutant superoxide dismutase-1-linked familial amyotrophic lateral sclerosis: Molecular mechanisms of neuronal death and protection. *J. Neurosci.* 17:8756–8766; 1998.
47. Gitter, B. D.; Cox, L. M.; Rydel, R. E.; May, P. C. Amyloid β peptide potentiates cytokine secretion by interleukin-1 β -activated human astrocytoma cells. *Proc. Natl. Acad. Sci. USA* 92:10738–10741; 1995.
48. Goto, J. J.; Gralla, E. B.; Valentine, J. S.; Cabelli, D. E. Reactions of hydrogen peroxide with familial amyotrophic lateral sclerosis mutant human copper-zinc superoxide dismutases studies by pulse radiolysis. *J. Biol. Chem.* 273:30104–30109; 1998.
49. Griffin, W. S.; Sheng, J. G.; Royston, M. C.; Gentleman, S. M.; McKenzie, J. E.; Graham, D. I.; Roberts, G. W.; Mrak, R. E. Glial-neuronal interactions in Alzheimer's disease: The potential role of a 'cytokine cycle' in disease progression. *Brain Pathol.* 8:65–72; 1998.
50. Harris, Z. L.; Klomp, L. W.; Gitlin, J. D. Aceruloplasminemia: An inherited neurodegenerative disease with impairment of iron homeostasis. *Am. J. Clin. Nutr.* 67:S972–S977; 1998.
51. Hart, P. J.; Liu, H.; Pellegrini, M.; Nersissian, A. M.; Gralla, E. B.; Valentine, J. S.; Eisenberg, D. Subunit asymmetry in the three-dimensional structure of a human CuZnSOD mutant found in familial amyotrophic lateral sclerosis. *Protein Sci.* 7:545–555; 1998.
52. Huang, X.; Atwood, C. S.; Hartshorn, M. A.; Multhaup, G.; Goldstein, L. E.; Scarpa, R. C.; Cuajungco, M. P.; Gray, D. N.; Lim, J.; Moir, R. D.; Tanzi, R. E.; Bush, A. I. The A β peptide of Alzheimer's disease directly produces hydrogen peroxide through metal ion reduction. *Biochemistry* 38:7609–7616; 1999.
53. Huang, X.; Cuajungco, M. P.; Atwood, C. S.; Hartshorn, M. A.; Tyndall, J. D. A.; Hanson, G. R.; Stokes, K. C.; Leopold, M.; Multhaup, G.; Goldstein, L. E.; Scarpa, R. C.; Saunders, A. J.; Lim, J.; Moir, R. D.; Glabe, C.; Bowden, E. F.; Masters, C. L.; Fairlie, D. P.; Tanzi, R. E.; Bush, A. I. Cu(II) potentiation of Alzheimer's A β neurotoxicity. *J. Biol. Chem.* 274:37111–37116; 1999.
54. Hu, J.; Van Eldik, L. J. Glial-derived proteins activate cultured astrocytes and enhance beta amyloid-induced glial activation. *Brain Res.* 842:46–54; 1999.
55. Julka, D.; Gill, K. D. Effect of aluminum on regional brain antioxidant defense status in Wistar rats. *Res. Exp. Med.* 196:187–194; 1996.
56. Kapaki, E.; Zournas, C.; Kanas, G.; Zambelis, T.; Kakami, A.; Papageorgiou, C. Essential trace element alterations in amyotrophic lateral sclerosis. *J. Neurol. Sci.* 147:171–175; 1997.
57. Kennedy, T.; Ghio, A. J.; Reed, W. S.; Samet, J.; Zagorski, J.; Quay, J.; Carter, J.; Dailey, L.; Hoidal, J. R.; Devlin, R. B. Copper-dependent inflammation and nuclear factor- κ B activation by particulate air pollution. *Am. J. Respir.* 19:366–378; 1998.
58. Kim, H. S.; Lee, S. H.; Kim, S. S.; Kim, Y. K.; Jeong, S. J.; Ma, J.; Han, D. H.; Cho, B. K.; Suh, Y. H. Post-ischemic changes in the

- expression of Alzheimer's APP isoforms in rat cerebral cortex. *Neuroreport* 9:533-537; 1998.
59. Kim, J. I.; Ju, W. K.; Choi, J. H.; Kim, J.; Choi, E. K.; Carp, R. I.; Wisniewski, H. M.; Kim, Y. S. Expression of cytokine genes and increased nuclear factor-kappa B activity in the brains of scrapie-infected mice. *Mol. Brain Res.* 73:17-27; 1999.
60. Kim, S.; Ponka, P. Effects of interferon-gamma and lipopolysaccharide on macrophage iron metabolism are mediated by nitric oxide-induced degradation of iron regulatory protein 2. *J. Biol. Chem.* 275:6220-6226; 2000.
61. Linder, M. C.; Hazegh-Azam, M. Copper biochemistry and molecular biology. *Am. J. Clin. Nutr.* 63:797S-811S; 1996.
62. Liochev, S. I.; Chen, L. L.; Hallewell, R. A.; Fridovich, I. The familial amyotrophic lateral sclerosis-associated amino acid substitutions E100G, G93A, and G93R do not influence the rate of inactivation of copper- and zinc-containing superoxide dismutase by H_2O_2 . *Arch. Biochem. Biophys.* 352:237-239; 1998.
63. Loeffler, D. A.; LeWitt, P. A.; Juneau, P. L.; Sima, A. A. F.; Nguyen, H. U.; Demaggio, A. J.; Brickman, C. M.; Brewer, G. J.; Dick, R. D.; Troyer, M. D.; Kanaley, L. Increased regional brain concentration of ceruloplasmin in neurodegenerative disorders. *Brain Res.* 738:265-274; 1996.
64. Lovell, M. A.; Robertson, J. D.; Teesdale, W. J.; Campbell, J. L.; Markesbery, W. R. Copper, iron and zinc in Alzheimer's disease senile plaques. *J. Neurol. Sci.* 158:47-52; 1998.
65. Lukiw, W. J.; Bazan, N. G. Strong nuclear factor - κ B-DNA binding parallels cyclooxygenase-2 gene transcription in aging and in sporadic Alzheimer's disease superior temporal lobe neocortex. *J. Neurosci. Res.* 53:583-592; 1998.
66. Mantyh, P. W.; Ghilardi, J. R.; Rogers, S.; DeMaster, E.; Allen, C. J.; Stimson, E. R.; Maggio, J. E. Aluminum, iron, and zinc ions promote aggregation of physiological concentrations of beta-amyloid peptide. *J. Neurochem.* 61:1171-1174; 1993.
67. Marcus, D. L.; Thomas, C.; Rodriguez, C.; Simberloff, K.; Tsai, J. S.; Strafaci, J. A.; Freedman, M. L. Increased peroxidation and reduced antioxidant enzyme activity in Alzheimer's disease. *Exp. Neurol.* 150:40-44; 1998.
68. Masliah, E.; Westland, C. E.; Rockenstein, E. M.; Abraham, C. R.; Mallory, M.; Veinberg, I.; Sheldon, E.; Mucke, L. Amyloid precursor protein protect neurons of transgenic mice against acute and chronic excitotoxic injuries in vivo. *Neuroscience* 78:135-146; 1997.
69. McLachlan, D. R. C.; Bergeron, C.; Smith, J. E.; Boomer, D.; Rifat, S. L. Risk for neuropathologically confirmed Alzheimer's disease and residual aluminum in municipal drinking water employing weighted residential histories. *Neurology* 46:401-405; 1996.
70. Meda, L.; Cassatella, M. A.; Szendrei, G. I.; Otvos, L.; Baron, P.; Villalba, M.; Ferrari, D.; Rossi, F. Activation of microglial cells by β -amyloid protein and interferon- γ . *Nature* 374:647-650; 1995.
71. Melov, S.; Schneider, J. A.; Day, B. J.; Hinerfeld, D.; Coskun, P.; Mirra, S. S.; Crapo, J. D.; Wallace, D. C. A novel neurological phenotype in mice lacking mitochondrial manganese superoxide dismutase. *Nat. Genet.* 18:159-163; 1998.
72. Mezziane, H.; Dodaart, J. C.; Mathes, C.; Little, S.; Clemens, J.; Paul, S. M.; Ungerer, A. Memory-enhancing effects of secreted forms of the β -amyloid precursor protein in normal and amnesic mice. *Proc. Natl. Acad. Sci. USA* 95:12683-12689; 1998.
73. Mezzetti, A.; Pierdomenico, S. D.; Costantini, F.; Romano, F.; De Cesare, D.; Cuccurullo, F.; Imbastro, T.; Riario-Sforza, G.; Di Giacomo, F.; Zuliani, G.; Fellin, R. Copper/zinc ratio and systemic oxidant load: Effect of aging and aging-related degenerative diseases. *Free Radic. Biol. Med.* 25:676-681; 1998.
74. Multhaup, G.; Schlicksupp, A.; Hesse, L.; Behr, D.; Ruppert, T.; Masters, C. L.; Beyreuther, K. The amyloid precursor protein of Alzheimer's disease in the reduction of copper (II) to copper (I). *Science* 271:1406-1409; 1996.
75. Multhaup, G.; Ruppert, T.; Schlicksupp, A.; Hesse, L.; Bill, E.; Pipkorn, R.; Masters, C. L.; Beyreuther, K. Copper-binding amyloid precursor protein undergoes a site-specific fragmentation in the reduction of hydrogen peroxide. *Biochemistry* 37:7224-7230; 1998.
76. Nunomura, A.; Perry, G.; Pappolla, M. A.; Wade, R.; Hirai, K.; Chiba, S.; Smith, M. A. RNA oxidation is a prominent feature of vulnerable neurons in Alzheimer's disease. *J. Neurosci.* 19:1959-1964; 1999.
77. Nunomura, A.; Perry, G.; Hirai, K.; Aliev, G.; Takeda, A.; Chiba, S.; Smith, M. A. Neuronal RNA oxidation in Alzheimer's disease and Down's syndrome. *Ann. NY Acad. Sci.* 893:362-364; 1999.
78. Nunomura, A.; Perry, G.; Pappolla, M. A.; Freidland, R. P.; Hirai, K.; Chiba, S.; Smith, M. A. Neuronal oxidative stress precedes amyloid- β deposition in Down syndrome. *J. Neuropathol. Exp. Neurol.* 59:1011-1017; 2000.
79. Odetti, P.; Angelini, G.; Dapino, D.; Zaccaro, D.; Garibaldi, S.; Dagna-Bricarelli, F.; Piombo, G.; Perry, G.; Smith, M.; Traverso, N.; Tabaton, M. Early glycoxidation damage in brains from Down's syndrome. *Biochem. Biophys. Res. Comm.* 243:849-851; 1998.
80. Pezzella, A.; d'Ischia, M.; Napolitano, A.; Misuraca, G.; Protta, G. Iron-mediated generation of the neurotoxin 6-hydroxydopamine quinone by reaction of fatty acid hydroperoxides with dopamine: A possible contributory mechanism for neuronal degeneration in Parkinson's disease. *J. Med. Chem.* 40:2211-2216; 1997.
81. Phelps, K. R.; Naylor, K.; Brien, T. P.; Wilbur, H.; Haqqie, S. S. Encephalopathy after bladder irrigation with alum: Case report and literature review. *Am. J. Med. Sci.* 318:181-185; 1999.
82. Riederer, P.; Sofic, E.; Rausch, W. D.; Schmidt, B.; Reynolds, G. P.; Jellinger, K.; Youdim, M. B. H. Transition metals, ferritin, glutathione, and ascorbic acid in parkinsonian brains. *J. Neurochem.* 52:515-520; 1989.
83. Rizzardini, M.; Chiesa, R.; Angeretti, N.; Lucca, E.; Salmona, M.; Forloni, G.; Cantoni, L. Prion protein fragment 106-126 differentially induces heme oxygenase-1 mRNA in cultured neurons and astroglial cells. *J. Neurochem.* 68:715-720; 1997.
84. Rogers, J.; Cooper, N. R.; Webster, S.; Schultz, J.; McGeer, P. L.; Styren, S. D.; Civin, W. H.; Brachova, L.; Bradt, B.; Ward, P.; Lieberburg, I. Complement activation by β -amyloid in Alzheimer's disease. *Proc. Natl. Acad. Sci. USA* 89:10016-10020; 1992.
85. Rogers, J.; O'Barr, S. Inflammatory mediators in Alzheimer's disease. In: Wasco, W.; Tanzi, R. E., eds. *Molecular mechanisms of dementia*. Totowa, NJ: Humana Press Inc.; 1997:177-198.
86. Rogers, J. T.; Leiter, L. M.; McPhee, J.; Cahill, C. M.; Zhan, S. S.; Potter, H.; Nilsson, L. N. G. Translation of the Alzheimer's amyloid precursor protein mRNA is up-regulated by interleukin-1 through 5'-untranslated region sequences. *J. Biol. Chem.* 274:6421-6431; 1999.
87. Russo, L. S.; Beale, G.; Sandroni, S.; Ballinger, W. E. Aluminum intoxication in undialysed adults with chronic renal failure. *J. Neurol. Neurosurg. Psychiatry* 55:697-700; 1991.
88. Sayre, L. M.; Zelasko, D. A.; Harris, P. L.; Perry, G.; Salomon, R. G.; Smith, M. A. 4-Hydroxynonenal-derived advanced lipid peroxidation end products are increased in Alzheimer's disease. *J. Neurochem.* 68:2092-2097; 1974.
89. Sayre, L. M.; Perry, G.; Harris, P. L. R.; Liu, Y.; Schubert, K. A.; Smith, M. A. In situ oxidative catalysis by neurofibrillary tangles and senile plaques in Alzheimer's disease: A central role for bound transition metals. *J. Neurochem.* 74:270-279; 2000.
90. Scali, C.; Prosperi, C.; Giovannelli, L.; Bianchi, L.; Pepeu, G.; Casamenti, F. Beta (1-40) amyloid peptide injection into the nucleus basalis of rats induces microglia reaction and enhances cortical gamma-aminobutyric acid release in vivo. *Brain Res.* 831:319-321; 1999.
91. Schipper, H. M. Astrocytes, brain aging, and neurodegeneration. *Neurobiol. Aging* 17:467-480; 1996.
92. Schipper, H. M.; Vininsky, R.; Brull, R.; Small, L.; Brawer, J. R. Astrocyte mitochondria: A substrate for iron deposition in the aging rat substantia nigra. *Exp. Neurol.* 152:188-196; 1998.
93. Schubert, D.; Behl, C.; Lesley, R.; Brack, A.; Dargusch, R.; Sagara, Y.; Kimura, H. Amyloid peptides are toxic via a common oxidative mechanism. *Proc. Natl. Acad. Sci. USA* 92:1989-1993.
94. Silei, V.; Fabrizi, C.; Venturini, G.; Salmona, M.; Bugiani, O.; Tagliavini, F.; Lauro, G. M. Activation of microglial cells by PrP and β -amyloid fragments raises intracellular calcium through L-type voltage sensitive calcium channels. *Brain Res.* 818:168-170; 1999.
95. Singh, R. J.; Karoui, H.; Gunther, M. R.; Beckman, J. S.; Mason, R. P.; Kalyanaram, B. Reexamination of the mechanism of hydroxyl radical adducts formed from the reaction between familial

- amyotrophic lateral sclerosis-associated Cu,Zn superoxide dismutase mutants and H₂O₂. *Proc. Natl. Acad. Sci. USA* 95:6675–6680; 1998.
96. Small, D. H. The role of the amyloid protein precursor (APP) in Alzheimer's disease. Does the normal function of APP explain the topography of neurodegeneration? *Neurosci. Res.* 23:795–806; 1998.
 97. Smith, M. A.; Taneda, S.; Richey, P. L.; Miyata, S.; Yan, S. D.; Stern, D.; Sayre, L. M.; Monnier, V. M.; Perry, G. Advanced Maillard reaction end products are associated with Alzheimer disease pathology. *Proc. Natl. Acad. Sci. USA* 91:5710–5714; 1994.
 98. Smith, M. A.; Sayre, L. M.; Anderson, V. E.; Beal, M. F.; Kowall, N. Oxidative damage in Alzheimer's. *Nature* 382:120–121; 1996.
 99. Smith, M. A.; Harris, P. L. R.; Sayre, L. M.; Beckman, J. S.; Perry, G. Widespread peroxynitrite-mediated damage in Alzheimer's disease. *J. Neurosci.* 17:2653–2657; 1997.
 100. Smith, M. A.; Harris, P. L. R.; Sayre, L. M.; Perry, G. Iron accumulation in Alzheimer disease is a source of redox-generated free radicals. *Proc. Natl. Acad. Sci. USA* 94:9866–9868; 1997.
 101. Smith, M. A.; Hirai, K.; Hsiao, K.; Pappolla, M. A.; Harris, P. L. R.; Siedlak, S. L.; Tabaton, M.; Perry, G. Amyloid- β deposition in Alzheimer transgenic mice is associated with oxidative stress. *J. Neurochem.* 70:2212–2215; 1998.
 102. Smith, M. A.; Wehr, K.; Harris, P. L. R.; Siedlak, S. L.; Connor, J. R.; Perry, G. Abnormal localization of iron regulatory protein (IRP) in Alzheimer disease. *Brain Res.* 788:232–236; 1998.
 103. Streit, W. J.; Walter, S. A.; Pennell, N. A. Reactive microgliosis. *Prog. Neurobiol.* 57:563–581; 1999.
 104. Veerhuis, R.; Janssen, I.; De Groot, C. J.; Van Muiswinkel, F. L.; Hack, C. E.; Eikelenboom, P. Cytokines associated with amyloid plaques in Alzheimer's disease brain stimulate human glial and neuronal cell cultures to secrete early complement proteins, but not C1-inhibitor. *Exp. Neurol.* 160:289–299; 1999.
 105. Wang, P.; Chen, H.; Qin, H.; Sankarapandi, S.; Becher, M. W.; Wong, P. C.; Zweier, J. L. Overexpression of human copper, zinc-superoxide dismutase (SOD1) prevents postischemic injury. *Proc. Natl. Acad. Sci. USA* 95:4556–4560; 1998.
 106. Wegrzyniak, B. H.; Dobrzanski, P.; Stoehr, J. D.; Wenk, G. L. Chronic neuroinflammation in rats reproduces components of the neurobiology of Alzheimer's disease. *Brain Res.* 780:294–303; 1998.
 107. Wiedau-Pazus, M.; Goto, J. J.; Rabizadeh, S.; Gralla, E. B.; Roe, J. A.; Lee, M. K.; Valentine, J. S.; Bredesen, D. E. Altered reactivity of superoxide dismutase in familial amyotrophic lateral sclerosis. *Science* 271:515–518; 1996.
 108. Williams, M. D.; Van Remmen, H.; Conrad, C. C.; Huang, T. T.; Epstein, C. J.; Richardson, A. Increased oxidative damage is correlated to altered mitochondrial function in heterozygous manganese superoxide dismutase knockout mice. *J. Biol. Chem.* 273:28510–28515.
 109. Xu, N.; Majidi, V.; Markesbery, W. R.; Ehmann, W. D. Brain aluminum in Alzheimer's disease using an improved GFAAS method. *Neurotoxicology* 13:735–743; 1992.
 110. Yang, E. Y.; Guo-Ross, S. X.; Bondy, S. C. The stabilization of ferrous iron by a toxic β -amyloid fragment and by an aluminum salt. *Brain Res.* 839:221–226; 1999.
 111. Yokel, R. A.; O'Callaghan, J. P. An aluminum-induced increase in GFAP is attenuated by some chelators. *Neurotoxicol. Teratol.* 20:55–60; 1997.
 112. Yoshida, T.; Tanaka, M.; Sotomatsu, A.; Hirai, S.; Okamoto, K. Activated microglia cause iron-dependent lipid peroxidation in the presence of ferritin. *Neuroreport* 9:1929–1933; 1998.
 113. Zhou, Y.; Richardson, J. S.; Mombourquette, M. J.; Weil, J. A. Free radical formation in autopsy samples of Alzheimer's and control cortex. *Neurosci. Lett.* 195:89–92; 1995.