UC San Diego

UC San Diego Previously Published Works

Title

Association of the selected dimensions of eudaimonic well-being with healthy survival to 85 years of age in older women.

Permalink

<https://escholarship.org/uc/item/50n4r2jp>

Journal

International psychogeriatrics, 26(12)

ISSN

1041-6102

Authors

Zaslavsky, Oleg
Rillamas-Sun, Eileen
Woods, Nancy Fugate
et al.

Publication Date

2014-12-01

DOI

10.1017/s1041610214001768

Peer reviewed

Association of the selected dimensions of eudaimonic well-being with healthy survival to 85 years of age in older women

Oleg Zaslavsky,¹ Eileen Rillamas-Sun,² Nancy Fugate Woods,³
Barbara B. Cochrane,^{2,3} Marcia L. Stefanick,⁴ Hilary Tindle,⁵ Lesley F. Tinker²
and Andrea Z. LaCroix^{2,6}

¹Faculty of Health Science and Social Welfare, University of Haifa, Haifa, Israel

²Public Health Sciences, Fred Hutchinson Cancer Research Center, Seattle, Washington, USA

³School of Nursing, University of Washington, Seattle, Washington, USA

⁴Stanford Prevention Research Center, School of Medicine, Stanford University, Palo Alto, California, USA

⁵Division of General Internal Medicine, University of Pittsburgh, Pennsylvania, USA

⁶Division of Epidemiology, School of Medicine, University of California, San Diego, California, USA

ABSTRACT

Background: Aspects of eudaimonic well-being, such as personal growth (PG) and purpose in life (PL), have been highlighted as important to older adults' health. We investigated the relationship of PG and PL with patterns of survival to the age of 85 years and older.

Methods: The sample included 8,880 women from the Women's Health Initiative cohort who reached 85 years of age by December 1, 2013, and for whom data on the PG and PL constructs were available. Women were classified into mutually exclusive outcomes: Healthy, Prevalent, Incident, Disabled, and Deceased. PG and PL were each assessed using a modified seven-item measure derived from the Psychological Well-Being scale.

Results: Women were most commonly classified as Healthy (38.2%, $n = 3,395$), followed by Incident (24.4%, $n = 2,163$), Disabled (19.0%, $n = 1,685$), Prevalent (14.3%, $n = 1,273$), and Deceased (4.1%, $n = 364$). Women with low PL and PG levels were more likely to have prevalent mobility disability and disease or incident death before the age of 85 years. Specifically, those who reported low levels of PG and PL had a 2.1- and 3.6-fold higher risk, respectively, of death.

Conclusions: These findings indicate that even among the oldest old, experience of purposeful life engagement and continuing PG may contribute to better health outcomes.

Key words: women's health, aging, well-being, functioning, purpose in life, personal growth

Introduction

A substantive body of research has highlighted the strong association between negative psychological states such as depression, anxiety, and psychological distress, and an increased risk of death or poor health among community-dwelling older adults (Cuijpers and Smit, 2002; Suls and Bunde, 2005). In contrast, there has been comparatively little

research linking positive psychological well-being with longer and high-functioning survival.

The concept of well-being refers to perceived optimal psychological functioning, with two schools of thought distinguishing between hedonic and eudaimonic approaches (Ryan and Deci, 2001). The hedonic approach focuses on happiness and defines well-being in terms of pleasure attainment and pain avoidance, whereas the eudaimonic approach focuses on self-actualization in accordance with one's daimon, or true self (Ryan and Deci, 2001). Prior research examined hedonic well-being (often termed subjective well-being) and life expectancy (Diener and Chan, 2011) and found that higher subjective well-being influences health and longevity in healthy populations. However,

Correspondence should be addressed to: Oleg Zaslavsky, PhD, Faculty of Health Sciences and Social Welfare, Department of Nursing, University of Haifa, Haifa 31905, Israel. Phone: +972-547832536; Fax: +972-48288017. Email: oleg.zaslavsky@fulbrightmail.org. Received 22 May 2014; revision requested 28 Jun 2014; revised version received 23 Jul 2014; accepted 24 Jul 2014. First published online 27 August 2014.

research on the impact of eudaimonic well-being (EW) on longevity and late-life functioning among community-dwelling older adults and especially among those in the oldest-old age group remains scarce.

Few observational studies suggest a link between life expectancy and eudaimonic markers of human flourishing such as sense of purpose in life (PL) and personal growth (PG; Vaillant, 2002; Buettner, 2008). In-depth interviews with high-functioning centenarians around the globe consistently demonstrated that PL and perceived PG were among the factors associated with the healthiest survival into advanced age (Buettner, 2008). Similar conclusions resonated in US-based Long-Life Observational Study of Vaillant (2002), which also underscored the importance of PL and perceived PG as longevity-associated strategies. Moreover, “meaningfulness” was one of the defining elements embedded in the Sense of Coherence construct, which in turn was linked to better psychological and physical health, and to longer survival in aged cohorts (Antonovsky, 1987; Wiesmann and Hannich, 2008; Lundman *et al.*, 2010; Nilsson *et al.*, 2010).

While there are many scales for assessing aspects of psychological well-being, Ryff’s scale is ideal for studying EW because it operationalizes the distinct eudaimonic dimensions of PG, PL, environmental mastery, autonomy, positive relations with others, and self-acceptance (Ryff and Keyes, 1995). Prior research examining possible biological underpinnings of eudaimonic constructs showed that individuals with higher levels of PG and PL demonstrated a slower increase in daily salivary cortisol, which is suggestive of favorable stress response, than those with lower levels of PG and PL (Ryff *et al.*, 2006). Similarly, PG levels were negatively correlated with total/high-density lipoprotein cholesterol ratio, considered protective against cardiovascular disease risks. Each of these biological markers has been implicated in the aging process (Varadhan *et al.*, 2008; Rahilly-Tierney *et al.*, 2011), suggesting a plausible physiological sequence.

Previous literature suggested that the PG and PL dimensions play an important role in later-life physical and emotional health in aged cohorts (Wood and Joseph, 2010; Friedman and Ryff, 2012); however, the association of these psychological factors in octo- and nonagenarian populations is limited. In this context, we examined Women’s Health Initiative (WHI) data to empirically estimate the association of PG and PL constructs with healthy survival to the age of 85 years and older. This study will be the first to

examine the effects of psychological well-being on living healthy into latest-life adulthood.

An examination of body mass index (BMI) as a predictor of healthy survival with and without disease and disability among women aged 85 years and older in the WHI was recently published (Rillamas-Sun *et al.*, 2013). Using outcomes modeled after this body of work, we aimed to understand how psychological factors relate to healthy survival and longevity in older women. Specifically, our objectives were:

1. To describe the distribution of PG and PL dimensions of EW in women at least 85 years of age.
2. To examine the relationship of PG and PL with healthy survival to at least 85 years of age without major chronic disease and mobility disability.

We hypothesized that individual measures of EW would vary widely in this older population of women. We also anticipated that higher levels of PG and PL would be associated with lower odds of disease, disability, and death.

Methods

Study sample

The multiethnic WHI Clinical Trial and Observational Study cohort of 161,808 postmenopausal women, aged 50 to 79 years at study entry, was enrolled between 1993 and 1998 across 40 US clinical centers. Details of the scientific rationale, study design, eligibility requirements, data collection protocols, and baseline characteristics of the cohort were previously published (Anderson *et al.*, 2003). In 2004–2005, WHI participants from the main study were invited to participate in the first extension study (2005–2010). A second extension study began in 2010, with completion expected in 2015. Mailings were sent annually to participants, who completed forms pertaining to incidence of clinical outcomes and changes in functioning. Adjudication by a study physician of cerebrovascular disease, cardiovascular disease, cancer (excluding non-melanoma skin cancer), and hip fractures were completed for women from the main WHI study and a subsample of women from the extension studies. In 2012, as part of the second extension study, participants completed a Lifestyle Questionnaire, which included a modified version of Ryff’s PG and PL scales (Ryff and Keyes, 1995). This analysis included women who were 85 years of age and older by the time of the latest clinical outcome evaluation on December 1, 2013, and who provided complete data on the PG and PL measures.

Outcome measures

The outcome measures were five mutually exclusive classifications modeled after the study by Rillamas-Sun *et al.* (2013):

1. Healthy: lived to the age of 85 years without major chronic disease or mobility disability.
2. Prevalent: lived to the age of 85 years and had at least one chronic disease at WHI baseline, but did not develop new disease or a mobility disability during study follow-up.
3. Incident: lived to the age of 85 years and developed at least one chronic disease, but did not develop mobility disability during study follow-up.
4. Disabled: lived to the age of 85 years and developed mobility disability, regardless of major chronic disease status.
5. Deceased: died before December 1, 2013, but after the collection of PG and PL measures in the second extension study.

Chronic diseases included cardiovascular disease, cerebrovascular disease, cancer (excluding non-melanoma skin cancer), type-2 diabetes, and hip fracture. These conditions were selected because they are highly prevalent in older women and greatly increase a woman's risk of death and disability. Disabling conditions, such as arthritis, were captured through identification of impaired mobility. Except for diabetes, which was identified by a self-report of doctor's diagnosis and medication use, all diseases were physician-adjudicated throughout most years of study follow-up. Participants' death records were centrally adjudicated by study physicians using hospital records, autopsy reports, and death certificates. Periodic checks of the National Death Index for all participants, including those lost to follow-up, were performed; the latest update was available through December 1, 2013. Mobility disability was defined as self-reported difficulty walking one block or climbing one flight of stairs or a self-reported need to use crutches, a walker, or a wheelchair to move on a level surface. As in Rillamas-Sun *et al.* (2013), we also excluded women who did not provide health information within 18 months of their 85th birth year and those who reported a mobility disability at baseline.

Eudaimonic constructs

PG and PL constructs were each assessed in 2012 using a modified seven-item measure derived from the Psychological Well-Being scale of Ryff and Keyes (1995; see Table S1, available as supplementary material attached to the electronic version of this paper at www.journals.cambridge.org/jid_IPG). Although each of the original subscales included 20 items designed

to measure these constructs, several shortened versions (ranging from 3 to 14 questions) have been developed and validated psychometrically (Ryff and Keyes, 1995; Abbott *et al.*, 2006). Participants rated their level of agreement with each item (e.g. "I have a sense of direction and purpose in life") in the past week on a five-point scale that ranged from 0 to 4. The construct scores were calculated by summing the score for the seven items, resulting in a possible range of 0 to 28 for each measure. Higher scores were suggestive of higher levels of PG and PL. The oldest-old women with at least five of the seven items non-missing for the PG ($n = 9,028$) and PL ($n = 8,988$) constructs were included in the analysis. The modified measures used for this analysis had acceptable reliability for the PG and PL scales (Cronbach's α 's of 0.73 and 0.65, respectively). These two scales' scores were divided into quartiles for analysis, based on their observed distribution.

Covariates

Data on demographic variables (race/ethnicity, family income, highest level of education) were obtained by self-report at the WHI baseline during the period 1993–1998. Self-reported frequency of alcohol consumption (Never; <1 drink per week; 1–4 drinks per week; >4 drinks per week) and total minutes of recreational physical activity (including walking, mild, moderate, and strenuous exercise) were collected in 2012. Additional variables, such as baseline obesity measured by BMI categories, depressive symptoms assessed using the Burnam adaptation of the Center for Epidemiological Studies Depression Scale short form (Burnam *et al.*, 1988), self-reported binary indicators (Y/N) of living alone, and baseline smoking status (*Never smoker*; *Past Smoker*; *Current smoker*) were also included as adjustment variables. Depressive symptoms and living status were collected concurrently with EW measures. Existing literature suggested that the above-mentioned factors are plausible confounders of the relationship between EW and healthy survival (Boyle *et al.*, 2009; 2010).

Statistical approach

We examined the distribution of the EW measures and other covariates by the outcome groups using descriptive statistics. Because the outcome was composed of five nominal categories, the association of the PG and PL constructs was examined using multinomial logistic regression. Multinomial logistic regression uses maximum likelihood to estimate the log odds of being in a given outcome category (Prevalent, Incident, Disabled, or

Deceased) compared with the reference category (Healthy), allowing for separate slope estimates (Long, 1997). The adjusted prediction of quartiles of PG and PL by the outcome categories and their 95% CIs were calculated and reported. The partially adjusted models controlled for age, race, and education (see Table S2, available as supplementary material attached to the electronic version of this paper at www.journals.cambridge.org/jid_IPG). The fully adjusted models controlled for age, race, education, income, alcohol use, depression, obesity, physical activity levels, smoking, and whether the participant lived alone. Tests for trend were performed by contrasting log-likelihood ratios of the models that differed by having either factorial or linear terms of PG or PL measures included. All *p* values reported are for two-tailed tests. The statistical software STATA, version 11.2 (StataCorp, College Station, TX), was used for the statistical analysis.

Results

Of the 9,986 women who were at least 85 years of age by December 1, 2013, 889 (8.9%) participants had missing information on EW measures and 196 (2.0%) had missing information on study outcomes, leaving a total analytical sample of 8,880 women. Age distribution was different between those with missingness on PG and PL measures and those with complete information (average age 88.5 (SD 2.5) and 88.0 (SD 2.3), $p < 0.001$ in excluded and analytical samples respectively) and similar between excluded and retained women with respect to study outcomes (average age 88.3 (SD 2.4) and 88.0 (SD 2.3), $p = 0.11$ in excluded and analytical samples respectively). A total of 3,395 women (38.2%) were classified as healthy, and 1,273 (14.3%) lived to the age of 85 years and maintained the same number of selected chronic conditions as at the WHI baseline. Of the remaining women, 2,163 (24.4%) developed at least one of the selected chronic conditions during study follow-up and before their 85th birth year; 1,685 (19.0%) lived to the age of 85 years and developed mobility disability without regard to disease status; and 364 (4.1%) died.

Table 1 displays univariate characteristics of the five outcome categories. Assessment of the EW indicators across outcome categories revealed a gradient decline in percentage of individuals reporting the highest quartile of PG and PL, ranging from 41.7% and 43.3% for those in the Healthy category to 2.4% and 1.9% for those in the Deceased group. In addition, women in the Healthy group were more likely to have higher levels of PG

and PL, whereas women in the Disabled group were more likely to have lower levels of PG and PL.

Table 2 shows the fully adjusted odds ratios (ORs) for the EW-related variables. Although not all ORs were statistically significant, consistently higher ORs for the Incident, Disabled, or Deceased categories versus the Healthy group were found in those reporting low levels of PG and PL as compared with higher levels. For example, those who reported the lowest levels of PG had 1.44 (95% CI = 1.17–1.76) odds of being in the Disabled rather than the Healthy category compared with their counterparts in the highest PG quartile. Likewise, women with the lowest PL levels had a 1.37 (95% CI = 1.14–1.64) odds of being in the Incident category relative to being in the Healthy group. Furthermore, women with the lowest levels of PL had a 3.6-fold higher risk of being in the Deceased versus the Healthy category, and, similarly, those who reported the lowest levels of PG had a 2.1-fold higher risk of dying relative to the reference category. Interestingly, ORs for Prevalent versus Healthy survival were not strongly associated with the EW levels, although a gradual increase in the odds of being in the Prevalent category could be clearly observed when contrasting higher with lower PL levels.

Figures 1 and 2 illustrate the estimated probabilities of an average study participant being in the Healthy, Prevalent, Incident, Disabled, or Deceased categories at different levels of PG and PL constructs when holding all other variables in the model at their means. The vertical lines are likelihood-ratio-based 95% CIs for the predicted probabilities. For PG, the probability of being in the Disabled or Deceased categories increases almost linearly as PG levels decline (p for linear trend < 0.001). A similar pattern was observed in the Incident and Disabled categories for the PL construct, although in the latter model log-likelihood estimates did not support the linear-trend assumption. Finally, the probabilities of being in the Healthy category increased directly as higher levels of the EW indicators were considered.

Discussion

We examined the association of PG and PL with survival to 85 years of age and older among almost 9,000 women, and found that these constructs of EW were independently associated with late-life risk of death and higher odds of having a mobility disability and major chronic morbidity before the age of 85 years. Older women who reported lower levels of PG and PL had respectively higher risk of death in the next two years of study

Table 1. Baseline and concurrent characteristics of the 8,880 participants 85 years of age and older in the Women's Health Initiative (WHI) study, by survival phenotype category

CHARACTERISTIC	HEALTHY ^a	PREVALENT ^b	INCIDENT ^c	DISABLED ^d	DECEASED ^e
N (%) of women	3,395 (38.2)	1,273 (14.3)	2,163 (24.4)	1,685 (19.0)	364 (4.1)
Age—mean (SD; years),	88.2 (2.3)	88.3 (2.4)	87.8 (2.2)	87.5 (1.8)	88.9 (2.7)
Personal growth—mean (SD; score)	19.6 (4.9)	19.4 (5.0)	19.3 (4.8)	18.5 (5.0)	17.8 (5.2)
PERSONAL GROWTH, N (%; n = 8,834)					
QRT1 (0–15)	765 (34.5)	300 (13.5)	524 (23.6)	504 (22.7)	125 (5.6)
QRT2 (16–19)	969 (37.5)	349 (13.5)	649 (25.1)	506 (19.6)	114 (4.4)
QRT3 (20–23)	785 (39.6)	304 (15.3)	481 (24.3)	339 (17.1)	74 (3.7)
QRT4 (24–28)	853 (41.7)	315 (15.4)	501 (24.5)	328 (16.0)	49 (2.4)
Purpose in Life—mean (SD; score)	17.8 (4.7)	17.5 (4.7)	17.4 (4.7)	16.6 (4.8)	15.4 (4.7)
PURPOSE IN LIFE, N (%; n = 8,794)					
QRT1 (0–14)	868 (33.4)	345 (13.3)	613 (23.6)	596 (23.0)	175 (6.7)
QRT2 (15–17)	818 (37.7)	314 (14.5)	564 (26.0)	405 (18.7)	68 (3.1)
QRT3 (18–21)	900 (40.1)	335 (14.9)	538 (24.0)	388 (17.3)	82 (3.7)
QRT4 (22–28)	773 (43.3)	268 (15.0)	427 (23.9)	284 (15.9)	33 (1.9)
WHITE RACE ^f , N (%; n = 8,855)					
Yes	3,155 (38.3)	1,174 (14.3)	2,005 (24.3)	1,550 (18.8)	305 (4.3)
No	233 (37.8)	95 (15.4)	151 (24.5)	129 (20.9)	8 (1.3)
EDUCATION ^f , N (%; n = 8,843)					
< High school	106 (35.3)	23 (7.7)	68 (22.7)	89 (29.7)	14 (4.7)
High school/GED	519 (36.6)	188 (13.2)	359 (25.3)	292 (20.6)	62 (4.4)
Some college	1,284 (37.5)	470 (13.7)	826 (24.1)	686 (20.1)	156 (4.6)
College grad	1,471 (39.8)	585 (15.8)	904 (24.4)	610 (16.5)	131 (3.5)
FAMILY INCOME ^f , N (%; n = 8,307)					
<\$20,000	498 (34.2)	189 (13.0)	348 (23.9)	358 (24.6)	63 (4.3)
\$20,000 to <\$50,000	2,215 (38.4)	850 (14.7)	1,404 (24.4)	1,066 (18.5)	230 (4.0)
\$50,000 or more	463 (42.6)	155 (14.3)	275 (25.3)	152 (14.0)	41 (3.8)
LIVE ALONE ^g , N (%; n = 8,057)					
Yes	1,911 (39.4)	697 (14.4)	1,206 (24.9)	874 (18.0)	163 (3.4)
No	1,169 (36.5)	464 (14.5)	753 (23.5)	657 (20.5)	163 (5.1)
SMOKING BEHAVIOR ^f , N (%; n = 8,768)					
Never smoker	2,029 (39.3)	746 (14.5)	1,260 (24.4)	926 (18.0)	199 (3.9)
Past smoker	1,270 (37.1)	502 (14.7)	831 (24.3)	674 (19.7)	148 (4.3)
Current smoker	49 (26.8)	21 (11.5)	50 (27.3)	54 (29.5)	9 (4.9)
ALCOHOL INTAKE ^g , N (%; n = 8,657)					
Never drinker	1,137 (34.0)	441 (13.2)	798 (23.8)	811 (24.2)	161 (4.8)
<1 drink/week	1,066 (38.8)	392 (14.3)	675 (24.6)	500 (18.2)	112 (4.1)
1–4 drinks/week	565 (41.9)	209 (15.5)	335 (24.9)	197 (14.6)	41 (3.0)
>4 drinks/week	587 (44.8)	217 (16.6)	294 (24.2)	138 (11.3)	41 (3.4)
DEPRESSION ^g , N (%; n = 8,880)					
Yes	448 (35.3)	175 (13.8)	299 (23.6)	281 (22.2)	65 (5.3)
No	2,947 (38.7)	1,098 (14.4)	1,864 (24.5)	1,404 (18.4)	299 (3.9)
BODY MASS INDEX CATEGORY ^f , N (%; n = 8,807)					
Underweight	34 (43.0)	11 (13.9)	22 (27.8)	9 (11.4)	3 (3.8)
Normal weight	1,528 (43.6)	566 (16.1)	865 (24.7)	417 (11.9)	132 (3.8)
Overweight	1,292 (38.3)	475 (14.1)	845 (25.0)	626 (18.6)	137 (4.1)
Obese	513 (27.8)	207 (11.2)	417 (22.6)	622 (33.7)	806 (4.7)
MINUTES OF PHYSICAL ACTIVITY ^g , N (%; n = 8,880)					
<Age-adjusted median	1,346 (30.9)	600 (13.8)	1,020 (23.4)	1,140 (26.1)	255 (5.9)
>Age-adjusted median	2,049 (45.3)	673 (14.9)	1,143 (25.3)	545 (12.1)	109 (2.4)

^aNo disease and no mobility disability.^bBaseline disease but not incident disease or mobility disability.^cIncident disease but not mobility disability.^dIncident mobility disability with or without disease.^eDid not survive to December 1, 2013.^fMeasure was collected at the WHI baseline during the period 1993–1998.^gMeasure was collected in 2012.

GED = tests of general educational development; QRT = quartile.

Table 2. Adjusted odds ratios (ORs) and 95% CIs of quartiles of "Personal Growth" ($N = 7,704$) and "Purpose in Life" ($N = 7,675$) constructs for Prevalent, Incident, Disabled, or Deceased categories at 85 years of age in Women's Health Initiative study participants^a

VARIABLE	PREVALENT ^b				INCIDENT ^c				DISABLED ^d				DECEASED ^e			
	OR (95% CI)	<i>p</i>	VALUE	<i>p</i> FOR TREND	OR (95% CI)	<i>p</i>	VALUE	<i>p</i> FOR TREND	OR (95% CI)	<i>p</i>	VALUE	<i>p</i> FOR TREND	OR (95% CI)	<i>p</i>	VALUE	<i>p</i> FOR TREND
PERSONAL GROWTH																
Continuous	1.00 (0.98–1.01)	0.47			0.99 (0.98–1.00)	0.07			0.98 (0.96–0.99)	0.001			0.95 (0.93–0.98)	<0.001		
score ^f																
QRT1	1.04 (0.85–1.27)	0.67			1.18 (0.98–1.42)	0.07			1.44 (1.17–1.76)	<0.001			2.10 (1.42–3.08)	<0.001		
QRT2	0.92 (0.76–1.13)	0.44			1.17 (0.99–1.38)	0.07			1.13 (0.93–1.38)	0.21			1.55 (1.06–2.28)	0.03		
QRT3	1.06 (0.86–1.31)	0.56		0.74	1.05 (0.88–1.26)	0.57	0.05		1.12 (0.91–1.38)	0.29		<0.001	1.33 (0.88–2.01)	0.18		<0.001
QRT4	Reference				Reference				Reference				Reference			
PURPOSE IN LIFE																
Continuous	0.98 (0.97–1.00)	0.14			0.98 (0.96–0.99)	<0.001			0.96 (0.94–0.97)	<0.001			0.92 (0.89–0.95)	<0.001		
score ^f																
QRT1	1.17 (0.95–1.45)	0.15			1.37 (1.14–1.64)	0.001			1.65 (1.35–2.02)	<0.001			3.55 (2.31–5.44)	<0.001		
QRT2	1.13 (0.91–1.40)	0.25			1.31 (1.09–1.56)	0.004			1.22 (0.99–1.50)	0.06			1.54 (0.96–2.48)	0.07		
QRT3	1.11 (0.91–1.37)	0.31		NA ^g	1.11 (0.93–1.32)	0.26		NA ^g	1.18 (0.96–1.44)	0.13		NA ^g	1.85 (1.17–2.92)	0.008		NA ^g
QRT4	Reference				Reference				Reference				Reference			

Note. Results are from multinomial logistic regression adjusted for age, race, education, income, alcohol use, depression, body mass index, physical activity, smoking status, and whether the participant lives alone. The higher value of the quartiles indicates higher personal growth and purpose in life.

^aReference category is those living to age 85 years or older with no disease and no mobility disability.

^bBaseline disease but not incident disease or mobility disability.

^cIncident disease but not mobility disability.

^dIncident mobility disability with or without disease.

^eDid not survive to December 1, 2013.

^fHigher score indicates more favorable status.

^gNull results for the test for linear trend.

QRT = quartile; OR = odds ratio.

Figure 1. (Colour online) Estimated adjusted probabilities of Healthy, Prevalent, Incident, Disabled, or Deceased categories of survival phenotype for 88-year-old woman among Women's Health Initiative participants ($N = 7,704$) by quartiles of Personal Growth construct. Categories denote: survived without major chronic disease and without disability (Healthy); survived with one or more major chronic diseases at baseline, but without disease or disability (Prevalent); survived and developed one or more major chronic morbidities, but not disability, during study follow-up (Incident); survived and developed mobility disability with or without disease (Disabled); and did not survive (Deceased). Results are from multinomial logistic regression adjusted for race, education, income, alcohol use, depression, body mass index, physical activity, smoking, and whether the participant lives alone. The higher value of the quartiles indicates higher Personal Growth. Error bars indicate likelihood-ratio-based 95% CIs.

follow-up than women who had higher levels of PG and PL. Furthermore, late-life survivors reporting lower levels of PG and PL had higher odds of having a mobility disability at 85 years of age or older compared with their counterparts who had a more favorably perceived psychological well-being. Among women who lived to late adulthood without mobility disability, having lower perceived levels of the selected EW constructs was associated with higher odds of having developed a major chronic morbidity than with the process of living with chronic conditions from baseline. Conversely, those who did not escape mobility disability were more likely to report lower levels of PG or PL than their able-bodied and healthier counterparts. To our knowledge this is the first study to examine the association between selected dimensions of the EW and patterns of survival in the oldest-old age group. However, one important caveat needs to be taken into consideration, as the limited prospective nature of the study precludes conclusive statements on the

directionality of the effects. It is possible to assume that women who developed chronic morbidity or disability during years of follow-up also experienced a decline in PG and PL given that EW measures were collected on average 15.5 years after baseline assessments.

The PL construct underscores one's perception of having meaning, purpose, and direction in life, whereas PG reflects the extent to which respondents make use of their personal talents and potential (Ryff and Keyes, 1995). High scores on these measures in latest adulthood suggest that psychological flourishing is possible even in the face of advanced aging processes.

Observational and empirical evidence suggest that psychological factors of human flourishing in latest adulthood are related to high-functioning survival. In the widely publicized book *The Blue Zones*, Buettner (2008) presented evidence that having a strong sense of PL was one of the main themes shared by a relatively large

Figure 2. (Colour online) Estimated adjusted probabilities of Healthy, Prevalent, Incident, Disabled, or Deceased categories of survival phenotype for 88-year-old woman among Women's Health Initiative participants ($N = 7,675$) by quartiles of Purpose in Life construct. Categories denote: survived without major chronic disease and without disability (Healthy); survived with one or more major chronic diseases at baseline, but without disease or disability (Prevalent); survived and developed one or more major chronic morbidities, but not disability, during study follow-up (Incident); survived and developed mobility disability with or without disease (Disabled); and did not survive (Deceased). Results are from multinomial logistic regression adjusted for race, education, income, alcohol use, depression, body mass index, physical activity, smoking, and whether the participant lives alone. The higher value of the quartiles indicates higher Purpose in Life. Error bars indicate likelihood-ratio-based 95% CIs.

cohort of high-functioning centenarians around the globe. Echoing these findings, US-based Long-Life Observational Study of Vaillant (2002) also highlighted that perceived PG and PL were among the factors associated with the healthiest survival into advanced age. In recent years cross-sectional and prospective studies have consistently demonstrated an association between PL and a reduced risk of all-cause mortality (Boyle *et al.*, 2009), reduced risk of Alzheimer disease, and mild cognitive impairment (Boyle *et al.*, 2010), reduced risk of developing functional dependence (Boyle *et al.*, 2010), and reduced risk of frailty syndrome (Andrew *et al.*, 2012). Similar studies examining the effects of PG on clinical outcomes are surprisingly scarce.

At present, mechanisms for the association of PL and PG with healthier survival are unknown. Physiological mechanisms for the relationship between EW and health have been hypothesized, including a direct enhancement

of neuroendocrine, immune, and cardiovascular systems functioning (Ryff *et al.*, 2004; Steptoe *et al.*, 2005) or indirect effect on health by buffering the effects of stress (Pressman and Cohen, 2005). Biological studies have provided partial support for these assertions, demonstrating that some aspects of psychological well-being are associated with important disease biomarkers such as salivary cortisol, the proinflammatory cytokine interleukin-6, and high-density lipoprotein cholesterol (Ryff *et al.*, 2006; Friedman *et al.*, 2007). Given the well-described association of low-grade inflammation with poor health and poor functioning in older adults, it is plausible that the beneficial effects of the eudaimonic markers of human flourishing on late-life health might be mediated through a physiological milieu. Thus, pending further research, one cautious interpretation is that eudaimonic indicators of well-being such as PL and PG might promote late-life physical health through favorable alteration of neuroendocrine

signaling and an enhancement of stress-resistance mechanisms.

Engagement in health-promoting behaviors is another sensible explanation for the link between psychological well-being and having better health. It is plausible to assume that those oldest old who have a favorable perceived psychological outlook also have a higher proclivity for health-promoting behaviors. However, in the current study, accounting for alcohol consumption, physical activity, smoking, and BMI did not substantially affect the association of PG and PL with the outcomes, suggesting that other factors may be involved (see Table S2, available as supplementary material attached to the electronic version of this paper at www.journals.cambridge.org/jid_IPG).

Our findings bear an important public health message. Psychological well-being is an integrative factor of one's physical health. Although it is still unclear whether the perceived sense of one's psychological flourishing is amenable to intervention in late adulthood, our findings highlight the importance of comprehensive psychological well-being assessment as one of mortality- and morbidity-sensitive measures. In addition, our results indicate that boosting EW in even the oldest adults – for instance, through community volunteer programs such as the Experience Corps (Carlson *et al.*, 2008) – might engage them in activities likely to promote their sense of PL and their PG (Greenfield and Marks, 2004).

Although we identified a relationship between lower levels of PG and PL and the risk of dying within two years post follow-up, we acknowledge that temporal ambiguity limits our ability to estimate risk associations of EW on the mobility disability and incident chronic disease outcomes. Because the evaluation of incident disability and disease occurred prior to the assessment of the PG and PL measures, it is reasonable to believe that women who developed these conditions might be more likely to report lower EW measures later in life. Yet, several studies report that levels of within-person EW are relatively stable over time, especially in older populations (Ryff, 1991; Springer *et al.*, 2011). This suggests that the women with lower levels of PG and PL in our sample may likely have had these low levels prior to their development of disability or disease. Thus, we believe there is a relationship between lower PG and PL levels and the risk of developing a mobility disability or major chronic disease; however, confirmation of these relationships in a prospective study is needed. The validation is especially pertinent given that the above-mentioned within-person stability in EW markers has yet to be confirmed among frail older adults and those with incident disability.

This study limited the scope of psychological constructs to measures of EW, even though other constructs, such as optimism, social support, and hedonic or emotional well-being, that have been measured in the WHI cohort might influence the relationship of PG and PL to physical health. In addition, PG and PL were only two of six constructs constituting Ryff's Psychological Well-Being scale (Ryff and Keyes, 1995). Although each of the six constructs can be used independently, the availability of the complete scale would offer a more comprehensive psychological profile (Ryff, 2014). Next, older women with low levels of functioning and health were less likely than women with higher levels of functioning to complete self-reported questionnaires. Loss to follow-up may underestimate the proportion of older women at risk of being in unfavorable survival categories. However, rather than determining the prevalence of survival groups, the aim of this analysis was to describe the association of the selected dimension of EW with patterns of survival to 85 years of age in older women. Finally, this oldest-old WHI cohort had a higher frequency of college education attainment than older women in the general population and was relatively homogeneous in terms of race; therefore, the cohort might not be fully representative. However, our models included adjustment for common indicators of prosperity, such as education and income, and statistically significant associations were still observed.

The study strengths are an assessment of PG and PL for a large group of older women. We also minimized misclassification bias of the outcomes by completing a detailed central adjudication performed by a trained clinical staff. Last, our rich database permitted examination of a wide array of likely confounders on the association of selected dimensions of EW with healthy survival.

In sum, these results add to a growing literature on the potential salutary effects of positive psychological functioning, and they suggest that even among oldest-old adults, experiences of purposeful life engagement and continuing PG may contribute to better health outcomes. Future prospective studies, however, are needed to better understand the dynamics between positive psychological factors and patterns of survival into advanced age.

Conflict of interest

The WHI program is funded by the National Heart, Lung and Blood Institute; the National Institute of Health; and the US Department of Health and Human Services through Contracts

N01WH22110, 24152, 32100-2, 32105-6, 32108-9, 32111-13, 32115, 32118-32119, 32122, 42107-26, 42129-32, and 44221. The funding agency had no role in the design or conduct of the study; collection, management, analysis, or interpretation of the data; or preparation, review, or approval of the manuscript.

Description of authors' roles

Dr O. Zaslavsky, Dr E. Rillamas-Sun, Dr B.B. Cochrane, Dr N.F. Woods, and Dr A.Z. LaCroix had full access to all of the data used in the study, and take responsibility for the integrity of the data and accuracy of the data analysis. Dr O. Zaslavsky, Dr E. Rillamas-Sun, Dr N.F. Woods, and Dr A.Z. LaCroix were involved with the study concept, design, interpretation of data, and preparation of manuscript. Dr O. Zaslavsky and Dr E. Rillamas-Sun carried out the data analysis and visualization. H. Tindle, M.L. Stefanick, and L.F. Tinker were involved with the preparation of manuscript.

Supplementary material

To view supplementary material for this article, please visit <http://dx.doi.org/10.1017/S1041610214001768>

Acknowledgments

The authors thank the WHI investigators and staff for their dedication and the study participants for making the program possible. A listing of WHI investigators can be found at <https://cleo.whi.org/researchers/Documents%20Write%20a%20Paper/WHI%20Investigator%20Short%20List.pdf>.

References

- Abbott, R. A., Ploubidis, G. B., Huppert, F. A., Kuh, D., Wadsworth, M. E. and Croudace, T. J.** (2006). Psychometric evaluation and predictive validity of Ryff's psychological well-being items in a UK birth cohort sample of women. *Health and Quality of Life Outcomes*, 4, 76. doi:10.1186/1477-7525-4-76.
- Anderson, G. L. et al.** (2003). Implementation of the Women's Health Initiative study design. *Annals of Epidemiology*, 13 (Suppl. 9), S5–S17. doi:10.1016/S1047-2797(03)00043-7.
- Andrew, M. K., Fisk, J. D. and Rockwood, K.** (2012). Psychological well-being in relation to frailty: a frailty identity crisis? *International Psychogeriatrics*, 24, 1347–1353. doi:10.1017/S1041610212000269.

- Antonovsky, A.** (1987). *Unraveling the Mystery of Health: How People Manage Stress and Stay Well*. San Francisco, CA: Jossey-Bass.
- Boyle, P. A., Barnes, L. L., Buchman, A. S. and Bennett, D. A.** (2009). Purpose in life is associated with mortality among community-dwelling older persons. *Psychosomatic Medicine*, 71, 574–579. doi:10.1097/PSY.0b013e3181a5a7c0.
- Boyle, P. A., Buchman, A. S., Barnes, L. L. and Bennett, D. A.** (2010). Effect of a purpose in life on risk of incident Alzheimer disease and mild cognitive impairment in community-dwelling older persons. *Archives of General Psychiatry*, 67, 304–310. doi:10.1001/archgenpsychiatry.2009.208.
- Buettner, D.** (2008). *The Blue Zones*. Washington, DC: National Geographic Society.
- Burnam, M. A., Wells, K. B., Leake, B. and Landsverk, J.** (1988). Development of a brief screening instrument for detecting depressive disorders. *Medical Care*, 26, 775–789. doi:10.1097/00005650-198808000-00004.
- Carlson, M. C. et al.** (2008). Exploring the effects of an “everyday” activity program on executive function and memory in older adults: Experience Corps®. *Gerontologist*, 48, 793–801. doi:10.1093/geront/48.6.793.
- Cuijpers, P. and Smit, F.** (2002). Excess mortality in depression: a meta-analysis of community studies. *Journal of Affective Disorders*, 72, 227–236. doi:10.1016/S0165-0327(01)00413-X.
- Diener, E. and Chan, M. Y.** (2011). Happy people live longer: subjective well-being contributes to health and longevity. *Applied Psychology: Health and Well-Being*, 3, 1–43. doi:10.1111/j.1758-0854.2010.01045.x.
- Friedman, E. M., Hayney, M., Love, G. D., Singer, B. H. and Ryff, C. D.** (2007). Plasma interleukin-6 and soluble IL-6 receptors are associated with psychological well-being in aging women. *Health Psychology*, 26, 305–313. doi:10.1037/0278-6133.26.3.305.
- Friedman, E. M. and Ryff, C. D.** (2012). Living well with medical comorbidities: a biopsychosocial perspective. *Journals of Gerontology. Series B, Psychological Sciences and Social Sciences*, 67, 535–544. doi:10.1093/geronb/67.5.535.
- Greenfield, E. A. and Marks, N. F.** (2004). Formal volunteering as a protective factor for older adults' psychological well-being. *Journals of Gerontology. Series B, Psychological Sciences and Social Sciences*, 59, S258–S264. doi:10.1093/geronb/59.5.S258.
- Long, J. S.** (1997). *Regression Models for Categorical and Limited Dependent Variables*. Thousand Oaks, CA: Sage.
- Lundman, B. et al.** (2010). Sense of coherence (SOC) related to health and mortality among the very old: the Umea 85+ study. *Archives of Gerontology and Geriatrics*, 51, 329–332. doi:10.1016/j.archger.2010.01.013.
- Nilsson, K. W., Leppert, J., Simonsson, B. and Starrin, B.** (2010). Sense of coherence and psychological well-being: improvement with age. *Journal of Epidemiology and Community Health*, 64, 347–352. doi:10.1136/jech.2008.081174.
- Pressman, S. D. and Cohen, S.** (2005). Does positive affect influence health? *Psychological Bulletin*, 131, 925–971. doi:10.1037/0033-2909.131.6.925.
- Rahilly-Tierney, C. R., Spiro, A. III, Vokonas, P. and Gaziano, J. M.** (2011). Relation between high-density

- lipoprotein cholesterol and survival to age 85 years in men (from the VA normative aging study). *American Journal of Cardiology*, 107, 1173–1177. doi:10.1016/j.amjcard.2010.12.015.
- Rillamas-Sun, E. et al.** (2013). Obesity and late-age survival without major disease or disability in older women. *JAMA Internal Medicine*, 174, 98–106. doi:10.1001/jamainternmed.2013.12051.
- Ryan, R. M. and Deci, E. L.** (2001). On happiness and human potentials: a review of research on hedonic and eudaimonic well-being. *Annual Review of Psychology*, 52, 141–166. doi:10.1146/annurev.psych.52.1.141.
- Ryff, C. D.** (1991). Possible selves in adulthood and old age: a tale of shifting horizons. *Psychology and Aging*, 6, 286–295. doi:10.1037/0882-7974.6.2.286.
- Ryff, C. D.** (2014). Psychological well-being revisited: advances in the science and practice of eudaimonia. *Psychotherapy and Psychosomatics*, 83, 10–28. doi:10.1159/000353263.
- Ryff, C. D. and Keyes, C. L.** (1995). The structure of psychological well-being revisited. *Journal of Personality and Social Psychology*, 69, 719–727. doi:10.1037/0022-3514.69.4.719.
- Ryff, C. D., Singer, B. H. and Love, G. D.** (2004). Positive health: connecting well-being with biology. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 359, 1383–1394. doi:10.1098/rstb.2004.1521.
- Ryff, C. D. et al.** (2006). Psychological well-being and ill-being: do they have distinct or mirrored biological correlates? *Psychotherapy and Psychosomatics*, 75, 85–95. doi:10.1159/000090892.
- Springer, K. W., Pudrovska, T. and Hauser, R. M.** (2011). Does psychological well-being change with age? Longitudinal tests of age variations and further exploration of the multidimensionality of Ryff's model of psychological well-being. *Social Science Research*, 40, 392–398. doi:10.1016/j.ssresearch.2010.05.008.
- Steptoe, A., Wardle, J. and Marmot, M.** (2005). Positive affect and health-related neuroendocrine, cardiovascular, and inflammatory processes. *Proceedings of the National Academy of Sciences*, 102, 6508–6512. doi:10.1073/pnas.0409174102.
- Suls, J. and Bunde, J.** (2005). Anger, anxiety, and depression as risk factors for cardiovascular disease: the problems and implications of overlapping affective dispositions. *Psychological Bulletin*, 131, 260–300. doi:10.1037/0033-2909.131.2.260.
- Vaillant, G.** (2002). *Aging Well: Surprising Guideposts to a Happier Life from the Landmark Harvard Study of Adult Development*. Boston, MA: Little, Brown.
- Varadhan, R. et al.** (2008). Higher levels and blunted diurnal variation of cortisol in frail older women. *Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, 63, 190–195. doi:10.1093/gerona/63.2.190.
- Wiesmann, U. and Hannich, H. J.** (2008). A salutogenic view on subjective well-being in active elderly persons. *Aging & Mental Health*, 12, 56–65. doi:10.1080/13607860701365998.
- Wood, A. M. and Joseph, S.** (2010). The absence of positive psychological (eudemonic) well-being as a risk factor for depression: a ten year cohort study. *Journal of Affective Disorders*, 122, 213–217. doi:10.1016/j.jad.2009.06.032.