

Lawrence Berkeley National Laboratory

Recent Work

Title

Energy and structure of dithiophosphinate complexes with trivalent actinides and lanthanides: Computational and experimental studies

Permalink

<https://escholarship.org/uc/item/5127p60v>

Authors

Zhang, Zhiyong

Tian, Guoxin

Rao, Linfeng

Publication Date

2004-06-01

Energy and Structure of Dithiophosphate Complexes with Trivalent Actinides and Lanthanides: Computational and Experimental Studies

Zhiyong Zhang, Guoxin Tian, Linfeng Rao

Lawrence Berkeley National Laboratory, Berkeley, California, U.S.A.

Separation of trivalent actinides from lanthanides has been a challenging subject of study in recent years due to the demand in the treatment of nuclear wastes. Bis(2,4,4-trimethylpentyl)dithiophosphinic acid, a commercially available dithiophosphinic acid (DTP), has been shown to be highly selective in the solvent extraction of An(III) over Ln(III). Various attempts have been made to reveal the fundamental chemical principles underlying the selectivity. In this study, Density Functional Theory (DFT) calculations were performed on the complexation of a smaller DTP, $(\text{CH}_3)_2\text{P}(\text{S})\text{SH}$, with Am(III) and lighter Ln(III) (La^{3+} , Nd^{3+} and Eu^{3+}), using Stuttgart small core relativistic effective core potentials and including the spin-orbit effects. The energy and structure of the hydrated and unhydrated complexes were calculated and compared. For the trivalent lanthanides, the energy of monohydrated DTP complexes, $\text{Ln}(\text{DTP})_4 \cdot \text{H}_2\text{O}$, is lower than that of unhydrated complexes, $\text{Ln}(\text{DTP})_4$. However, for americium(III), the unhydrated complex, $\text{Am}(\text{DTP})_4$, has lower energy than the hydrated one. The computational results are consistent with experimental observations by spectroscopic techniques including FT-IR, TRLFS and ES/MS, suggesting that the difference in the degree of hydration between the Ln(III) and An(III) complexes contributes to the selectivity of DTP for An(III) over Ln(III) in solvent extraction.

This work was supported by the Director, Office of Science, Office of Basic Energy Sciences, Division of Chemical Sciences, under U.S. Department of Energy Contract No. DE-AC03-76SF0098 at the Lawrence Berkeley National Laboratory.