UC San Diego

UC San Diego Previously Published Works

Title

Theory-Guided Development of Fertility Care Implementation Strategies for Adolescent and Young Adult Cancer Survivors.

Permalink

<https://escholarship.org/uc/item/513745sf>

Journal

Journal of adolescent and young adult oncology, 10(5)

ISSN

2156-5333

Authors

Dornisch, Anna
Yang, Emily H
Gruspe, Jamie
et al.

Publication Date

2021-10-01

DOI

10.1089/jayao.2020.0156

Peer reviewed

Theory-guided development of fertility care implementation strategies for adolescent and young adult cancer survivors

Running title: Designing oncofertility care implementation

Anna Dornisch, MAS,^{1*} Emily H. Yang,^{1*} Jamie Gruspe, MPH,¹ Erin R. Roesch, MD,^{1#} Paula Aristizabal, MD MAS,^{1,2} Greg A. Aarons, PhD,³ Teresa Helsten, MD,¹ Michelle B. Takemoto, PhD MPH,⁴ Sally A. D. Romero, PhD MPH,¹ Bonnie N. Kaiser, PhD MPH,⁵ H. Irene Su, MD MSCE^{1,6}

¹Moore's Cancer Center, University of California, San Diego

²Rady Children's Hospital – San Diego

³Department of Psychiatry, University of California, San Diego

⁴ServiceNow, Inc.

⁵Department of Anthropology and Global Health Program, University of California, San Diego

⁶Department of Obstetrics, Gynecology and Reproductive Sciences, University of California, San Diego

*AD and EHY contributed equally to this manuscript

#Cleveland Clinic

Corresponding Author:

H. Irene Su, MD MSCE

Department of Obstetrics, Gynecology and Reproductive Sciences and Moore's Cancer Center

University of California, San Diego

3855 Health Sciences Drive

Dept 0901

La Jolla, CA 92093-0901

858-822-5986

hisu@health.ucsd.edu

Funding: NIH UL1TR001442, NIH TL1TR001443

Author contribution: Design (ER, PA, GAA, MBT, TH, HIS), data collection (AD, EHY, JG, ER, PA, MBT, HIS), data analysis (AD, EHY, JG, SADR, BNK, HIS), manuscript preparation (AD, EHY, HIS), manuscript review (JG, ER, PA, GAA, MBT, TH, SADR, BNK)

Abstract

Purpose: Oncofertility care at cancer diagnosis remains under-implemented across oncology and fertility care settings, with limited tools to scale up effective implementation strategies. Using implementation science theory, we systematically assessed factors that influence oncofertility care implementation and map scalable strategies, particularly electronic health record (EHR)-enabled ones, that fit adult and pediatric oncology care contexts.

Methods: Using purposeful sampling, we recruited healthcare providers and female, reproductive-aged survivors of adolescent and young adult cancers (AYA survivors) from a comprehensive cancer center and a freestanding children's hospital to semi-structured interviews and focus groups. Using thematic analysis combining inductive codes with deductive codes using the Consolidated Framework for Implementation Research (CFIR), we characterized barriers and facilitators to care and designed responsive strategies. Two coders independently coded each transcript.

Results: We recruited 19 oncology and fertility providers and 9 cancer survivors. We identified barriers and facilitators to oncofertility care in the CFIR domains of individual, inner setting, outer setting, and process, allowing us to conceptualize oncofertility care to encompass three core components (screening, referral, and fertility preservation counseling) and map five strategies to these components that fit an adult and a children's context and bridge oncology and fertility practices. The strategies were screening using a best practice advisory, referral order, telehealth fertility counseling, provider audit and feedback, and provider education. All but provider education were EHR tools with embedded efficiencies.

Conclusions: An implementation science approach systematically assessed oncofertility care and mapped strategies to provide a theory-based approach and scalable EHR tools to support wider dissemination.

Key words: oncofertility, implementation science, CFIR, fertility preservation

Background

Unmet reproductive health care needs are highly prevalent among the nearly 400,000 reproductive-aged, female survivors of AYA cancers (AYA survivors) in the U.S.¹ They undergo treatments such as radiation, chemotherapy, surgery, and/or endocrine therapy, which may adversely impact future fertility.^{2,3} AYA survivors often want to have their own families; infertility from cancer treatment significantly impairs quality of life.^{4,5} Because oncofertility care – fertility counseling and preservation procedures – at cancer diagnosis and post-treatment can decrease the risk of infertility,^{6,7} clinical guidelines from oncology and fertility societies recommend that oncologists discuss infertility risk with patients and offer fertility preservation options or referrals to reproductive specialists throughout the cancer continuum.^{6,8-10}

Despite clinical guidelines, uptake of fertility counseling at cancer diagnosis remains highly variable. In the adult setting, 2015-2019 Quality of Oncology Practice Initiative data showed that 44% of patients younger than age 50 were counseled on reproductive risks across 400 practices.¹¹ Continually moderate implementation of fertility counseling is attributed to heterogeneous barriers and facilitators.¹² Examples include inadequate recognition of reproductive health needs by patients and providers,^{13,14} unclear role expectations of oncology versus fertility providers,¹⁵⁻¹⁸ lack of clear referral pathways,¹⁹⁻²¹ disparities in counseling between males and females,²² and lack of access to fertility programs, particularly in pediatric oncology settings.²³ These research and quality improvement efforts have been limited by scope and methodology, motivating a systematic approach to assess health system barriers, map scalable strategies to address them, and describe the design process for future dissemination and adaptation.

Electronic health records (EHRs) may facilitate strategies enabling fertility counseling. EHR systems can set rule-based reminders to staff and/or providers, automate referral pathways, generate reports of fertility referral and counseling, collect patient-reported information through a patient portal, and support telehealth. For widely used EHRs, functionalities are shareable, and mobile apps can upgrade

universally. Moreover, connectivity via smartphones is over 80% for AYAs, regardless of socioeconomic status, facilitating reach via such EHR apps.²⁴

To date, an implementation science approach – the study of methods to promote integration of evidence-based practices into routine health care – has not been undertaken to address the know-do gap (i.e., gap between what we know and what we do) in fertility counseling.²⁵ Thus, guided by the Consolidated Framework for Implementation Research (CFIR),²⁶ we systematically assessed barriers and facilitators to fertility counseling at cancer diagnosis and use of EHR tools as strategies to integrate fertility counseling into two oncology programs, one adult and one pediatric, and to bridge oncology and fertility programs. We focused on the female AYA survivor population because of an observed gap in care in our context, compared with male survivors. We compared adult versus pediatric and inpatient versus outpatient settings. We then designed a multi-component implementation strategy to fit these clinical contexts.

Methods

The study was approved by IRBs at University of California San Diego (UCSD) and Rady Children's Hospital. Researchers were female oncologists, reproductive endocrinologists, implementation scientists, and medical students. Between October 2018-May 2019, we enrolled reproductive-aged female survivors of AYA cancers and healthcare providers of this population to participate in semi-structured interviews or focus groups. Study settings were Moores Cancer Center, an adult comprehensive cancer center, and Rady Children's Hospital, a freestanding children's hospital affiliated with UCSD. At Moores, disease teams included hematologic malignancies, gastrointestinal, neuro-oncology, breast oncology, and radiation oncology. At Rady Children's, disease teams included liquid tumor, solid tumor, bone marrow transplant, and the survivorship clinic. There is an institutional fertility program at Moores, but not at Rady Children's.

We developed CFIR-based guides for provider interviews and AYA survivor focus groups to assess multi-level ecologic factors impacting fertility counseling and explore use of EHR-based implementation strategies. The guides encompassed questions on CFIR domains (intervention, individual, inner setting, outer setting, process) and relevant domain-specific constructs..²³

We conducted 19 health care provider semi-structured interviews, 8 at the adult program and 11 at the pediatric program. Oncologists, advanced practice providers, nurses and pharmacists from each disease team were approached for participation, because clinic processes varied by disease team. Program clinical and quality leaders were also recruited. After obtaining informed consent, we conducted 1-hour long interviews in person or via video calls of each participant with 2-3 investigators.

We conducted 4 focus groups with 2-3 reproductive-aged AYA survivor participants each. They were recruited from the investigators' prior research studies on reproductive health in AYA survivors..^{27, 28} Among participants who agreed to be contacted for future studies, we restricted to individuals treated at either of the two oncology programs and younger than 45. We purposefully sampled survivors at higher and lower risks of infertility (e.g., sarcoma survivors vs. thyroid cancer survivors) and at variable times since treatment in order to perspectives that may be impacted by these factors. Participants received recruitment emails, the study team answered questions, and consents were signed and returned prior to video focus groups. Focus groups were 1-hour long via video calls with 2-3 investigators.

Audio recording and note-taking occurred during interviews and focus groups. Participants were compensated with e-giftcards. Recruitment was stopped when data saturation was achieved.

Data analysis

We conducted thematic analysis facilitated by MaxQDA software..²⁹ In addition to deductive themes (e.g., CFIR constructs^{26, 30}), we identified inductive themes, those arising from the data, using the following steps: 1) two independent coders (AD, JG) read the transcripts, becoming familiar with the text and developing initial codes by consensus, 2) they coded three transcripts iteratively and refined the

codebook, 3) the final codebook was determined by consensus (AD, JG, HIS), 4) all data were coded, and 5) data were summarized by theme, with systematic comparison of pediatric versus adult settings and inpatient versus outpatient settings.³¹

Results

We contacted 19 providers and 72 AYA survivors for participation. Ten physicians (8 medical oncologists, 1 surgical oncologist, and 1 radiation oncologist), 2 advanced practice providers, 1 pharmacist, and 6 nurses participated. Median (range) of years in practice was 23 (29) years. Our sample had 8 adult providers and 11 pediatric providers, with 16 women and 3 men. Among the 9 AYA survivors, mean age was 33.1 (SD 6.8) years, and their cancer diagnoses included thyroid, cervical and bone cancers, leukemia and lymphomas. Table 1 summarizes barriers and facilitators to fertility counseling by CFIR domains (i.e., individual characteristics, inner context, outer context, process) and related constructs.

Individual characteristics

Providers had variable content knowledge about the existence of oncofertility care guidelines, fertility risks of cancer treatments, and fertility preservation procedures. Even for oncology providers who addressed fertility, depth of content knowledge was sometimes perceived to be lacking: “*[I] feel as though as I am inadequately trained in this arena. Having said all of that, I do feel like we are addressing the issue and at least putting it on the table, but I don’t think we’re doing enough*” (Advanced practice provider). This content knowledge gap limited self-efficacy; access to knowledge on fertility risks, which may be derived from a number of resources, was felt to be key to improving self-efficacy. Beyond content knowledge, oncology providers lacked knowledge on how fertility counseling and referral are operationalized in their clinic.

From the perspective of AYA survivors, fertility counseling would inform their decisions about timing of cancer treatment, and lack of knowledge about the option of fertility preservation procedures

prevented AYA survivors from undertaking them: *“I think before I would have liked to know that. In my opinion, I think doctors being doctors push treatment right away. And if I had known, like ... if I would've been able to preserve or do something for fertility, I think I would've chosen to wait on treatment and done that”* (AYA survivor).

While most providers expressed that oncofertility care is relevant to AYA survivors, a common theme was the belief that the timing of counseling at cancer diagnosis can be complicated by overwhelmed patients, competing oncology workup and/or treatment initiation needs, unclear cancer treatment plans that preclude informing patients about their reproductive risks, and poor prognosis: *“I don't know if they're going to need chemo because for various reasons, and they're going to surgery first, so I don't bring it up at all. And for the patients that I'm sending for neoadjuvant, I just assume the oncologist is going to do that unless they ask me specifically. If the patient asks me about it then I'll put the referral in”* (Surgical oncologist). As suggested by this oncologist, regardless of if/when a provider brings up fertility, patient-driven requests are a useful cue to action.

Preferences varied on which provider, i.e., physician (oncology or fertility), advanced practice providers, or nursing, conducts the primary fertility counseling. Oncology physicians reporting self-efficacy prefer to undertake primary counseling themselves, while other providers opt for automated referrals for all survivors to fertility specialists. There was consensus that nursing has the potential for fertility education, but would require significant additional oncofertility education.

Inner context

Discussions of clinic characteristics important to delivering counseling focused on the implementation climate and readiness. EHR tools were suggested and/or endorsed by providers as compatible with automating screening for fertility needs in clinic or before visits via patient portal, referral pathways between oncology and fertility, and accrual of fertility care metrics. For providers, advantages were alleviating personnel workload, systematically selecting the targeted population (e.g. via setting age parameters and type of encounter for automated fertility needs screens) and generating

shareable tools among organizations using the same EHR platform. EHR documentation, particularly with discrete fields, allows efficient collection of quality metrics for feedback and accreditation: *“It’s also important to have the resource of being able to pull metrics because otherwise you have this warm fuzzy feeling in your heart that we’re doing super well, but then the data shows that that was an erroneous warm fuzzy feeling”* (Quality leader).

Another provider theme centered on adapting available resources, including paper or EHR screening tools, EHR note templates, quality metrics reporting, and quality improvement processes for fertility counseling implementation. In addition, all levels of providers noted a lack of fertility educational materials that were sufficiently in-depth and with patient-friendly content: *“Recently, for example, I had a patient... we didn’t have the resources here. We had to try and pull resources and then go and discuss with the patients. I don’t think as pediatric oncologists we have been trained to do that.”*

Outer context

External factors influence clinic and individual provider delivery of oncofertility care. An oncology physician noted, *“[At] national meetings [oncofertility] is always a big topic, something that’s on our radar a lot.”* Additionally, advocacy and funding organizations use oncofertility care as a quality metric and a requirement for receiving funding:

One of [a pediatric oncology non-profit’s] big requirements is to have this fertility preservation talk at the very beginning, before starting chemotherapy ...They require that to get their sponsorship, we have to have formal protocol in place. (Pediatric oncologist)

Insurance coverage for fertility preservation services was repeatedly discussed as a barrier by both providers and patients. Even if insurance coverage for fertility counseling is available to patients, procedures may not be covered, and authorization processes are burdensome. For example, after a fertility preservation referral order has been placed in one oncology program, insurance authorization is required and obtained by a central or provider team-specific authorization unit. The authorization is a

physical letter, and to bypass the delay, the provider team always made additional calls to confirm referrals were being authorized.

Lastly, delivery of oncofertility care is complex due to involvement of different organizations, i.e., oncology clinics, fertility clinics, non-profit organizations, and insurance companies. Oncology programs do not always have a fertility program within the institution. Oncology and fertility clinics also need to join a network for fertility preservation financial support from non-profit organizations such as Livestrong. Both provider and AYA survivor participants noted a strong need to bridge these different organizations systematically and seamlessly. As an example, there was great reported burden on and lack of a streamlined, replicable process to enable oncology and fertility financial authorization staff and patients to interact with insurers to determine coverage.

Process

Many providers reflected on the types of engagement needed for effective implementation. Leadership engagement at three distinct levels were proposed: organizational level that sets oncofertility care as an institutional goal (e.g., selection of the Quality Oncology Practice Initiative quality metric³²), cancer team leaders that endorse care goals for the team, and a fertility champion (often cited as nurses) within each team. Providers viewed engagement of multiple champions as a facilitator because *“people feel resistance when it’s like this is one person’s project”* (Oncologist). Moreover, engaging AYA survivors as their own advocates would effectively cue their providers.

Co-planning implementation strategies with stakeholders was endorsed for fit and buy-in. Adaptation of processes will be needed to fit clinic context, e.g. which appointments are appropriate for introducing fertility needs screens. Several providers suggested piloting implementation strategies with engaged teams before full-scale implementation.

Three core components emerged as central to planning fertility counseling implementation (Figure 1). First, patients need to be screened for fertility needs. Second, a fertility care referral needs to be placed, if appropriate. Third, patients need access to a fertility specialist for additional fertility

counseling and appropriate fertility preservation strategies. Different types of individuals can participate in each stage.

Differences between pediatrics and adult settings

Several themes unique to pediatrics emerged. For adolescents, providers and AYA survivors similarly discussed that the timing of fertility counseling before, concomitant or after consultation with their parents/guardians should be guided by the family. Some providers questioned whether adolescents would be interested in talking about future fertility, believing it was best to talk to the parents only. Pediatric providers also discussed at what age it is appropriate to talk about fertility care: “*Like would the kid want to know about having children at 15 or would you talk just to the parents?*” (Advanced practice provider). These concerns were echoed in the focus groups, as several patients, when asked if they initiated oncofertility conversations with their providers, shared that fertility was not on their mind when diagnosed with cancer in adolescence. Additionally, complexity of privacy laws with regard to whether adolescents and/or parents/guardians have access to their patient portal (through which telehealth and screening questionnaires may be delivered) emerged as a barrier to EHR patient portal strategies.

Investigators’ mapping of strategies to fertility needs screening, referral, and specialist access

Based on barriers and facilitators reported by participants, the investigators then designed multiple strategies for the three core components, selected strategies and identified implementors. Table 2 specifies the five selected strategies for the two oncology programs: 1) automatic fertility needs screen using a best practice advisory (BPA), 2) an opt-out fertility referral pathway through the EHR system EPIC, 3) adding an option to conduct fertility counseling via telehealth, 4) audit and feedback to providers, and 5) conducting educational meetings. Table 3 specifies other strategies that were considered but not selected. Importantly, strategies apply to both female and male patients.

Key differences between the inpatient and outpatient environment were identified in mapping strategies. EHR tool specifications and pathways require different programming logic between the inpatient and outpatient setting. Inpatient insurance authorizations are not required for consultations, so

providers hypothesized that this would improve access to risk counseling by fertility specialists.

However, female fertility preservation procedures of oocyte or embryo banking generally occur in outpatient settings, which would limit the access of hospitalized patients.

Discussion

Guided by an implementation science framework, we conducted a qualitative study with oncology providers and AYA survivors from one adult and one pediatric oncology setting to systematically identify barriers and facilitators to fertility counseling at cancer diagnosis and co-design strategies with stakeholders. We report this systemized approach to inform teams seeking to develop or adapt their implementation of oncofertility care as part of routine oncology care.

Using CFIR domains and constructs allowed systematic assessment of the facilitators and barriers that influence fertility counseling implementation and change of processes. CFIR offered a pragmatic structure for our multi-level problem and a large number of domains and constructs that we could query.²⁶ While our qualitative guides encompassed questions based on a larger set of constructs, ultimately, the number of key relevant constructs (Table 1) was smaller and can guide the environmental scan of other clinical settings. Compatibility, feedback, available resources and planning constructs were particularly important in designing specific strategies for the two oncology programs.

Findings led to conceptualizing three core components of oncofertility care. Then, an array of implementation strategies targeting these components were evaluated, considering both our findings, implementation strategies from the Expert Recommendations for Implementing Change project,³³ and a priori research on components of fertility care models.¹² The goals were to select strategies that fit our context and may be applied across adult and pediatric settings; we aimed to minimize the number of strategies to limit complexity and system burden. The process of describing the strategies (naming, defining, operationalizing the actor, action, action targets, temporality, dose, implementation outcomes addressed and theoretical justification) clarified which ones were not feasible (e.g., high quality patient

educational materials on fertility risk) or did not fit (e.g., patient portal for screening).³⁴ The findings aim to enable planning of fertility counseling implementation efforts.

Across domains, we found variability in beliefs about whose role it was to provide fertility counseling, concordant with prior reports.¹⁴ For some providers, this concern was due to lack of self-efficacy about addressing fertility, an individual characteristic, while others attributed to inadequate time or incompatibility with a clinical visit, an inner setting characteristic. We also found that many types of providers could perform fertility needs screening, while only physicians and advanced practice providers could place referrals. Hence, specifying the actors of an implementation strategy was necessary.

We sought stakeholder input on EHR tools as implementation strategies. In designing and building EHR-tools, we could modify existing EHR tools such as the reporting tool for audit and feedback and the patient-friendly portal that enables secure video visits. Because of the limited number of EHR software systems, these tools could be widely disseminated and scalable. We also found that EHR tools may not be acceptable due to provider fatigue, feasible due to privacy laws, or appropriate due to low uptake of patient portal apps. Building these tools can be time- and labor-intensive, but a successful build in one clinical program may be disseminated to others.

This work advances prior research on oncofertility care delivery. Multiple reports highlight gaps in and interventions on provider knowledge and confidence, but show that while interventions increase providers' knowledge of fertility preservation, they largely do not increase care delivery.³⁵⁻³⁷ This is because barriers are multi-dimensional, e.g., need for inter-organizational networks and engaging oncology team on implementation procedures, as we have shown in the current study. A recent scoping review identified 9 core domains for oncofertility care.³⁸ We found that individual studies in this review contribute to 1-2 domains, e.g. communication with patients and training providers, but lacked a systemized interrogation of a clinical system and development of a coordinated, system-based intervention that we propose in our work. Detailing our approach to evaluating two clinical systems and

stakeholders and developing a coordinated intervention will enable other programs to adapt this more specific approach to their context.

Limitations include that we were limited to two oncology programs. While they were selected to reflect pediatric and adult oncology, their academic and existing single fertility clinic referral site limit generalizability. Additionally, our sampling strategy for health care providers and AYA survivors, like other qualitative work, was purposeful and not random, risking selection bias as the healthcare providers and AYA survivors who took part in our study may have an interest in fertility care. While the developed strategies target both sexes, female AYA survivors were specifically recruited because of the gap in care in our settings. It is possible that we did not capture additional barriers for male and childhood cancer survivors and their parents/guardians. As we developed strategies, we were limited by available resources. There is a strong need for the creation of educational resources to support fertility risk discussion, where current tools lack specificity and/or are too time intensive. Financial costs are a significant barrier that cannot be overcome with clinic-based implementation strategies alone.

In summary, we describe a systematic approach to design implementation strategies for fertility counseling at an adult and children's oncology program. We contribute data on the most salient CFIR constructs to assess and specifications on a set of potential strategies that may be adapted and deployed to improve the fertility care of adolescents and young adults with cancer.

Acknowledgments: We wish to thank Kris Brodsho for guidance on EHR capabilities and Nicole Stadnick, PhD for implementation science expertise.

Preprint: doi: <https://doi.org/10.1101/2020.06.30.20143891>

Author Disclosure Statement: Dr. Takemoto works for ServiceNow, Inc., which did not sponsor, support, or have oversight of this research.

References

1. Miller KD, Nogueira L, Mariotto AB, et al. Cancer treatment and survivorship statistics, 2019. *CA Cancer J Clin*. 2019;69(5):363-385.
2. van Dorp W, Haupt R, Anderson RA, et al. Reproductive Function and Outcomes in Female Survivors of Childhood, Adolescent, and Young Adult Cancer: A Review. *J Clin Oncol*. 2018;36(21):2169-2180.
3. Barton SE, Najita JS, Ginsburg ES, et al. Infertility, infertility treatment, and achievement of pregnancy in female survivors of childhood cancer: a report from the Childhood Cancer Survivor Study cohort. *Lancet Oncol*. 2013;14(9):873-881.
4. Gorman JR, Su HI, Roberts SC, Dominick SA, Malcarne VL. Experiencing reproductive concerns as a female cancer survivor is associated with depression. *Cancer*. 2015;121(6):935-942.
5. Gorman JR, Usita PM, Madlensky L, Pierce JP. Young breast cancer survivors: their perspectives on treatment decisions and fertility concerns. *Cancer Nurs*. 2011;34(1):32-40.
6. Oktay K, Harvey BE, Partridge AH, et al. Fertility Preservation in Patients With Cancer: ASCO Clinical Practice Guideline Update. *J Clin Oncol*. 2018;36(19):1994-2001.
7. Fallat ME, Hutter J, Bioethics AAoPCo, Hematology/Oncology AAoPSo, Surgery AAoPSo. Preservation of fertility in pediatric and adolescent patients with cancer. *Pediatrics*. 2008;121(5):e1461-1469.
8. American Society for Reproductive M, American College of O, Gynecologists' Committee on Gynecologic P. Prepregnancy counseling: Committee Opinion No. 762. *Fertil Steril*. 2019;111(1):32-42.
9. Network NCC. NCCN Clinical Practice Guidelines in Oncology: Adolescent and Young Adult (AYA) Oncology, Version 1.2021-September 10, 2020. https://www.nccn.org/professionals/physician_gls/default.aspx#age. Published 2020. Accessed November 1, 2020.
10. Patel A, Schwarz EB, Society of Family P. Cancer and contraception. Release date May 2012. SFP Guideline #20121. *Contraception*. 2012;86(3):191-198.
11. Patel P, Kohn TP, Cohen J, Shiff B, Kohn J, Ramasamy R. Evaluation of Reported Fertility Preservation Counseling Before Chemotherapy Using the Quality Oncology Practice Initiative Survey. *JAMA Netw Open*. 2020;3(7):e2010806.
12. Anazodo A, Laws P, Logan S, et al. How can we improve oncofertility care for patients? A systematic scoping review of current international practice and models of care. *Hum Reprod Update*. 2019;25(2):159-179.
13. Gorman JR, Bailey S, Pierce JP, Su HI. How do you feel about fertility and parenthood? The voices of young female cancer survivors. *J Cancer Surviv*. 2012;6(2):200-209.
14. Quinn GP, Vadaparampil ST, Lee JH, et al. Physician referral for fertility preservation in oncology patients: a national study of practice behaviors. *J Clin Oncol*. 2009;27(35):5952-5957.
15. Johnson RH, Kroon L. Optimizing fertility preservation practices for adolescent and young adult cancer patients. *J Natl Compr Canc Netw*. 2013;11(1):71-77.
16. Duffy C, Allen SM, Dube C, Dickersin K. Oncologists' confidence in knowledge of fertility issues for young women with cancer. *Journal of Cancer Education*. 2012;27(2):369-376.
17. Overbeek A, Van den Berg M, Louwé L, et al. Practice, attitude and knowledge of Dutch paediatric oncologists regarding female fertility. *Neth J Med*. 2014;72(5):264-270.
18. Ghorbani B, Madahi P, Shirazi E, Ardekani HS, Kamali K. Iranian oncologists' attitude towards fertility preservation in a sample group. *Journal of Reproduction & Infertility*. 2011;12(1):33.

19. Loren AW, Brazauskas R, Chow EJ, et al. Physician perceptions and practice patterns regarding fertility preservation in hematopoietic cell transplant recipients. *Bone Marrow Transplant*. 2013;48(8):1091-1097.
20. Clayman ML, Harper MM, Quinn GP, Reinecke J, Shah S. Oncofertility resources at NCI-designated comprehensive cancer centers. *J Natl Compr Canc Netw*. 2013;11(12):1504-1509.
21. Warner E, Yee S, Kennedy E, et al. Oncofertility knowledge, attitudes, and practices of Canadian breast surgeons. *Annals of surgical oncology*. 2016;23(12):3850-3859.
22. Köhler TS, Kondapalli LA, Shah A, Chan S, Woodruff TK, Brannigan RE. Results from the survey for preservation of adolescent reproduction (SPARE) study: gender disparity in delivery of fertility preservation message to adolescents with cancer. *J Assist Reprod Genet*. 2011;28(3):269-277.
23. Rabah DM, Wahdan IH, Merdawy A, Abourafe B, Arafa MA. Oncologists' knowledge and practice towards sperm cryopreservation in Arabic communities. *Journal of Cancer Survivorship*. 2010;4(3):279-283.
24. Center PR. Mobile Fact Sheet. <http://www.pewinternet.org/fact-sheet/mobile/>. Published 2018. Accessed October 10, 2018, 2018.
25. Implementation Science. <https://cancercontrol.cancer.gov/IS/>. Updated February 5, 2020. Accessed May 17, 2020.
26. Damschroder LJ, Aron DC, Keith RE, Kirsh SR, Alexander JA, Lowery JC. Fostering implementation of health services research findings into practice: a consolidated framework for advancing implementation science. *Implementation science : IS*. 2009;4:50.
27. Su HI, Stark S, Kwan B, et al. Efficacy of a web-based women's health survivorship care plan for young breast cancer survivors: a randomized controlled trial. *Breast Cancer Res Treat*. 2019;176(3):579-589.
28. Su HI, Kwan B, Whitcomb BW, et al. Modeling variation in the reproductive lifespan of female adolescent and young adult cancer survivors using AMH. *J Clin Endocrinol Metab*. 2020;105(8):2740-2751.
29. M. P. Qualitative evaluation and research methods, 2nd ed. *Thousand Oaks, CA: Sage Publications*. 1990.
30. Damschroder LJ, Lowery JC. Evaluation of a large-scale weight management program using the consolidated framework for implementation research (CFIR). *Implement Sci*. 2013;8:51.
31. Ulin PR RE, Tolley EE. Qualitative methods in public health: a field guide for applied research. In. 1st ed. San Francisco: Jossey-Bass2005.
32. McNiff K. The quality oncology practice initiative: assessing and improving care within the medical oncology practice. *J Oncol Pract*. 2006;2(1):26-30.
33. Powell BJ, Waltz TJ, Chinman MJ, et al. A refined compilation of implementation strategies: results from the Expert Recommendations for Implementing Change (ERIC) project. *Implement Sci*. 2015;10:21.
34. Proctor EK, Powell BJ, McMillen JC. Implementation strategies: recommendations for specifying and reporting. *Implementation Science*. 2013;8(1):139.
35. Zhang HF, Jiang QH, Huang GY, et al. The Educational Program for Healthcare Providers Regarding Fertility Preservation for Cancer Patients: a Systematic Review. *Journal of cancer education : the official journal of the American Association for Cancer Education*. 2020.
36. Quinn GP, Vadaparampil ST. Fertility preservation and adolescent/young adult cancer patients: physician communication challenges. *J Adolesc Health*. 2009;44(4):394-400.
37. Forman EJ, Anders CK, Behera MA. A nationwide survey of oncologists regarding treatment-related infertility and fertility preservation in female cancer patients. *Fertil Steril*. 2010;94(5):1652-1656.

38. Anazodo A, Laws P, Logan S, et al. The Development of an International Oncofertility Competency Framework: A Model to Increase Oncofertility Implementation. *The oncologist*. 2019;24(12):e1450-e1459.

Figure 1: Fertility care pathway involves a fertility care needs screen, a fertility referral, and fertility counseling with multiple potential actors in each step.

Figure 1

Table 1

Construct	
Individual	
Knowledge and Beliefs	Facilitators: <ul style="list-style-type: none"> • Variation in which oncology visit and which provider should address fertility • Patient-driven requests for fertility care would prompt provider actions
	Barriers: <ul style="list-style-type: none"> • Provider content knowledge gap: Treatment-related fertility risks, fertility preservation procedures, fertility care clinical guidelines • Provider operational knowledge gap on how to refer to fertility care • Provider belief that fertility discussions are not appropriate when patients are overwhelmed, cancer workup/treatments more pressing, cancer treatment plans unknown or prognosis is poor
Self-Efficacy	Facilitators: <ul style="list-style-type: none"> • Resources to support knowledge gaps on treatment-specific fertility risks: pharmacists, fertility specialists, or reliable tool • Resources to support nurse-led education of survivors
Inner Setting	
Implementation Climate	Facilitators: <ul style="list-style-type: none"> • Feedback: Implementation metrics of each clinical team preferred over individual provider • Feedback: Peer pressure through public audit/feedback • Compatibility: EHR-enabled, automated screening tools, referral pathways, collection quality metrics
	Barriers: <ul style="list-style-type: none"> • Compatibility: Heterogeneous EHR templates and systems • Compatibility: Automated screening protocols lacks flexibility (e.g. patient overwhelmed, treatment plan unknown, too close to treatment to allow for fertility preservation procedures)
Readiness for Implementation	Facilitators: <ul style="list-style-type: none"> • Access to knowledge and information: Expertise on fertility risks and procedures • Access: Devices and Internet connectivity for in clinic telehealth access to off-site specialists • Available resources: EHR system to support screening, referrals, quality metrics, patient portal • Available resources: Existing patient screening tools, templated notes or pathways, quality improvement processes that can be adapted for fertility care
	Barriers: <ul style="list-style-type: none"> • Available resources: Heterogeneity in personnel resources to support screening (navigator, social work), risk counseling (pharmacist), in person translation • Available resources: Lack of educational materials with depth for patients
Outer Setting	
External Policy & Incentives	Facilitators: <ul style="list-style-type: none"> • Fertility care as a quality metric from accreditation organizations • Clinical guidelines from oncology societies recommend fertility care • Systemized delivery of fertility care as a requirement of AYA foundation funding • Insurance coverage for fertility preservation services
	Barrier: <ul style="list-style-type: none"> • Insurance approval of fertility preservation is inconsistent, time consuming and complex
Peer Pressure	Facilitator: <ul style="list-style-type: none"> • High quality, systemized fertility care can set an oncology program apart from others
Cosmopolitanism (networks)	Barrier: <ul style="list-style-type: none"> • Inter-organizational networks (between oncology and fertility clinics or fertility care funding organizations and clinics) are lacking
Implementation Process	
Engaging	Facilitators: <ul style="list-style-type: none"> • Oncology team: Identify a nursing champion to support team-specific implementation procedures • Organization: Engage leadership, e.g. cancer cabinet or quality committee, to select fertility care as an institutional goal

	<ul style="list-style-type: none"> • Patients educated about fertility care can prompt their oncology providers
Planning	<p>Facilitators:</p> <ul style="list-style-type: none"> • Plan for providing implementation metrics/feedback to providers • Adapt processes to fit each clinic, e.g. determine when/which appointments appropriate for fertility screening of a newly diagnosed cancer patient, order of face-to-face consultations with adolescents and parents (together, tandem) • Plan for resources needed, e.g. Internet access, language translators, provider educational session • Stage implementation scale up: First pilot in a limited setting, e.g. a few oncology teams

Table 2

Strategy	Actor	Action	Action Target(s)	Temporality	Dose	Implementation & Services Outcomes	Justification
Mandate Change	Cancer center leadership	Select fertility counseling as cancer center quality goal	- Oncology teams	Once	Once	- Uptake of fertility counseling by AYA survivors at diagnosis and in survivorship	1. Prioritizes implementation 2. Access to quality improvement team and resources
Automate Fertility Needs Screen: Remind Clinicians	EHR system EPIC	Best Practice Advisory (BPA) pops up to 1) remind clinicians about fertility counseling, 2) shortcut to referral order	- Physicians - APPs	Trigger criteria: new oncology visit, age (<42 females, <50 males), cancer diagnosis in EHR	Each provider will see BPA maximum of one time; after referral, no other providers will see BPA for 2 years	- Screen all newly diagnosed and 2-year post-treatment AYA survivors for fertility needs	1. Compatible with EHR 2. Addresses oncology provider content & operational knowledge gap
Automate Fertility Referral Between Clinics	EHR system EPIC	Fertility specialist referral order with cancer treatment plan automatically placed in a STAT fertility scheduler queue for insurance authorization and contacting patient within 72 hours	- Oncology physicians and APPs - Fertility clinic schedulers ownership of insurance authorization & patient contact - Fertility specialists know proposed cancer treatments	Each referral order	Once per referral	- Fewer patients lost to care between two clinics - Insurance authorization & scheduling efficiency - More precise fertility risk counseling by fertility specialists	1. Compatible with EHR 2. Addresses oncology providers operational knowledge gap
Fertility Counseling: Add Service Sites	Fertility specialist, EHR system and patient portal	Televideo fertility counseling using EHR provider tool and EHR patient portal	- AYA survivors at diagnosis and post-treatment	Offered to patient by fertility scheduler after fertility referral order placed	An initial 30-60-minute fertility counseling visit on treatment-related reproductive risks and fertility preservation options	- Uptake of fertility counseling by AYA survivors - Patient-centered, timely visits	1. Compatible with EHR 2. Addresses geographic and time (multiple visits to multiple providers) barriers

Audit & Feedback	EHR system, quality team	Metrics on screening, referral, counseling	<ul style="list-style-type: none"> - Oncology teams - Fertility teams 	After initiation of screening & automated referral pathways	Monthly reports by individual provider and by clinic team	- Fidelity of BPA screening and referral pathway	1. Compatible with EHR-based screening and referral 2. Peer pressure
Conduct Educational Meetings	Implementation team, fertility specialist	20-minute educational session on fertility content and operationalizing implementation strategies	<ul style="list-style-type: none"> - Cancer center cabinet and quality committee - Oncology physicians and APPs at oncology team meeting - Fertility clinic schedulers and administrators 	After preparation of strategies	Once per group	- Increase acceptability of implementation strategies	1. Addresses oncology provider content & operational knowledge gap

Table 3

Fertility Care Pathway Order	Strategy	Facilitators & Barriers	Justification
Fertility Needs Screen	Involve patients: AYA survivors screened through questionnaire sent via EHR patient portal MyChart	Facilitators: <ul style="list-style-type: none"> • Patient time: not limited by in person visit time • EHR patient portal uptake high in adults, low in pediatrics 	Available Resource Compatibility
		Barriers: <ul style="list-style-type: none"> • Heterogeneity of questionnaires used by providers • Cannot be used in inpatient setting • Privacy laws do not allow parents access to patient portal in children $\geq$ age 12 • No Spanish version of patient portal • Wording appropriate to children and adolescents 	Compatibility & Available Resources Compatibility External Policy & Incentives Adaptability Complexity
	Involve patients: Paper questionnaire in waiting room	Facilitators: <ul style="list-style-type: none"> • Does not require patient to enroll in patient portal • Could be given in multiple languages 	Available Resource Adaptability
		Barriers: <ul style="list-style-type: none"> • Wording appropriate to children and adolescents • Available patient time to consider screening questions limited • Provider may miss the paper questionnaire 	Complexity Available Resource Compatibility
	Revise professional roles: Dedicated fertility navigator	Facilitator: <ul style="list-style-type: none"> • One provider that does all the steps: screening and referral 	Complexity
		Barriers: <ul style="list-style-type: none"> • Cost • Depending on volume, difficult to capture all new diagnoses 	Available Resources Compatibility
	Technical assistance – EHR templates to document screens	Barrier: <ul style="list-style-type: none"> • Heterogeneity of note templates limits automation and uniform uptake 	Complexity, compatibility
Fertility Referral	Remind clinicians: EHR inbox messages on new patients	Barrier: <ul style="list-style-type: none"> • Providers overwhelmed by inbox messages 	Compatibility
	Automated referral for all new patients	Facilitator: <ul style="list-style-type: none"> • Has one less step required of oncology providers 	Complexity
		Barrier: <ul style="list-style-type: none"> • Leads to unnecessary consults 	Relative Advantage
	Remind clinicians: added referral order	Barrier: <ul style="list-style-type: none"> • Heterogeneity of order sets used by different disease teams 	Complexity

	to admission order set	<i>Would not work in outpatient setting</i>	<i>Compatibility</i>
--	---------------------------	---	----------------------