UC Irvine

UC Irvine Previously Published Works

Title

Molecular mapping of deletion breakpoints on chromosome 4 of *Drosophila melanogaster*.

Permalink

<https://escholarship.org/uc/item/51c7d0p6>

Journal

Chromosoma, 112(8)

ISSN

0009-5915

Authors

Podemski, Lynn
Sousa-Neves, Rui
Marsh, J Lawrence
et al.

Publication Date

2004-06-01

DOI

10.1007/s00412-004-0286-4

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Lynn Podemski · Rui Sousa-Neves ·
J. Lawrence Marsh · John Locke

Molecular mapping of deletion breakpoints on chromosome 4 of *Drosophila melanogaster*

Received: 28 January 2004 / Revised: 1 April 2004 / Accepted: 2 April 2004 / Published online: 8 June 2004
© Springer-Verlag 2004

Abstract As part of our effort to induce and identify mutations in all genes on chromosome 4 of *Drosophila melanogaster*, we have mapped the breakpoints of eight chromosome 4 deficiencies relative to the predicted genes along this chromosome. Although the approximate locations of *Df(4)G*, *Df(4)C3*, *Df(4)M101-62f*, *Df(4)M101-63a*, *Df(4)J2*, *Df(4)O2*, *Df(4)C1-10AT*, and *Df(4)B2-2D* are known (some from cytological observations and others predicted from *P* element locations), the extents of these deletions have not been mapped with respect to the predicted genes identified by the *Drosophila* Genome Project. Polymerase chain reaction primers were designed to amplify the predicted exons of all chromosome 4 genes, and homozygous embryos for each deficiency were identified and their DNA used to test for the presence or absence of these exons. By testing for the inability to amplify various exons along the length of the chromosome, we were able to determine which predicted genes are missing in each deficiency. The five deficiencies, *Df(4)G*, *Df(4)C3*, *Df(4)C1-10AT*, and *Df(4)B2-2D* (all terminal deletions), and *Df(4)M101-62f* (a proximal interstitial deletion), enabled us to partition the gene-containing, right arm of chromosome 4 into five regions. Region A [uncovered by *Df(4)M101-62f*] contains the proximal-most 21 genes; region B [uncovered by *Df(4)B2-2D*] contains the next 12 genes; region C [uncovered by *Df(4)B2-2D* and *Df(4)C1-10AT*] contains the next 17 genes; region D [uncovered by *Df(4)B2-2D*, *Df(4)C1-10AT*, and *Df(4)C3*] contains the next 21 genes; and region E [uncovered by *Df(4)B2-2D*, *Df(4)C1-10AT*, *Df(4)C3*, and *Df(4)G*] contains the distal-most ten genes. By using *Df(4)M101-62f*, *Df(4)B2-2D*, *Df(4)C1-10AT*, *Df(4)C3*, and *Df*

(4)G in complementation tests, we can assign newly induced recessive lethal mutations to one of the five regions on chromosome 4. This will substantially reduce the amount of DHPLC analysis required to match each mutation to a predicted transcript on chromosome 4.

Introduction

The fourth chromosome of *Drosophila melanogaster* is the smallest of the chromosomes. It is about 5 Mb in length, and approximately 4 Mb of that comprises satellite repeats and does not appear to contain any genes (Locke and McDermid 1993). The gene-containing region is approximately 1.2 Mb in length, and forms a polytenized region with cytological band designations 101E–102F8. Chromosome 4 has several unusual features that set it apart from the other autosomes and point to its heterochromatic nature. Unlike the other chromosomes, there is no crossing over in females during normal meiosis (Bridges 1935). The fourth chromosome often shows a poorly banded appearance in salivary gland polytene chromosome spreads similar to regions of β -heterochromatin. Chromosome 4 contains low and middle copy repetitive elements that are typically found in β -heterochromatin (Miklos et al. 1988; Carmena and Gonzalez 1995; Pimpinelli et al. 1995). The recent annotation of the *Drosophila* genome shows that the density of transposable elements on chromosome 4 is significantly higher than that of the other arms. The average density of transposable elements on the *X*, *2L*, *2R*, *3L* and *3R* arms is only 10–15 elements per megabase (million bases), while the density on chromosome 4 is 82 elements per megabase. The difference between the densities is made up almost exclusively in LINE-like and TIR elements, and these elements are distributed all along the chromosome, both between and within genes (Kaminker et al. 2002). Chromosome 4 is also highly enriched for the repetitive element DINE-1, which has been localized almost exclusively to chromosome 4 and centric heterochromatic regions (Locke et al. 1999). Heterochromatic protein 1

L. Podemski · J. Locke (✉)
Department of Biological Sciences, University of Alberta,
Edmonton, AB, Canada, T6G 2E9
e-mail: john.locke@ualberta.ca

R. Sousa-Neves · J. L. Marsh
Department of Developmental and Cell Biology, University of
California, Irvine,
Irvine, CA, 92697, USA

(HP1), an important constituent of heterochromatin, has been shown to bind extensively to chromosome 4 (James et al. 1989). A chromosome binding protein, Painting of Fourth (POF), has been shown to localize exclusively to chromosome 4, and it has been speculated that this protein may be involved in the regulation of genes in a heterochromatinized environment (Larsson et al. 2001). In addition, *P* element transgenes inserted into chromosome 4 frequently show the variegated expression (position-effect variegation) typical of genes inserted into heterochromatic boundaries (Wallrath and Elgin 1995; Wallrath et al. 1996). Mapping studies of variegating and non-variegating transgene inserts into chromosome 4 suggest that the polytenized regions of this chromosome comprise interspersed euchromatic and heterochromatic domains (Sun et al. 2000).

The lack of crossing over during meiosis makes the construction of a conventional genetic map of chromosome 4 impossible. Hochman (1976) undertook an extensive mutational screen on chromosome 4, in which he was able to identify 37 vital loci and 6 visible mutations. From a statistical analysis he also estimated the presence of another ~30 loci, bringing the total gene number on chromosome 4 to approximately 70. He was, however, only able to map his mutations based on complementation to two chromosome 4 deletions: *Df(4)M101-1* (101EF;102B6-7) and *Df(4)G* (102E2;102F8). He assumed the majority of the mutants (26) fell into the large region between these two deletions.

There are relatively few chromosome 4 deficiencies available from the stock center. Most of these deficiencies were created by X-ray mutagenesis or, more recently, FLPase mediated events (Ahmad and Golic 1998; Kramps

et al. 2002). Two of the deficiencies we discuss here, *Df(4)B2-2D* and *Df(4)C1-10AT*, were created in a novel *P* element screen (Sousa-Neves et al., unpublished). These deficiencies have been mapped approximately on the basis of cytology and using only a few chromosome 4 genes in complementation analysis. Release 3.1 of the *Drosophila* genome sequence (Adams et al. 2000; Celniker et al. 2002; Misra et al. 2002) has identified 82 putative loci on chromosome 4. We are currently undertaking a large mutational screen, using ethylmethyl sulfonate (EMS), in which we hope to create mutations in every locus on chromosome 4. We then propose to match each mutation to a chromosome 4 transcript using DHPLC analysis and confirm the locations by sequencing. To streamline our identification and matching process, we have used polymerase chain reaction (PCR) amplification to map molecularly the putative transcripts to the available chromosome 4 deficiencies. We have been able to map the extents of these deletions, and hence divide the chromosome into five regions based on the molecular mapping of five chromosome 4 deficiencies. Complementation analysis can thus be used to map each of our new EMS-induced mutations to one of the five groups, which is the first step in the localization of the mutations to their respective genes. This set of molecularly defined deletions will also be of value to other *Drosophila* researchers who wish to map chromosome 4 mutations.

Fig. 1 An example of a polymerase chain reaction (PCR) experiment used to determine molecular breakpoints of a deficiency. DNA from homozygous *Df(4)G* embryos was tested with PCR primers from four genes: CG2999 (*unc-13*), CG11081 (*plexA*), CG11062 (*activin-beta*), and CG11027 (*Arf102F*). Oregon R DNA was used as a positive control for PCR amplification, and a PCR negative control (no DNA template) was also used to ensure that the amplified bands were dependent upon the addition of template DNA. The PCR products from *unc-13* and *plexA* were successfully amplified in the *Df(4)G* DNA lanes, indicating that these two regions are present in *Df(4)G* DNA. The products from the *activin-beta* and *Arf102F* primers were not amplified in the *Df(4)G* lanes, indicating that these two regions are deleted in the *Df(4)G* DNA

Table 1 A summary of the polymerase chain reaction (PCR) primer pairs used to determine the location of the *Dff(4)* breakpoints. The breakpoint of each *Dff(4)* is bounded by a PCR primer pair on the proximal (left) and distal (right) side of the break. Note that *Dff(4)O2* is the only deficiency with two identified breakpoints. The primer pairs are identified by their computed gene number (CGxxxxx) and the strand orientation forward (F) or reverse (R)

Deficiency <i>Dff(4)</i>	PCR primer pairs at the proximal breakpoint (proximal pair; distal pair)	Primer sequences (5'-3')	PCR primer pairs at the distal breakpoint (proximal pair; distal pair)	Primer sequences (5'-3')
<i>M101-62f</i>	Could not be located		CG1674-F4A/CG1674-R4A	CTGTTTACTCGCATGTGTTA GTTTCATAGGCACAAATACAT
<i>B2-2D</i>	CG1748-F2/CG1748-R2	AAGCATTGAGCATTAACCAA ATGAAGGCACACGAACTAT	CG1674-F5A/CG1674-R5A	TCATTTCACGTCGATTAAACAC TTTCTGTGTTTCTTGATTGTTTC
	CG2380-F6/CG2380-R6	GCCCAAGCCGAAAGAGTTTAA CCAAATGAGTAACGGGTAGC		
<i>C1-104T</i>	CG11533-F1/CG11533-R1	TAGCAACCTGAGTCTTATAAT CAGCAGAGCATCGAAGGTAT		
	CG1449-F1/CG1449-R1	TGGTAATGACGACGGCTCTC CTGTATAATGGCACTGGAGTA		
<i>J2</i>	CG1838-F3/CG1838-R3	CTGTATAATGGCACTGGAGTA TTCAGCATTCATCAACGGTTT		
	CG1464-F1/CG1464-R1	TCACCATCTGCGCAAAATTAC CGGATAAAGACGATGATGCC		
<i>C3</i>	CG32019-F9/CG32019-R9	GATCAGAAAGTCGCGGAGTC ACCTCTGAACTAGTGATTAATC		
	CG32019-F42/CG32019-R42	TGGCTGTGATGTTAGATGCTG TGCAAGTCGATAGTGTACATT		
<i>O2</i>	CG32019-F9/CG32019-R9	GATCAGAAAGTCGCGGAGTC ACCTCTGAACTAGTGATTAATC	CG11091-F1/CG11091-R1	GGAAAGTGTAGAATCTGTTG ATTCAACCTATTTCACAGTGT
	CG32019-F42/CG32019-R42	TGGCTGTGATGTTAGATGCTG TGCAAGTCGATAGTGTACATT	CG11186-F1/CG11186-R1	CGTAGTTCTCTGTTTAGTTAA TGATGCTAAACAACACTGAACAC
<i>G</i>	CG11064-F15/CG11064-R15	AACCTCCGATGTAATTTCAT GTTGCCCGCTTTCTGCTAT		
	CG11062-F3/CG11062-R3	ACGGAGAAAGGTAACATTA GGCAAGCTGTAAAAGTAATT		

Materials and methods

Stocks of *D. melanogaster*

We tested the following chromosome 4 deficiency stocks: *Df(4)M101-62f*, *Df(4)M101-63a*, *Df(4)C3*, *Df(4)G*, *Df(4)O2*, *Df(4)J2*, *Df(4)C1-10AT*, and *Df(4)B2-2D*. Some of the listed chromosome 4 deficiency stocks are no longer available from the stock center. For example, we were unable to obtain *Df(4)M101-1* and *Df(4)38* because our genetic analysis of the stocks revealed that they no longer contained the deficient chromosomes.

Crosses and identification of homozygous embryos

Deficiency stocks were mated to a *ci¹ey^R* stock. Virgin *Df(4)/ci¹ey^R* females were mated to *Df(4)/ci¹ey^R* males and allowed to lay eggs on agar plates. After approximately 24 h, single embryos were picked from the plates into tubes, and the DNA was extracted from each embryo by squashing in 10 μ l of 10 mM TRIS-Cl, pH 8, 1 mM

EDTA, pH 8, 200 μ g/ml Proteinase K solution. The tubes were incubated at 37°C for 45 min, and heated at 99°C for 10 min. Each embryo preparation was tested first for the presence of the *ci¹* chromosome using primers JS1 (ATACATATGTTTCATTACGG) and JS3 (TACTCAGTTCAAATCTTGTG), and second for DNA quality using a PCR primer pair known to be present on the deficient chromosome. Embryo preparations that produce a band in the DNA quality test, but fail to produce a band in the *ci¹* test must therefore be homozygous for the deficiency.

Molecular mapping

Homozygous deficient embryo DNA was used to test for the presence or absence of primers specific to genes located on chromosome 4. Primer pairs for nearly all the putative or known gene exons on chromosome 4 (based on Release 3.1) were synthesized and PCR amplification was performed using a selection of primer pairs and homozygous deficient embryo DNA as a template. A PCR product of the expected band size indicated that the primers (and hence that exon) were still present in the deficiency DNA, while the lack of a band indicated that the gene was

Fig. 2a–h Polytene chromosome squashes from the salivary glands of chromosome 4 deficiency chromosomes heterozygous with cytologically normal chromosomes. **a** The cytological location of the *Df(4)M101-62f* breakpoints. This chromosome is deleted for the region including 101E–102A6. **b** The breakpoint locations of *Df(4)B2-2D*. The deleted chromosome points to the left and the cytologically normal *ci^D* chromosome points to the right. The chromosome is deleted for bands 102B2 to the telomere. **c** The *Df(4)C1-10AT* chromosome lying adjacent to two *ci^Dspa^{pol}* chromosomes. The deletion removes the region from 102C2 to the telomere. **d** The *Df(4)J2* deletion, which includes the terminal portion of the chromosome starting at 102D2. **e** The *Df(4)C3* deletion, which is cytologically identical to *Df(4)J2*, and removes the region from 102D2 to the telomere. **f** The deleted region of *Df(4)G*, which removes the region from 102E2 to the telomere. **g** The interstitial deletion of *Df(4)O2*, which removes the region from 102D1 to 102D4. **h** The *Df(4)M101-63a* deficiency, which is cytologically normal since no band aberration can be seen

completely or partially deleted. An example of this method showing two genes present and two genes absent in *Df(4)G* DNA is shown in Fig. 1. Once an approximate molecular location for the deletion had been established, primer pairs specific to each exon of the genes near the breakpoints were used for PCR amplification to map the location of the deletion breakpoint in finer detail. A list of the relevant primers is shown in Table 1. Interstitial breakpoints were mapped to a location within a particular exon of a gene, or to the region between two genes. If the most distal exon of the last gene on chromosome 4, *CG18026* (*caps*), was shown to be absent, the deficiency was deemed a terminal deletion.

Cytology

Polytene chromosome squashes were made from dissected salivary glands of third instar larvae of the eight *Df(4)* containing stocks (Fig. 2a–h). The glands were stained in lacto-aceto-orcein before squashing to enhance the visibility of the bands. Photographs of the slides were taken under phase contrast microscopy. Most of the *Df(4)*s were stocked over a cytologically normal fourth chromosome, with the exception of *Df(4)B2-2D* and *Df(4)C1-10AT*, which were kept most easily as fourth chromosome trisomics. *Df(4)B2-2D* is stocked with two *ci^D* chromosomes and *Df(4)C1-10AT* is stocked with two *ci^Dspa^{pol}* chromosomes.

Results

The mapping of each *Df(4)* is described below.

Df(4)M101-62f

This deficiency is the proximal-most interstitial deletion of chromosome 4. Using methodology like that shown in Fig. 1, we mapped the distal breakpoint of this deletion to a 946 bp region of DNA between exons 4 and 5 in gene *CG1674*. Cytological observations place this breakpoint just proximal to band 102B1 (Fig. 2a). The proximal breakpoint location is not clear from our cytological observations. Our molecular breakpoint analysis has placed it somewhere proximal to the first exon of *CG17923*, the second most-proximal predicted gene on chromosome 4.

The most proximal predicted gene on chromosome 4, *CG32013*, appears to be present in DNA from homozygous *Df(4)M101-62f* embryos; however, all attempts to amplify a PCR product that spans the expected deletion (*CG32013* to the *CG1674* breakpoint) from homozygous deficient embryo DNA were unsuccessful. In addition, PCR amplifications using *CG32013*-specific primer pairs produced bands of the expected sizes with DNA from embryos that lack chromosome 4 entirely [null-4s derived from *C(4)RM/0* parents]. This paradoxical result can be explained in several ways.

First, gene *CG32013* is repetitive in nature, and thus the primer pairs for *CG32013* are able to amplify regions on other chromosomes besides chromosome 4. BLASTN comparison (Altschul et al. 1990) of the *CG32013* sequence with the genome showed significant homology of some regions of the *CG32013* gene sequence to centromeric regions of other chromosomes. In addition,

hybridization of *CG32013* probes to genomic DNA blots resulted in multiple bands (data not shown), indicating that there are repetitive sequences in *CG32013*. However, we could not find any additional whole copies of the *CG32013* gene using a BLASTN search. We sequenced the null-4 PCR products amplified with *CG32013* primer pairs and found the products to be exactly identical to the released genome sequence predicted for chromosome 4. Therefore, multiple copies of *CG32013*, if they exist, must be highly conserved, but not in the current genome sequence.

Another explanation for the ability to amplify regions of *CG32013* from null-4 embryo DNA is that gene *CG32013* is not actually located on chromosome 4 as shown in the Release 3.1 sequence. We attempted to verify the Release 3.1 location of gene *CG32013* by PCR amplifying the region between *CG32013* and the adjacent gene *CG17923*, which we know to be on chromosome 4 because of its absence in *Df(4)M101-62f* and null-4s. As PCR templates, we used genomic DNA isolated from five different *D. melanogaster* stocks, including the same stock used to produce the RPCI-98 BAC library (Hoskins et al. 2002). In addition to the genomic DNA templates, DNA from clone BACR05L22, the proximal-most BAC on chromosome 4 based on the physical map of Locke et al. (2000), was also used as a PCR template. The proximal end of this clone corresponds to the proximal end of the Release 3.1 sequence. The PCRs using the genomic DNA all failed to produce the band predicted from the Release 3.1 sequence, but the BAC clone DNA did produce the predicted band. We also performed Southern blot analysis of this region using the five genomic DNAs and BACR05L22 DNA. A hybridization probe was produced by PCR amplification of Oregon R DNA using a primer pair specific to gene *CG17923*. The probe was hybridized to genomic DNA digested with *EcoR* I or *Hind* III. The Release 3.1 sequence predicts both digests should result in a single fragment (2.8 kb for *EcoR* I, and 9.0 kb for *Hind* III) spanning the *CG32013*–*CG17923* genes. The BAC clone produced the predicted bands in both the *EcoR* I and *Hind* III digest lanes; however, the genomic DNAs from the five *Drosophila* stocks all produced a common, single band of a different size (1.4 kb for *EcoR* I, 4.7 kb for *Hind* III). No hybridization was seen at the sizes predicted from the Release 3.1 sequence. A second probing was done, using a hybridization probe from gene *CG32013*. This probe should produce identical bands to the first probing if the two genes are indeed adjacent. The BAC digests showed hybridization to the same, predicted, single bands for the *EcoR* I and *Hind* III digests, while this probe resulted in hybridization to 8–12 bands in the *EcoR* I and *Hind* III lanes of all five stocks, indicating it contained repetitive sequences. None of these bands was the predicted size, and none was the same size as the band produced from the *CG17923* probe, indicating that the two genes are not located within the same restriction fragment as predicted in Release 3.1. Both the PCR and Southern data indicate these two genes cannot be adjacent on chromosome 4. Since the restriction digest pattern of the

BAC clone differs from the digest pattern of genomic DNA in this region, we must also conclude that neither clone BACR05L22 nor the sequence data from Release 3.1 is representative of the *Drosophila* genome in the region proximal to gene *CG17923*. The Southern blot results, combined with the PCR results from *nullu-4* embryos, indicate that gene *CG32013* is likely not located on chromosome 4 and that BACR05L22 is chimeric with the gene *CG32013*-containing sequence being ligated adjacent to *CG17923* on the end of a chromosome 4-bearing fragment.

On the basis of these findings and other PCR tests, we conclude that *Df(4)M101-62f* is deleted or disrupted for the following genes: *CG17923*, *CG32011*, *CG32010*, *CG32009*, *CG17245* (*plexB*), *CG2125* (*ci*), *CG2168* (*RpS3A*), *CG17964* (*pan*), *CG32005*, *CG32008* (*Ank*), *CG32000*, *CG32006*, *CG31997*, *CG2219*, *CG2316*, *CG31998*, *CG1587* (*Crk*), *CG31999*, *CG1629* (*yellow-h*), and *CG1674*. This group defines region A of our complementation map of chromosome 4 (Fig. 3).

Df(4)B2-2D

This deficiency is the largest terminal deletion of chromosome 4 (Fig. 2b). The proximal breakpoint of the deletion is located at cytological position 102B2, which

we have located between the most distal exon of *CG1748* (*RhoGAP102A*) and the most proximal exon of *CG2380* (*NfI*). This deletion consists of the following genes: *CG2380* (*NfI*), *CG2381* (*Syt7*), *CG1836* (*Rad23*), *CG2177*, *CG2165* (*BEST:CK01140*), *CG1710* (*Hcf*), *CG2052*, *CG2041* (*Igs*), *CG1495* (*CamKI*), *CG2009* (*bip2*), *CG11533*, plus those genes deleted in *Df(4)C1-10AT*. This group defines region B of our chromosome 4 complementation map (Fig. 3).

Df(4)C1-10AT

This deficiency is a terminal deletion of chromosome 4 (Fig. 2c). The proximal breakpoint is located at cytological position 102C2, between the most distal exon of *CG11533*, and the proximal exon of *CG1449* (*zfh2*). This deletion consists of the following genes: *CG1449* (*zfh2*), *CG1981* (*ThdI*), *CG1507* (*Pur-alpha*), *CG1970*, *CG1862* (*Ephrin*), *CG1909*, *CG11260*, *CG1922* (*oncut*), *CG1511* (*Eph*), *CG1901* (*mav*), *CG1732*, *CG9935*, *CG31992*, *CG1783*, *CG11360*, *CG1838* (*myoglianin*), *CG1464* (*ey*), plus those genes deleted in *Df(4)C3*. It is possible that *CG11533* may also be disrupted if the deletion removes the regulatory region of this gene. This group defines region C (Fig. 3).

Fig. 3 A summary map of chromosome 4 deletions. The chromosome has been divided into five regions designated as A, B, C, D, and E based on the breakpoint locations of five chromosome 4 deficiencies [*Df(4)M101-62f*, *Df(4)B2-2D*, *Df(4)C1-10AT*, *Df(4)C3*, and *Df(4)G*]. The breakpoints of two other deficiencies [*Df(4)J2* and *Df(4)O2*] are also shown, but they have not been used to partition

chromosome 4. The cytological locations of the breakpoints are indicated. The inset boxes show the locations of the deleted regions in more detail. The CG numbers refer to the gene identifiers from the annotated sequence Release 3.1 of the *Drosophila* genome (Adams et al. 2000; Celniker et al. 2002; Misra et al. 2002)

Df(4)J2

This deficiency is a terminal deletion. The proximal breakpoint has been located to a 10.8 kb region between the distal exon of *CG1838* (*myoglianin*) and the proximal exon of *CG1464* (*ey*) (Fig. 3). Cytological observations have shown the breakpoint to be located at 102D2 (Fig. 2d). This deficiency will not be used to define a complementation group on chromosome 4 because of its similarity to *Df(4)C3*.

Df(4)C3

This deficiency is a terminal deletion of chromosome 4. The proximal breakpoint is located at cytological position 102D2 (Fig. 2e), which we have positioned within the distal portion of the gene *CG32019* (*bt*). This deletion consists of the following genes: *CG32019* (*bt*), *CG1793* (*Arc70*), *CG11231*, *CG11153* (*Sox102F*), *CG11152*, *CG11148*, *CG2999* (*unc-13*), *CG11224*, *CG32021*, *CG10811* (*eIF-4G*), *CG11144* (*Glu-RA*), *CG32016*, *CG11093*, *CG11091* (*sphinx*), *CG11186* (*toy*), *CG11081* (*plexA*), *CG11077*, *CG11076*, *CG11154* (*ATPsyn-beta*), *CG18069* (*CamKII*), *CG32018*, *CG11064* (*RfaBP*), plus those deleted in *Df(3)G*. This group defines region D of the complementation map (Fig. 3).

Df(4)G

This deficiency is the most distal terminal deletion of chromosome 4 (Fig. 2f). In addition to the deletion, we found that this deficiency contains a second site mutation in the gene *CG32019* (*bt*). The proximal deletion breakpoint is located at cytological position 102E2, which we have positioned within a 4916 bp region between the last exon of *CG11064* (*RfaBP*), and the most proximal exon of *CG11062* (*activin-beta*). This deletion includes all distal genes, and thus consists of the following genes: *CG32019* (*bt*, due to a second site mutation), *CG11062* (*activin-beta*), *CG11049* (*sv*), *CG11059* (*calsyntenin-1*), *CG11027* (*Arf102F*), *CG11155*, *CG32017*, *CG17461* (*Kif3C*), *CG17743* (*pho*), *CG32015*, *CG18026* (*caps*). This group defines region E (Fig. 3).

Df(4)O2

This deficiency is an interstitial deletion located at 102D1–D4 (Fig. 2g). We have mapped the proximal breakpoint to a region within the distal half of the gene *CG32019* (*bt*). The distal breakpoint is located within a 12.6 kb region between the distal exon of *CG11091* (*sphinx*) and the proximal exon of *CG11186* (*toy*). The genetic data indicate that *Df(4)O2* is deficient for *CG11186* (*toy*), so it is likely that the regulatory region of *toy* is deleted (Kammermeier et al. 2001).

Df(4)M101-63a

This chromosomal aberration genetically fails to complement only *ci* recessive alleles and *RpS3A* [*Minute(4)101* or *CG2168*]. One breakpoint has been localized to a 263 bp region within the *RpS3A* gene, which accounts for the lack of complementation to *Minute(4)101* mutations. The sequences on either side of the breakpoint are still present in DNA from homozygous *Df(4)M101-63a* embryos, indicating that this aberration is not a deletion. Thus the lesion in *RpS3A* is due to either an insertion into the *RpS3A* gene, or an inversion with one breakpoint located in the *RpS3A* gene. We attempted to obtain sequence past the breakpoint by single-primer PCR and inverse PCR, but these attempts were unsuccessful. Cytological observations do not show any aberration in the banding pattern (Fig. 2h), indicating that the inverted or inserted region must be relatively small in cytological terms.

Discussion

The current annotation (Release 3.1) of the *Drosophila* genome indicates that chromosome 4 contains 82 putative gene transcripts, interspersed with repeat elements. We have used these sequences to design PCR primers for exons of these genes. Amplification by PCR from homozygous deficient embryo DNA has allowed us molecularly to map the breakpoints of eight chromosome 4 deficiencies. The putative genes have been partitioned to the five regions of the chromosome specified by the breakpoint locations of five of the eight deficiencies.

Chromosome 4 deficiencies *B2-2D*, *C1-10AT*, *C3*, *J2*, and *G* are all terminal deletions of chromosome 4. *Df(4)M101-62f* is a interstitial deletion of the proximal region of chromosome 4, and *Df(4)O2* is an interstitial deletion of chromosome 4 in the 102D region. *Df(4)M101-63a* is either an inversion of the chromosome with one breakpoint in *RpS3A*, or an insertion into this same region. One gene, *CG1748* (*RhoGAP102A*), could not be placed within any of the five groups. It lies within the approximately 35 kb region between the distal breakpoint of *Df(4)M101-62f* and the proximal break of *Df(4)B2-2D*.

We have demonstrated that there is an error in the Release 3.1 genome sequence in the region proximal to gene *CG17923*. This error was likely due to the sequencing of the BAC clone BACR05L22, which we have shown to have different restriction fragments than the same region of genomic DNA extracted from five strains, including the isogenized stock used to build the BAC library. The ability to amplify PCR products using primers specific to gene *CG32013* on DNA from nullo-4 embryos indicates that this putative gene is, in fact, located on a chromosome other than the fourth. Because of this error, there may be genes located in the region proximal to *CG17923* that have not been included in this study owing to their absence from the Release 3.1 sequence.

The molecular mapping of these deficiencies will be a valuable resource for *Drosophila* researchers working on

chromosome 4 genes. The inability to make a conventional genetic map owing to the lack of recombination on the fourth chromosome has retarded the investigation of chromosome 4 genes. Complementation mapping to the deficiencies studied here will allow researchers easily to delimit a mutation to one specific block of genes. The chromosome 4 deficiency map will also be a very important resource for our own chromosome 4 mutational screen. The ability to locate the mutations to specific blocks of genes by complementation mapping will significantly reduce the number of DHPLC runs needed to find base substitutions, and will reduce the number of false identifications of mutants due to second site non-lethal base pair changes.

Acknowledgements We thank Scott Hanna for excellent technical assistance and advice, W. Gehring for providing the *Df(4)O2* and *Df(4)J2* stocks, Travis Thompson for providing the *D(4)C3* and *D(4)G* stocks and the Bloomington Stock Center for providing the *Df(4)M101-63a* stock. This work was funded with grants from the Canadian Institutes of Health Research (CIHR) and the Natural Sciences and Engineering Research Council (NSERC) of Canada.

References

- Adams MD, Celniker SE, Holt RA, Evans CA, Gocayne JD et al (2000) The genome sequence of *Drosophila melanogaster*. *Science* 287:2185–2195
- Ahmad K, Golic K (1998) The transmission of fragmented chromosomes in *Drosophila melanogaster*. *Genetics* 148:775–792
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *J Mol Biol* 215:403–410
- Bridges CB (1935) The mutants and linkage data of chromosome four of *Drosophila melanogaster*. *Biol Zh (Moscow)* 4:401–420
- Carmena M, Gonzalez C (1995) Transposable elements map in a conserved pattern of distribution extending from beta-heterochromatin to centromeres in *Drosophila melanogaster*. *Chromosoma* 103:676–684
- Celniker SE, Wheeler DA, Kronmiller B, Carlson JW, Halpern A et al (2002) Finishing a whole-genome shotgun: release 3 of the *Drosophila melanogaster* euchromatic genome sequence. *Genome Biol* 3:RESEARCH0079. Epub
- Hochman B (1976) The fourth chromosome of *Drosophila melanogaster*. In: Ashburner M, Novitski E (eds) *The genetics and biology of Drosophila*. Academic, London, pp 903–928
- Hoskins RA, Nelson CR, Berman BP, Laverty TR, George RA et al (2002) Annotation of the *Drosophila melanogaster* euchromatic genome: a systematic review. *Genome Biol* 3:RESEARCH0083. Epub
- James TC, Eissenberg JC, Craig C, Deitrich V, Hobson A, Elgin SC (1989) Distribution patterns of HP1, a heterochromatin-associated nonhistone chromosomal protein of *Drosophila*. *Eur J Cell Biol* 50:170–180
- Kaminker JS, Bergman CM, Kronmiller B, Carlson J, Svirskaas R, Patel S, Frise E, Wheeler DA, Lewis SE, Rubin GM, Ashburner M, Celniker SE (2002) The transposable elements of the *Drosophila melanogaster* euchromatin: a genomics perspective. *Genome Biol* 3:RESEARCH0084. Epub
- Kammermeier L, Leemans R, Hirth F, Flister S, Wenger U, Walldorf U, Gehring W, Reichert H (2001) Differential expression and function of the *Drosophila Pax6* genes *eyeless* and *twin of eyeless* in embryonic central nervous system development. *Mech Dev* 103:71–78
- Kramps T, Peter O, Brunner E, Nellen D, Froesch B, Chatterjee S, Murone M, Züllig S, Basler K (2002) Wnt/Wingless signaling requires BCL9/legless-mediated recruitment of pygopus to the nuclear β -catenin-TCF complex. *Cell* 109:47–60
- Larsson J, Chen D, Rasheva V, Rasmuson-Lestander A, Pirrotta V (2001) Painting of fourth, a chromosome-specific protein in *Drosophila*. *Proc Natl Acad Sci U S A* 98:6273–6278
- Locke J, McDermid H (1993) Analysis of *Drosophila* chromosome four using pulsed field gel electrophoresis. *Chromosoma* 102:718–723
- Locke J, Howard LT, Aippersbach N, Podemski L, Hodgetts RB (1999) The characterization of DINE-1, a short, interspersed repetitive element present on chromosome 4 and in the centric heterochromatin of *Drosophila melanogaster*. *Chromosoma* 108:356–366
- Locke J, Podemski L, Aippersbach N, Kemp H, Hodgetts R (2000) A physical map of the polytenized region (101EF–102F) of chromosome 4 in *Drosophila melanogaster*. *Genetics* 155:1175–1183
- Miklos GL, Yamamoto MT, Davies J, Pirrotta V (1988) Microcloning reveals a high frequency of repetitive sequences characteristic of chromosome 4 and the beta-heterochromatin of *Drosophila melanogaster*. *Proc Natl Acad Sci U S A* 85:2051–2055
- Misra S, Crosby MA, Mungall CJ, Matthews BB, Campbell KS et al (2002) Annotation of the *Drosophila melanogaster* euchromatic genome: a systematic review. *Genome Biol* 3:RESEARCH0083. Epub
- Pimpinelli S, Berloco M, Fanti L, Dimitri P, Bonaccorsi S, Marchetti E, Caizzi R, Caggese C, Gatti M (1995) Transposable elements are stable structural components of *Drosophila melanogaster* heterochromatin. *Proc Natl Acad Sci U S A* 92:3804–3808
- Sun F-L, Cuaycong M, Craig C, Wallrath L, Locke J, Elgin SC (2000) The fourth chromosome of *Drosophila melanogaster*: interspersed euchromatic and heterochromatic domains. *Proc Natl Acad Sci U S A* 97:5340–5345
- Wallrath LL, Elgin SC (1995) Position effect variegation in *Drosophila* is associated with an altered chromatin structure. *Genes Dev* 9:1263–1277
- Wallrath LL, Guntur VP, Rosman LE, Elgin SC (1996) DNA representation of variegating heterochromatic P-element inserts in diploid and polytene tissues of *Drosophila melanogaster*. *Chromosoma* 104:519–527