UC Irvine

UC Irvine Previously Published Works

Title

Estimate of late 1974 stratospheric concentration of gaseous chlorine compounds (CIX)

Permalink

<https://escholarship.org/uc/item/51n216dm>

Journal

Geophysical Research Letters, 2(6)

ISSN

0094-8276

Authors

Cicerone, RJ
Stedman, DH
Stolarski, RS

Publication Date

1975

DOI

10.1029/GL002i006p00219

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

ESTIMATE OF LATE 1974 STRATOSPHERIC CONCENTRATION
OF GASEOUS CHLORINE COMPOUNDS (ClX)

R. J. Cicerone and D. H. Stedman

Department of Atmospheric and Oceanic Science
The University of Michigan, Ann Arbor, Michigan 48105

R. S. Stolarski

NASA Johnson Space Center, Houston, Texas 77058

Abstract. We estimate the total concentration [ClX] versus altitude of Cl-containing molecules, principally HCl, Cl and ClO. The predicted present stratospheric concentrations appear to be due to photolysis near 30 km of man-made CF_2Cl_2 and CFCI_3 , and CCl_4 from a ground-level source, man-made or natural. Based on preliminary measurements of CH_3Cl at ground we estimate its contribution to stratospheric ClX through the major known loss mechanism for CH_3Cl , abstraction by OH. Ground-level sources of HCl and Cl_2 are large but tropospheric processes preclude significant impact on the stratosphere. Our computed profiles are consistent with first HCl concentrations measured by Lazrus et al. [1974, 1975] in the 15-27 km region.

Molina and Rowland [1974a; 1975] have proposed that the increasing use of man-made chlorofluoromethanes may represent a serious threat to atmospheric O_3 . Subsequent studies by Cicerone et al. [1974], Crutzen [1974], and Wofsy et al. [1975] have not refuted this proposal; instead they have offered additional documentation and quantitative estimates of the size and timetable of the effect on O_3 . The potential seriousness of this problem might require this increase to cease before definitive proof of O_3 depletion can be obtained. Therefore, we focus attention on some key intermediate species for which measurements are needed. In particular, we estimate the concentration of ClX (Cl + HCl + ClO) now in the stratosphere due to several sources. We show that ground-level ClX or Cl_2 sources (natural or anthropogenic) do not contribute appreciably to the stratosphere. With certain plausible assumptions we estimate the present stratospheric ClX profile, to be used as a baseline against which to assess future measurements. Thus, future stratospheric increases of ClX can be estimated with known rates of injection by photodissociation of chlorofluoromethanes or other chlorocarbons or by explosive volcanoes. Reliable measurements of ClX are obviously necessary and preferred over theoretical estimates but available data is sketchy and scarce.

Ground-level sources of HCl and Cl_2 include the conversion of sea-salt aerosol chlorides to HCl, volcanic and fumarolic emissions, diffusion through the earth's crust, and industrial processes. These have been discussed by many authors including Eriksson [1960], Valach [1967], Duce

[1969], Chesselet et al. [1972], Stolarski and Cicerone [1974], Ryan and Mukherjee [1974], and Wofsy and McElroy [1974]. The latter two references also concluded that HCl is by far the dominant Cl-containing form (barring organic compounds) in the troposphere, that is, $[\text{HCl}] = [\text{ClX}]$, where [] denotes concentration (cm^{-3}). Ground-level and low altitude measurements by Junge [1957], Duce et al. [1965], Duce [1969] and Chesselet et al. [1972] cluster around an HCl volume mixing ratio $f(\text{HCl}) = 10^{-9}$ but these may be upper limits because each apparatus may have permitted particle-to-gas conversion in its inlet system [Stedman et al., 1975]. We note that Farmer et al. [1974] reported from ground-level spectral observations that $f(\text{HCl}) < 3 \times 10^{-10}$ averaged over the lowest scale volume.

Measurements of HCl at all tropospheric altitudes are not available but Ryan and Mukherjee [1974] and Stedman et al. [1975] show that the tropospheric HCl profile should be proportional to the H_2O vapor profile. The H_2O vapor mixing ratio, $f(\text{H}_2\text{O})$, at 10 km is 10^{-3} to 10^{-2} times its ground-level value of 10^{-2} [U.S. Standard Atmosphere Supplements, 1966]. It decreases further between 10 km and a nominal tropopause height, 15 km. Thus, $f(\text{HCl})$ at 10 km is no more than 10^{-11} even if $f(\text{HCl})$ at ground is as high as 10^{-9} . Just as we have an H_2O -dry stratosphere (observations show $f(\text{H}_2\text{O}) \approx 3$ ppm; Mastenbrook [1974]) we would also expect a "ClX-dry" stratosphere ($f(\text{ClX}) < 10^{-11}$) if the only sources of ClX were at ground.

Ryan and Mukherjee's conclusion that tropospheric HCl is controlled by H_2O is based on their analysis of microscopic or molecular processes involving the thermodynamics of $\text{HCl}(\text{H}_2\text{O})_n$ polymers. Stedman et al. [1975] have reached the same conclusion for several highly soluble gases through an argument that does not invoke the existence of such polymers. They assumed that HCl gas and water vapor diffuse onto water droplets at rates proportional to their respective concentrations. From thermodynamic data they showed that the HCl vapor pressure around rain droplets is considerably less than ambient tropospheric HCl partial pressure, when the rain droplets form by condensation of all available H_2O and HCl in a given volume. Their model predicts a globally averaged Cl^- molarity of rainfall that is consistent with Duce's [1969] observation that rain droplets are considerably undersaturated in HCl. The theory is not so clear

for ice crystals, but from snow and ice composition data Stedman *et al.* argue that one can expect $f(\text{HCl})$ to decrease rapidly with altitude in the troposphere, just like $f(\text{H}_2\text{O})$.

The conclusion is that ground-level sources of HCl and Cl_2 are inconsequential for the stratosphere despite the large size (perhaps 10^{14} g/year or $10^{10} \text{ cm}^{-2} \text{ sec}^{-1}$) of the HCl source due to sea-salt conversion [Duce, 1969; Chesselet *et al.*, 1972; Ryan and Mukherjee, 1974]. Clearly if ground-level sources of HCl are this large the tropospheric lifetime of gaseous HCl must be relatively short. Wofsy [1974] and Wofsy and McElroy [1974] have estimated this lifetime to be 1-9 days. Chesselet *et al.* [1972] have inferred that HCl in overland air may have a lifetime < 1 day. H_2O control of HCl would give HCl the same lifetime (~ 1 week) as H_2O , through the rapid and dynamic processes of H_2O evaporation, condensation, freezing, sublimation and transport [Stedman *et al.*, 1975].

There is, of course, the possibility of other tropospheric sources of ClX, such as the release of Cl atoms from chlorocarbons, man-made or natural. Molina and Rowland [1974a, 1975] and Cicerone *et al.* [1974] and papers they cite show that CF_2Cl_2 and CFCl_3 probably do not release much Cl in the troposphere. CCl_4 produced at ground-level is also not likely to decompose in the troposphere. Chlorocarbons with C-H bonds should be more susceptible to attack by tropospheric OH. Lovelock [1974] has detected the presence of CHCl_3 , CH_3CCl_3 , and $\text{CHCl}=\text{CCl}_2$, albeit in small amounts, in the atmosphere. Other such compounds, e.g., CH_3Cl , may be present in larger amounts [R. Rasmussen, 1974, private communication], but unless they can provide tropospheric sources of HCl large enough to overcome the H_2O control of HCl (the required source strength would vary with altitude; near ground 10^{15} g/year might suffice) we expect a tropospheric ClX profile like that sketched in Figure 1, discussed later.

To estimate the present and future concentration profiles in the stratosphere we have considered the photolysis of CF_2Cl_2 , CFCl_3 and CCl_4 and the reaction of atmospheric OH with CH_3Cl produced at ground. Direct stratospheric injection of Cl_2 and HCl by explosive volcanoes and the projected Space Shuttle may also be important at specific times and places but estimates of temporal and global averages of injection rates [Stolarski and Cicerone, 1974; Ryan and Mukherjee, 1974; Cadle, 1975] yield relatively small masses compared to the man-made chlorofluoromethane sources.

We calculated the altitude profiles of ClX concentrations due to CF_2Cl_2 , CFCl_3 , CCl_4 and CH_3Cl in the following manner: for CF_2Cl_2 and CFCl_3 there appears little doubt that no natural sources exist. In addition, the anthropogenic release rates at ground-level are reasonably well-known [see e.g., Rowland and Molina, 1975; McCarthy, 1974]. For the time-dependent calculation of ClX concentrations we used the vertical profile of eddy-diffusion coefficient, K , deduced by Hunten [Hunten, 1975 and unpublished manuscript, 1974]. For photolysis rates of CF_2Cl_2 and CFCl_3 we used data from Rowland and Molina [1975], and procedures summarized by Cicerone

et al. [1974]. For the CCl_4 calculations we assumed a steady state profile and photolysis as the only sink [with photodissociation cross sections of Molina and Rowland, 1974b]. The identities and strengths of CCl_4 sources are not completely clear. The steady state assumption on CCl_4 is partially justified by arguments and data presented by Molina and Rowland [1974b]. If natural sources dominate one would expect steady state, but measured CCl_4 mixing ratios show an apparent increase from about 7×10^{-11} in 1972 [Lovelock *et al.*, 1973; Wilkniiss *et al.*, 1973] to about 1.2×10^{-10} in 1974 [Lovelock, 1974; Wilkniiss *et al.*, 1975]. It may not be safe to interpret these numbers as a uniform global increase, but if so, they probably imply a large anthropogenic (non U.S.) input since 1972. CCl_4 released from the U.S. alone has been reasonably steady for about 20 years and M. J. Molina [private communication, 1974] has shown that the U.S. output of the last 15 years can account for about 50% of the presently measured CCl_4 content of the atmosphere. This is contrary to Lovelock's [1974] interpretation of CCl_4 release statistics. Natural sources of CCl_4 are suggested by certain features of the measured concentrations [Lovelock, 1974].

Our calculations yield the ClX mixing ratio profiles of Figure 1. We show separately the contributions from CF_2Cl_2 and CFCl_3 together and from CCl_4 . A suggested tropospheric profile (consistent with H_2O control) due to ground-level sources of Cl-containing inorganic compounds is also indicated. We also show the HCl data points of Lazrus *et al.* [1974, 1975]. Two upper limits on HCl near 15 km by Farmer *et al.* [1974] are not shown because he has re-interpreted the data; he now estimates $f(\text{HCl})$ to be about 10^{-9} above 16 km, based on the total content in the zenith direction [Farmer, 1975]. A striking feature of our

Figure 1. Computed ClX (HCl + ClO + Cl) mixing ratios versus altitude for late 1974. The ClX contribution from CF_2Cl_2 and CFCl_3 photolysis was calculated from the known history of anthropogenic emissions, that from CCl_4 as described in the text. Recent HCl measurements are shown by X's. Nearly all ClX is HCl below 25 km. A nominal tropospheric HCl profile is also shown, reflecting H_2O control.

Figure 2. Computed ClX (HCl + ClO + Cl) mixing ratios versus altitude, due to a ground-level source of CH₃Cl, and the computed CH₃Cl profile for late 1974. The total ClX mixing ratio due to CF₂Cl₂, CFC1₃, CCl₄ and CH₃Cl is also shown, as are recent HCl measurements (X's) by Lazrus et al. [1974; 1975].

computed profiles is the positive upward gradient in ClX mixing ratio between 15 and 25 or 30 km. Such a profile shape is expected from a source of ClX in the middle stratosphere where CF₂Cl₂, CFC1₃ and CCl₄ photolyze [Molina and Rowland, 1974a,b; Cicerone et al., 1974; Crutzen, 1974; Wofsy et al., 1975]. This profile shape is shown clearly by the Lazrus et al. HCl values, and it indicates why HCl measurements near 15 km are difficult [Farmer et al., 1974]. It should be mentioned that ClX is almost all HCl below 20 km, according to present models [see e.g., Wofsy and McElroy, 1974; Crutzen, 1974; Molina and Rowland, 1974b]. The ClX due to CCl₄ is from 3 Cl atoms removed from each CCl₄ molecule, and a ground-level CCl₄ mixing ratio fixed at 10⁻¹⁰. We infer that the ground flux of CCl₄ is about 6 x 10⁵ cm⁻² sec⁻¹ over the globe. Almost all of these molecule enter the stratosphere.

A Cl concentration of 10⁵ cm⁻³ at 75 km was inferred from Cl⁻ measurements by a rocket-borne mass spectrometer between 71 and 77 km [Arnold and Krankowsky, 1971]. This mesospheric Cl concentration is consistent with our profiles in Figures 1 and 2 but it is also possible that meteoric sources contribute to the mesosphere. Further, the inference of Cl concentrations from Cl⁻ measurements is risky: the Cl⁻ might have been from contaminants, charge-exchange and recombination rate constants are poorly known, NO₃⁻ should not exchange charge with Cl as they assumed, and a reaction they neglected, H + Cl⁻ → HCl + e, has a constant of 1 x 10⁻⁹ cm³ sec⁻¹ [Ferguson, 1975, private communication]. Finally, it should be noted that the Lazrus et al. data were gathered with chemically impregnated filter paper. Efficiency of collection of gaseous HCl may be less than 100%, further calibration may demand that these HCl values be increased [Lazrus, 1975, private communication]. Possible particle-to-gas conversion could cause opposing errors.

In Figure 2 we display our computed vertical profile of CH₃Cl, along with its contribution of stratospheric ClX and the total ClX to be expected from CH₃Cl and the CF₂Cl₂, CFC1₃ and CCl₄ sources, and the Lazrus et al. data. Our computations for CH₃Cl are based on an assumed steady state vertical profile of CH₃Cl with a mixing ratio of 4 x 10⁻¹⁰ at ground level, as suggested by recent measurements [R. Rasmussen, 1974; J. E. Lovelock, 1975, private communication]. The only significant sink for CH₃Cl appears to be the abstraction reaction CH₃Cl + OH → CH₂Cl + H₂O with immediate CH₂Cl decomposition by uncertain chemical reactions probably involving O₂. The rate for this reaction was estimated as 4.8 x 10⁻¹² exp(-1400/T) from Davis' et al. [1975] room temperature rate and a (guessed) activation energy of 2.8 kcal. For an OH profile we took model results from McElroy et al. [1974]. The present simple model implies a globally averaged ground-level source of 2.6 x 10⁸ CH₃Cl molecules cm⁻² sec⁻¹. This source is about 10 x the direct industrial production rate in the U.S. in 1972. The flux of CH₃Cl upward through the 15 km level is about 1.7 x 10⁶ cm⁻² sec⁻¹ and leads to more HCl than has been measured when it is added to the sources for Figure 1. Presumably, a natural source exists for CH₃Cl, but its identity is unknown. Lovelock et al. [1973] and Zafirou [1974] have discussed possible natural sources of CH₃I.

Substantial agreement appears between the Lazrus et al. HCl measurements and the ClX prediction in Figure 1, where the CF₂Cl₂, CFC1₃ and CCl₄ sources are included. Without CCl₄, our model predicts about one-third as much ClX as observed by Lazrus et al. Similarly, calculations for CF₂Cl₂ and CFC1₃ alone, by Wofsy et al. [1975], found one-half as much ClX as Lazrus et al. If the Figure 1 total ClX profile is taken as a guide one would expect ClO concentrations of about 3 x 10⁷ cm⁻³ between 30 and 35 km in late 1974 according to present schemes of ClX chemistry. Figure 2 shows that the uncertain CH₃Cl source implied by Rasmussen's and Lovelock's measurements may have carried as many Cl atoms to the 30 km level as CF₂Cl₂, CFC1₃ and CCl₄ had by 1974. It must be emphasized, however, that CF₂Cl₂ and CFC1₃ released after the mid-1960's will soon result in more Cl atoms injected near 30 km than the (presumed) steady state CCl₄ and CH₃Cl contributions. If CF₂Cl₂ and CFC1₃ emissions were kept constant at 1973 levels, their tropospheric mixing ratios would increase by about a factor of 10 and stratospheric ClX mixing ratios to about 4 x 10⁻³ when steady state were reached, in 50 to 100 years [Crutzen, 1974; Cicerone et al., 1975; Rowland and Molina, 1975; Wofsy et al., 1975].

Acknowledgments. We thank E. E. Ferguson, A. L. Lazrus, M. J. Molina, R. S. Narcisi and F. S. Rowland for helpful discussion, NASA for financial support through Grant NGR-23-006-616 and the Climatic Impact Assessment Program for NSF Grant DES 74-11478.

References

- Arnold, F. and D. Krankowsky, Negative ions in the lower ionosphere: A comparison of a model computation and a mass-spectrometer measurement, *J. Atmos. Terr. Phys.*, **33**, 1693-1702, 1971.
- Cadle, R. D., Volcanic emissions of halides and sulfur compounds to the troposphere and stratosphere, *J. Geophys. Res.*, **80**, in press, 1975.
- Chesselet, R., J. Morelli and P. Buat-Menard, Some aspects of the geochemistry of marine aerosols, in *The Changing Chemistry of the Oceans*, ed. D. Dyrssen and D. Jagner, Wiley, New York, pp. 93-114, 1972.
- Cicerone, R. J., R. S. Stolarski and S. Walters, Stratospheric ozone destruction by man-made chlorofluoromethanes, *Science*, **185**, 1165-1167, 1974.
- Crutzen, P. J., Estimates of the possible future ozone reductions from continued use of fluorochloro-methanes (CF_2Cl_2 , CFCl_3), *Geophys. Res. Lett.*, **1**, 205-208, 1974.
- Davis, D., P. Hogan and Y. Oh, University of Maryland Gas Kinetics Laboratory Report No. 2, 1975.
- Duce, R. A., On the source of gaseous chlorine in the marine atmosphere, *J. Geophys. Res.*, **74**, 4597-4599, 1969.
- Duce, R., J. Winchester and T. Van Nahl, Iodine, bromine and chlorine in the Hawaiian marine atmosphere, *J. Geophys. Res.*, **70**, 1775-1800, 1965.
- Eriksson, E., The yearly circulation of chloride and sulfur in nature: meteorological, geochemical and pedological implications, *Tellus*, **12**, 63-109, 1960.
- Farmer, C. B., Spectroscopic detection of chlorine compounds in the stratosphere, paper presented at 4th Conf. on CIAP, Cambridge, Massachusetts, Feb. 7, 1975.
- Farmer, C., O. Raper, R. Toth and R. Schindler, Recent results of aircraft infrared observations of the stratosphere, *Proc. Third Conf. on CIAP, DOT-TSC-OST-74-15*, 234-245, 1974.
- Hunten, D. M., Residence times of aerosols and gases in the stratosphere, *Geophys. Res. Lett.*, **2**, 26-28, 1975.
- Junge, C., Chemical analysis of aerosol particles and of gas traces on the island of Hawaii, *Tellus*, **9**, 528-537, 1957.
- Lazrus, A., B. Gandrud, W. Sedlacek and R. Woodard, Measurements of stratospheric halogens, paper presented at Amer. Geophys. Union Fall Meeting, San Francisco, Dec. 12, 1974.
- Lazrus, A., B. Gandrud, R. Woodard and W. Sedlacek, Measurements of stratospheric halogens and nitric acid, paper presented at 4th Conf. on CIAP, Cambridge, Massachusetts, Feb. 7, 1975.
- Lovelock, J., Atmospheric halocarbons and stratospheric ozone, *Nature*, **252**, 292-294, 1974.
- Lovelock, J., R. Maggs, and R. Wade, Halogenated hydrocarbons in and over the Atlantic, *Nature*, **241**, 194-196, 1973.
- Mastenbrook, H. J., Water-vapor measurements in the lower stratosphere, *Can. J. Chem.*, **52**, 1527-1531, 1974.
- McCarthy, R. L., Fluorocarbons in the atmosphere, *Trans. Am. Geophys. Union*, **56**, 1153, 1974.
- McElroy, M. B., S. C. Wofsy, J. E. Penner and J. C. McConnell, Atmospheric ozone: Possible impact of stratospheric aviation, *J. Atmos. Sci.*, **31**, 287-303, 1974.
- Molina, M. J. and F. S. Rowland, Stratospheric sink for chlorofluoromethanes: Chlorine atom catalyzed destruction of ozone, *Nature*, **249**, 810-812, 1974a.
- Molina, M. J. and F. S. Rowland, Predicted present stratospheric abundances of chlorine species from photodissociation of carbon tetrachloride, *Geophys. Res. Lett.*, **1**, 309-312, 1974b.
- Rowland, F. S. and M. J. Molina, Chlorofluoromethanes in the environment, *Rev. Geophys. Space Phys.*, **13**, in press, 1975.
- Ryan, J. A. and N. Mukherjee, Estimate of tropospheric HCl cycle, MDAC-WD paper No. 2290, McDonnell-Douglas Company, Huntington Beach, California, 1974.
- Stedman, D., W. Chameides and R. Cicerone, The vertical distribution of soluble gases in the troposphere, submitted to *Geophys. Res. Lett.*, 1975.
- Stolarski, R. S. and R. J. Cicerone, Stratospheric chlorine: A possible sink for ozone, *Can. J. Chem.*, **52**, 1610-1615, 1974.
- U. S. Atmosphere Supplements, U. S. Government Printing Office, Washington, D. C., 1966.
- Valach, P., The origin of the gaseous form of natural atmospheric chlorine, *Tellus*, **9**, 509-516, 1967.
- Wilkniss, P., J. Swinnerton, D. Bressan, R. Lamontagne and R. Larson, CO , CCl_4 , Freon-11, CH_4 and Rn-222 concentrations at low altitude over the Arctic Ocean in January, 1974, *J. Atmos. Sci.*, **32**, 158-162, 1975.
- Wilkniss, P., R. Lamontagne, R. Larson, J. Swinnerton, C. Dickson and T. Thompson, Atmospheric trace gases in the southern hemisphere, *Nature*, **245**, 45-47, 1973.
- Wofsy, S., M. McElroy and N. Sze, Freon consumption: Implications for atmospheric ozone, *Science*, **187**, 535-537, 1975.
- Wofsy, S. C. and M. B. McElroy, HO_x , NO_x and ClO_x : Their role in atmospheric photochemistry, *Can. J. Chem.*, **52**, 1582-1591, 1974.
- Wofsy, S. C., On gaseous chlorine and the role of HCl-water aerosols in the Earth's lower atmosphere, unpublished manuscript, Harvard University, 1974.
- Zafiriou, O. C., Photochemistry of halogens in the marine atmosphere, *J. Geophys. Res.*, **79**, 2730-2732, 1974.

(Received March 10, 1975

Accepted April 3, 1975)