

UCLA

UCLA Previously Published Works

Title

The role of APOBEC3-induced mutations in the differential evolution of monkeypox virus.

Permalink

<https://escholarship.org/uc/item/52g0g0pj>

Journal

Virus Evolution, 9(2)

ISSN

2057-1577

Authors

Li, Xiangting
Habibipour, Sara
Chou, Tom
[et al.](#)

Publication Date

2023

DOI

10.1093/ve/vead058

Copyright Information

This work is made available under the terms of a Creative Commons Attribution-NonCommercial License, available at <https://creativecommons.org/licenses/by-nc/4.0/>

Peer reviewed

The role of APOBEC3-induced mutations in the differential evolution of monkeypox virus

Xiangting Li,^{1,†,‡} Sara Habibipour,^{2,†} Tom Chou,^{1,3,*,§} and Otto O. Yang²

¹Department of Computational Medicine, UCLA, Los Angeles, CA, United States, ²Departments of Medicine and Microbiology, Immunology, and Molecular Genetics, UCLA, Los Angeles, CA, United States and ³Department of Mathematics, UCLA, Los Angeles, CA, United States

[†]These authors contributed equally to this work.

[‡]<https://orcid.org/0000-0001-5238-7364>

[§]<https://orcid.org/0000-0003-0785-6349>

*Corresponding author: E-mail: tomchou@ucla.edu

Abstract

Recent studies show that newly sampled monkeypox virus (MPXV) genomes exhibit mutations consistent with Apolipoprotein B mRNA Editing Catalytic Polypeptide-like3 (APOBEC3)-mediated editing compared to MPXV genomes collected earlier. It is unclear whether these single-nucleotide polymorphisms (SNPs) result from APOBEC3-induced editing or are a consequence of genetic drift within one or more MPXV animal reservoirs. We develop a simple method based on a generalization of the General-Time-Reversible model to show that the observed SNPs are likely the result of APOBEC3-induced editing. The statistical features allow us to extract lineage information and estimate evolutionary events.

Keywords: APOBEC3; monkeypox virus; synonymous mutations; phylogenetics; DNA substitution model.

Introduction

Monkeypox (MPX), or Mpox, is a viral zoonotic disease that has been reported sporadically in western and central Africa. It is caused by the orthopoxvirus MPXV, which is related to variola virus, the past cause of the smallpox pandemic. MPX resides in a yet unidentified animal reservoir, and whether it has achieved endemicity in humans is unclear (EUCDC, 2019; USCDC, 2022; WHO, 2022).

In 1970, the first human case of monkeypox was reported in the Democratic Republic of the Congo (DRC) (Bremner et al., 1980). Since then, it has been suggested that MPX may have become endemic in the DRC and spread to several other Central and West African countries (Bunge et al., 2022). Outside of Africa, there were sporadic cases of MPX reported from 2003 to 2021 until the recent 2022 worldwide outbreak (EUCDC, 2019; USCDC, 2022; WHO, 2022). These previous sporadic cases were shown to be related to travel to Africa (Bunge et al., 2022). While cases in the most recent outbreak did not involve travel to or from Africa, the sequences from the 2022 outbreak were found to be genetically clustered with 2018–2019 cases (Isidro et al., 2022). Moreover, O'Toole and Rambaut first reported signs of apolipoprotein B mRNA Editing Catalytic Polypeptide-like3 (APOBEC3)-mediated editing in those sequences (O'Toole and Rambaut, 2022a; O'Toole and Rambaut, 2022b; O'Toole and Rambaut, 2022c).

APOBEC3 is a group of human enzymes of the innate immune system capable of inhibiting certain types of viruses through

deaminating cytosine to uracil, causing a G-to-A mutation in the complement strand when it is synthesized (Sadeghpour et al., 2021). Most human APOBEC3 molecules tend to deaminate TC dinucleotides, except APOBEC3G, which prefers CC dinucleotides (Beale et al., 2004; Yu et al., 2004; Stenglein et al., 2010; Hultquist et al., 2011; Burns et al., 2013). Compared to MPXV sequences collected in the 1970s, the single-nucleotide polymorphisms (SNPs) in the 2022 outbreak were found to be biased for the pattern 'TC → TT' and 'GA → AA' (O'Toole and Rambaut, 2022a; O'Toole and Rambaut, 2022b; O'Toole and Rambaut, 2022c). O'Toole and Rambaut hypothesized that the observed SNPs in the 2022 outbreak were the result of APOBEC3-induced editing and that they are evidence of within-human evolution of MPXV (O'Toole and Rambaut, 2022a; O'Toole and Rambaut, 2022b; O'Toole and Rambaut, 2022c).

Subsequent phylogenetic analysis of the genomes collected between 2017 and 2022 revealed significantly high frequencies of APOBEC3-associated SNPs, further supporting the hypothesis of within-human evolution of MPXV (Isidro et al., 2022; Gigante et al., 2022). In addition, all the 2022 sequences appeared to cluster together and little evidence of APOBEC3-related mutations was found prior to 2016. In the context of those studies, the APOBEC3-associated SNPs were defined as specific 'dinucleotide mutation' of the form 'TC → TT' and 'GA → AA.'

Because DNA evolution is a complex process involving mutation and selection, there may be a number of different explanations for the higher fraction of APOBEC3-associated SNPs observed

in later sequences; we wish to quantify the influence of APOBEC3 editing by factoring out the effects of these other mechanisms. If the average mutation rates *per site* of the TC → TT type are intrinsically higher than other types of mutations, independent of APOBEC3 activity, then a high fraction of APOBEC3-associated SNPs could be explained without invoking APOBEC3 involvement. Here, we take ‘intrinsically’ to mean APOBEC3-like mutations (You, 2000; Hussein, 2005; Budden and Bowden, 2013) caused by mutagens unrelated to APOBEC3 such as UV radiation that induce ‘C → T’ mutations (Drouin and Therrien, 1997). Besides the mutagens, the observed mutations themselves may confer higher fitness to MPXV, for instance, by enabling the virus to evade immune surveillance and infect cells more effectively and by exploiting tRNA abundance via codon usage biases (Parvathy, Udayasuriyan and Bhadana, 2021). Mutations that are indistinguishable from those induced by APOBEC3 activity can thus be selected even in the absence of APOBEC3.

To distinguish APOBEC3-induced mutations from other high-frequency mutations, we can establish a baseline by investigating early (pre-2016) MPXV genomes. However, time-varying mutation rates, uncertainty in sampling time points, and a limited number of early sequences complicate ancestor sequence reconstruction and mutation event timing. To circumvent these uncertainties, we develop a method to quantify ‘relative’ mutation rates and find (1) that even in genomes collected before 2016, the TC → TT mutation rate relative to other types is higher than average and (2) that TC → TT mutations are even more abundant in genomes collected after 2016. Mutagens induced by UV radiation may account for the higher mutation rate of TC → TT in the pre-2016 genomes, while our hypothesis is that the acceleration of TC → TT mutations in the later (post-2016) genomes arise from human-specific APOBEC3 editing.

To factor out selection, we examine the evolution of synonymous SNPs, which do not alter the amino acid sequence and should have minimal influence on the viral fitness. However, even within different synonymous mutations, codon usage bias may arise; a previous study on the codon usage bias of monkeypox virus collected before 2016 concluded that the genome-wide frequency of any given nucleotide at the third codon position was not dependent on nucleotides at the first or second position (Karamathil et al., 2018), suggesting that natural selection did not favor particular codons in the MPXV genomes. We count the number of synonymous mutations and synonymous APOBEC3-relevant mutations by first identifying all mutations at all sites, then identifying specific mutations that are synonymous. We also consider the preceding nucleotide, which may or may not be within the same codon, to count the number of APOBEC-relevant dinucleotide mutations such as TC → TT.

Our model allows for variation in the ‘raw’ or absolute mutation rate across different genomic sites, lineages, and generations. For example, one specific lineage of the virus may acquire a beneficial mutation at a specific time, allowing the virus to replicate faster and thus increase the subsequent mutation rates in this faster-growing lineage. However, to quantitatively understand the evolution of synonymous SNPs, we make a key assumption that the ‘ratio’ of rates of any two types of mutation is the same across all sites, lineages, and across time even though the raw rates may vary across sites, lineages, and time. Our method is designed to capture common patterns of ‘relative’ mutation rates shared by all members of a lineage sharing a common ancestor.

The assumption of constant mutation rate ratios underpins Markov models of DNA nucleotide substitution, such as the JC69

(Jukes and Cantor, 1969) and GTR (Tavaré, 1986) models. Evolution models used in maximum likelihood and Bayesian inference phylogenetic methods also implicitly rely on the constant rate ratio assumption (Felsenstein, 1981; Felsenstein, 2001; Nguyen et al., 2014) as typical phylogenetic methods are derived based on relative rates of mutations and do not involve absolute time. Thus, our key assumption of constant relative mutation rates is commonly used. Under such an assumption, the number of the observed mutations of a given type will be proportional to the number of another type in the same way across genomes from different lineages and generations. On the other hand, molecular clock theory posits that the ‘total’ number of neutral mutations increases linearly with elapsed physical time or number of generations, without making any assumptions about the relative rates or probabilities of the different mutation types.

This similarity inspires us to integrate phylogenetic models with principles derived from molecular clocks. We aim to correlate the number of synonymous mutations with a conceptualized relative time scale. We refer to the linear proportionality of the number of a specific mutation to the total number of synonymous mutations as the ‘relative molecular clock’.

In the next (Materials and Methods) section, we formalize and validate our ideas by first introducing a complete DNA substitution model with raw rates and absolute time. The raw rates ν will cancel each other when we consider ratios of mutations in our final results. We also introduce an additional assumption of independence between sites, which allows us to easily model the distribution of mutations using a simple binomial distribution and perform statistical hypothesis testing based on this distribution. In particular, our method does not require any knowledge of the ancestor sequence.

Using representative MPXV genomes available through the National Center for Biotechnology Information (NCBI), we demonstrate that our relative molecular clocks can describe the distribution of the fraction of APOBEC3-associated SNPs to the total number of synonymous SNPs. The genomes collected before and after 2016 clearly separate into two groups, with each group exhibiting a shared relative rate of APOBEC3-associated mutations. We surmise that these two groups of genomes were derived from two different evolutionary environments. The genomes collected before 2016 were likely direct descendants from the animal reservoir, while the genomes collected after 2016 were likely derived from the human host. Given that MPXV jumped from the animal reservoir to humans (Huang et al., 2022), we can infer statistical properties such as the number of synonymous mutations between the common ancestor of the post-2016 genomes (which has not yet undergone APOBEC3 editing) and a reference genome. This statistically inferred common ancestor can be older than the last common ancestor of the same group and provide a better estimate of the time of zoonotic MPXV transmission. Based on the inferred relative rates and additional independence assumption, we also developed a statistical test to determine whether a given genome can be considered as the common ancestor of the post-2016 genomes. Evolution of other viruses with APOBEC3-induced editing can also be analyzed using the models presented here.

Materials and Methods

DNA evolution model

We construct a relative clock by using the number of synonymous SNPs. To relate our method to both molecular clock and phylogenetic models, we start with a formal substitution model with




Figure 1. Schematic of the stochastic evolution model. (A) There are different potential synonymous mutation sites along the genome. Some potential mutations are APOBEC3-induced (red), some are reverse-APOBEC3-induced (blue), and some are neither (green). The transitions (from open circles to hash-filled circles) are indicated by arrows. Transitions of each type occur independently at their type-specific transition rate. (B) An example of two lineages (i) and (j) arising from a common ancestor that shared a common evolutionary path until time $t^{(ij)}$. Each black dot represents a mutation. An open circle represents the time a lineage is sampled. The waiting time between two successive mutations are independent and randomly distributed with the same total rate $N\bar{v}$. The specific type of mutation is also randomly chosen according to the relative rates of the different mutation types. (C) Counts of mutations using the common ancestor as a reference. (D) Counting the numbers of mutations of lineage (i) at time $t^{(i)}$ using the lineage (j) at time $t^{(j)}$ as a reference. Since sequences share part of their evolutionary paths starting with the common ancestor, mutations α_1 and α_2 are not identified by using lineage (j) as a reference. The subsequent mutations of lineage (j) count toward $n^{(ij)}$. For completeness, we included all mutations instead of only synonymous ones in the schematic sample sequences in (C) and (D).

raw mutation rates and absolute time. For the sake of simplicity, we first consider a variant of the GTR model (Tavaré, 1986) that accounts for the local dinucleotide context and mutation rates that are homogeneous across the genome, all lineages, and all generations. We then discuss how relaxing these homogeneity assumptions, by proper rescaling of absolute time, can still lead to a constant-‘relative’-mutation-rate structure.

Single-site DNA evolution models typically use a 4×4 matrix whose elements describe substitutions among the four nucleotides. For example, the entry \mathbf{V}_{AC} might describe the $A \rightarrow C$ mutation rate, defined by the expected number of mutations per site per unit time. The rate matrix \mathbf{V} can be decomposed into the product of the mutation probabilities per site per replication cycle and the raw replication rate. The JC69 model assumes that each available mutation is equally likely to occur and uses a single parameter v that describes the overall mutation rate. Therefore, in JC69, $\mathbf{V} = v\mathbf{Q}$, where \mathbf{Q} is a normalized substitution matrix $\mathbf{Q}_{XY} = \frac{1}{3}$ if $X \neq Y$ and $\mathbf{Q}_{XX} = -1$. Similarly, the rate matrix \mathbf{V} in the GTR model can be decomposed into $\mathbf{V} = v\mathbf{Q}$, where v and \mathbf{Q} are both parameters of the model.

To reformulate the DNA evolution model to include local dinucleotide context, we extend \mathbf{Q} to a 16×16 matrix with entries describing the normalized rate of mutations between each pair of dinucleotides. In the following subsections, we allow v to be dependent on genomic sites x , lineage (i), and time t , while \mathbf{Q} is a constant matrix to be shared by all mutation sites, lineages, and generations. Under this generalization, the mutation rate matrix for lineage (i) at time t and site x is $\mathbf{V}^{(i)}(x, t) = v^{(i)}(x, t)\mathbf{Q}$. Because the

mutation rate separates from the constant substitution matrix \mathbf{Q} , the ratios of mutation rates are independent of the mutation rate prefactor $v^{(i)}(x, t)$.

Several factors can influence the substitution matrix, including the DNA replication and proofreading mechanisms, the environmental factors such as UV, pH, and temperature that may induce spontaneous base substitution, and other proteins that actively edit the DNA sequence, such as the APOBEC3 family of proteins. As an orthopoxvirus, MPXV carries its own replication proteins and replicates in the cytoplasm of the host cell (Peng et al., 2023). Within the human host, the environmental factors and host proteins are relatively constant. Therefore, we conclude that the ratio of mutation rates are mostly conserved, and our key assumption is reasonable.

Here, we focus primarily on synonymous SNPs. Historically, synonymous SNPs can be considered as neutral mutations and are often associated with the molecular clock (Fitch and Langley, 1976). However, selection pressure related to codon usage (Wallace, Airoidi and Drummond, 2013) can differentially affect synonymous mutations. Our model does not require neutrality, but it does require that mutations of the same type occur with similar rates relative to other mutations. Statistically, different synonymous mutations of the same pattern, e.g. $TA \rightarrow TT$, are considered as identical. This allows us to count the number of synonymous mutations of the same type and infer the relative rates of these mutations. Our consideration mostly involves DNA substitution models rather than codon substitution models by Goldman and Yang (1994); Yang (2014). We identify synonymous

dinucleotide mutations by the context along the whole sequence rather than only through synonymous sites.

Now suppose there is a common ancestor sequence S_0 in which there are a total of N possible synonymous mutations, i.e. roughly $N/3$ codons since most codons have three synonymous alternatives, a total of N_{+a} possible synonymous APOBEC3-induced mutations (TC \rightarrow TT and GA \rightarrow AA), and a total of N_{-a} possible synonymous reverse-APOBEC3 mutations (TT \rightarrow TC and AA \rightarrow GA). To quantify N and N_{+a} , each possible nucleotide change that retains the subsequent amino acid is counted as a separate synonymous mutation. As shown in Fig. 1(A), the basic idea is to assume that each mutation of the same type along the genome is independent and identically distributed with a type-specific rate.

Synonymous APOBEC3-induced mutations are defined as synonymous mutations that match the pattern TC \rightarrow TT or its reverse complement GA \rightarrow AA. The reverse-APOBEC3 synonymous mutations are defined as synonymous mutations that match the pattern TT \rightarrow TC or its complement AA \rightarrow GA. Together, these two types of synonymous mutations will be denoted ‘APOBEC3-relevant’ in the following sections. The remaining possible number of synonymous mutations $N_0 = N - N_{+a} - N_{-a}$ are non-APOBEC3-relevant.

We assume that all possible synonymous mutations of the same type $\sigma \in \{0, +a, -a\}$ have the same mutation rate v_σ and that each mutation occurs independently of others. For example, the mutation rate for synonymous APOBEC3-induced mutations is v_{+a} , and the mutation rate for synonymous reverse-APOBEC3 mutations is v_{-a} . The average mutation rate for synonymous non-APOBEC3-induced mutations is v_0 . The average mutation rate across all possible synonymous mutations is thus

$$\bar{v} = \frac{N_{+a}v_{+a} + N_{-a}v_{-a} + N_0v_0}{N}. \quad (1)$$

Table 1 summarizes the notation used in the following sections.

The different types of mutations invoked in our model are listed in Table 2.

Single-lineage evolution

First consider a single lineage of MPXV genomes evolving from the common ancestor S_0 starting at time 0. In terms of the synonymous mutations, the lineage observed at time t is described by a sequence $\alpha(t) = (\alpha_1, \dots, \alpha_{n(t)})$, where each $\alpha_k \in \{1, \dots, N\}$ labels the location (which also defines the mutation type) of the k^{th} (chronological) synonymous mutation, relative to some reference genome (e.g. S_0). Here, $n(t)$ describes the total number of synonymous mutations that have occurred up to time t . An example of two lineages $\alpha^{(i)}$ and $\alpha^{(j)}$ is shown in Fig. 1(B).

To simplify the analysis, we also adopt an *infinite sites assumption* (Ma et al., 2008) in which the total number $n^{(i)}(t)$ of synonymous mutations of lineage $\alpha^{(i)}$ satisfies $n^{(i)}(t) \ll N$. For the MPXV genomes we investigated, the number n of observed synonymous mutations is around 100, and the total number N of possible synonymous mutations is around 100,000. Therefore, we can safely assume that all observed mutations $\alpha_k^{(i)}$ occurs on different genomic sites for different lineages (i) and chronological order k . Multiple mutations and back mutations occurring on the same site are rare and neglected. We note that the infinite sites assumption is not fundamental to our calculation. To drop this assumption, we need to consider the relation between the number of *observed* mutations and the number of *actual* mutations. For example, if one back mutation occurs, the observed number of mutations will be two less than the actual number of mutations. Given the total number of actual mutations, one can calculate

Table 1. Notation used in our model and analysis.

Name	Symbol	Typical values for MPXV
No. of possible synonymous mutations (of a given type)	$N, (N_{\pm a})$	100,000
No. of observed synonymous mutations (of a given type)	$n, (n_{\pm a})$	100
Mutation rate of a single nucleotide (of a given type)	$v, (v_{\pm a})$	–
A specific mutation out of N possible mutations	α	$\{1, \dots, N\}$
No. of observed synonymous mutations up to time t	$n(t)$	100
Sequence of mutations of i^{th} lineage	$\alpha^{(i)}(t) = (\alpha_1^{(i)}, \dots, \alpha_{n(t)}^{(i)})$	–
k^{th} (chronological) mutation in lineage (i)	$\alpha_k^{(i)}$	$\{1, \dots, N\}$
No. of observed mutations of genome (i) relative to (j)	$n^{(ij)}$	100
Sampling time of lineage or genome (i)	$t^{(i)}$	–
Emergence time of last common ancestor of lineages (i), (j)	$t^{(ij)}$	–
No. of mutations shared by lineages (i) and (j)	$k^{(ij)}$	–

Table 2. A list of the different types of synonymous mutations considered in our model.

Mutation type	Symbol	Synonymous mutation type
Index	σ	$\{0, +a, -a\}$
APOBEC3-induced	n_{+a}	TC \rightarrow TT, GA \rightarrow AA
reverse-APOBEC3	n_{-a}	TT \rightarrow TC, AA \rightarrow GA
APOBEC3-relevant	$n_a = n_{+a} + n_{-a}$	TC \leftrightarrow TT, GA \leftrightarrow AA
synonymous	n	all types of synonymous mutations
non-APOBEC3-relevant	$n - n_a$	all but APOBEC3-relevant mutations
AC \rightarrow AT	–	AC \rightarrow AT, GT \rightarrow AT
A \rightarrow C	–	A \rightarrow C, T \rightarrow G

the expected number of observed mutations. Inversely, when the number of observed mutations is known, we can obtain an estimate of the number of actual mutations by the number that produces a matching expected number of observed mutations. In the current setting, the chances of back mutations and multiple mutations on the same nucleotide are small. Therefore, the difference between the number of observed mutations and the number of actual mutations is also very small.

The average synonymous mutation rate \bar{v} (Eq. 1) comes into play when we measure the number of mutations $n^{(i)}(t)$ of the lineage $\alpha^{(i)}$ at time t . The expected number of mutations is then given by

$$\mathbb{E}[n^{(i)}(t)] = N\bar{v}t, \quad (2)$$

and the expected number of mutations of type σ is given by $N\sigma v_\sigma t$. To obtain a probability distribution of different types of mutations, we assume that mutation events on different dinucleotide sites are independent, allowing for at least a dinucleotide level of site dependence.

This independent-site assumption is also widely adopted in the literature (Felsenstein, 1981; Felsenstein, 2001; Nguyen et al., 2014) and reflects the stochastic nature of molecular processes and the feature that DNA replication and proofreading manifest themselves locally at nucleotide sites. Under this independence assumption, we have the following conditional probability:

$$\mathbb{P}(\alpha^{(i)} = (\alpha_1^{(i)}, \dots, \alpha_n^{(i)}) | n^{(i)}(t) = n) = \prod_{k=1}^n \frac{N\sigma_k v_{\sigma_k}}{N\bar{v}}, \quad (3)$$

where $\sigma_k \in \{+, -, 0\}$ defines the type of the k^{th} mutation, which is implicitly defined by the sequence $(\alpha_1^{(i)}, \dots, \alpha_n^{(i)})$.

Now, let n_{+a} denote the number of synonymous APOBEC3-induced mutations (TC \rightarrow TT and GA \rightarrow AA) in the lineage α , as exemplified in Figure 1(C). Similarly, we let n_{-a} , n_0 denote the number of synonymous reverse-APOBEC3 mutations (TT \rightarrow TC and AA \rightarrow GA) and synonymous non-APOBEC3 mutations, respectively. When the ancestor can be used as the reference sequence, only the number of APOBEC3-induced mutations is considered. However, as we discuss in the next section, when the ancestor sequence is unknown, the forward APOBEC3 mutation on the test sequence and reverse-APOBEC3 mutations on the reference sequence can not be distinguished. Conditioned on a total of $n(t) = n$ mutations, the probability of n_{+a} synonymous APOBEC3-induced mutations arising is

$$\mathbb{P}(n_{+a}(t) = n_{+a} | n(t) = n) = \binom{n}{n_{+a}} \left(\frac{N_{+a}v_{+a}}{N\bar{v}} \right)^{n_{+a}} \left(1 - \frac{N_{+a}v_{+a}}{N\bar{v}} \right)^{n-n_{+a}}. \quad (4)$$

In other words, the random variable $n_{+a}(t)$ given $n(t)$ follows a binomial distribution with parameter n and probability $N_{+a}v_{+a}/(N\bar{v})$.

If the ancestor sequence S_0 is known, we can infer that for all samples $\alpha^{(i)}(t^{(i)})$, the number of synonymous APOBEC3-induced mutations $n_{+a}^{(i)}(t^{(i)})$ follows a binomial distribution with parameters $n^{(i)}$ and $\frac{N_{+a}v_{+a}}{N\bar{v}}$. If $n^{(i)}$ is sufficiently large, we expect that

$$n_{+a}^{(i)}(t^{(i)}) \approx \frac{N_{+a}v_{+a}}{N\bar{v}} n^{(i)}(t^{(i)}), \quad \text{for } 1 \ll n^{(i)} \ll N. \quad (5)$$

Eq. (5) reveals a simple linear relationship followed by all samples. Unfortunately, we do not know the ancestor sequence, and in order to extract evolution information from data, we must compare the evolution across multiple lineages.

Multi-lineage evolution

We now consider multiple lineages $\alpha^{(i)}(t^{(i)})$ evolved from the same ancestor sequence S_0 and sampled at time $t^{(i)}$. The statistics are straightforward if different lineages evolved independently. However, in reality, some lineages are related to each other by sharing parts of their evolutionary paths, as depicted in Fig. 1(B).

Shared evolutionary paths change how distinct mutations are enumerated. Suppose that two lineages $\alpha^{(i)}$ and $\alpha^{(j)}$ share the first $k^{(ij)}$ mutations and diverge at the $k^{(ij)} + 1$ -st mutation. After divergence, they acquire mutations independently. Among a total of $n^{(i)}(t^{(i)}) + n^{(j)}(t^{(j)})$ mutations in these two lineages, there are $k^{(ij)}$ pairs of identical mutations. The subsequent random mutations are mutually independent of each other.

To construct a systematic notation, for any two lineages $\alpha^{(i)}$ and $\alpha^{(j)}$, we define $k^{(ij)}$ to be the largest integer such that $\alpha_\ell^{(i)} = \alpha_\ell^{(j)}$ for all $\ell \leq k^{(ij)}$. For unrelated lineages, $k^{(ij)} \equiv 0$.

Let $A = \{\alpha_j^{(i)}\}$ denote the set of all mutations across all lineages. Under our working assumptions, the ℓ_1^{th} mutation $\alpha_{\ell_1}^{(i)}$ in lineage (i) and ℓ_2^{th} mutation $\alpha_{\ell_2}^{(j)}$ in lineage (j) are identical if and only if they have identical chronological order ($\ell_1 = \ell_2$) in the common evolution history before the two lineages diverged ($\ell_1 < k^{(ij)}$ and $\ell_2 < k^{(ij)}$). When the condition $\ell_1 = \ell_2 < k^{(ij)}$ is not satisfied, the mutations are unrelated. Unrelated mutations are independent and identically distributed. Examples of related mutations (α_1, α_2) and unrelated mutations $(\alpha_3, \alpha_3^{(j)}, \alpha_4, \alpha_4^{(j)})$ are shown in Figure 1(B).

Now, pick any two lineages $\alpha^{(i)}$ and $\alpha^{(j)}$ and define the *relative number of mutations* $n^{(ij)}$ of (i) with respect to (j) as the number of synonymous mutations identified when (j) is assumed to be the ancestor sequence, i.e. when the reference genome is (j). Figure 1(D) provides an example of $n^{(ij)}$. Let $t^{(ij)}$ be the time when two lineages $\alpha^{(i)}$ and $\alpha^{(j)}$ diverged, as indicated in Fig. 1(B). Then, $k^{(ij)} = n^{(i)}(t^{(ij)}) = n^{(j)}(t^{(ij)})$. Under our infinite sites assumption,

$$\begin{aligned} n^{(ij)} &\approx n^{(i)}(t^{(i)}) - n^{(i)}(t^{(ij)}) + n^{(j)}(t^{(j)}) - n^{(j)}(t^{(ij)}) \\ n_{+a}^{(ij)} &\approx n_{+a}^{(i)}(t^{(i)}) - n_{+a}^{(i)}(t^{(ij)}) + n_{-a}^{(j)}(t^{(j)}) - n_{-a}^{(j)}(t^{(ij)}) \\ n_{-a}^{(ij)} &\approx n_{-a}^{(i)}(t^{(i)}) - n_{-a}^{(i)}(t^{(ij)}) + n_{+a}^{(j)}(t^{(j)}) - n_{+a}^{(j)}(t^{(ij)}) \end{aligned} \quad (6)$$

If $1 \ll n^{(ij)} \ll N$, substituting Eq. (5) into the right-hand side of Eq. (6), we find order of magnitude relationships between the number of mutations between lineage (i) and reference lineage (j) and the times of sampling and lineage divergence:

$$\begin{aligned} n^{(ij)} &\approx N\bar{v}(t^{(i)} + t^{(j)} - 2t^{(ij)}) \\ n_{+a}^{(ij)} &\approx N_{+a}v_{+a}(t^{(i)} - t^{(ij)}) + N_{-a}v_{-a}(t^{(j)} - t^{(ij)}) \\ n_{-a}^{(ij)} &\approx N_{-a}v_{-a}(t^{(i)} - t^{(ij)}) + N_{+a}v_{+a}(t^{(j)} - t^{(ij)}). \end{aligned} \quad (7)$$

Therefore, upon defining $n_a^{(ij)} = n_{+a}^{(ij)} - n_{-a}^{(ij)}$, we find the asymptotic expression:

$$n_a^{(ij)} \approx \frac{N_{+a}v_{+a} + N_{-a}v_{-a}}{N\bar{v}} n^{(ij)}, \quad \text{for } 1 \ll n^{(ij)} \ll N. \quad (8)$$

Due to the inability to distinguish APOBEC3-induced mutations on genome i from reverse-APOBEC3 mutations on genome j, the number of observed APOBEC3-induced mutations $n_{+a}^{(ij)}$ of genome i with respect to genome j is the sum of $N_{+a}v_{+a}(t^{(i)} - t^{(ij)})$ and $N_{-a}v_{-a}(t^{(j)} - t^{(ij)})$. This quantity depends on $(t^{(i)} - t^{(ij)})$ and $(t^{(j)} - t^{(ij)})$, the branch lengths of both lineage (i) and lineage (j). Therefore, the ratio $n_{+a}^{(ij)}/n^{(ij)}$ will also depend on the branch lengths and is not a constant when $N_{+a}v_{+a} \neq N_{-a}v_{-a}$. To avoid this complication, we must consider both the forward and reverse mutations, which leads to the desired linearity in Eq. (8). Conditioned on $n^{(ij)} = n$, $n_a^{(ij)}$ still follows the binomial distribution with mean and

probability parameters n and $p = (N_{+a}v_{+a} + N_{-a}v_{-a})/(N\bar{v})$:

$$\begin{aligned} \mathbb{P}(n_a^{(ij)}(t^{(i)}) = n_a^{(ij)} | n^{(ij)} = n) \\ = \binom{n}{n_a^{(ij)}} \left(\frac{N_{+a}v_{+a} + N_{-a}v_{-a}}{N\bar{v}} \right)^{n_a^{(ij)}} \\ \times \left(1 - \frac{N_{+a}v_{+a} + N_{-a}v_{-a}}{N\bar{v}} \right)^{n - n_a^{(ij)}}. \end{aligned} \quad (9)$$

When an *unrelated* genome ℓ is used as a reference, the proportionality indicated in Eq. (8) is no longer valid. In this case, because of unrelatedness and the infinite sites assumption, we have $n^{(i\ell)} = n^{(ij)} + \Delta n^{(j\ell)}$ and $n_a^{(i\ell)} = n_{a,ij} + \Delta n_a^{(j\ell)}$, where $\Delta n^{(j\ell)}$ and $\Delta n_a^{(j\ell)}$ are two values that are independent of i . Consequently, $n_a^{(i\ell)}$ and $n^{(i\ell)}$ are not proportional to each other but follow the linear relationship

$$n_a^{(i\ell)} \approx \frac{N_{+a}v_{+a} + N_{-a}v_{-a}}{N\bar{v}} n^{(i\ell)} + \left[\Delta n_a^{(j\ell)} - \frac{N_{+a}v_{+a} + N_{-a}v_{-a}}{N\bar{v}} \Delta n^{(j\ell)} \right]. \quad (10)$$

The fraction $(N_{+a}v_{+a} + N_{-a}v_{-a})/(N\bar{v})$ and constant $\Delta n_a^{(j\ell)} - (N_{+a}v_{+a} + N_{-a}v_{-a})/(N\bar{v}) \Delta n^{(j\ell)}$ are parameters to be obtained from data. If genomes (j) and (ℓ) have evolved from the same ancestor with the same mutation rates, the constant term is zero. Nonzero constant term partially measures the degree to which the mutation rates of the two lineages are different.

Relaxing relative clock assumptions

So far in our relative clock construction, we assumed that the raw mutation rates are constant over genomic locations, lineages, and time. Although this assumption is not realistic, phylogenetic-like models do not require specification of raw mutation rates. Since our hybrid approach shares features with phylogenetic methods, we can relax the constant mutation rate assumption while retaining key relative rate information. The assumption of constant mutation rates across lineages and generations can be relaxed by properly reinterpreting the time variable t , while the assumption of constant mutation rates across genomic locations can be relaxed by changing how total mutation rates are calculated.

For time-varying overall mutation rates, $v(t)$, we define a new nonlinear measure of time $\tau := \int_0^t v(t)dt$, over which the mutation rate is again constant. Interpreted biologically, τ is interpreted biologically as being proportional to the number of DNA replication cycles since the last common ancestor. It has been found that the mutation rate is positively correlated with DNA replication frequency and negatively correlated with the generation time (Li et al., 1996; Bromham and Penny, 2003; Moorjani et al., 2016). Under those circumstances, we replace the absolute time t in the previous sections with a pseudo-time τ that measures the number of generations since the last common ancestor, which is universal across all different lineages. Since Eqs. (5) and (8) are independent of the time variable, they are invariant for the new time τ .

Incorporating possible heterogeneity in mutation rates across different sites requires a more complex argument. Genomic sites with an overall higher mutation rate contribute most significantly to the mutations observed between lineages. Instead of simply multiplying v_{σ_k} by the number of sites N_{σ_k} available for the mutation of type σ_k , we need to calculate the total mutation rate of type σ_k via a sum $\sum_{k=1}^{N_{\sigma_k}} v_{\sigma_k,s}$ that weights the mutation rates at each site s . As the mutations accumulate, indirect and complex cellular interactions may also change the rate of mutation. However, under an infinite sites approximation, only a very small

fraction of the sites have mutated. Therefore, the total mutation rates and their ratios remain constant for a very long time. In the unlikely case where a small number of sites experience extremely high mutation rates, while all other sites have negligible mutation rates, the infinite sites assumption is effectively violated and the ratios of the total mutation rates may no longer remain constant. We conclude that the spatial homogeneity assumption can be effectively relaxed provided the distribution of mutation rates are not too disperse. After relaxation of these assumptions and proper reinterpretation, our previous calculations and equations hold.

Statistical tests for changes in the mutation rate

Although time-dependent changes in *separable* mutation rates $v(t)$ can be taken into account and do not affect mutation rate ratios, our modeling approach allows one to statistically test for inseparable shifts in the mutation rate where ratios of mutation rates change. A model in which mutation rates change in time is simulated in Appendix ‘Hypothetical scenarios’ in the Supporting Information (SI) and shown in Fig. S1(A). Specifically, Eq. (9) (the distribution of the count of a specific mutation type, conditioned on known relative mutation rates and the total number of synonymous mutations) enables the calculation of a 95 per cent prediction interval. Instances where samples fall outside this interval suggest a significant alteration in the mutation rate. We provide an example of this test in the Results and in Figure S1(B).

Eq. (9) can be further used to test whether a focal genome and a set of known genomes share the same relative mutation rates. To enhance resolution, the focal genome is used as a reference and its coordinates are set to the origin $(n, n_a) = (0, 0)$. The null hypothesis that the reference genome and the set of other genomes (i) exhibit the same relative mutation rate can be tested by how well $(n^{(i)}, n_a^{(i)})$ represent draws from the binomial distribution given in Eq. (9). For example, one can apply least-squares regression on the data points $(n^{(i)}, n_a^{(i)})$ for each genome i . Under the null hypothesis, the n_a -intercept of the regression line should be at $b \equiv n_a(n=0) = 0$. A t -test statistic given by $b/SE(B)$ can then be used, providing a sensitive test of the null hypothesis.

Selection effects and model validation via simulation

Our theoretical analysis predicts a linear relationship between $n_a^{(ij)}$ and $n^{(ij)}$, which we test by implementing two types of simulations.

Mutation-only processes

First, we simulate DNA evolution using independent and identically distributed (i.i.d.) GTR processes. For simplicity, we first exclude selection pressure and assume equal frequencies and equal mutation rates of all four nucleotides, aligning our model with the JC69 limit (Jukes and Cantor, 1969).

For each trial, we randomly sample 10^6 nucleotides from the equilibrium distribution of $\{A, T, C, G\}$ to construct an initial sequence. For each nucleotide position, we then simulate a Markov mutation process using the JC69 transition matrix and a variant of the Gillespie algorithm (Gillespie, 1977). During the simulation of the substitution model, we count the total number of mutations and plot the associated numbers of $TC \rightarrow TT$ and $A \rightarrow C$ mutations in Figure 2(A) and (B), respectively. Across all four independent trajectories shown, the linear relationship between the number of all mutation types and the total mutation number




Figure 2. Simulations of different scenarios. (A, B) Four representative trajectories of an iid substitution process of the JC69 limit. A total of 10^6 nucleotides are used for each simulation and both TC → TT and A → C mutations are linearly correlated with the total number of point mutations.

holds, even up to $\sim 10^4$ mutations. This justifies and implies broad applicability of the infinite sites assumption.

Mutation-selection processes

In a second set of simulations, we factor in selection pressure, using a simple asexual reproduction logistic growth model with a shared carrying capacity K . In this simulation, we track the population of genes through the numbers of background synonymous mutations $n - n_a$, synonymous APOBEC3-driven mutations n_a , and 'hidden' (nonsynonymous) beneficial mutations n_h . The population undergoes a birth-death process defined by birth and death rates

$$\beta(n_a, n_h) = \beta_0 s(n_a, n_h), \quad (11)$$

$$\mu(n_a, n_h) = \mu_0 + \beta(n_a, n_h) \frac{N_G(t)}{K}, \quad (12)$$

where $N_G(t)$ is the total population of genomes, K is the carrying capacity, and the selection coefficient is given by

$$s(n_a, n_h) = 1 + \sigma_a n_a + \sigma_h n_h. \quad (13)$$

The selection coefficient depends linearly on the numbers of mutations n_a and n_h through the selection strengths and σ_a and σ_h . Genomes that have larger $s(n_a, n_h)$ reproduce faster.

At birth, one daughter has a probability to acquire an additional mutation of specified type. Mutation probabilities are chosen such that the total rates of each type of mutation are identical. To connect with the real-world evolutionary process, after each unit of time (corresponding to the average time between replication of a single genome), a sample of the population is drawn and genomes with specific number $n - n_a$ of synonymous non-APOBEC3 mutations $n - n_a$ are collected, along with their associated numbers of synonymous APOBEC3 mutations n_a . These $(n_a, n - n_a)$ points are collected across all sampled time points and plotted in Figs. 3(A) and (C). From the infinite sites approximation, Eq. 2, and the constant relationship between different types of mutation, $n - n_a$, is tightly associated with time.

We assign different selection coefficients σ_a and σ_h to understand how selection on the synonymous APOBEC3 mutations and other beneficial mutations may alter the linear relationship between n_a and n . In a mutation-only process (negligible σ_a and σ_h), the expected number of mutations of each type is the same as the others, e.g., $n_a \approx (n - n_a)$, as indicated by the

blue trajectory in Fig. 3(A). In Figure 3(B), we plot, for each value of σ_a indicated, the trajectory slopes for 25 different values of $\log_{10} K$ uniformly sampled within $\log_{10} K \in [2, 4]$ or $K \in [10^2, 10^4]$. The increased mutation rate with larger population size is consistent with prior theoretical understanding of selection effects (Lewontin et al., 2000; Ingvarsson, 2008). Even when the selection for APOBEC3 mutations is strong ($\sigma_a \geq 0.01$) and the carrying capacity is large ($K > 10^3$), the effective APOBEC3 mutation rate is at most doubled, as shown in Figure 3(B).

If the selection of the hidden beneficial mutations is large (say, $\sigma_h \gtrsim \sigma_a$), the ratio $n_a/(n - n_a) \approx 1$, as shown by the red trajectory in Figure 3(C) and the marked red values of slopes (diamonds) in Fig. 3(D). Here, the hidden beneficial mutations drive selection, leading again to comparable n_a and $n - n_a$.

Summarizing, our simulations verify (1) that the infinite sites hypothesis holds up to $\sim 10^3$ mutations in a genome of 10^6 bp, (2) that the linear relationship between the number of different types of mutations holds even under selection, (3) that the effective mutation rate of APOBEC3-driven mutations can be increased by selection only if they are relatively strongly selected for ($\sigma_a \geq 0.01$), and (4) that strong selection for other beneficial mutations can mask the increase in the effective mutation rate of APOBEC3-driven mutations due to selection.

Sequence analysis and alignments

We obtained the sequences of 237 MPXV genomes from the NCBI database as listed in Table S1. To date, there are hundreds of MPXV genomes available. Dates of the sequences are inferred from the collection date and other contextual information. However, a majority of them are samples collected from the most recent outbreak. Thus, a large number of them are closely related, and using our proposed analysis, contribute little information to the evolutionary history of MPXV. The sequences were aligned using the affine gap model from the BioAlignments package (Christensen et al., 2022) in JuliaLang (Bezanson et al., 2017). Alignment results are manipulated using the BioSequences and GenomicAnnotations packages in JuliaLang.

The dinucleotide and synonymous mutations were identified by first enumerating all mutations in the alignment, in the context of whole-genome sequences. Then, we identify the synonymous mutations by comparing the amino acid sequences of the genes. These considerations are not restricted to the third codon position.




Figure 3. Simulations of scenarios that include mutation and selection following Eqs. 11–13. We used $\beta_0 = 1$, $\mu_0 = 0.5$ and assumed that with each birth, one daughter has equal probability 0.1 of acquiring each type of mutation. The total initial numbers of possible mutations N_i for synonymous non-APOBEC3 mutations, synonymous APOBEC3 mutations, and hidden nonsynonymous mutations are set to 10^5 , 10^3 , and 10^3 , respectively. (A) Numbers of synonymous APOBEC3-relevant mutations n_a associated with the numbers of synonymous non-APOBEC3-relevant mutations $n - n_a$ from samples of simulation trajectories. Here, no selection for hidden mutations ($\sigma_h = 0$) is present. (B) Each diamond symbol represents the slope of an independently simulated selection-mutation process, as shown in (A). Slopes for different σ_a and 25 uniformly sampled values of $\log K$ are shown. (A) and (B) show that large population size K and strong selection coefficient σ_a can lead to a larger effective mutation rate of APOBEC3-driven mutations. By design, under weak selection strength σ_a , the effective mutation rate of APOBEC3-driven mutations is identical to the non-APOBEC3-driven mutations. (C) Values of n_a associated with $n - n_a$ under different strengths σ_h of selection of hidden beneficial mutations. Strong selection of hidden beneficial mutations can restore the increased effective mutation rate of APOBEC3-driven mutations to the same level as the pure mutation processes. (D) For large $\sigma_h > \sigma_a$, the expansion of n_a is limited (red diamonds).

For sequences without gene annotations, we used NC_063383 as the template to annotate the genes. Then, these annotations were used to generate pairwise alignment results and identify synonymous mutations. The statistical tests were developed based on likelihood ratio tests for binomial distributions.

Appendix ‘Lineage Analysis’ in the Supporting Information (SI) provides details on the inference of a tree based on ordering of genome mutations. Our result is shown in Fig. S3(A) and compares well with the phylogenetic tree shown in Figure S3(B) which as derived from using standard NCBI BLAST (King et al., 2010). However, BLAST does not correctly resolve subtle ancestral relationships between recently collected genomes, which our method does.

Results

Statistically distinct subgroups with different evolutionary features

For the reasons discussed in **Multi-lineage evolution**, when the phylogenetic tree is not resolved, we cannot distinguish APOBEC3-induced mutations from their reverse mutation. If the phylogenetic tree is established, these two directions of substitutions can be distinguished from established methods such as

the UNREST model (Yang, 1994). Here, APOBEC3-relevant mutations are defined as the mutations matching patterns $TC \leftrightarrow TT$ or $GA \leftrightarrow AA$, as detailed in Table 2. The number of these synonymous mutations is $n_a = n_{+a} + n_{-a}$. For a time-scaled phylogenetic tree construction method, ‘root-to-tip’ regression is used to estimate the evolutionary rate of the virus, represented by the slope of the regression line (Rambaut et al., 2016). In the absence of a resolved phylogenetic tree, a linear fit of two types of mutations in our relative clock is essentially a ‘tip-to-tip’ regression.

Successful application of the relative molecular clock to pre-2016 genomes

We obtained six MPXV genomes collected before 2016 from GenBank. The genome KP849470 was arbitrarily chosen as the reference genome against which other genomes are aligned. For each genome (i), we calculate the relative number of synonymous mutations $n^{(i)KP849470}$ and number of APOBEC3-relevant mutations $n_a^{(i)KP849470}$. For simplicity, we omit the reference genome in the superscript and write $n^{(i)}$ and $n_a^{(i)}$, respectively. Each genome is then plotted on the (n, n_a) plane, where they fall near a line which we find by using a least-squares fit and Eq. (8). As shown in Fig. 4(A), the data points are well fitted by the linear model,




Figure 4. Relative molecular clocks for different types of mutations with respect to an arbitrarily chosen reference genome, instead of time, exhibit linearity. (A) The number of synonymous APOBEC3-relevant mutations (TC → TT) is plotted against the number of synonymous mutations for the genomes collected before 2016. (B) The number of synonymous APOBEC3-relevant mutations (TC → TT) is plotted against the number of synonymous mutations for the genomes collected after 2016.

allowing us to conclude the pre-2016 MPXV genomes have evolved with a constant set of relative mutation rates.

Given the shared relative mutation rates, we can use the linear fit as a baseline to compare the APOBEC3-relevant mutation rates. A total of $N \sim 100,000$ possible synonymous mutations were identified on the reference genome KP849470, with $N_{+a} \sim 3,000$ of them APOBEC3-induced, and $N_{-a} \sim 8,000$ of them reverse-APOBEC3-induced. In total, APOBEC3-relevant mutations account for about 11 per cent of all possible synonymous mutations. By contrast, the slope of the linear fit in Fig. 4(A) is 0.157 ± 0.007 , which represents the fraction of observed APOBEC3-relevant mutations n_a/n . This fraction is significantly larger than the fraction of possible APOBEC3-relevant mutations $N_a/N \approx 0.11$, suggesting that the APOBEC3-relevant mutation rate is higher than that of the average mutation already extant in the pre-2016 MPXV genomes. The ratio $n_a^{(i)}/n \sim 0.157$ serves as a better baseline to test whether the APOBEC3-relevant mutation rate is higher in the post-2016 MPXV genomes.

Successful application of the relative molecular clock to post-2016 genomes

We obtained 226 MPXV genomes from GenBank collected after 2016. Genome MK783028 was chosen to be the reference genome against which other genomes are aligned. The relative number of synonymous mutations $n^{(i)MK783028}$ and number of APOBEC3-relevant mutations $n_a^{(i)MK783028}$ are computed and each genome is plotted in the (n, n_a) plane. The results are shown in Fig. 4(B). These points also fall near a line, indicating that the post-2016 MPXV genomes also evolved with a constant set of relative mutation rates.

The total number of possible mutations of different types on the reference genome MK783028 is similar to that of KP849470. However, the slope n_a/n of the linear fit in Fig. 4(B) is 0.912 ± 0.002 , which is significantly larger than the baseline slope 0.157 ± 0.007 of the pre-2016 genomes.

Interpretation of different APOBEC3-relevant mutation rates

In conclusion, our key assumption of constant ratios of mutation rates is separately applicable to the two subgroups of genomes

(distinguished by their year of collection). The mutation rate ratios are particular to each group. Because we have factored out the selection pressure by considering only synonymous mutations, selection is unlikely to account for the difference in the two mutation rate ratios. If this difference is due to random neutral mutations within the animal hosts, we would expect that the ratio of the number of APOBEC3-relevant mutations to the number of all synonymous mutations is the same for both groups of genomes resulting in similar slope in Figure 4. In addition, since the total number of possible mutations of different types on the reference genome is similar for both groups of genomes, the difference in the observed frequencies of APOBEC3-relevant mutations can only be explained by the difference in the relative APOBEC3-relevant mutation rate.

The SNP variants across the pre-2016 genomes that often belong to different clades likely reflect the evolution of the virus in unknown animal reservoirs. By contrast, the genomes in the post-2016 group are clustered into a single clade, and the associated SNP variants likely reflect the evolution of the virus in the human host. Therefore, we conclude that the APOBEC3-relevant mutation frequency $n_a/N_a > n/N$ is relatively higher in the pre-2016 group ($n_a/N_a > n/N$), possibly due to some intrinsic or extrinsic physical or chemical mutagens like UV radiation, as mentioned previously. The APOBEC3-relevant mutation rate is extremely high in the post-2016 group, which could be a result of persistent APOBEC3-editing in the human host.

APOBEC3 and other human-specific environmental factors may also impose selection pressure that favors certain mutations. For example, tRNA abundance may be different in the human host compared to the animal reservoirs. However, such effects are generally expected to be gene-specific, exhibiting a stronger effect on highly expressed genes. Our analysis in Appendix ‘Distribution of APOBEC3-induced mutations’ (see SI) does not support this hypothesis as we do not find exceptionally high APOBEC3-relevant per-site mutation rates in specific genes (see Fig. S4). Neither are mutation types correlated across genes (see Fig. S5). Moreover, our simulations show that even a strong selection coefficient favoring APOBEC3-relevant mutations can at most double the APOBEC3-relevant mutation rate, which is much smaller than the observed difference between the pre-2016 and post-2016 groups




Figure 5. (A) The number of observed APOBEC3-relevant mutations n_a of all genomes plotted against the number of observed synonymous mutations n , using the genome KP849470 as a reference. The two genomes KJ642617 and MK783029 are highlighted since they are the closest genomes to the intersection of the two lines corresponding to the pre-2016 and post-2016 genomes. (B) The number of observed APOBEC3-relevant mutations n_a , using genome KJ642617 as the reference, plotted against the number of observed synonymous mutations n . The open circle represents the location of the hypothesized most recent common ancestor (MRCA) of the post-2016 genomes in the animal reservoir. The inset shows that the line of the post-2016 group does not intersect the origin, suggesting that KJ642617 is not a common ancestor of the post-2016 group. (C) The number of observed APOBEC3-relevant mutations n_a plotted against the number of observed synonymous mutations n , using MK783029 as reference. The inset shows that the line of the pre-2016 group does not pass through the origin, suggesting that MK783029 has already undergone some APOBEC3-editing after its ancestor's transmission to human.




Figure 6. The number of synonymous APOBEC3-induced mutations plotted against the collection date of the post-2016 genomes, using KJ642617 as the reference. The red diamond represents the genomes within the lineage with the longest branch length, starting with MK783028 and ending with ON694329.

(see Figure 3). Other beneficial nonsynonymous mutations could further reduce the increase in the ratio. Therefore, we conclude that the observed difference in APOBEC3-relevant mutation rate is mostly likely due to a difference between the APOBEC3-editing activity in animal reservoirs and the human host.

General applicability of relative molecular clocks and bias for APOBEC3-relevant mutations

We have shown that our recalibrated molecular clock method is applicable to both pre- and post-2016 genomes in the context

of APOBEC3-relevant mutations versus all synonymous mutations. In fact, the relative molecular clock method is applicable to other types of synonymous mutations as well. For example, we considered the A \rightarrow C type of synonymous mutations versus synonymous non-APOBEC3-relevant mutations in Fig. S6. The linearity is still observed and is consistent in both pre- and post-2016 genomes. This observation suggests not only that the relative molecular clock method is generally applicable.

The ancestor of post-2016 MPXV genomes

The geometry of pre- and post-2016 genomes on the same plane

By choosing a common reference genome for the two groups of genomes, we can visualize them on the same (n, n_a) -plane. The geometric relationship between the two groups of genomes reveal information about the evolutionary history of the virus in these two groups of hosts.

When KP849470 is used as the reference genome, the post-2016 genomes are still distributed along a line in the (n, n_a) -plane, as shown in Fig. 5(A). However, since the post-2016 genomes have undergone a different evolutionary history, the genome KP849470 should not be co-linear with the post-2016 genomes. In other words, the line representing the post-2016 genomes should not pass through the origin, but the lines of the pre- and post-2016 genomes will intersect at some point. Genomes at this intersection will share a common evolutionary history in both the animal reservoir and the human host. Since the post-2016 genomes may come from the animal reservoir, the intersection point represents the most recent common ancestor (MRCA) of the post-2016 genomes in the animal reservoir.

The genomes KJ642617 and MK783029 lie approximately at the intersection of the two lines when the genome KP849470 is used as reference. They are thus good candidates for the MRCA of the post-2016 genomes. Small deviations from the precise intersection point of the two lines can be explained by the stochastic nature of mutations.




Figure 7. Different evolutionary environments shape the shared features of pre- and post-2016 groups of MPXV genomes. Pre-2016 cross-species transmission did not persist in human hosts. The blue dots represent the shared features due to evolution in the animal reservoir. The red dots represent the shared features due to evolution in the human hosts. We hypothesize an unobserved stage of continuous and persistent evolution in the human hosts between the 2017 cross-species transmission and the 2022 global outbreak.

Statistical tests to determine the MRCA of the post-2016 MPXV genomes

In order to minimize the effects of stochasticity and improve precision, we determine whether genomes KJ642617 and MK783029 are the MRCA of interest by choosing them as the reference genome (setting their positions to $(n, n_a) = (0, 0)$), respectively. Next, we determine how well the two lines corresponding to the pre-2016 and post-2016 genomes intersect at the origin.

Using least-squares regression, we found that when KJ642617 is used as the reference genome (Fig. 5(B)), the n -intercept of the linear fit of the post-2016 genomes is -9.27 ± 0.34 , but when MK783029 is used as reference (Fig. 5(C)), the n -intercept of the pre-2016 genomes is 2.80 ± 0.55 . In both cases, the P -values of the null hypothesis that the n -intercept is zero are less than 0.01. Thus, neither KJ642617 nor MK783029 is representative of the MRCA of the post-2016 MPXV genomes.

However, when we apply the combined P -value test described previously, based on the conditional probability formula Eq. (9), the P -values for the null hypotheses are both greater than 0.05. This discrepancy reflects the fact that the observed number of APOBEC3-relevant mutations has a sharper distribution concentrated around the mean value than the distribution predicted in theory. This sharper distribution might result from multiple factors, including the close phylogenetic relationship between genomes and selection pressure.

Statistical inference of the MRCA in the animal host

Although neither KJ642617 nor MK783029 are good candidates for the MRCA in the animal hosts, we can infer the number of synonymous mutations with respect to a given reference genome by analyzing the intersection of the linear fits to the pre- and post-2016 genomes.

To estimate the time at which the MRCA first emerges, the reference genome should be chosen from the pre-2016 group. Only if the reference genome is chosen from the pre-2016 group will the critical ancestor have a fewer number of synonymous mutations than other post-2016 genomes. Such a choice would allow us to linearly extrapolate the strict molecular clock fit to the post-2016 genomes and find the emergence time of the MRCA.

Although we can use any pre-2016 genome as a reference, KJ642617 is the most similar to the post-2016 genomes, as measured by the number of observed synonymous mutations. For mutations that occur in a Poisson-like process, the effects of stochasticity measured by standard deviation is proportional to the square root of total number of mutations. Therefore, using KJ642617 as the reference reduces the number of observed mutations and therefore the noise, providing a more precise estimate. As shown in Fig. 5(A), ordinary least-squares regression provides an estimate of the 95 per cent confidence interval $n_{\text{ancestor}} \in$

[11.61, 12.76] of the number of synonymous mutations of the critical ancestor with respect to KJ642617.

Molecular clock of the post-2016 group

The knowledge of the relative number of synonymous mutations of the critical ancestor is helpful for timing its emergence assuming a strict molecular clock is present during its subsequent evolution.

In order to understand whether mutation rates have changed over time, we apply in Appendix ‘Lineage analysis’ (see SI) the root-to-tip regression, using the genetic distance defined by the number of synonymous mutations and calibration data defined by manually identified collection dates of the genomes, over a phylogenetic tree constructed by asymmetry of TC \rightarrow TT mutations. The total rate $N\bar{v}$ of synonymous mutations for the post-2016 genomes is estimated to be around 5 per year. Their number of synonymous mutations $n^{(i)}(t^{(i)})$ are plotted against the collection date, as shown in Fig. 6.

Multiple factors such as differences between emergence and collection times, cross-lineage and cross-generation differences in the mutation rates, and different geographic locations can result in deviations from the linear relationship between the number of mutations and the collection time predicted by the strict molecular clock. Consequently, unlike the relative clocks shown in Figs. 4 and 5, many genomes seem to have a significant deviation from the predictions of a strict clock. Relatedness between genomes will make the observed number of mutations correlated and not truly independent, rendering statistics such as R^2 and P -values biased and less reliable. More specifically, a large fraction of samples collected in 2022 have almost identical number of synonymous mutations. Any fitting method that passes through the center of these samples will have a high R^2 .

In order to minimize the deviation from predictions of a strict molecular clock, we considered the *principal lineage* defined by the longest branch length identified by the phylogenetic method described in the next section, shown in ‘Lineage analysis’ in the SI and Fig. S3. For the twenty-two genomes sampled along this the principal lineage, the linear fit between the number of synonymous mutations to collection date yields a slope of 5.17 synonymous mutations per year. As shown in Fig. 6, this fit is quite good with $R^2 = 0.997$.

Weighted least square fitting is preferred for dating the MRCA

When the strict molecular clock assumption of constant mutation rates is valid, the precise number of synonymous mutations with respect to time is given by a constant-rate Poisson process. After a given time t , both the mean and the variance in the number of synonymous mutations is proportional to t .

To account for the change in variance, we use a weighted least-squares method to fit the data. The weights are inversely proportional to the variance in the number of mutations and thus inversely proportional to time t . To convert collection dates into time t , we need to have a primary estimate of the emergence time of the MRCA. This is obtained by ordinary least-squares fitting of the data and is set to 01 October 2016. However, due to sampling bias, the variance of the number of mutations for the whole dataset is not proportional to time t , so we restricted our analysis to the genomes within the principal lineage of MPXV.

The linear relationship between the mean number of synonymous mutations and time suggests that no major change in the mutation rate occurred along the principal lineage, including during the unsampled period between 2019 and 2021. A few genomes collected in 2022 have a different number of synonymous mutations. This deviation may be due to cross-lineage variations. More observations are needed to test whether the mutation rate has changed since the 2022 global outbreak.

We then extrapolated the linear fit obtained via weighted least squares and found a 95 per cent confidence interval of the time of emergence of the ancestor $t^{(\text{ancestor})} \in [2016-10-12, 2017-05-05]$. The main source of uncertainty is the uncertainty in the weighted least-squares fit, where the weighting accounts for possible correlations between different samples. This estimate is close to that of a previously published estimate (O'Toole and Rambaut, 2022c).

Discussion and Conclusions

We developed a simple molecular clock-based method for analyzing relative mutation rates that is applicable to a broad class of DNA evolution models. Molecular clock theory assumes constant a evolutionary rate across lineages and generations. However, this assumption is often violated due to differences in environment and evolutionary mechanisms. Using a less restrictive assumption of constant ratios of mutation rates, we developed a method to recalibrate a molecular clock using, instead of time, another molecular clock. If the relative rates of different types of mutations are constant, different lineages from the same ancestor will have the same relative numbers of these different types of mutations. When we applied our analysis to monkeypox virus (MPXV), we find that there are two distinct groups of MPXV genomes, those collected before 2016 and those after. Each group adheres to the recalibrated molecular clock prediction, with unique relative mutation rates, indicating different evolutionary pressures potentially caused by variations in animal and human hosts. The post-2016 group of sequences share biased hypermutations of the pattern TC \rightarrow TT, characteristic of human APOBEC3-induced editing. Imposing an additional assumption of independence between sites, we are able to statistically characterize the initial common ancestor of lineages appearing after 2016. Further invoking an infinite sites assumption mathematically simplifies the relation between different types of mutations and the overall analysis. These assumptions allowed us to predict a mean number of mutations that increases linearly with time (Eqs. 2, 7, and 14), as well as a linear relationship between the mutations of a specific type to the total number of mutations (Eqs. 5 and 8), respectively.

Applying our analysis to MPXV samples shows that they evolved in a way that is quantitatively consistent with our assumptions and that there are two distinct epochs of mutations characterized by differences in the rates of the specific dinucleotide mutation TC \rightarrow TT, typical of APOBEC3 editing. We ruled out several alternative possibilities for the high APOBEC3-relevant mutation frequencies such as selection or drift in animal hosts or

differences in the number of available mutation sites. Dynamic biases in mutation rates driven by physical or chemical mutagens are also unlikely since the pre-2016 group of MPXV genomes represent a baseline that incorporates the effects of mutagens other than human-specific enzymes.

Additionally, our analysis not only identified the statistical departure but also showed that the relative mutation rate of TC \rightarrow TT is preserved in the pre-2016 and the post-2016 sequences, respectively, as shown in Fig. 4. The tight linear proportionality observed is also worth reporting and further investigation. It may reflect a relatively homogeneous environment for different individuals in the host populations. It can also allow one to discern subtle changes in relative mutation rates. In particular, our analysis suggests that the KJ642617 sequence is not a direct ancestor of the post-2016 group, despite it being very close. Similarly, the MK783029 sequence in the post-2016 group is very likely to have undergone a few APOBEC3-editing events since the MRCA of the post-2016 group.

We conclude that the most likely scenario is that pre-2016, MPXV genomes evolved in the animal host. The post-2016 group of MPXV genomes has undergone persistent and continuous human APOBEC3 editing after zoonotic transmission circa late 2016/early 2017. This scenario is implied in Fig. 7.

Unlike applications of DNA evolution models to phylogenetic inference (Posada, 2013), our method provides a simple and vivid goodness-of-fit measure by checking whether different genomes share the same linear relationship between different types of mutations. While high APOBEC3-relevant mutation rates have been observed and reported in previous studies, our evolutionary model-based approach delineates shared features within and between the pre- and post-2016 groups of MPXV genomes. Our model also allows us to identify and characterize the critical ancestor of the post-2016 group, which may contain information about post-2016 epidemics. Quantification of the molecular clock of MPXV suggested that they have a constant mutation rate in the principal lineage of the post-2016 group. This allowed us to estimate the time of the ancestral zoonotic transmission leading to the post-2016 MPXV genomes.

We further validated strong APOBEC3-editing in post-2016 genomes by comparing the lineage analysis results with the results from the BLAST Tree algorithm. Both trees show similar clusters of genomes, but our method provides better resolution of the subtle phylogenetic relations (\simeq) between genomes collected in 2017 and 2018.

We have not explored the quantitative interpretations of connections between the rate of synonymous mutations and the rate of nonsynonymous mutations; nonetheless, our linear model and analysis can be directly applied to other viruses experiencing APOBEC3-driven evolution. For example, the synonymous mutation rate of SARS-Cov-2 was found to be relatively stable, while its nonsynonymous mutation rate varied over time (Neher, 2022). There have been some research into the relationship between the mutation rates of synonymous and nonsynonymous mutations (Zhang, 2005; Bloom, 2014; Spielman and Wilke, 2016) but they have focused on improving the inference of the phylogenetic tree. Low levels of nonsynonymous mutations have already been observed in Feline parvovirus and Canine parvovirus and are related to strong selection (Wang et al., 2022).

The idea of inferring evolutionary insights from mutation ratios, such as the nonsynonymous to synonymous mutation ratio (dN/dS), is well established and has been used to infer selective pressures on protein-coding regions (Zhang, 2005; Bloom, 2014; Spielman and Wilke, 2016). Other methods have been developed

to identify site-specific mutation rate shifts (Yang, 1994; Murrell et al., 2012; Duchemin et al., 2022). However, these methods often involve phylogenetic tree reconstruction using computationally intensive approaches like maximum likelihood or Bayesian inference, and they lack an intuitive goodness-of-fit measure. In contrast, our method bypasses the need for phylogenetic trees, utilizes only sequence alignments against a common reference, and offers an intuitive way to measure fit to data. We can also evaluate the probability of nonsynonymous mutations per replication, but this can depend on selection since past nonsynonymous mutations would have likely changed the fitness of the genome. A sequence with more potentially beneficial mutations is more likely to exhibit a higher effective rate of nonsynonymous mutations, while excess deleterious mutations lead to slower nonsynonymous mutation. While the basic mutation rate is not directly affected by individual mutations, beneficial mutations are more likely to survive and spread, whereas deleterious ones tend to be eliminated. The effective mutation rate, influenced by selection, provides insight into the evolutionary landscape. Therefore, APOBEC3-driven (nonsynonymous) mutations may disrupt the dynamic mutation-selection balance, which may profoundly influence the evolution of molecular phenotypes under stabilizing selection (Rouzine, Brunet and Wilke, 2008; Goyal et al., 2012; Nourmohammad, Schiffels and Lässig, 2013). Finding beneficial mutations is mathematically similar to the problem of a high-dimensional random walk searching for target sites. The interplay between target (beneficial mutations) search and APOBEC3 editing may be an interesting direction of for future analysis. Combining these features with previously developed virus dynamics models could yield a more realistic picture of virus evolution in human (Kreger, Komarova and Wodarz, 2021; Lord and Bonsall, 2021).

Data availability

The computer code produced in this study are available at:

- Analysis computer scripts: Github (github.com/hsianktin/monkeypox).

This study includes no data deposited in external repositories.

Supplementary data

Supplementary data are available at *Virus Evolution* online.

Acknowledgements

T.C. acknowledges support from the National Institutes of Health through grant R01HL146552.

Conflict of interest: The authors have no competing interests.

Author Contributions

X.L. conceived and constructed the evolution model and performed the statistical analysis. S.H. and O.Y. conceived the initial problem and constructed the virus phylogeny. X.L., S.H., and T.C. wrote the manuscript. T.C. and O.Y. assisted in editing the manuscript.

References

- D. T. D. T. G. (1977) 'Exact stochastic simulation of coupled chemical reactions', *The Journal of Physical Chemistry*, 81: 2340–2361.
- Beale, R. C. L. et al. (2004) 'Comparison of the differential context-dependence of DNA deamination by APOBEC enzymes: Correlation with mutation spectra in vivo', *Journal of Molecular Biology*, 337: 585–596.
- Bezanson, J. et al. (2017) 'Julia: A fresh approach to numerical computing', *SIAM Review*, 59: 65–98.
- Bloom, J. D. (2014) 'An experimentally informed evolutionary model improves phylogenetic fit to divergent lactamase homologs', *Molecular Biology and Evolution*, 31: 2753–69.
- Breman, J. G. et al. (1980) 'Human monkeypox, 1970-79', *Bulletin of the World Health Organization*, 58: 165.
- Bromham, L. and D. Penny (2003) 'The modern molecular clock', *Nature Reviews Genetics*, 4: 216–24.
- Budden, T. and N. Bowden (2013) 'The role of altered nucleotide excision repair and UVB-induced DNA damage in melanomagenesis', *International Journal of Molecular Sciences*, 14: 1132–1151.
- Bunge, E. M. et al. (2022) 'The changing epidemiology of human monkeypox—a potential threat? a systematic review', *PLoS Neglected Tropical Diseases*, 16: e0010141.
- Burns, M. B. et al. (2013) 'APOBEC3b is an enzymatic source of mutation in breast cancer', *Nature*, 494: 366–370.
- Centers for Disease Control & Prevention USCDC. (2022) About monkeypox. <https://www.cdc.gov/poxvirus/monkeypox/about/index.html>, accessed 16 Oct 2022.
- Christensen, T. A. et al. (2022) 'Tim Holy, Morten Piibeleht, and tanhevg', *Biojulia/bioalignments.jl*, 3.0.0. <https://zenodo.org/record/7166110>, accessed 16 Oct 2022.
- Drouin, R. and J.-P. Therrien (1997) 'UVB-induced cyclobutane pyrimidine dimer frequency correlates with skin cancer mutational hotspots in p53', *Photochemistry and Photobiology*, 66: 719–726.
- Duchemin, L. et al. (2022) Evaluation of methods to detect shifts in directional selection at the genome scale', *Molecular Biology and Evolution*, 40: msac247.
- European Centre for Disease Prevention & Control EU CDC. (2019) Risk assessment: Monkeypox multi-country outbreak. <https://www.ecdc.europa.eu/en/publications-data/risk-assessment-monkeypox-multi-country-outbreak>, accessed 16 Oct 2022.
- Felsenstein, J. (1981) 'Evolutionary trees from DNA sequences: A maximum likelihood approach', *Journal of Molecular Evolution*, 17: 368–376.
- Felsenstein, J. (2001) 'Taking variation of evolutionary rates between sites into account in inferring phylogenies', *Journal of Molecular Evolution*, 53: 447–455.
- Fitch, W. M. and C. H., Langley (1976) *Molecular Anthropology*. Evolutionary rates in proteins: Neutral mutations and the molecular clock, Springer US, pp. 197–219.
- Gigante, C. M. et al. (2022) 'Multiple lineages of monkeypox virus detected in the United States, 2021–2022', *Science*, 378: 560–565.
- Goldman, N. and Z., Yang (1994) 'A codon-based model of nucleotide substitution for protein-coding DNA sequences', *Molecular Biology and Evolution*, 11: 725–736.
- Goyal, S. et al. (2012) 'Dynamic mutation-selection balance as an evolutionary attractor', *Genetics*, 191: 1309–1319.
- Huang, Y., L., Mu and W., Wang Monkeypox: epidemiology, pathogenesis, treatment and prevention', *Signal Transduction and Targeted Therapy*, 7: 2022.
- Hultquist, J. F. et al. (2011) 'Human and rhesus APOBEC3d, APOBEC3f, APOBEC3g, and APOBEC3h demonstrate a conserved capacity to restrict Vif-deficient HIV-1', *Journal of Virology*, 85: 11220–11234.
- Hussein, M. R. (2005) 'Ultraviolet radiation and skin cancer: molecular mechanisms', *Journal of Cutaneous Pathology*, 32: 191–205.
- Igor, M. R., E., Brunet and O. W., Claus (2008) 'The traveling-wave approach to asexual evolution: Müller's ratchet and speed of adaptation', *Theoretical Population Biology*, 73: 24–46.

- Ingvarsson, P. K. (2008) 'Molecular evolution of synonymous codon usage in populus', *BMC Evolutionary Biology*, 8: 307.
- Isidro, J. et al. (2022) 'Phylogenomic characterization and signs of microevolution in the 2022 multi-country outbreak of monkeypox virus', *Nature Medicine*, 28: 1569–1572.
- Jian, M. et al. (2008) 'The infinite sites model of genome evolution', *Proceedings of the National Academy of Sciences*, 105: 14254–14261.
- Jukes, T. H. and C. R., Cantor (1969) *Mammalian Protein Metabolism. Evolution of protein molecules*, Elsevier, pp. 21–132.
- Karumathil, S. et al. (2018) 'Evolution of synonymous codon usage bias in West African and Central African strains of Monkeypox virus', *Evolutionary Bioinformatics*, 14: 117693431876136.
- King, S. et al. (2010) BLAST tree: Fast filtering for genomic sequence classification, 2010 *IEEE International Conference on Bioinformatics and BioEngineering*, IEEE.
- Kreger, J., N. L., Komarova and D., Wodarz (2021) 'A hybrid stochastic-deterministic approach to explore multiple infection and evolution in HIV', *PLOS Computational Biology*, 17: 1–26.
- Lewontin, R. C. and K. V., Krimpas et al. (2000) *Evolutionary Genetics: From Molecules to Morphology*. Vol.1, Cambridge University Press.
- Lord, J. S. and M. B., Bonsall (2021) 'The evolutionary dynamics of viruses: virion release strategies, time delays and fitness minima', *Virus Evolution*, 7: veab039.
- Moorjani, P. et al. (2016) 'Variation in the molecular clock of primates', *Proceedings of the National Academy of Sciences*, 113: 10607–10612.
- Murrell, B. et al. (2012) 'Modeling HIV-1 drug resistance as episodic directional selection', *PLoS Computational Biology*, 8: e1002507.
- Neher, R. A. (2022) Contributions of adaptation and purifying selection to SARS-CoV-2 evolution, <https://www.biorxiv.org/content/10.1101/2022.08.22.504731v1>, accessed 16 Oct 2022.
- Nguyen, L. -T. et al. (2014) 'IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies', *Molecular Biology and Evolution*, 32: 268–274.
- Nourmohammad, A., S., Schiffels and M., Lässig (2013) 'Evolution of molecular phenotypes under stabilizing selection', *Journal of Statistical Mechanics: Theory and Experiment*, 01: 01012.
- O'Toole, Á. Rambaut, A. (2022a) Initial observations about putative APOBEC3 deaminase editing driving short-term evolution of MPXV since 2017. <https://virological.org/t/initial-observations-about-putative-apobec3-deaminase-editing-driving-short-term-evolution-of-mpxv-since-2017/830>, accessed 16 Oct 2022.
- O'Toole, Á. Rambaut, A. (2022b) Update to observations about putative APOBEC3 deaminase editing in the light of new genomes from USA. <https://virological.org/t/update-to-observations-about-putative-apobec3-deaminase-editing-in-the-light-of-new-genomes-from-usa/847>, accessed 16 Oct 2022.
- O'Toole, Á. Rambaut, A. (2022c) An APOBEC3 molecular clock to estimate the date of emergence of hMPXV. <https://virological.org/t/an-apobec3-molecular-clock-to-estimate-the-date-of-emergence-of-hmpxv/885>, accessed 16 Oct 2022.
- Peng, Q. et al. (2023) 'Structure of monkeypox virus DNA polymerase holoenzyme', *Science*, 379: 100–105.
- Posada, D. (2013) 'Phylogenetic models of molecular evolution: Next-generation data, fit, and performance', *Journal of Molecular Evolution*, 76: 351–352.
- Qin, Y. et al. (2004) 'Single-strand specificity of APOBEC3G accounts for minus-strand deamination of the HIV genome', *Nature Structural & Molecular Biology*, 11: 435–442.
- Rambaut, A. et al. (2016) 'Exploring the temporal structure of heterochronous sequences using TempEst (formerly path-o-gen)', *Virus Evolution*, 2: vew007.
- Sadeghpour, S. et al. (2021) 'Human APOBEC3 variations and viral infection', *Viruses*, 13: 1366.
- Spielman, S. J. and C. O., Wilke (2016) 'Extensively parameterized mutation-selection models reliably capture site-specific selective constraint', *Molecular Biology and Evolution*, 33: 2990–3002.
- Stenglein, M. D. et al. (2010) 'APOBEC3 proteins mediate the clearance of foreign DNA from human cells', *Nature Structural & Molecular Biology*, 17: 222–229.
- Tavaré, S. (1986) 'Some probabilistic and statistical problems in the analysis of DNA sequences', *Lect Math Life Sci (Am Math Soc)*, 17: 57–86.
- Thankeswaran Parvathy, S., V., Udayasuriyan and V., Bhadana (2021) 'Codon usage bias', *Molecular Biology Reports*, 49: 539–565.
- Wallace, E. W. J., E. M., Airoidi and D., Allan Drummond (2013) 'Estimating selection on synonymous codon usage from noisy experimental data', *Molecular Biology and Evolution*, 30: 1438–1453.
- Wang, X. et al. (2022) 'Low intrahost and interhost genetic diversity of carnivore protoparvovirus 1 in domestic cats during a feline panleukopenia outbreak', *Viruses*, 14: 1412.
- Wen-Hsiung, L. et al. (1996) 'Rates of nucleotide substitution in primates and rodents and the generation-time effect hypothesis', *Molecular Phylogenetics and Evolution*, 5: 182–187.
- World Health Organization WHO. (2022) Monkeypox fact sheet. <https://www.who.int/news-room/fact-sheets/detail/monkeypox>, accessed 16 Oct 2022.
- Yang, Z. (1994) 'Estimating the pattern of nucleotide substitution', *Journal of Molecular Evolution*, 39: 105–111.
- Yang, Z. (2014) *Molecular Evolution*. Oxford University Press Oxford, .
- You, Y. -H. (2000) 'Cyclobutane pyrimidine dimers form preferentially at the major p53 mutational hotspot in UVB-induced mouse skin tumors', *Carcinogenesis*, 21: 2113–7.
- Zhang, J. (2005) 'Evaluation of an improved branch-site likelihood method for detecting positive selection at the molecular level', *Molecular Biology and Evolution*, 22: 2472–2479.