

# UCSF

## UC San Francisco Previously Published Works

### Title

Sexually Dimorphic Neurons in the Ventromedial Hypothalamus Govern Mating in Both Sexes and Aggression in Males

### Permalink

<https://escholarship.org/uc/item/53234547>

### Journal

Cell, 153(4)

### ISSN

0092-8674

### Authors

Yang, Cindy F  
Chiang, Michael C  
Gray, Daniel C  
[et al.](#)

### Publication Date

2013-05-01

### DOI

10.1016/j.cell.2013.04.017

Peer reviewed



Published in final edited form as:

Cell. 2013 May 9; 153(4): 896–909. doi:10.1016/j.cell.2013.04.017.

## Sexually dimorphic neurons in the ventromedial hypothalamus govern mating in both sexes and aggression in males

Cindy F. Yang<sup>1,2</sup>, Michael Chiang<sup>2</sup>, Daniel C. Gray<sup>3</sup>, Mahalakshmi Prabhakaran<sup>2</sup>, Maricruz Alvarado<sup>2</sup>, Scott A. Juntti<sup>1</sup>, Elizabeth K. Unger<sup>4</sup>, James A. Wells<sup>3,5</sup>, and Nirao M. Shah<sup>1,2,4</sup>

<sup>1</sup>Program in Neuroscience, University of California, San Francisco, San Francisco, CA 94158

<sup>2</sup>Dept. of Anatomy, University of California, San Francisco, San Francisco, CA 94158

<sup>3</sup>Program in Chemistry and Chemical Biology, University of California, San Francisco San Francisco, CA 94158

<sup>4</sup>Program in Biomedical Sciences, University of California, San Francisco San Francisco, CA 94158

<sup>5</sup>Depts. of Pharmaceutical Chemistry and Cellular and Molecular Pharmacology, University of California, San Francisco, San Francisco, CA 94158

### SUMMARY

Sexual dimorphisms in the brain underlie behavioral sex differences, but the function of individual sexually dimorphic neuronal populations is poorly understood. Neuronal sexual dimorphisms typically represent quantitative differences in cell number, gene expression, or other features, and it is unknown if these dimorphisms control sex-typical behavior in one sex exclusively or in both sexes. The progesterone receptor (PR) controls female sexual behavior, and we find many sex differences in number, distribution, or projections of PR-expressing neurons in the adult mouse brain. We have ablated one such PR-expressing neuronal population located in the ventromedial hypothalamus (VMH) using a novel genetic strategy. Ablation of these neurons in females greatly diminishes sexual receptivity. Strikingly, the corresponding ablation in males reduces mating and aggression. Our findings reveal the functions of a molecularly-defined, sexually dimorphic neuronal population in the brain. Moreover we show that sexually dimorphic neurons can control distinct sex-typical behaviors in both sexes.

### INTRODUCTION

Males and females show sex differences in many behaviors, including mating and aggression, that result from sexually dimorphic development or activation of the underlying neural circuits. Gonadal sex hormones exert a profound influence on vertebrate sex-typical

© 2013 Elsevier Inc. All rights reserved.

\*Correspondence: nms@ucsf.edu.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

behaviors by controlling sex differences in the brain (Cooke et al., 1998; Dewing et al., 2003; Gagnidze et al., 2010; Jazin and Cahill, 2010; McCarthy and Arnold, 2011; Morris et al., 2004; Simerly, 2002; De Vries, 1990; Xu et al., 2012; Yang et al., 2006). Most behaviors and neural circuits are shared between the sexes such that sexually dimorphic neuronal clusters represent a small fraction of the neurons within larger brain regions. It has therefore been difficult to discern which dimorphic, hormone-responsive neurons in the brain control each of the various sex differences in physiology and behavior. In addition, neuronal sex differences usually represent quantitative rather than all-or-nothing dimorphisms in gene expression or cytological features. It is presently unclear whether such groups of dimorphic neurons regulate gender-typical behaviors in one or both sexes.

Progesterone controls female reproduction, including sexual receptivity, by signaling via its cognate receptor, PR (Levine et al., 2001; Mani et al., 1997). PR is widely distributed in the brain, and the PR+ neurons that regulate sexual receptivity remain to be identified unambiguously (Blaustein and Feder, 1979; Olster and Blaustein, 1990; Quadros et al., 2008). The VMH, which contains a small pool of PR+ neurons in its ventrolateral division (VMHvl), is well characterized for its relevance to female mating in mammals (Blaustein, 2008; Cohen and Pfaff, 1992; Flanagan-Cato, 2011; Rubin and Barfield, 1983). Studies with c-Fos suggest that many VMHvl neurons, including a subset of PR+ neurons, are activated following female mating (Flanagan-Cato et al., 2006). However, lesions or manipulations of neuronal activity of the VMH can lead to no change, a decrease, or an increase in female sexual behavior (Goy and Phoenix, 1963; Kow et al., 1985; Leedy and Hart, 1985; Mathews and Edwards, 1977a, 1977b; Musatov et al., 2006; Pfaff and Sakuma, 1979a, 1979b; Robarts and Baum, 2007; La Vaque and Rodgers, 1975). Some studies also report a concurrent increase in body weight, suggesting a complex role of this region in feeding and mating (King, 2006; Musatov et al., 2007). This phenotypic diversity is likely due to manipulations that variably affect the heterogeneous neuronal subsets within the VMH (Kurrasch et al., 2007), adjacent brain regions, and fibers of passage. Given these challenges, the identity and function of VMHvl neurons that specifically influence female mating remain unclear.

In accord with the notion that the VMHvl influences female sexual behavior, the VMHvl exhibits quantitative cell and molecular sex differences (Dugger et al., 2007; Grgurevic et al., 2012; Matsumoto and Arai, 1983, 1986; Patisaul et al., 2008; Wu et al., 2009; Xu et al., 2012). Intriguingly, lesions or manipulations of neural activity of the VMH or the surrounding neurons have long suggested an important role of this region in controlling aggression (Hess and Akert, 1955; Kruk et al., 1979; Reeves and Plum, 1969; Wheatley, 1944). In fact, this region is activated during male aggression, and correspondingly, electrical activation or inhibition elicits or inhibits fighting, respectively (Kollack-Walker and Newman, 1995; Lin et al., 2011; Veening et al., 2005). However, as with VMH neurons that regulate female receptivity, the identity of VMH neurons that influence aggression is unknown. In principle, these behaviors may be regulated by a single set or by non-overlapping sets of neurons.

We utilized genetic strategies in mice to visualize PR+ neurons and to assess their contributions to mating and aggression. We find many sex differences in PR+ neurons in the adult brain, including in the VMHvl. We have developed a Cre-loxP strategy to ablate any

molecularly defined neuronal population via targeted viral delivery of a genetically engineered caspase. Using this approach, we have ablated PR<sup>+</sup> VMHvl neurons in adult females and observe a dramatic reduction in sexual receptivity. The corresponding ablation in males reduces mating and territorial aggression. Thus our results define a role of PR<sup>+</sup> VMHvl neurons in sex-typical behaviors. Moreover, we establish that a discrete, sexually dimorphic neuronal population influences sexually dimorphic behaviors in both sexes.

## RESULTS

### Visualizing PR expression in the mouse brain

We wished to identify PR<sup>+</sup> neurons at high cellular resolution. We inserted an *IRES-PLAP-IRES-nuclear LacZ (PL)* reporter into the 3' UTR of *PR* using gene targeting (Figures 1A, B). As described previously (Shah et al., 2004), this cassette permits expression of placental alkaline phosphatase (PLAP), which labels neuronal processes, and nuclear targeted  $\beta$ -galactosidase ( $\beta$ -gal) in PR<sup>+</sup> cells. This strategy maintains the expression and function of PR and permits examination of PR<sup>+</sup> neurons in otherwise wildtype (WT) mice. Accordingly, and in contrast to *PR*<sup>-/-</sup> mice (Chappell et al., 1997; Lydon et al., 1995), *PR*<sup>PL/PL</sup> females were similar to WT females in fecundity and also maintained normal sex hormone titers (Figure 1C–G).

In the forebrain, we observed  $\beta$ -gal activity in pools of neurons in specific hypothalamic nuclei, posterodorsal medial amygdala (MeApd), medial division of the posteromedial bed nucleus of the stria terminalis (BNSTmpm), various cortical areas, basal ganglia, and dentate gyrus (Figures 1H–M, 2, S1). This distribution of cells mirrors the expression pattern of PR mRNA in adjacent sections (Figure 1H–M). In regions such as the basal ganglia with low level PR expression that precludes visualization by in situ hybridization, we can detect PR message by RT-qPCR (Figure S1A). The distribution of  $\beta$ -gal<sup>+</sup> cells is in accord with histological and pharmacological studies (Becker, 1999; Blaustein and Feder, 1979; Olster and Blaustein, 1990; Quadros et al., 2008). In the case of the basal ganglia, our studies localize PR expression to sparsely distributed neurons across the rostrocaudal axis (Figure S1B–D). In addition, we find unreported PR<sup>+</sup> neuronal pools scattered within the basal forebrain (Figure 2), an observation confirmed by RT-qPCR from this region (Figure S1A). The ~1 week  $t_{1/2}$  of  $\beta$ -gal in neurons precludes detection of PR mRNA changes across the 4–6 day estrous cycle (Allen, 1922; Smith et al., 1995). However, the long  $t_{1/2}$  and superb signal:noise of  $\beta$ -gal labeling allows sensitive detection of PR expression. Together, the *PR*<sup>PL</sup> reporter mouse confirms and extends previous reports of PR expression in the mouse brain.

### Widespread sex differences in the distribution and cell number of PR<sup>+</sup> neurons

We observed previously unreported as well as known sex differences in PR<sup>+</sup> cells in the adult *PR*<sup>PL</sup> brain (Figures 2, S2A, and Table S1). We found more PR<sup>+</sup> cells in the female preoptic area (POA), the adjacent anteroventral periventricular hypothalamic nucleus (AVPV), arcuate nucleus, and VMHvl (Figure 2A–L). The VMHvl contains cells expressing the estrogen receptor alpha (ER $\alpha$  or *Esr1*) (Xu et al., 2012), and we find that >92% PR<sup>+</sup> neurons co-label for ER $\alpha$  in both sexes (Figure S2B). We asked whether PR<sup>+</sup> VMHvl

neurons expressed *Cckar*, a GPCR required for sexual receptivity and expressed in the female but essentially absent in the male VMHvl (Xu et al., 2012). We observed that  $67\% \pm 3$  (Mean  $\pm$  SEM) of PR+ VMHvl cells co-labeled with *Cckar* whereas  $96\% \pm 0.2$  of *Cckar*+ VMHvl cells were PR+ ( $n = 3$  *PR<sup>PL/PL</sup>* females, 500 cells analyzed/brain) (Figure S2C–E). Thus PR+ neurons represent the vast majority of VMHvl neurons that express *Cckar*, a gene required for female mating.

We observed many clusters of PR+ cells (~15–40 cells/cluster) in the male but not female basal forebrain (Figure 2M–P). Together with a sex difference in androgen receptor expression in this region (Shah et al., 2004), our findings suggest an unappreciated role of the basal forebrain in responding to sex hormones. We also found more PR+ cells in the male BNSTmpm and MeApd (Figure 2Q–X). This increased PR expression is surprising because there is little circulating progesterone in males; our findings are nevertheless consistent with studies indicating a role of PR in male behaviors (Phelps et al., 1998; Schneider et al., 2005, 2009; Witt et al., 1995). As suggested previously (Mani et al., 1994a; Power et al., 1991; Tsutsui, 2012), PR may function in a progesterone-independent manner or locally synthesized progesterone may activate PR in males. Consistent with these sex differences in PR expression, the POA, BNSTmpm, MeApd, arcuate nucleus, and VMHvl have been implicated in sex differences in behavior or physiology (Cooke et al., 1998; Morris et al., 2004; Simerly, 2002), and PR+ neurons in these regions could contribute to such sexually dimorphic output.

We find that the dimorphic PR+ cells co-label with pan-neuronal markers (Figure S2F). However, within any given brain region expressing PR dimorphically, only a subset of neurons is PR+. Even within the VMHvl, only  $49\% \pm 4$  of NeuN+ cells co-label with PR ( $n = 3$  brains,  $10^3$  NeuN+ cells analyzed for PR/brain). There is a sex difference in the soma size of thionin-labeled neurons within the rat VMHvl (Dugger et al., 2007). However, there was no such sex difference in PR+ VMHvl neurons (Figure S2G), suggesting a species difference or that other VMHvl neurons account for this dimorphism. The sex differences in PR expression cannot result solely from sex differences in neuronal numbers. Indeed, no sex difference in neuronal number has been reported in the basal forebrain or VMHvl, and in the POA and arcuate nucleus, which contain more neurons in males (Gorski et al., 1980; Leal et al., 1998), we find more PR+ neurons in females. Finally, the 3–4 fold more PR+ neurons in the male BNSTmpm and MeApd exceeds the <2-fold more neurons in these regions in males (Morris et al., 2008; Shah et al., 2004; Wu et al., 2009). Thus, our studies confirm known sex differences (POA, VMHvl, arcuate nucleus, MeApd) (Blaustein et al., 1980; Brown et al., 1996; Grgurevic et al., 2012; Kudwa et al., 2009; Quadros et al., 2002) and reveal new sexual dimorphisms in PR expression (basal forebrain, BNSTmpm) in the mammalian brain.

### Visualizing sex differences in projections of PR+ neurons

We determined whether sexually dimorphic PR+ neurons projected to distinct locations in the two sexes. Consistent with PR expression in interconnected regions such as the POA, BNST, MeA, and VMHvl, we observed a rich distribution of PLAP+ fibers in the *PR<sup>PL/PL</sup>* forebrain (data not shown) that precluded identification of dimorphic projection patterns.

We devised a genetic strategy to visualize the projections of any subset of PR<sup>+</sup> neurons. We first targeted an *IRES-Cre recombinase* cassette to the 3' UTR of *PR* (Figures 3A, S3A, B). As expected, these *PR-IRES-Cre (PR<sup>Cre</sup>)* mice, like *PR<sup>PL</sup>* mice, are viable and fertile, and Cre expression mirrors that of PR in the brain (Figure S3C–F). We also designed a lentiviral vector that expresses PLAP in a Cre-dependent manner (Lenti-lxlap, Figures 3A, S3G). This lentivirus is replication-incompetent and integrates into the host genome, properties that restrict PLAP expression to Cre<sup>+</sup> cells for the life of the cells. This virus infects cells in both WT and *PR<sup>Cre</sup>* mice, but we only observe PLAP expression in *PR<sup>Cre</sup>* mice (Figure 3B–E).

The VMH has been implicated in sex-specific behaviors, and we therefore traced the projections of PR<sup>+</sup> VMHvl neurons in adults. We initially determined that we could visualize maximal expression of PLAP 7–8 days following delivery of Lenti-lxlap into the VMH (CFY, unpublished observations). Such injections revealed the soma and local arbors of PR<sup>+</sup> VMHvl neurons (Figure 3F–I). In contrast to the wide-ranging projections of the entire VMH (Saper et al., 1976; Krieger et al., 1979), we observed PLAP<sup>+</sup> projections of PR<sup>+</sup> VMHvl neurons in the AVPV and adjacent periventricular area, POA, and periaqueductal gray (PAG) (Figure 3J–U). Unlike PR<sup>+</sup> VMHvl projections in the guinea pig (Ricciardi and Blaustein, 1994), mouse PR<sup>+</sup> VMHvl neurons did not appear to project appreciably to the BNST or MeA, suggesting subtle species differences in these cells. Although we observed similar localization of PLAP<sup>+</sup> projections of PR<sup>+</sup> VMHvl neurons in both sexes (Figures 3J–W, S3H, and Table S2), there was a striking, previously unreported 7-fold increase in PLAP<sup>+</sup> fibers in the female AVPV (Figure 3J–M). This sex difference cannot solely result from the dimorphism (~30%) in PR<sup>+</sup> VMHvl cell number. In fact, we even observed the dimorphic AVPV projection in *PR<sup>Cre</sup>* females in whom a few PR<sup>+</sup> VMHvl neurons had been infected. Thus, more PR<sup>+</sup> female VMHvl neurons project to the AVPV or their axonal termini arborize more extensively. The AVPV is thought to control ovulation, and the PAG can regulate sexual receptivity in females (Sakuma and Pfaff, 1979; Simerly, 2002). In summary, PR<sup>+</sup> VMHvl neurons project to a subset of VMH targets, their efferents are sexually dimorphic, and each of their targets can influence sexually dimorphic behaviors or physiology.

### A novel genetic approach to ablate adult neurons in vivo

We determined the requirement of PR<sup>+</sup> VMHvl neurons in sex-typical behaviors by targeting Cre-dependent, virally encoded toxins to the VMHvl of *PR<sup>Cre</sup>* mice. Initial studies suggested that virally encoded diphtheria toxin A or tBid (Jiang and Wang, 2004; Maxwell et al., 1986) were partially effective in ablating PR<sup>+</sup> neurons in vivo even though they were effective in tissue culture cells (CFY, unpublished observations). We therefore employed a genetically engineered caspase-3, a caspase whose activation commits a cell to apoptosis, to kill adult neurons in vivo (Figure 4A) (Gray et al., 2010). Endogenous caspase-3 normally exists as pro-caspase-3, and apoptotic signals activate upstream caspases that cleave pro-caspase-3 into its active form (Figure 4A). Our designer pro-caspase-3, pro-taCasp3, lacks the cleavage site for upstream caspases and encodes a cleavage site for the heterologous enzyme Tobacco Etch Virus protease (TEVp). Provision of TEVp activates pro-taCasp3 into the apoptosis-inducing taCasp3. We generated an adeno-associated virus (AAV) to drive

expression of pro-taCasp3 and TEVp in a Cre-dependent manner (Figures 4B, S4A) (Atasoy et al., 2008). This virus (AAV-flex-taCasp3-TEVp) utilizes the T2A peptide encoding sequence to ensure bi-cistronic expression of pro-taCasp3 and TEVp. Importantly, taCasp3 triggers cell-autonomous apoptosis, thereby minimizing toxicity to adjacent non-Cre+ cells (Gray et al., 2010).

Infection of HEK293T cells with this virus leads to rapid Cre-dependent cell death (Figure 4C, D). We next tested whether this virus could ablate adult PR+ neurons by stereotaxically targeting it to the VMHvl of adult *PR<sup>+/PL</sup>* or *PR<sup>Cre/PL</sup>* mice. PR+ VMHvl neurons appeared unaffected in controls but were essentially completely lost in *PR<sup>Cre/PL</sup>* mice 2–4 weeks following viral delivery (Figures 4E, F, S4B). We tested whether the taCasp3-encoding AAV targeted to the VMHvl diffused to and ablated PR+ cells in distant hypothalamic regions. We therefore enumerated PR+ cells along the rostrocaudal extent of the hypothalamus in a cohort of virally injected control and *PR<sup>Cre</sup>* mice. This analysis revealed no difference in PR+ cell counts between *PR<sup>Cre</sup>* and control females (number of PR+ cells: Control,  $619 \pm 60$  and *PR<sup>Cre</sup>*,  $679 \pm 150$ ;  $n = 5/\text{cohort}$ ,  $p = 0.7$ ). Thus taCasp3-mediated ablation appears restricted to the vicinity of the injection site. We observed local spread of the virus to the arcuate and present these findings below. In separate experiments, we found that stereotaxic delivery of the taCasp3-encoding virus ablated Cre+ neurons in different brain regions (CFY, EKV, and MC, unpublished observations), indicating that we have devised a general strategy for targeted ablation of Cre+ cells.

### The dimorphic PR+ VMHvl cluster of neurons regulates female sexual behavior

We tested the role of PR+ VMHvl neurons in female mating. We targeted AAV-flex-taCasp3-TEVp bilaterally to the VMHvl of adult *PR<sup>Cre</sup>* and control females (Figure 5A). To assure optimal sexual receptivity, females were ovariectomized at the time of viral injection and, following recovery, hormonally primed to be in estrus when tested with WT males.

We observed a marked diminution of female sexual behavior in such *PR<sup>Cre</sup>* females (Figure 5B–G, Movies S1 and S2). As in many vertebrates, female mating in mice is stereotyped and includes permitting the male to approach and mount and dorsiflexing the neck and back (lordosis) upon sensory stimulation to the dorsum (Harvey, 1951; McGill, 1962). This allows the males to intromit (penetrate, as determined by his thrust pattern) and attempt ejaculation. *PR<sup>Cre</sup>* females rejected mount attempts by kicking or running away (Figure 5B), thereby reducing the fraction of mounts that progressed to intromission (receptivity index, Figure 5C). In sharp contrast to controls, *PR<sup>Cre</sup>* females walked around during intromission, lordosed rarely, and with a >20-fold reduction in lordosis duration (Figures 5D–F). This reduced sexual behavior of *PR<sup>Cre</sup>* females affected the WT male partner's performance (Figure 5H–J). Males were interested in both *PR<sup>Cre</sup>* and control females, initiating anogenital sniffing, mounting, and intromission equivalently, but were less successful in ejaculating with the former (Figures 5H, S5A, S5B). Accordingly, males intromitted only briefly with *PR<sup>Cre</sup>* females even though they mounted the females more and for longer duration (Figure 5I, J). Correspondingly the total duration of intromission per assay was also reduced (Control,  $279 \text{ s} \pm 41$  and *PR<sup>Cre</sup>*,  $121 \text{ s} \pm 19$ ;  $n = 10$ ,  $p = 3 \times 10^{-3}$ ). In summary,

targeted ablation of adult PR+ VMHvl neurons leads to a significant diminution in female mating.

We next assessed the ablation of PR+ VMHvl cells in these *PR<sup>Cre</sup>* females. We observed that most ( $97\% \pm 1$ ;  $n = 10$  control and 16 *PR<sup>Cre</sup>* females) PR+ VMHvl neurons were ablated upon injection of the taCasp3-encoding AAV into *PR<sup>Cre</sup>* females (Figure 5G). Co-injection of this AAV and a constitutively expressed EGFP-encoding AAV revealed spread to the adjacent arcuate nucleus, which contains PR+ neurons (Figure 2I–L) and controls feeding and the estrous cycle (Atasoy et al., 2012; Simerly, 2002). Consistent with the lack of estrous cycle or body weight phenotypes in *PR<sup>Cre</sup>* mice (see below and Figure S5), our injections spared most PR+ arcuate neurons in *PR<sup>Cre</sup>* females ( $74\% \pm 12$  of controls). There was no correlation in the extent of loss of PR+ arcuate neurons and reduced sexual receptivity ( $R^2 = 5 \times 10^{-3}$ ,  $p = 0.8$ ). Moreover, we found that *PR<sup>Cre</sup>* females ( $n = 7$ ) in whom the number of PR+ arcuate neurons was indistinguishable from controls also rejected males and displayed reduced sexual receptivity (Rejections/assay: Controls,  $1 \pm 1$  and *PR<sup>Cre</sup>* females,  $35 \pm 7$ ;  $p = 6 \times 10^{-5}$ ,  $n = 7$ ; Receptivity Index: Controls, 0.5 and *PR<sup>Cre</sup>* females  $0.2 \pm 0.1$ ,  $p = 3 \times 10^{-3}$ ,  $n = 7$ ). Thus PR+ VMHvl neurons are required for normal female sexual behavior.

We tested the specificity of the behavioral deficit in *PR<sup>Cre</sup>* females following ablation of PR + VMHvl neurons. Despite their reduced sexual receptivity, these mice sniffed and groomed males normally (Figure S5C, D) (Groom duration: Control,  $2 \text{ s} \pm 1$  and *PR<sup>Cre</sup>*,  $5 \text{ s} \pm 1$ ,  $n = 10$ ,  $p = 0.3$ ). There were no overt deficits in tests of anxiety, motivated behavior, motor coordination, and locomotor activity (Figure S5E–H). In contrast to the weight gain subsequent to a VMH lesion (Dhillon et al., 2006; Hetherington and Ranson, 1940; King, 2006; Majdic et al., 2002), *PR<sup>Cre</sup>* females maintained body weight similar to controls upon ablation of PR+ VMHvl neurons (Figure S5I). Thus, we have partitioned the VMHvl to reveal that PR+ VMHvl neurons are required for normal levels of female sexual receptivity but not for all social or other behaviors and physiology.

In separate studies we ablated PR+ VMHvl neurons but left the ovaries intact to examine whether other female-typical behaviors are regulated by these neurons. This ablation did not disrupt the estrous cycle as assayed by vaginal cytology (Figure S5J). To test for maternal behaviors, we obtained litters from *PR<sup>Cre</sup>* and control females by co-housing them with WT males. Similar to control females, *PR<sup>Cre</sup>* females displayed various elements of maternal care toward their litters, including pup retrieval and aggression toward unfamiliar intruders in their cage (Figure S5K–O). Our results therefore show that ablation of PR+ VMHvl neurons reduced female sexual displays without overt disruption of other female-typical behaviors and physiology.

PR regulates female mating (Lydon et al., 1995), and our findings suggest that it functions in the VMHvl to do so, consistent with prior work (Mani et al., 1994a, 1994b; Ogawa et al., 1994; Pollio et al., 1993). Cckar is also required for female mating (Xu et al., 2012). Most Cckar+ VMHvl neurons are PR+ (Figure S2C–E), resulting in a near-complete loss of these cells upon ablation of PR+ VMHvl neurons (Figure 5K, L). It is possible that PR or Cckar act elsewhere to control female mating, and these genes only mark a pool of VMHvl



neurons that controls this behavior. We favor a more parsimonious model in which PR and Cckar function in the VMHvl to regulate female mating. In any event, our findings show that PR+, Cckar+ VMHvl neurons are essential for high, WT levels of female sexual behavior.

### The dimorphic PR+ VMHvl cluster of neurons regulates mating and aggression in males

The VMHvl has been implicated in regulating female mating and male fighting. PR+ neurons represent ~50% of VMHvl neurons and these regulate female mating (Figure 5), but fighting could be controlled by PR+ or PR- VMHvl cells. We tested whether PR+ VMHvl neurons regulate male behaviors by ablating them with the taCasp3-encoding AAV (Figure 6A). *PR<sup>Cre</sup>* and control males were allowed to recover for 4 weeks following viral delivery, singly housed, and tested for mating and fighting.

*PR<sup>Cre</sup>* and control males initiated mounting intruder females equivalently, but *PR<sup>Cre</sup>* males were less likely to intromit (Figures 6B and S6A). The reduced intromissions likely resulted from the fewer mounts exhibited by *PR<sup>Cre</sup>* males (Figure 6C). Even when these males intromitted, there was a decrease in the number and duration of intromissions (Figures 6C, D, S6B). The decreased intromission count was significant ( $n = 16/\text{cohort}$ ;  $p = 5 \times 10^{-3}$ ) even when normalized to the fewer mounts. Thus, ablation of male PR+ VMHvl neurons leads to specific deficits in consummatory elements of mating. This phenotype is not accompanied by deficits in presumptively appetitive behaviors such as sniffing (Figure 6B, S6C–E), sex discrimination, or territory marking. There was no difference between *PR<sup>Cre</sup>* and control males in sex discrimination as shown by predominantly female-directed ultrasonic vocalization (Figure 6E) (Nyby et al., 1977). Both *PR<sup>Cre</sup>* and control males also marked their territory equivalently (Figure 6F, G) (Desjardins et al., 1973; Kimura and Hagiwara, 1985). Together, PR+ VMHvl neurons are essential for the normal display of male sexual behavior.

We tested whether ablation of PR+ VMHvl neurons disrupted aggression toward a WT male intruder. *PR<sup>Cre</sup>* males exhibited a >2-fold reduction in the probability of initiating aggression compared to controls (Figure 7A). Even when *PR<sup>Cre</sup>* males fought, they attacked less, for a shorter duration, and with a longer inter-attack interval (Figure 7B–D). Male fighting includes tail rattles and overt attacks such as biting. Control and *PR<sup>Cre</sup>* residents rattled their tails equivalently, but *PR<sup>Cre</sup>* males bit the intruders over 3-fold less (Figure 7E). Thus ablation of PR+ VMHvl neurons significantly reduces male aggression.

We assessed the ablation of PR+ VMHvl neurons in males tested behaviorally. Most of these neurons ( $95\% \pm 1$ ;  $n = 14$  control and 35 *PR<sup>Cre</sup>* males) were ablated in *PR<sup>Cre</sup>* males (Figure 7F–H) whereas PR+ arcuate neurons were largely spared ( $92\% \pm 12$  of controls). There was no correlation in the extent of loss of PR+ neurons in the arcuate and the reduced mating or fighting (mating,  $R^2 = 4 \times 10^{-4}$ ,  $p = 0.9$ ; fighting,  $R^2 = 2 \times 10^{-2}$ ,  $p = 0.7$ ). *PR<sup>Cre</sup>* males ( $n = 15$ ) in whom the number of PR+ arcuate neurons was indistinguishable from controls also exhibited deficits in mating and fighting (Percent males intromitting: Controls, 67% and *PR<sup>Cre</sup>* males, 27%,  $n = 15$ ,  $p = 0.02$ ; Percent males attacking: Controls, 75% and *PR<sup>Cre</sup>* males, 20%,  $n = 15$ ,  $p = 1 \times 10^{-3}$ ). Taken together, our findings demonstrate that PR+ VMHvl neurons control the normal display of male mating and fighting.

We tested the specificity of the deficits in *PR<sup>Cre</sup>* males following ablation of PR+ VMHv1 neurons. Despite deficits in mating and fighting, these males sniffed and groomed intruders in a WT manner (Figures 6B, 7A, S6C–E, S7A–C). *PR<sup>Cre</sup>* males performed at WT levels in assays of anxiety, motivated behavior, motor coordination, and locomotor activity (Figure S7D–G). These males maintained normal body weight, and there was no change in the weight of gonads, seminal vesicles, and serum testosterone titers (Figure S7H–J). Thus, PR+ VMHv1 neurons are specifically required in males for the high, WT levels of mating and aggression.

## DISCUSSION

We have identified a small, sexually dimorphic cluster of ~2,000 PR+ hypothalamic neurons that is essential for the normal display of sexual receptivity in females and sexual and aggressive behaviors in males. Our findings directly demonstrate that sexually dimorphic neurons in the brain influence dimorphic behaviors. Moreover these PR+ neurons are functionally bivalent in that they regulate distinct dimorphic behaviors in the two sexes.

### Control of social behaviors by the VMH

Experimental studies and clinical observations have suggested that the VMH or adjacent hypothalamic regions regulate aggression and female mating (Bard, 1928; Blaustein, 2008; Clemente and Chase, 1973; Colpaert and Wiepkema, 1976; Grossman, 1972; Hess and Akert, 1955; Kow et al., 1985; Kruk et al., 1979; Lin et al., 2011; Olivier and Wiepkema, 1974; Pfaff and Sakuma, 1979a, 1979b; Reeves and Plum, 1969; Swaab, 2003; La Vaque and Rodgers, 1975; Wheatley, 1944). Despite intense scrutiny, the neurons that control these behaviors remained unidentified. In fact, whether separate or overlapping neuronal groups control these innate behaviors was also unknown. Our studies reveal the molecular identity of the long sought-after neurons in or around the VMH that influence male fighting and female mating. While other neighboring neurons may also influence these behaviors, we show that PR+ VMHv1 neurons are required for the normal display of mating in females and fighting in males. These PR+ neurons also regulate male mating. Non-targeted inhibition of neurons in this region disrupts male fighting but not mating (Lin et al., 2011), suggesting partial inactivation or incomplete targeting of the neurons that regulate male mating. By contrast, our ablation of the PR+ VMHv1 population revealed a role for these cells in male mating. Generalized arousal systems may feed into the VMH to enhance social interactions (Schober et al., 2011). We do not observe altered locomotor activity, sensorimotor coordination, or general social interactions in mice lacking PR+ VMHv1 neurons, suggesting that these neurons are unlikely to exert a major influence on neural pathways that increase such arousal. In summary, we show that PR+ VMHv1 neurons are required for the normal display of mating in both sexes and fighting in males. Given the conservation of genes and neuroanatomy across placental mammals, these VMHv1 neurons may regulate mating and aggression in many mammals, including humans.

### Distributive neural control of sexually dimorphic behaviors

It is curious that ablation of a highly restricted, molecularly defined set of neurons results in deficits in male mating and fighting. These PR+ neurons may integrate social cues relevant

to both behaviors, allowing males to mate or fight appropriately. Such dual control could also reflect further diversity within PR+ VMHvl neurons such that subsets of these neurons regulate one or the other behavior. In fact, in vivo recordings and c-Fos studies (Lin et al., 2011) reveal male VMHvl neurons that are activated during encounters with both sexes as well as neurons that appear responsive to either male or female encounters.

We find that different components of male behaviors require distinct neuronal populations. Males lacking PR+ VMHvl neurons have a male behavioral repertoire: they distinguish between the sexes with vocalizations (Stowers et al., 2002), attack males, and mate with females. Moreover, these males mark territory like WT males, thereby providing an objective indicator that their internal representation of sexual identity is masculine. Nevertheless these males display specific deficits in mating and fighting, indicating that ablation of PR+ VMHvl neurons dissociates the repertoire of masculine behaviors. Such partial behavioral deficits could reflect compensatory mechanisms activated upon the loss of these neurons. However, acute inactivation of the VMH mimics the behavioral deficits we observe (Lin et al., 2011), suggesting a minimal role of compensatory mechanisms. Thus, male mating and fighting are encoded in a distributive or redundant manner in the brain. Similarly, ablation of these neurons reduces female sexual receptivity without overtly disrupting estrous cyclicity or maternal care, indicating that these behaviors and physiology may also be controlled by distinct neuronal groups. Together, our findings show that sex-typical behaviors are represented distributively, and different neuronal populations in the underlying neural circuit control specific components of these behaviors. In fact genes such as *Cckar* also control these behaviors in a modular manner; for instance, *Cckar*<sup>-/-</sup> females show reduced sexual receptivity without alterations in other behaviors or physiology (Xu et al., 2012). Thus modular control of sexually dimorphic behaviors across multiple levels, including genes and neurons, may be a general organizational principle of the underlying neural circuits.

### **Control of sex-typical behaviors by sexually dimorphic VMHvl neurons**

Studies in diverse animals have defined the relevance of particular brain regions to sex-typical behaviors (Brenowitz, 1991; Cooke et al., 1998; Ferveur et al., 1995; Kelley, 1997; Konishi, 1989; Morris et al., 2004). However, within a brain region only specific subsets of neurons are sexually dimorphic (Ng et al., 2009; De Vries and Panzica, 2006; Xu et al., 2012), and with rare exceptions in invertebrates (Kohatsu et al., 2011; von Philipsborn et al., 2011), the function of sexually dimorphic neurons is unknown. Ablation of the ~2000 sexually dimorphic PR+ VMHvl neurons, a fraction of the ~10<sup>8</sup> neurons in the mouse brain, results in specific deficits in complex social behaviors. Such specificity likely results from manipulation of a molecularly defined subset of neurons. Indeed PR+ neurons represent only ~50% of VMHvl neurons that, in turn, represent a fraction of VMH neurons.

The mechanisms whereby sexually dimorphic neurons control dimorphic behaviors are poorly understood. It is possible that PR+ VMHvl neurons represent unrelated cell types in the two sexes, as evidenced by the sex differences in cell number and distribution, projection targets, and expression of *Cckar*. This is unlikely because PR+ VMHvl neurons also share many features, including location, projection targets, gene expression (PR, ER $\alpha$ ), and

developmental lineage (Grgurevic et al., 2012). Thus, it appears that a common pool of PR+ VMHvl neurons is present in both sexes, but their sex differences may allow them to transform synaptic inputs in a sex-specific manner or to relay either male or female-specific input to drive sexually dimorphic behavioral output.

Most behaviors are common to the sexes, suggesting that each sex possesses the motor pathways to display dimorphic behaviors of the opposite sex. Most sex differences in the brain represent quantitative and not all-or-none cellular or molecular sex differences. It is unknown whether these shared but dimorphic neurons regulate sex-typical behaviors in both sexes. Alternately, such neurons may regulate a dimorphic output in one sex, and in the other sex, they may be functionally vestigial, subserve a non-dimorphic function, or suppress a function of the opposite sex (De Vries and Boyle, 1998). We show that PR+ VMHvl neurons are functionally bivalent in the sense that they control sex-typical behaviors in both males and females. This dual function may prove adaptive if such neurons can generate a dimorphic behavior of the opposite sex in the appropriate context; in addition, bivalence may permit facile interchange of sex-typical behaviors between the sexes during speciation. Such flexibility may underlie the rapid evolution of sexually dimorphic traits (Darwin, 1871), including behaviors such as allocation of parental care and social dominance hierarchies. Given such evolutionary considerations, it remains to be seen whether all sexually dimorphic neuronal populations control sex-typical behaviors in both sexes.

## EXPERIMENTAL PROCEDURES

### Viruses

**AAV-flex-taCasp3-TEVp**—The plasmid encoding AAV-flex-taCasp3-TEVp (Figure S4A) was generated using routine subcloning. High titer virus of serotype 2/1 ( $3 \times 10^{12}$  IU/mL) was generated from the plasmid at the UNC, Chapel Hill Vector Core.

**Lenti-IxIplap**—The plasmid encoding this VSVG pseudo-typed lentivirus was generated using standard subcloning (Figure S3G). High titer virus ( $\sim 10^8$  IU/mL) was generated using standard protocols (Barde et al., 2001).

### Stereotaxic surgery

The virus was stereotaxically delivered under anesthesia to the VMHvl (Coordinates: rostrocaudal,  $-1.48$  mm; mediolateral,  $\pm 0.78$  mm; depth,  $5.8$  mm; see also Supplemental Procedures) (Paxinos and Franklin, 2003). Injections of taCasp3-encoding AAV were spiked (9:1) with constitutive EGFP-encoding AAV to verify accuracy of the injection placement in control and *PR<sup>Cre</sup>* mice.

### Behavior

Testing was performed as described previously (Juntti et al., 2010; Wu et al., 2009; Xu et al., 2012) (see also Supplemental Procedures). To test for sexual receptivity, females were castrated and, subsequent to estrus induction with estrogen and progesterone, inserted singly into the home cage of a sexually experienced WT male. Lordosis was defined as the female

holding still with a dorsiflexed neck while being intromitted. Each experimental cohort included a set of control and *PR<sup>Cre</sup>* mice.

### Supplemental Experimental Procedures

Details regarding animals, histology, data analyses, and the procedures described above can be found in the Supplemental Information. All animal studies were in accordance with IACUC protocols at UCSF.

### Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

### ACKNOWLEDGMENTS

We thank C. Saper for sharing reagents; A. Lasek and U. Heberlein for a practical on stereotaxis; R. Axel for discussions; T. Clandinin, H. Ingraham, S. Lomvardas, C. Saper, and Shahlab members for comments on the manuscript; and N. Agarwal, A. Wang, and M. Borius for technical support. This work was supported by a Genentech Graduate Fellowship (CFY); NSF Graduate Fellowship (SAJ); NIH (EKU, F31NS078959); NARSAD, Program in Biomedical Breakthrough Research, Ellison Medical Foundation, and NIH (R01NS049488, R01NS083872, and DP1MH099900) (NMS).

### REFERENCES

- Allen E. The oestrous cycle in the mouse. *American Journal of Anatomy*. 1922; 30:297–371.
- Atasoy D, Aponte Y, Su HH, Sternson SM. A FLEX switch targets Channelrhodopsin-2 to multiple cell types for imaging and long-range circuit mapping. *J. Neurosci*. 2008; 28:7025–7030. [PubMed: 18614669]
- Atasoy D, Betley JN, Su HH, Sternson SM. Deconstruction of a neural circuit for hunger. *Nature*. 2012; 488:172–177. [PubMed: 22801496]
- Bard P. A diencephalic mechanism for the expression of rage with special reference to the sympathetic nervous system. *American Journal of Physiology*. 1928; 84:490–515.
- Barde, I.; Salmon, P.; Trono, D. In *Current Protocols in Neuroscience*. John Wiley & Sons, Inc.; 2001. Production and Titration of Lentiviral Vectors.
- Becker JB. Gender differences in dopaminergic function in striatum and nucleus accumbens. *Pharmacol. Biochem. Behav*. 1999; 64:803–812. [PubMed: 10593204]
- Blaustein JD. Neuroendocrine regulation of feminine sexual behavior: lessons from rodent models and thoughts about humans. *Annu Rev Psychol*. 2008; 59:93–118. [PubMed: 17678443]
- Blaustein JD, Feder HH. Cytoplasmic progesterin-receptors in guinea pig brain: Characteristics and relationship to the induction of sexual behavior. *Brain Research*. 1979; 169:481–497. [PubMed: 571753]
- Blaustein JD, Ryer HI, Feder HH. A sex difference in the progesterin receptor system of guinea pig brain. *Neuroendocrinology*. 1980; 31:403–409. [PubMed: 7192804]
- Brenowitz EA. Altered perception of species-specific song by female birds after lesions of a forebrain nucleus. *Science*. 1991; 251:303–305. [PubMed: 1987645]
- Brown TJ, Yu J, Gagnon M, Sharma M, MacLusky NJ. Sex differences in estrogen receptor and progesterin receptor induction in the guinea pig hypothalamus and preoptic area. *Brain Res*. 1996; 725:37–48. [PubMed: 8828584]
- Bunting M, Bernstein KE, Greer JM, Capecchi MR, Thomas KR. Targeting genes for self-excision in the germ line. *Genes Dev*. 1999; 13:1524–1528. [PubMed: 10385621]
- Chappell PE, Lydon JP, Conneely OM, O'Malley BW, Levine JE. Endocrine defects in mice carrying a null mutation for the progesterone receptor gene. *Endocrinology*. 1997; 138:4147–4152. [PubMed: 9322923]

- Clemente CD, Chase MH. Neurological substrates of aggressive behavior. *Annu. Rev. Physiol.* 1973; 35:329–356. [PubMed: 4581768]
- Cohen RS, Pfaff DW. Ventromedial hypothalamic neurons in the mediation of long-lasting effects of estrogen on lordosis behavior. *Prog. Neurobiol.* 1992; 38:423–453. [PubMed: 1589577]
- Colpaert FC, Wiepkema PR. Effects of ventromedial hypothalamic lesions on spontaneous intraspecies aggression in male rats. *Behav Biol.* 1976; 16:117–125. [PubMed: 943158]
- Cooke B, Hegstrom CD, Villeneuve LS, Breedlove SM. Sexual differentiation of the vertebrate brain: principles and mechanisms. *Front Neuroendocrinol.* 1998; 19:323–362. [PubMed: 9799588]
- Darwin, C. *The Descent of Man, and Selection in Relation to Sex.* Princeton University Press; 1871.
- Desjardins C, Maruniak JA, Bronson FH. Social rank in house mice: differentiation revealed by ultraviolet visualization of urinary marking patterns. *Science.* 1973; 182:939–941. [PubMed: 4745598]
- Dewing P, Shi T, Horvath S, Vilain E. Sexually dimorphic gene expression in mouse brain precedes gonadal differentiation. *Brain Res. Mol. Brain Res.* 2003; 118:82–90. [PubMed: 14559357]
- Dhillon H, Zigman JM, Ye C, Lee CE, McGovern RA, Tang V, Kenny CD, Christiansen LM, White RD, Edelman EA, et al. Leptin Directly Activates SF1 Neurons in the VMH, and This Action by Leptin Is Required for Normal Body-Weight Homeostasis. *Neuron.* 2006; 49:191–203. [PubMed: 16423694]
- Dugger BN, Morris JA, Jordan CL, Breedlove SM. Androgen receptors are required for full masculinization of the ventromedial hypothalamus (VMH) in rats. *Horm Behav.* 2007; 51:195–201. [PubMed: 17123532]
- Ferveur JF, Störtkuhl KF, Stocker RF, Greenspan RJ. Genetic feminization of brain structures and changed sexual orientation in male *Drosophila*. *Science.* 1995; 267:902–905. [PubMed: 7846534]
- Flanagan-Cato LM. Sex differences in the neural circuit that mediates female sexual receptivity. *Front Neuroendocrinol.* 2011; 32:124–136. [PubMed: 21338620]
- Flanagan-Cato LM, Lee BJ, Calizo LH. Co-localization of midbrain projections, progesterone receptors, and mating-induced fos in the hypothalamic ventromedial nucleus of the female rat. *Hormones and behavior.* 2006; 50:52–60. [PubMed: 16546183]
- Gagnidze K, Pfaff DW, Mong JA. Gene expression in neuroendocrine cells during the critical period for sexual differentiation of the brain. *Prog. Brain Res.* 2010; 186:97–111. [PubMed: 21094888]
- Gorski RA, Harlan RE, Jacobson CD, Shryne JE, Southam AM. Evidence for the existence of a sexually dimorphic nucleus in the preoptic area of the rat. *J. Comp. Neurol.* 1980; 193:529–539. [PubMed: 7440781]
- Goy RW, Phoenix CH. Hypothalamic regulation of female sexual behaviour; establishment of behavioural oestrus in spayed guinea-pigs following hypothalamic lesions. *J. Reprod. Fertil.* 1963; 5:23–40. [PubMed: 13949643]
- Gray DC, Mahrus S, Wells JA. Activation of specific apoptotic caspases with an engineered small-molecule-activated protease. *Cell.* 2010; 142:637–646. [PubMed: 20723762]
- Grgurevic N, Büdefeld T, Spanic T, Tobet SA, Majdic G. Evidence that sex chromosome genes affect sexual differentiation of female sexual behavior. *Horm Behav.* 2012; 61:719–724. [PubMed: 22483977]
- Grossman SP. Aggression, avoidance, and reaction to novel environments in female rats with ventromedial hypothalamic lesions. *Journal of Comparative and Physiological Psychology.* 1972; 78:274–283. [PubMed: 5062000]
- Harvey, W. *Exercitationes de generatione animalium* (London: Typis Du-Gardianis). In: Willis, R., translator. *The Works of William Harvey.* Miami, USA: HardPress Publishing, 2012; 1651. p. 186-187.
- Hess WR, Akert K. Experimental data on role of hypothalamus in mechanism of emotional behavior. *AMA Arch Neurol Psychiatry.* 1955; 73:127–129.
- Hetherington AW, Ranson SW. Hypothalamic lesions and adiposity in the rat. *The Anatomical Record.* 1940; 78:149–172.
- Jazin E, Cahill L. Sex differences in molecular neuroscience: from fruit flies to humans. *Nat. Rev. Neurosci.* 2010; 11:9–17. [PubMed: 20019686]

- Jiang X, Wang X. Cytochrome C-mediated apoptosis. *Annu. Rev. Biochem.* 2004; 73:87–106. [PubMed: 15189137]
- Juntti SA, Tollkuhn J, Wu MV, Fraser EJ, Soderborg T, Tan S, Honda S-I, Harada N, Shah NM. The androgen receptor governs the execution, but not programming, of male sexual and territorial behaviors. *Neuron.* 2010; 66:260–272. [PubMed: 20435002]
- Kelley DB. Generating sexually differentiated songs. *Curr. Opin. Neurobiol.* 1997; 7:839–843. [PubMed: 9464984]
- Kimura T, Hagiwara Y. Regulation of urine marking in male and female mice: effects of sex steroids. *Horm Behav.* 1985; 19:64–70. [PubMed: 4038966]
- King BM. The rise, fall, and resurrection of the ventromedial hypothalamus in the regulation of feeding behavior and body weight. *Physiol. Behav.* 2006; 87:221–244. [PubMed: 16412483]
- Kohatsu S, Koganezawa M, Yamamoto D. Female contact activates male-specific interneurons that trigger stereotypic courtship behavior in *Drosophila*. *Neuron.* 2011; 69:498–508. [PubMed: 21315260]
- Kollack-Walker S, Newman SW. Mating and agonistic behavior produce different patterns of Fos immunolabeling in the male Syrian hamster brain. *Neuroscience.* 1995; 66:721–736. [PubMed: 7644033]
- Konishi M. Birdsong for neurobiologists. *Neuron.* 1989; 3:541–549. [PubMed: 2701843]
- Kow LM, Harlan RE, Shivers BD, Pfaff DW. Inhibition of the lordosis reflex in rats by intrahypothalamic infusion of neural excitatory agents: evidence that the hypothalamus contains separate inhibitory and facilitatory elements. *Brain Res.* 1985; 341:26–34. [PubMed: 4041787]
- Krieger MS, Conrad LC, Pfaff DW. An autoradiographic study of the efferent connections of the ventromedial nucleus of the hypothalamus. *J. Comp. Neurol.* 1979; 183:785–815. [PubMed: 762273]
- Kruk MR, Van der Poel AM, De Vos-Frerichs TP. The induction of aggressive behaviour by electrical stimulation in the hypothalamus of male rats. *Behaviour.* 1979; 70:292–322. [PubMed: 574764]
- Kudwa AE, Harada N, Honda S-I, Rissman EF. Regulation of progesterone receptors in medial amygdala: estradiol, phytoestrogens and sex. *Physiol. Behav.* 2009; 97:146–150. [PubMed: 19258019]
- Kurrasch DM, Cheung CC, Lee FY, Tran PV, Hata K, Ingraham HA. The neonatal ventromedial hypothalamus transcriptome reveals novel markers with spatially distinct patterning. *J. Neurosci.* 2007; 27:13624–13634. [PubMed: 18077674]
- Leal S, Andrade JP, Paula-Barbosa MM, Madeira MD. Arcuate nucleus of the hypothalamus: effects of age and sex. *J. Comp. Neurol.* 1998; 401:65–88. [PubMed: 9802701]
- Leedy MG, Hart BL. Female and male sexual responses in female cats with ventromedial hypothalamic lesions. *Behavioral Neuroscience.* 1985; 99:936–941. [PubMed: 3843309]
- Levine JE, Chappell PE, Schneider JS, Sleiter NC, Szabo M. Progesterone receptors as neuroendocrine integrators. *Front Neuroendocrinol.* 2001; 22:69–106. [PubMed: 11259133]
- Lin D, Boyle MP, Dollar P, Lee H, Lein ES, Perona P, Anderson DJ. Functional identification of an aggression locus in the mouse hypothalamus. *Nature.* 2011; 470:221–226. [PubMed: 21307935]
- Lydon JP, DeMayo FJ, Funk CR, Mani SK, Hughes AR, Montgomery CA Jr, Shyamala G, Conneely OM, O'Malley BW. Mice lacking progesterone receptor exhibit pleiotropic reproductive abnormalities. *Genes Dev.* 1995; 9:2266–2278. [PubMed: 7557380]
- Majdic G, Young M, Gomez-Sanchez E, Anderson P, Szczepaniak LS, Dobbins RL, McGarry JD, Parker KL. Knockout Mice Lacking Steroidogenic Factor 1 Are a Novel Genetic Model of Hypothalamic Obesity. *Endocrinology.* 2002; 143:607–614. [PubMed: 11796516]
- Mani SK, Allen JM, Clark JH, Blaustein JD, O'Malley BW. Convergent pathways for steroid hormone- and neurotransmitter-induced rat sexual behavior. *Science.* 1994a; 265:1246–1249. [PubMed: 7915049]
- Mani SK, Blaustein JD, Allen JM, Law SW, O'Malley BW, Clark JH. Inhibition of rat sexual behavior by antisense oligonucleotides to the progesterone receptor. *Endocrinology.* 1994b; 135:1409–1414. [PubMed: 7925102]
- Mani SK, Blaustein JD, O'Malley BW. Progesterone receptor function from a behavioral perspective. *Horm Behav.* 1997; 31:244–255. [PubMed: 9213138]

- Mathews D, Edwards DA. Involvement of the ventromedial and anterior hypothalamic nuclei in the hormonal induction of receptivity in the female rat. *Physiol. Behav.* 1977a; 19:319–326. [PubMed: 607243]
- Mathews D, Edwards DA. The ventromedial nucleus of the hypothalamus and the hormonal arousal of sexual behaviors in the female rat. *Hormones and Behavior.* 1977b; 8:40–51. [PubMed: 863393]
- Matsumoto A, Arai Y. Sex difference in volume of the ventromedial nucleus of the hypothalamus in the rat. *Endocrinol. Jpn.* 1983; 30:277–280. [PubMed: 6662066]
- Matsumoto A, Arai Y. Male-female difference in synaptic organization of the ventromedial nucleus of the hypothalamus in the rat. *Neuroendocrinology.* 1986; 42:232–236. [PubMed: 3754037]
- Maxwell IH, Maxwell F, Glode LM. Regulated expression of a diphtheria toxin A-chain gene transfected into human cells: possible strategy for inducing cancer cell suicide. *Cancer Res.* 1986; 46:4660–4664. [PubMed: 3460697]
- McCarthy MM, Arnold AP. Reframing sexual differentiation of the brain. *Nat. Neurosci.* 2011; 14:677–683. [PubMed: 21613996]
- McGill TE. Sexual behavior in three inbred strains of mice. *behavior.* 1962; 19:341–350.
- Morris JA, Jordan CL, Breedlove SM. Sexual differentiation of the vertebrate nervous system. *Nat. Neurosci.* 2004; 7:1034–1039. [PubMed: 15452574]
- Morris JA, Jordan CL, Breedlove SM. Sexual dimorphism in neuronal number of the posterodorsal medial amygdala is independent of circulating androgens and regional volume in adult rats. *J. Comp. Neurol.* 2008; 506:851–859. [PubMed: 18076082]
- Musatov S, Chen W, Pfaff DW, Kaplitt MG, Ogawa S. RNAi-mediated silencing of estrogen receptor {alpha} in the ventromedial nucleus of hypothalamus abolishes female sexual behaviors. *Proc. Natl. Acad. Sci. U.S.A.* 2006; 103:10456–10460. [PubMed: 16803960]
- Musatov S, Chen W, Pfaff DW, Mobbs CV, Yang X-J, Clegg DJ, Kaplitt MG, Ogawa S. Silencing of estrogen receptor alpha in the ventromedial nucleus of hypothalamus leads to metabolic syndrome. *Proc. Natl. Acad. Sci. U.S.A.* 2007; 104:2501–2506. [PubMed: 17284595]
- Ng L, Bernard A, Lau C, Overly CC, Dong H-W, Kuan C, Pathak S, Sunkin SM, Dang C, Bohland JW, et al. An anatomic gene expression atlas of the adult mouse brain. *Nat. Neurosci.* 2009; 12:356–362. [PubMed: 19219037]
- Nyby J, Wysocki CJ, Whitney G, Dizinno G. Pheromonal regulation of male mouse ultrasonic courtship (*Mus musculus*). *Anim Behav.* 1977; 25:333–341. [PubMed: 889149]
- Ogawa S, Olazábal UE, Parhar IS, Pfaff DW. Effects of intrahypothalamic administration of antisense DNA for progesterone receptor mRNA on reproductive behavior and progesterone receptor immunoreactivity in female rat. *J. Neurosci.* 1994; 14:1766–1774. [PubMed: 8126569]
- Olivier B, Wiepkema PR. Behaviour changes in mice following electrolytic lesions in the median hypothalamus. *Brain Res.* 1974; 65:521–524. [PubMed: 4472010]
- Olster DH, Blaustein JD. Immunocytochemical colocalization of progesterone receptors and beta-endorphin or enkephalin in the hypothalamus of female guinea pigs. *J. Neurobiol.* 1990; 21:768–780. [PubMed: 2144316]
- Patisaul HB, Fortino AE, Polston EK. Sex differences in serotonergic but not gamma-aminobutyric acidergic (GABA) projections to the rat ventromedial nucleus of the hypothalamus. *Endocrinology.* 2008; 149:397–408. [PubMed: 17947355]
- Paxinos, G.; Franklin, KBJ. Academic Press; 2003. *The Mouse Brain in Stereotaxic Coordinates: Compact Second Edition, Second Edition.*
- Pfaff DW, Sakuma Y. Deficit in the lordosis reflex of female rats caused by lesions in the ventromedial nucleus of the hypothalamus. *J. Physiol. (Lond.).* 1979a; 288:203–210. [PubMed: 469716]
- Pfaff DW, Sakuma Y. Facilitation of the lordosis reflex of female rats from the ventromedial nucleus of the hypothalamus. *J. Physiol. (Lond.).* 1979b; 288:189–202. [PubMed: 469715]
- Phelps SM, Lydon JP, O'Malley BW, Crews D. Regulation of male sexual behavior by progesterone receptor, sexual experience, and androgen. *Horm Behav.* 1998; 34:294–302. [PubMed: 9878278]
- Von Philipsborn AC, Liu T, Yu JY, Masser C, Bidaye SS, Dickson BJ. Neuronal control of *Drosophila* courtship song. *Neuron.* 2011; 69:509–522. [PubMed: 21315261]




- Pollio G, Xue P, Zanisi M, Nicolin A, Maggi A. Antisense oligonucleotide blocks progesterone-induced lordosis behavior in ovariectomized rats. *Brain Res. Mol. Brain Res.* 1993; 19:135–139. [PubMed: 8361336]
- Power RF, Mani SK, Codina J, Conneely OM, O'Malley BW. Dopaminergic and ligand-independent activation of steroid hormone receptors. *Science.* 1991; 254:1636–1639. [PubMed: 1749936]
- Quadros PS, Pfau JL, Goldstein AYN, De Vries GJ, Wagner CK. Sex differences in progesterone receptor expression: a potential mechanism for estradiol-mediated sexual differentiation. *Endocrinology.* 2002; 143:3727–3739. [PubMed: 12239082]
- Quadros PS, Schlueter LJ, Wagner CK. Distribution of progesterone receptor immunoreactivity in the midbrain and hindbrain of postnatal rats. *Dev Neurobiol.* 2008; 68:1378–1390. [PubMed: 18712784]
- Reeves AG, Plum F. Hyperphagia, rage, and dementia accompanying a ventromedial hypothalamic neoplasm. *Arch. Neurol.* 1969; 20:616–624. [PubMed: 5769839]
- Ricciardi KH, Blaustein JD. Projections from ventrolateral hypothalamic neurons containing progesterin receptor- and substance P-immunoreactivity to specific forebrain and midbrain areas in female guinea pigs. *J. Neuroendocrinol.* 1994; 6:135–144. [PubMed: 7519513]
- Roberts DW, Baum MJ. Ventromedial hypothalamic nucleus lesions disrupt olfactory mate recognition and receptivity in female ferrets. *Horm Behav.* 2007; 51:104–113. [PubMed: 17011561]
- Rubin BS, Barfield RJ. Induction of estrous behavior in ovariectomized rats by sequential replacement of estrogen and progesterone to the ventromedial hypothalamus. *Neuroendocrinology.* 1983; 37:218–224. [PubMed: 6621803]
- Sakuma Y, Pfaff DW. Facilitation of female reproductive behavior from mesencephalic central gray in the rat. *Am. J. Physiol.* 1979; 237:R278–284. [PubMed: 495775]
- Saper CB, Swanson LW, Cowan WM. The efferent connections of the ventromedial nucleus of the hypothalamus of the rat. *J. Comp. Neurol.* 1976; 169:409–442. [PubMed: 61975]
- Schneider JS, Burgess C, Sleiter NC, DonCarlos LL, Lydon JP, O'Malley B, Levine JE. Enhanced Sexual Behaviors and Androgen Receptor Immunoreactivity in the Male Progesterone Receptor Knockout Mouse. *Endocrinology.* 2005; 146:4340–4348. [PubMed: 16002522]
- Schneider JS, Burgess C, Horton TH, Levine JE. Effects of progesterone on male-mediated infant-directed aggression. *Behav. Brain Res.* 2009; 199:340–344. [PubMed: 19146882]
- Schober J, Weil Z, Pfaff D. How generalized CNS arousal strengthens sexual arousal (and vice versa). *Horm Behav.* 2011; 59:689–695. [PubMed: 20950622]
- Shah NM, Pisapia DJ, Maniatis S, Mendelsohn MM, Nemes A, Axel R. Visualizing sexual dimorphism in the brain. *Neuron.* 2004; 43:313–319. [PubMed: 15294140]
- Simerly RB. Wired for reproduction: organization and development of sexually dimorphic circuits in the mammalian forebrain. *Annu. Rev. Neurosci.* 2002; 25:507–536. [PubMed: 12052919]
- Smith RL, Geller AI, Escudero KW, Wilcox CL. Long-term expression in sensory neurons in tissue culture from herpes simplex virus type 1 (HSV-1) promoters in an HSV-1-derived vector. *J. Virol.* 1995; 69:4593–4599. [PubMed: 7609023]
- Stowers L, Holy TE, Meister M, Dulac C, Koentges G. Loss of sex discrimination and male-male aggression in mice deficient for TRP2. *Science.* 2002; 295:1493–1500. [PubMed: 11823606]
- Swaab, D. *The Human Hypothalamus: Basic and Clinical Aspects Part I: Nuclei of the Human Hypothalamus.* Elsevier; 2003. The ventromedial nucleus (VMN; nucleus of Cajal); p. 239–242.
- Tsutsui K. Neurosteroid biosynthesis and action during cerebellar development. *Cerebellum.* 2012; 11:414–415. [PubMed: 22198872]
- La Vaque TJ, Rodgers CH. Recovery of mating behavior in the female rat following VMH lesions. *Physiol. Behav.* 1975; 14:59–63. [PubMed: 1171475]
- Veening JG, Coolen LM, De Jong TR, Joosten HW, De Boer SF, Koolhaas JM, Olivier B. Do similar neural systems subserve aggressive and sexual behaviour in male rats? Insights from c-Fos and pharmacological studies. *Eur. J. Pharmacol.* 2005; 526:226–239. [PubMed: 16263109]
- De Vries GJ. Sex Differences in Neurotransmitter Systems. *Journal of Neuroendocrinology.* 1990; 2:1–13. [PubMed: 19210390]

- De Vries GJ, Boyle PA. Double duty for sex differences in the brain. *Behav. Brain Res.* 1998; 92:205–213. [PubMed: 9473673]
- De Vries GJ, Panzica GC. Sexual differentiation of central vasopressin and vasotocin systems in vertebrates: different mechanisms, similar endpoints. *Neuroscience.* 2006; 138:947–955. [PubMed: 16310321]
- Wheatley MD. The hypothalamus and affective behavior in cats: a study of the effects of experimental lesions, with anatomic correlations. *Arch NeurPsych.* 1944; 52:296–316.
- Witt DM, Young LJ, Crews D. Progesterone modulation of androgen-dependent sexual behavior in male rats. *Physiol. Behav.* 1995; 57:307–313. [PubMed: 7716208]
- Wu MV, Manoli DS, Fraser EJ, Coats JK, Tollkuhn J, Honda S-I, Harada N, Shah NM. Estrogen masculinizes neural pathways and sex-specific behaviors. *Cell.* 2009; 139:61–72. [PubMed: 19804754]
- Xu X, Coats JK, Yang CF, Wang A, Ahmed OM, Alvarado M, Izumi T, Shah NM. Modular genetic control of sexually dimorphic behaviors. *Cell.* 2012; 148:596–607. [PubMed: 22304924]
- Yang X, Schadt EE, Wang S, Wang H, Arnold AP, Ingram-Drake L, Drake TA, Lusis AJ. Tissue-specific expression and regulation of sexually dimorphic genes in mice. *Genome Res.* 2006; 16:995–1004. [PubMed: 16825664]

**HIGHLIGHTS**

- Widespread adult sex differences in progesterone receptor (PR) expressing neurons
- Ventrolateral compartment of ventromedial hypothalamus (VMHvl) contains PR+ neurons
- PR+ VMHvl neurons exhibit cellular and molecular dimorphisms between the sexes
- PR+ VMHvl neurons essential for normal mating in both sexes and fighting in males



**Figure 1. Visualizing PR+ neurons in the mouse brain**

(A) Generating the PR<sup>PL</sup> allele. ACN is a self-excising neomycin selection cassette (Bunting et al., 1999). Orange rectangles are exons and the red line in the 3' exon denotes the stop codon.

(B) PCR to detect homologous recombination at the PR locus. Primers used to detect integration of the 5' (F1, R1) and 3' (F2, R2) arms of the targeting vector. ACN precludes detection of the 3' recombination event in ES cells.

(C, D) No difference between WT and PR<sup>PL/PL</sup> females in litter size and frequency.

(E–G) No difference in titers of sex hormones between WT and  $PR^{PL/PL}$  adults.

(H–M) Boxed areas in Nissl-stained coronal sections (Paxinos and Franklin, 2003) through the adult brain depict locations of the regions shown in panels to the right. PR expression in  $PR^{PL/+}$  female as labeled by  $\beta$ -gal activity mirrors expression of PR mRNA in adjacent sections.

Scale bars = 50  $\mu$ m.

Mean  $\pm$  SEM; n = 12/genotype (C–G); n = 3 (H–M).


See also Figure S1 and Table S3.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript



**Figure 2. Sexual dimorphism in PR-expression in the adult brain**

Boxed areas in Nissl-stained coronal sections through the adult brain depict regions of *PR<sup>PL/PL</sup>* mice labeled for  $\beta$ -gal activity in the panels to the right.

(A–L) More PR+ cells in the female AVPV/POA, VMHvl, and arcuate nucleus.

(M–X) More PR+ cells in the male basal forebrain, BNSTmpm, and MeApd.

(Y) Representation of sexually dimorphic PR expression in different brain regions as projected on a mid-sagittal section. c, caudal, d, dorsal, r, rostral, v, ventral.

Scale bars = 50  $\mu$ m (C, K) and 100  $\mu$ m (G, O, S, W). Inset scale bars = 25  $\mu$ m.


Mean  $\pm$  SEM; n = 4/sex; \*p < 0.04, \*\*p < 0.01.  
See also Figure S2 and Table S1.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript



**Figure 3. PR+ VMHvl neurons project in a sexually dimorphic manner**

(A) Strategy to visualize projections of PR+ neurons.

(B–E) Lenti-lxIplap targeted to the VMH infects cells in *PR<sup>Cre/+</sup>* and WT mice as visualized by EGFP+ cells. Only a few cells are PR+ in this region so there is no apparent difference in the number of EGFP+ cells in *PR<sup>Cre</sup>* and WT mice. PLAP+ soma and local arbors of VMHvl neurons are only observed in *PR<sup>Cre</sup>* mice.

(F–U) Boxed areas in Nissl-stained coronal sections depict regions shown in panels to the right. Lenti-lxIplap targeted to the VMHvl of adult *PR<sup>Cre/+</sup>* mice labels PLAP+ soma and




local arbors of VMHvl neurons (F–I). The lentiviral titer limits the number of infected Cre+ neurons and does not highlight the sex difference in the number of these neurons. The variable multiplicity of infection can lead to apparent size differences in PLAP-labeled soma. However, there is no sex difference in the soma size of these neurons (Figure S2G). PR+ VMHvl neurons project to the AVPV, POA, and PAG (J–U). There are more PLAP+ projections to the AVPV in females (J–M).

(V, W) Schematic summarizing projections of PR+ VMHvl neurons. No difference in anatomical extent of projections in different regions, but female AVPV receives more innervation from these neurons.

Scale bars = 100  $\mu\text{m}$  (C), 50  $\mu\text{m}$  (H,P,T), 25  $\mu\text{m}$  (L).

Mean  $\pm$  SEM; n = 7/sex; \*p < 0.001.

See also Figure S3, Tables S2, S3.



**Figure 4. Genetic strategy to ablate neurons in a Cre-dependent manner**  
 (A) Intramolecular cleavage of endogenous pro-caspase-3 by upstream caspases activates caspase-3, which then induces apoptosis. This intramolecular cleavage site has been replaced by a TEV linker domain (black bar) in inactive taCasp3 (pro-taCasp3) such that only TEV protease activates taCasp3, which then induces apoptosis.  
 (B) Viral strategy to ablate PR+ neurons conditionally.  
 (C, D) Cell death 1 week following infection of Cre:EGFP+ HEK293T cells with AAV-flex-taCasp3-TEVp. n = 3 experiments.


(E, F) Ablation of PR+ VMHvl neurons in a *PR<sup>PL/Cre</sup>* but not *PR<sup>PL/+</sup>* female injected with AAV-flex-taCasp3-TEVp. n = 10/experimental group.  
Scale bar = 100  $\mu$ m (C, D) and 25  $\mu$ m (E, F).  
See also Figure S4.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript



**Figure 5. PR+ VMHvl neurons regulate female sexual receptivity**

(A) Experimental design to test the role of PR+ VMHvl neurons in female behaviors. Mating was tested with ovariectomized females primed to be in estrus. Other behaviors were tested with gonadally-intact females.

(B–J) PR<sup>Cre</sup> and control females were injected with AAV-flex-taCasp3-TEVp and tested for sexual behavior with WT males.

(B) PR<sup>Cre</sup> females spend more time rejecting male mating attempts, walking away when the male approaches.

(C–E) *PR<sup>Cre</sup>* females display lower receptivity index (mounts leading to intromission/total mounts) and reduced number and duration of lordosis events.

(F) *PR<sup>Cre</sup>* females spend more time moving about and being unreceptive during intromission.

(G) Fewer than 20% of PR+ neurons remain in the VMHvl of *PR<sup>Cre</sup>* females, who reject male mating attempts more than control females.

(H) Males sniff and initiate mating equivalently with *PR<sup>Cre</sup>* and WT females but ejaculate in fewer assays with *PR<sup>Cre</sup>* females.


(I) Males mount *PR<sup>Cre</sup>* females more but without a corresponding increase in intromission.

(J) Males mount *PR<sup>Cre</sup>* females longer, but intromit for shorter duration.

(K, L) Ablation of PR+ VMHvl neurons in *PR<sup>Cre</sup>* females results in loss of Cckar expression.

Mean  $\pm$  SEM; n = 10/experimental group (B–J); n = 3 (K, L); \*p < 0.02, \*\* p < 0.005. Scale bar = 50  $\mu$ m.

See also Figure S5, Table S3, Movies S1, S2.



**Figure 6. PR+ VMHvl neurons regulate male sexual behavior**

(A) Experimental design to test the role of PR+ VMHvl neurons in male behaviors. (B–G) PR<sup>Cre</sup> and control males were injected with AAV-flex-taCasp3-TEVp and tested for mating, ultrasonic vocalizations toward male or female intruders, and territory marking. (B) PR<sup>Cre</sup> males intromit females in fewer assays. (C, D) PR<sup>Cre</sup> males mount and intromit females less and have shorter bouts of intromissions. (E) Both PR<sup>Cre</sup> and control males emit more vocalizations to females. (F, G) No difference between PR<sup>Cre</sup> and control males in the number and distribution of urine

marks. % marks in center =  $100 * (\# \text{ urine marks not abutting cage perimeter} / \# \text{ of all urine marks})$ .

Mean  $\pm$  SEM; n = 24/experimental group (B–D, F,G), n = 5/experimental group (E); \*p < 0.008, \*\*p < 0.001.


See also Figures S6 and S7.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript



**Figure 7. PR+ VMHvl neurons regulate male aggression**

(A–F) *PR<sup>Cre</sup>* and control resident males were injected with AAV-flex-taCasp3-TEVp targeted to the VMHvl and tested for aggression toward a WT male intruder.

(A) All residents sniff intruders equivalently, but *PR<sup>Cre</sup>* males attack less.

(B–D) When *PR<sup>Cre</sup>* males fight, they attack less, for a shorter duration, and with longer intervals between attacks.

(E) *PR<sup>Cre</sup>* males bite less.



(F) Fewer than 20% of PR+ neurons remain in the VMHvl of *PR<sup>Cre</sup>* males, who attack intruders less.

Mean  $\pm$  SEM; n = 24/experimental group; \*p < 0.04, \*\*p = 0.009.

(G, H) Ablation of PR+ VMHvl neurons in a *PR<sup>PL/Cre</sup>* male injected with AAV-flex-taCasp3-TEVp. Scale bar = 25  $\mu$ m.

See also Figure S7.