

**UCLA**

**UCLA Previously Published Works**

**Title**

Support for a Single Underlying Dimension of Self-Reported Health in a Sample of Adults with Low Back Pain in the United States

**Permalink**

<https://escholarship.org/uc/item/5396f9pr>

**Authors**

Hays, Ron D

Rodriguez, Anthony

Qureshi, Nabeel

et al.

**Publication Date**

2024-05-14

**DOI**

10.1007/s11482-024-10327-8

Peer reviewed



# Support for a Single Underlying Dimension of Self-Reported Health in a Sample of Adults with Low Back Pain in the United States

Ron D. Hays<sup>1</sup> · Anthony Rodriguez<sup>2</sup> · Nabeel Qureshi<sup>3</sup> · Chengbo Zeng<sup>4</sup> · Maria Orlando Edelen<sup>2,4</sup>

Received: 21 December 2023 / Accepted: 7 May 2024  
© The Author(s) 2024

## Abstract

There is increasing interest in measuring “whole person” health and deriving an overall summary score. Underlying physical and mental health dimensions have been found consistently in prior studies of self-reported health, but it is unclear whether a single underlying health factor is supported across health domains. We examine the dimensionality of nine domains from the Patient-Reported Outcomes Measurement Information System (PROMIS®)-29 + 2 profile measure, the PROMIS social isolation scale, the Personal Wellbeing Index, and the EQ-5D-5L preference score in a sample of 1256 adults with back pain in the United States: mean age was 55 (range 18–94), 52% female, 74% non-Hispanic White, 61% were married or living with a spouse, and the highest level of education completed for 35% of the sample was a high school degree or general education diploma. The sample reported substantially more pain intensity, pain interference, and worse physical function than the U.S. general population. Product-moment correlations among the measures ranged from 0.25 to 0.83 (median correlation = 0.52). A bifactor model showed that a general health factor accounted for most of the covariation among measures, but physical function, pain interference, and pain intensity loaded slightly more on the physical health group factor than on the general health factor. The study provides some support for combining multiple aspects of self-reported health into an overall indicator of whole-person health.

**Keywords** Health dimensions · Factor analysis · Social health · Well-being · PROMIS®

The World Health Organization (2002) defined health as functioning and well-being in physical, mental, and social health. Generic self-report measures are designed to be generally relevant and to be used to assess different subgroups of people (Patrick & Deyo, 1989). There is an emerging interest in the concept of whole-person health:

---

Extended author information available on the last page of the article

“engaging with the whole person, not just the physical body but the emotional, mental and spiritual aspects as well is critical to healing” (Kligler, 2022). An overall whole-person score may provide a useful summary of perceived health. For example, Yin et al. (2016) found that a single factor accounted for the covariation among four items included in the United States Behavioral Risk Factor Surveillance System and recommended the use of a single summary score to monitor population trends. However, the reasonableness of an overall whole-person health measure needs to be evaluated in comprehensive patient-reported health measures (Chi et al., 2023).

Prior work has consistently found multiple correlated underlying dimensions for self-reported health. There is extension evidence in support of underlying physical and mental health (Essink-Bot et al., 1997; Hays et al., 2009; Hays & Stewart, 1990). For example, a confirmatory factor analysis of a 64-item measure administered to individuals with HIV supported a physical health factor defined by physical function, role function, freedom from pain, disability days, and quality of sex life, and a mental health dimension defined by the overall quality of life, emotional well-being, hopefulness, lack of loneliness, will to function, quality of family life, quality of friendships, and cognitive function/distress, with physical and mental health correlated 0.31 with one another (Hays et al., 1995). Correlations of 0.62 and 0.66, respectively, between physical and mental health factors were found for the RAND-36 (Farivar et al., 2007; Hays et al., 1998). Rasch-derived physical and mental health scores from the SF-36 correlated 0.74 (Chang et al., 2007). Finally, the correlation between Patient-Reported Measurement and Information System (PROMIS)-29 profile physical and mental health summary scores was 0.69 (Hays et al., 2018). In summary, prior work indicates that self-reported health is multidimensional with substantial correlations among the underlying dimensions.

We extend the work of Yin et al. (2016) by evaluating the extent to which an overall health summary score is supported in a comprehensive set of multi-item health measures. Generic health profile measures yield scores on multiple domains. In this study, we include widely used measures from the PROMIS project, a United States National Institutes of Health Roadmap initiative to create item banks for use in the general population and for different medical conditions that have been touted as state-of-the-science instruments (Cella et al., 2019; Kaplan & Hays, 2022). We supplement the PROMIS measures with the Personal Wellbeing Index, a subjective well-being measure endorsed by the World Health Organization and the Organization for Economic Cooperation and Development (Cummins et al., 2003). Generic health preference-based measures yield a single score anchored at 0 = dead and 1 = perfect health. We include the latest version of the most widely used health preference-based measure in the world (Brazier et al., 2017).

The inclusion of an important profile measure in the United States, a worldwide subjective well-being measure, and the most widely used preference-based measure makes it possible to thoroughly evaluate the plausibility of a summary indicator of whole-person health. Because low back pain is common and regarded as the leading cause of years lived with disability worldwide (Hoy et al., 2014; Wu et al., 2020), this study focuses on adults with low back pain.

## Methods

We administered surveys in English to members of KnowledgePanel®, an online panel that relies on probability-based sampling methods for recruitment and provides a representative sample of non-institutionalized adults 18 and older residing in the United States (Ipsos, 2018). Data was collected at baseline and six months later for a subsample with back pain. The PROMIS-29 + 2 and the EQ-5D-5L were administered at both time points. Personal well-being and the PROMIS social isolation scale were only included in the 6-month survey. The analyses reported here are limited to the 6-month data to assess the dimensionality of these health measures.

At baseline, the survey vendor (Ipsos) sent an email invitation to 7224 KnowledgePanel members on September 22, 2022, and gave them 10 days to complete the survey. Email reminders were sent to non-responders on Day 3 of the field period. Additional reminders were sent to the remaining non-responders every 3 days for up to 10 days. Upon completion, respondents received an entry into the KnowledgePanel sweepstakes. Fifty-seven percent ( $n=4117$ ) completed the survey and 19 who reported having one or two of the fake health conditions included in the survey to identify careless or insincere respondents (Hays et al., 2023) were excluded, resulting in a baseline sample of 4098. The 6-month survey was offered only to those who reported back pain and did not endorse a fake health condition on the baseline survey. Seventy-nine percent of the eligible baseline respondents completed a 6-month survey ( $n=1256$ ).

## Measures

### PROMIS

The PROMIS ontology describes multiple domains within dimensions of physical, mental, and social health (Cella & Hays, 2022). Physical health is divided into physical symptoms and physical function, mental health is represented by affect, behaviors, and cognition, and social health includes social function and social relationships (e.g., family, and friends).

The PROMIS-29 (Cella et al., 2019) is the most widely used PROMIS profile measure. It assesses pain intensity using a single 0–10 numeric rating item, and seven health domains (physical function, fatigue, pain interference, depression, anxiety, ability to participate in social roles and activities, and sleep disturbance) using 4 polytomous (5 response categories) items per domain. Support for the reliability and validity of the PROMIS-29 has been shown in several prior studies (Cook et al., 2021; McMullen et al., 2022; Pecorelli et al., 2023; Peipert et al., 2018). In addition to the PROMIS-29, the study included the 2-item cognitive function scale from the PROMIS-29 + 2. The cognitive function scale had a reliability of 0.77 in a prior study (Hays et al., 2023). The PROMIS 4-item social isolation scale was also administered (Hahn et al., 2014).

All PROMIS measures are scored on a T-score metric with a mean of 50 and SD of 10 in the general U.S. population for all except sleep disturbance where the score is relative to a combination of the general population and clinical patients.

### Personal Well-Being

To ensure adequate representation of subjective well-being (Cummins et al., 2004), we administered 10 items developed by the International Wellbeing Group (2013). Eight of the items used a 0 (No satisfaction at all) to 10 (Completely satisfied) response scale and asked “How satisfied are you with”:

- 1) your standard of living?
- 2) your health?
- 3) what you are achieving in life?
- 4) your personal relationships?
- 5) how safe you feel?
- 6) feeling part of your community?
- 7) your future security?
- 8) your spirituality or religion?

A ninth question asked, “Overall, how satisfied are you with your life as a whole these days?” (0=Not satisfied at all, 10=Completely satisfied). The final question asked, “Overall, to what extent do you feel the things you do in your life are worthwhile?” (0=Not at all worthwhile, 10=Completely worthwhile). Seven of the items are used in scoring the Personal Wellbeing Index (Cummins et al., 2003). Test-retest (intraclass correlation) reliability of 0.84 for a 1–2-week interval was found (Lau & Cummins, 2005). We created a 10-item average personal well-being score with a 0–10 possible score range.

### EQ-5D-5L

The EQ-5D-5L items refer to “Your health today” and assess mobility, self-care, usual activities, pain/discomfort, and anxiety/depression with five response options (*no problems, some problems, moderate problems, severe problems, and extreme problems*), with 3125 possible health states (Herdman et al., 2011). The EQ-5D-5L U.S. weights we used were derived using Tobit modeling of time trade-off preference elicitation (Pickard et al., 2019). Extensive support for the reliability and validity of the EQ-5D-5L has been reported (Feng et al., 2021). The EQ-5D-3L was included along with the SF-36 to evaluate the dimensionality of self-reported health in a prior study (Essink-Bot et al., 1997). Because the EQ-5D-5L preference-based score combines information from physical, mental, and social health, we hypothesized that it would be a good indicator of overall health and be useful in assessing the reasonableness of the overall whole-person health dimension. Palimaru and Hays (2017) found that 69% of the variance in overall quality of life was accounted for by the EQ-5D-3L and PROMIS global health items.

## Demographic Variables and Medical Conditions

We assessed gender, age, education, race/ethnicity, marital status, whether working full time and the presence of 20 medical conditions (see Table 1).

### Subjects

Table 1 shows that 52% of the back pain sample that completed the 6-month survey were female. The mean age was 55 (18–94 range). Thirty-five percent of the sample had a high school education or less. The majority were non-Hispanic White (74%), with 10% Hispanic, and 8% non-Hispanic Black. Most were married (61%). Thirty-six percent were working full-time. The most common medical conditions reported other than back pain were allergies (58%), hypertension and high cholesterol (47% each), and arthritis (46%).

### Analysis Plan

We provide means, standard deviations, and internal consistency reliability (Cronbach, 1951) estimates for the PROMIS scales and personal well-being scale, means, standard deviations, and test-retest (stability) estimates over 6-months (baseline to 6-months later) for the pain intensity item and EQ-5D-5L. In addition, we report product-moment correlations among the measures. Then we estimate confirmatory factor analysis models.

We evaluated one-factor, two-factor, three-factor, and bifactor models (Reise et al., 2007; Rodriguez et al., 2016). For the one-factor model, loadings for all measures were estimated on the single factor. For the two-factor model, physical health was defined by physical function, pain interference, and pain intensity while mental health was defined by depression, anxiety, social isolation, personal well-being, cognitive function, and sleep disturbance. Fatigue, the ability to participate in social roles and activities, and the EQ-5D-5L were allowed to load on both physical and mental health. For the three-factor model, physical health was defined by the same domains as for the two-factor model. Mental health was defined by depression, anxiety, personal well-being, cognitive function, and sleep disturbance. Social health was indicated by the ability to participate in social roles and activities, and social isolation. Fatigue was allowed to load on both physical and mental health. The EQ-5D-5L was allowed to load on physical, mental, and social health. The bifactor model included all 12 measures loading on the general factor, 3 loadings on the physical health group factor (physical function, pain interference, pain intensity), and 4 loadings on the mental health group factor (depression, personal well-being, anxiety, social isolation). We assess model fit using the comparative fit index (CFI) and the root mean square error of approximation (RMSEA). CFI values of about 0.95 or higher and RMSEA values of about 0.06 or lower are indicators of a good practical fit of the model to the data (Hu & Bentler, 1999; MacCallum et al., 1996). In addition, we report the explained common variance for the general factor. We

**Table 1** Characteristics of the Sample ( $n = 1256$ )

| Variable  | %        |
|---|----------|
| Female  | 52       |
| Age   |          |
| 18–29   | 10       |
| 30–44   | 20       |
| 45–59   | 23       |
| 60–94   | 47       |
| Education   |          |
| Did not graduate high school                          | 7        |
| High school degree or general education diploma (GED) | 28       |
| Some college or AA degree                             | 29       |
| Bachelor's degree                                     | 20       |
| Masters' degree or higher                             | 16       |
| Race/ethnicity  |          |
| Non-Hispanic White                                    | 74       |
| Hispanic  | 10       |
| Non-Hispanic Black                                    | 8        |
| Non-Hispanic reporting 2 or more races                | 4        |
| Non-Hispanic reporting another race                   | 4        |
| Marital status  |          |
| Married or living with a spouse                       | 65       |
| Never married   | 16       |
| Divorced  | 11       |
| Widowed   | 6        |
| Separated   | 2        |
| Working full time                                     | 36       |
| Allergies   | 58       |
| Hypertension  | 47       |
| High cholesterol                                      | 47       |
| Arthritis   | 46       |
| Trouble sleeping                                      | 42       |
| Neck pain   | 39       |
| Sciatica  | 33       |
| Anxiety   | 29       |
| Depression  | 29       |
| Stomach   | 25       |
| Trouble hearing                                       | 23       |
| Trouble seeing  | 22       |
| Diabetes  | 19       |
| Asthma  | 17       |
| Dermatitis  | 14       |
| Cancer  | 13       |
| Chronic obstructive pulmonary disease                 | 8        |
| Heart disease   | <b>8</b> |

**Table 1** (continued)

| Variable     | % |
|--------------|---|
| Heart attack | 3 |
| Stroke       | 3 |
| Angina       | 2 |

also assess the fit of the bifactor model in age, gender, education, and race/ethnicity subgroups. Finally, to evaluate the potential effects of including the EQ-5D-5L on the loadings for the other indicators in the factor analyses, we estimated the bifactor model excluding the EQ-5D-5L.

Analyses were conducted using SAS 9.4 (SAS Institute, Cary, NC).

## Results

Descriptive statistics for the measures are shown in Table 2. The sample reported substantially more pain intensity, pain interference, and worse physical function than the United States general population. Internal consistency reliability coefficients ranged from 0.82 to 0.95 and 6-month test-retest correlates were 0.68 for the pain intensity item and 0.77 for the EQ-5D-5L. As seen in Table 3, the

**Table 2** Mean, Standard Deviation, and Reliability of the Measures

| Scale   | Mean | Standard Deviation | Reliability |
|---|------|--------------------|-------------|
| Fatigue   | 51.6 | 9.7                | 0.94        |
| Ability to participate in social roles and activities | 51.9 | 9.2                | 0.95        |
| Depression  | 50.2 | 9.8                | 0.94        |
| Personal well-being                                   | 6.8  | 2.2                | 0.94        |
| Anxiety   | 50.7 | 9.7                | 0.91        |
| Social isolation                                      | 47.6 | 10.7               | 0.93        |
| Sleep disturbance                                     | 51.6 | 8.6                | 0.88        |
| EQ-5D-5L  | 0.77 | 0.23               | 0.77*       |
| Cognitive function                                    | 51.3 | 8.2                | 0.82        |
| Pain interference                                     | 54.4 | 8.8                | 0.96        |
| Pain intensity item                                   | 56.5 | 9.8                | 0.68*       |
| Physical function                                     | 45.8 | 8.9                | 0.94        |

All measures except for personal well-being and the EQ-5D-5L are reported on a T-score metric (mean = 50 in the U.S. general population). The possible range of the personal well-being score was 0 to 10. The mean EQ-5D-5L in the U.S. for an online sample (Jiang et al., 2021) was 0.80 (SD = 0.24). Internal consistency reliability is reported for the 10 multi-item scales and 6-month test-retest reliability (stability) is reported for the EQ-5D-5L and the pain intensity item (indicated with \* above)



**Table 3** Product-Moment Correlations Among PROMIS-29+2, EQ-5D-5L, Personal Well-Being, and Social Isolation Measures

|                             | Fatigue | Social roles and activities | Depression | Personal well-being | Anxiety | Social isolation | Sleep disturbance | EQ-5D-5L | Cognitive function | Pain interference | Pain intensity | Physical function |
|-----------------------------|---------|-----------------------------|------------|---------------------|---------|------------------|-------------------|----------|--------------------|-------------------|----------------|-------------------|
| Fatigue                     | 1.00    |                             |            |                     |         |                  |                   |          |                    |                   |                |                   |
| Social roles and activities | -.70    | 1.00                        |            |                     |         |                  |                   |          |                    |                   |                |                   |
| Depression                  | 0.62    | -.59                        | 1.00       |                     |         |                  |                   |          |                    |                   |                |                   |
| Personal well-being         | -.57    | 0.57                        | -.69       | 1.00                |         |                  |                   |          |                    |                   |                |                   |
| Anxiety                     | 0.58    | -.55                        | 0.83       | -.60                | 1.00    |                  |                   |          |                    |                   |                |                   |
| Social isolation            | 0.58    | -.53                        | 0.69       | -.66                | 0.61    | 1.00             |                   |          |                    |                   |                |                   |
| Sleep disturbance           | 0.58    | -.53                        | 0.51       | -.45                | 0.50    | 0.45             | 1.00              |          |                    |                   |                |                   |
| EQ-5D-5L                    | -.56    | 0.66                        | -.57       | 0.42                | -.50    | -.42             | -.42              | 1.00     |                    |                   |                |                   |
| Cognitive function          | -.47    | 0.50                        | -.52       | 0.52                | -.53    | -.43             | -.43              | 0.44     | 1.00               |                   |                |                   |
| Pain interference           | 0.52    | -.68                        | 0.39       | -.36                | 0.36    | 0.32             | 0.41              | -.69     | -.36               | 1.00              |                |                   |
| Pain intensity              | 0.43    | -.52                        | 0.35       | -.25                | 0.31    | 0.25             | 0.38              | -.63     | -.29               | 0.73              | 1.00           |                   |
| Physical function           | -.45    | 0.64                        | -.32       | 0.25                | -.29    | -.25             | -.33              | 0.71     | 0.32               | -.71              | -.58           | 1.00              |

Social roles and activities = Ability to participate in social roles and activities

absolute value of product-moment correlations among the measures ranged from 0.25 (social isolation with physical function and pain intensity) to 0.83 (depressive symptoms and anxiety). The median of the absolute value of the correlations was 0.52.

Because of the large sample size, all confirmatory factor analysis models were rejectable statistically at  $p < .0001$ . The one-factor model did not fit the data:  $\chi^2$  (54 df) = 2794.19 (CFI = 0.74, RMSEA = 0.20). The two-factor model estimated a 0.50 correlation between physical and mental health. That model fit the data better than the one-factor model, but it still was below the thresholds of acceptable practical fit:  $\chi^2$  (50 df) = 759.51 (CFI = 0.93, RMSEA = 0.11). The three-factor model fit the data less well than the two-factor model:  $\chi^2$  (48 df) = 1059.96 (CFI = 0.90, RMSEA = 0.13). Moreover, the estimated correlation between mental and social health was 0.97. The bifactor model with a general health factor and physical health and mental health group factors provided an acceptable fit:  $\chi^2$  (36 df) = 170.51 (CFI = 0.99, RMSEA = 0.05). Model fit was like that of the overall sample in all the age, gender, education, and race/ethnic subgroups (Table 4).

Table 5 shows the 12-factor loadings on a general factor, 3 loadings on the physical health group factor, 4 loadings on the mental health group factor, and 10 correlated uniqueness estimates. Loadings on the general health factor were larger than on a group factor for 9 of the 12 measures, but physical function, pain interference, and pain intensity loaded slightly more on physical health rather than the general health factor. The general factor explained 78% of the common variance. The bifactor model that excluded the EQ-5D-5L fit the data equally well (CFI = 0.99, RMSEA = 0.05) and yielded standardized factor loadings that were virtually identical to the model with all 12 indicators (Table 5).

**Table 4** Fit of Bifactor Model in Overall Sample and Age, Gender, Education, and Race/ethnicity Subgroups

| Sample                            | Chi-square | Comparative Fit Index | Root Mean Square Error of Approximation |
|-----------------------------------|------------|-----------------------|---|
| Overall sample                    | 170.51     | 0.99                  | 0.05                                    |
| Age                               |            |                       |   |
| 18–59 years old                   | 109.80     | 0.99                  | 0.06                                    |
| 60–94 years old                   | 104.78     | 0.99                  | 0.06                                    |
| Gender                            |            |                       |   |
| Females                           | 106.86     | 0.99                  | 0.05                                    |
| Males                             | 106.76     | 0.98                  | 0.06                                    |
| Education                         |            |                       |   |
| High school education or less     | 82.32      | 0.99                  | 0.05                                    |
| More than a high school education | 120.72     | 0.99                  | 0.05                                    |
| Race/ethnicity                    |            |                       |   |
| Non-Hispanic White                | 156.10     | 0.98                  | 0.06                                    |
| Another race/ethnicity            | 170.51     | 0.99                  | 0.05                                    |

**Table 5** Standardized Confirmatory Factor Loading Matrix for PROMIS-29+2, EQ-5D-5L, Personal Well-being, and Social Isolation Measures from the Bifactor Model (Estimates from Model Excluding EQ-5D-5L shown within parentheses)

| Scale   | General Health | Physical Health | Mental Health |
|---|----------------|-----------------|---------------|
| Fatigue   | -.81 (-.81)    |                 |               |
| Ability to participate in social roles and activities | 0.78 (0.78)    |                 |               |
| Depression  | -.77 (-.77)    |                 | -.38 (-.35)   |
| Personal well-being                                   | 0.75 (0.75)    |                 | 0.30 (0.31)   |
| Anxiety   | -.72 (-.71)    |                 | -.28 (-.25)   |
| Social isolation                                      | -.69 (-.68)    |                 | -.44 (-.48)   |
| Sleep disturbance                                     | -.69 (-.69)    |                 |               |
| EQ-5D-5L  | 0.68           |                 |               |
| Cognitive function                                    | 0.63 (0.63)    |                 |               |
| Pain interference                                     | -.62 (-.62)    | -.70 (-.70)     |               |
| Pain intensity  | -.52 (-.52)    | -.57 (-.57)     |               |
| Physical function                                     | 0.52 (0.52)    | 0.55 (0.56)     |               |

Blank cells indicate that the loading was not estimated. CFI=0.99 and RMSEA=0.05 for both the model with all 12 indicators and the model with 11 indicators (excluding EQ-5D-5L). For the model with 12 indicators, 10 correlated uniqueness estimates were estimated: physical function with EQ-5D-5L ( $r=0.36$ ), pain intensity with EQ-5D-5L ( $r=-0.27$ ), pain interference with EQ-5D-5L ( $r=-0.26$ ), the ability to participate in social roles and activities with physical function ( $r=0.22$ ), pain interference ( $r=-0.19$ ), EQ-5D-5L ( $r=0.13$ ), pain intensity ( $r=-.11$ ), and fatigue ( $r=-.07$ ), and anxiety with depressive symptoms ( $r=0.17$ ) and cognitive function ( $r=-0.06$ )

## Discussion

This study of adults with back pain shows that the nine measures from the PROMIS-29+2, the PROMIS social isolation scale, the personal well-being measure, and the EQ-5D-5L are substantially intercorrelated and, at a higher level, represent a single underlying dimension of health. The findings extend the Yin et al. (2016) research of a single factor for the four items to a more comprehensive collection of self-report measures of health. The results suggest that it is reasonable to combine PROMIS measures into a single score using factor-scoring coefficients as weights or a preference-based score (Dewitt et al., 2018),

Based on the two-factor model, one might have concluded, consistent with previous research, that there are two dimensions: physical and mental health. However, by specifying a bifactor model, we were able to evaluate the extent to which the 12 measures represented a general health concept. There was a strong indication of the presence of a general health factor as well as unique information shared among three physical health indicators (physical function, pain interference, and pain intensity) that a National Institutes of Health (NIH) Pain Consortium steering committee research task force proposed combining into an Impact Stratification Score for assessing adults with chronic low back pain (Deyo et al., 2014; Hays et al., 2021). The largest loading on the general health factor was

observed for the PROMIS fatigue scale and 10 of the 12 indicators had standardized loadings (absolute value) of 0.62 or larger.

Consistent with most prior work, we failed to identify a separate social health factor despite including the PROMIS social isolation scale and the ability to participate in social roles and activities scale. One prior confirmatory factor analysis of PROMIS measures suggested three underlying factors: physical health represented by PROMIS measures of physical function, pain interference, pain behavior, and fatigue; mental health represented by anger, anxiety, depression, and fatigue; and social health represented by social role performance and social role satisfaction (Carle et al., 2015). However, the social health factor correlated more strongly with physical and mental health (0.67 and 0.68, respectively) than physical and mental health correlated with one another ( $r = 0.57$ ), suggesting that social health was not a distinct dimension.

This study has limitations. The generalizability of the results is uncertain given the sample was limited to adult members of KnowledgePanel with back pain who responded to a baseline and follow-up survey. In addition, a wide range of measures was administered, but it is possible that stronger support for a social health factor would be obtained if additional measures had been included. Nonetheless, the study provides important information about the dimensionality of self-reported health.

Future research is needed to put this work in the context of emerging interest in whole-person health. For example, the U.S. Veterans Health Administration emphasizes the importance of whole-person health as part of comprehensive health care (Kligler, 2022). A deeper understanding of the dimensionality of health measures can contribute substantially to efforts to define and measure whole-person health. The reasonableness of single summary indexes such as the whole health index must also be evaluated further (Chi et al., 2023).

**Acknowledgments** We thank Rebecca Weir and Mary Slaughter for the preparation of the dataset.

**Authors Contribution** RDH conducted the analysis and wrote the drafts. All coauthors provided feedback on drafts and approved the final submission.

**Funding** This work was supported by the National Center for Complementary and Integrative Health (NCCIH) Grant No. 1R01AT010402-01A1.

**Data Availability** The data set analyzed for this study is publicly available from the ICPSR database repository number openicpsr-198049.

## Declarations

**Ethics Approval** This study was approved by the RAND Human Subjects Protection Committee (2019-0651-AM02).

**Competing Interests** The authors have no competing interests to declare that are relevant to the content of this article.

**Consent** All respondents provided electronic consent before starting the survey.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References


- Brazier, J., Ratcliffe, J., Salomon, J., & Tsuchiya, A. (2017). *Measuring and valuing health benefits for economic evaluation* (2nd ed.). Oxford University Press. <https://doi.org/10.1093/med/9780198725923.001.0001>
- Carle, A. C., Riley, W., Hays, R. D., & Cella, D. (2015). Confirmatory factor analysis of the patient reported outcomes measurement information system (PROMIS) adult domain framework using item response theory scores. *Medical Care*, *53*(10), 894–900.
- Cella, D., Choi, S. W., Condon, D. M., Schalet, B., Hays, R. D., Rothrock, N. E., et al. (2019). PROMIS® adult health profiles: Efficient short-form measures of seven health domains. *Value in Health*, *22*(5), 537–544. <https://doi.org/10.1016/j.jval.2019.02.004>
- Cella, D., & Hays, R. D. (2022). A patient reported outcome ontology: Conceptual issues and challenges addressed by the patient-reported outcomes measurement information system® (PROMIS®). *Patient Related Outcome Measures*, *13*, 189–197. <https://doi.org/10.2147/PROM.S371882>
- Chang, C.-H., Wright, B. D., Cella, D., & Hays, R. D. (2007). The SF-36 physical and mental health factors were confirmed in cancer and HIV/AIDS patients. *Journal of Clinical Epidemiology*, *60*(1), 68–72. <https://doi.org/10.1016/j.jclinepi.2006.07.001>
- Chi, W. C., Overhage, J. M., Sponholtz, T. P., Nguyen, B. T., Brady, P. M., Falconi, A. P., & Johnson, M. P. (2023). The whole health index: A practical, valid, and reliable tool to measure whole-person health and manage population health. *New England Journal of Medicine Catalyst*. <https://doi.org/10.1056/CAT.23.0015>
- Cook, C. E., Rethorn, Z. D., Chiarotto, A., Garcia, A. N., & Gottfried, O. (2021). Construct validity and item response theory analysis of the PROMIS-29 v2.0 in recipients of lumbar spine surgery. *Spine*, *46*(24), 1721–1728. <https://doi.org/10.1097/BRS.00000000000004085>
- Cronbach, L. J. (1951). Coefficient alpha and the internal structure of tests. *Psychometrika*, *16*, 297–334. <https://doi.org/10.1007/BF02310555>
- Cummins, R. A., Eckersley, R., Pallant, J., Van Vugt, J., & Misajon, R. (2003). Developing a national index of subjective wellbeing: The Australian Unity wellbeing index. *Social Indicators Research*, *64*(2), 159–190. <https://doi.org/10.1023/A:1024704320683>
- Cummins, R. A., Lau, A. L., & Stokes, M. (2004). HRQOL and subjective well-being: Noncomplementary forms of outcome measurement. *Expert Review of Pharmacoeconomics & Outcomes Research*, *4*(4), 413–420. <https://doi.org/10.1586/14737167.4.4.413>
- Dewitt, B., Feeny, D., Fischhoff, B., Cella, D., Hays, R. D., Hess, R., Pilkonis, P. A., Revicki, D. A., Roberts, M. S., Tsevat, J., Lan, Y., & Hanmer, J. (2018). Estimation of a preference-based summary score for the patient-reported outcomes measurement information system: The PROMIS®-preference (PROPr) scoring system. *Medical Decision Making*, *38*(6), 683–698. <https://doi.org/10.1177/0272989X18776637>
- Deyo, R. A., Dworkin, S. F., Amtmann, D., Andersson, G., Borenstein, D., Carragee, E., Carrino, J., Chou, R., Cook, K., DeLitto, A., Goertz, C., Khalsa, P., Loeser, J., Mackey, S., Panagis, J., Rainville, J., Tosteson, T., Turk, D., Von Korf, M., & Weiner, D. K. (2014). Report of the NIH task force on research standards for chronic low back pain. *Pain Medicine*, *15*(8), 1249–1267. <https://doi.org/10.1111/pme.12538>
- Essink-Bot, M. L., Krabbe, P. F., Bonsel, G. J., & Aaronson, N. K. (1997). An empirical comparison of four generic health status measures. The Nottingham health profile, the medical outcomes study 36-item short-form health survey, the COOP/WONCA charts, and the EuroQol instrument. *Medical Care*, *35*(5), 522–537. <https://doi.org/10.1097/00005650-199705000-00008>
- Farivar, S. S., Cunningham, W. E., & Hays, R. D. (2007). Correlated physical and mental health summary scores for the SF-36 and SF-12 health survey, V.I. *Health and Quality of Life Outcomes*, *5*, 54. <https://doi.org/10.1186/1477-7525-5-54>

- Feng, Y. S., Kohlmann, T., Janssen, M. F., & Buchholz, I. (2021). Psychometric properties of the EQ-5D-5L: A systematic review of the literature. *Quality of Life Research, 30*(3), 647–673. <https://doi.org/10.1007/s11136-020-02688-y>
- Hahn, E. A., DeWalt, D. A., Bode, R. K., Garcia, S. F., DeVellis, R. F., Correia, H., & Cella, D. (2014). New English and Spanish social health measures will facilitate evaluating health determinants. *Health Psychology, 33*(5), 490–499. <https://doi.org/10.1037/hea0000055>
- Hays, R. D., Bjorner, J., Revicki, D. A., Spritzer, K. L., & Cella, D. (2009). Development of physical and mental health summary scores from the patient-reported outcomes measurement information system (PROMIS) global items. *Quality of Life Research, 18*(7), 873–880. <https://doi.org/10.1007/s11136-009-9496-9>
- Hays, R. D., Cunningham, W. E., Ettl, M. K., Beck, C. K., & Shapiro, M. F. (1995). Health-related quality of life in HIV disease. *Assessment, 2*(4), 363–380. <https://doi.org/10.1177/1073191195002004006>
- Hays, R. D., Edelen, M. O., Rodriguez, A., & Herman, P. (2021). Support for the reliability and validity of the National Institutes of Health impact stratification score in a sample of active-duty U.S. military personnel with low back pain. *Pain Medicine, 22*(10), 2185–2190. <https://doi.org/10.1093/pm/pnab175>
- Hays, R. D., Prince-Embury, S., & Chen, H.-Y. (1998). *RAND-36 health status inventory*. The Psychological Corporation.
- Hays, R. D., Qureshi, N., Herman, P. M., Rodriguez, A., Kapteyn, A., & Edelen, M. O. (2023). Effects of excluding those who report having “Syndromitis” or “Chekalism” on data quality: Longitudinal health survey of a sample from Amazon’s mechanical Turk. *Journal of Medical Internet Research, 25*, e46421. <https://doi.org/10.2196/46421>
- Hays, R. D., Spritzer, K. L., Schalet, B. D., & Cella, D. (2018). PROMS@-29 v2.0 profile physical and mental health summary scores. *Quality of Life Research, 27*(7), 1885–1891. <https://doi.org/10.1007/s11136-018-1842-3>
- Hays, R. D., & Stewart, A. L. (1990). The structure of self-reported health in chronic disease patients. *Psychological Assessment, 2*(1), 22–30. <https://doi.org/10.1037/1040-3590.2.1.22>
- Herdman, M., Gudex, C., Lloyd, A., Janssen, M., Kind, P., Parkin, D., Bonnel, G., & Badia, X. (2011). Development and preliminary testing of the new five-level version of EQ-5D (EQ-5D-5L). *Quality of Life Research, 20*(10), 1727–1736. <https://doi.org/10.1007/s11136-011-9903-x>
- Hoy, D., March, L., Brooks, P., Blyth, F., Woolf, A., Bain, C., Williams, G., Smith, E., Vos, T., Barendregt, J., Murray, C., Burstein, R., & Buchbinder, R. (2014). The global burden of low back pain: Estimates from the global burden of disease 2010 study. *Annals of the Rheumatic Diseases, 73*(6), 968–974. <https://doi.org/10.1136/annrheumdis-2013-204428>
- Hu, L. T., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural Equation Modeling, 6*(1), 1–55. <https://doi.org/10.1080/10705519909540118>
- International Wellbeing Group. (2013). *Personal wellbeing index* (5th ed.). Australian Centre on Quality of Life, Deakin University <https://www.acqol.com.au/uploads/pwi-a/pwi-a-english.pdf>. Accessed March 2024
- Ipsos. (2018). Be sure with KnowledgePanel. [https://www.ipsos.com/sites/default/files/18-11-53\\_Overview\\_v3.pdf](https://www.ipsos.com/sites/default/files/18-11-53_Overview_v3.pdf). Access February 2024.
- Jiang, R., Janssen, M. F. B., & Pickard, A. S. (2021). US population norms for the EQ-5D-5L and comparison of norms from face-to-face and online samples. *Quality of Life Research, 30*(3), 803–816. <https://doi.org/10.1007/s11136-020-02650-y>
- Kaplan, R. M., & Hays, R. D. (2022). Health-related quality of life measurement in public health. *Annual Review of Public Health, 43*, 355–373. <https://doi.org/10.1146/annurev-publhealth-052120-012811>
- Kligler, B. (2022). Whole health in the veterans health administration. *Global advances in Health and Medicine, 11*, 2164957X221077214. <https://doi.org/10.1177/2164957X221077214>
- Lau, A. L. D., & Cummins, R. A. (2005). *Test-retest reliability of the personal wellbeing index*. [Unpublished manuscript]. The Hong Kong Polytechnic University.
- MacCallum, R. C., Browne, M. W., & Sugawara, H. M. (1996). Power analysis and determination of sample size for covariance structure modeling. *Psychological Methods, 1*(2), 130–149. <https://doi.org/10.1037/1082-989X.1.2.130>
- McMullen, K., Bamer, A., Ryan, C. M., Schneider, J. C., Gibran, N., Stewart, B. T., Mroz, T., Wolf, S., & Amtmann, D. (2022). Validation of PROMIS-29 domain scores among adult burn survivors: A National Institute on Disability, Independent Living, and Rehabilitation Research burn model system study. *Journal of Trauma and Acute Care Surgery, 92*(1), 213–222. <https://doi.org/10.1097/TA.0000000000003365>

- Palimaru, A., & Hays, R. D. (2017). Associations of health-related quality of life with overall quality of life in the patient-reported outcomes measurement information system (PROMIS®) project. *Applied Research in Quality of Life*, 12(2), 241–250. <https://doi.org/10.1007/s11482-017-9515-x>
- Patrick, D. L., & Deyo, R. A. (1989). Generic and disease-specific measures in assessing health status and quality of life. *Medical Care*, 27(3), S217–S232. <https://doi.org/10.1097/00005650-198903001-00018>
- Pecorelli, N., Guarneri, G., Vallorani, A., Limongi, C., Licinio, A.W., Di Salvo, F., Tamburrino, D., Partelli, S., Crippa, S., & Falconi, M. (2023). Validation of the PROMIS-29 questionnaire as a measure of recovery after pancreatic surgery. *Annals of surgery*, 278(5), 732–739. [10.1097/SLA.0000000000006020](https://doi.org/10.1097/SLA.0000000000006020).
- Peipert, B. J., Goswami, S., Yount, S. E., & Sturgeon, C. (2018). Health-related quality of life in MEN1 patients compared with other chronic conditions and the United States general population. *Surgery*, 163(1), 205–211. <https://doi.org/10.1016/j.surg.2017.04.030>
- Pickard, A. S., Law, E. H., Jiang, R., Pullenayegum, E., Shaw, J. W., Xie, F., Oppe, M., Boye, K. S., Champman, R. H., Gong, C. L., Balch, A., & Busschback, J. J. V. (2019). United States valuation of EQ-5D-5L health states using an international protocol. *Value in Health*, 22(8), 931–941. <https://doi.org/10.1016/j.jval.2019.02.009>
- Reise, S. P., Morizot, J., & Hays, R. D. (2007). The role of the bifactor model in resolving dimensionality issues in health outcomes measures. *Quality of Life Research*, 16(Suppl 1), 19–31. <https://doi.org/10.1007/s11136-007-9183-7>
- Rodriguez, A., Reise, S. P., & Haviland, M. G. (2016). Evaluating bifactor models: Calculating and interpreting statistical indices. *Psychological Methods*, 21(2), 137–150. <https://doi.org/10.1037/met0000045>
- World Health Organization. (2002). Towards a common language for functioning, disability and health ICF. <https://cdn.who.int/media/docs/default-source/classification/icf/icfbeginnersguide.pdf>. Accessed March 2024.
- Wu, A., March, L., Zheng, X., Huang, J., Wang, X., Zhao, J., Blyth, F. M., Smith, E., Buchbinder, R., & Hoy, D. (2020). Global low back pain prevalence and years lived with disability from 1990 to 2017: Estimates from the global burden of disease study 2017. *Annals of Translational Medicine*, 8(6), 299. <https://doi.org/10.21037/atm.2020.02.175>
- Yin, S., Njai, R., Barker, L., Siegel, P. Z., & Liao, Y. (2016). Summarizing health-related quality of life (HRQOL): Development and testing of a one-factor model. *Population Health Metrics*, 14, 22. <https://doi.org/10.1186/s12963-016-0091-3>

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## Authors and Affiliations

Ron D. Hays<sup>1</sup>  · Anthony Rodriguez<sup>2</sup> · Nabeel Qureshi<sup>3</sup> · Chengbo Zeng<sup>4</sup> · Maria Orlando Edelen<sup>2,4</sup>

✉ Ron D. Hays  
drhays@ucla.edu

<sup>1</sup> UCLA Division of General Internal Medicine & Health Services Research, UCLA, Department of Medicine, Los Angeles, CA, USA

<sup>2</sup> Behavioral and Policy Sciences, RAND Corporation, Boston, MA, USA

<sup>3</sup> Behavioral and Policy Sciences, RAND Corporation, Santa Monica, CA, USA

<sup>4</sup> Patient Reported Outcomes, Value and Experience (PROVE) Center; Department of Surgery, Brigham and Women's Hospital, Boston, MA, USA