

UCSF

UC San Francisco Previously Published Works

Title

Tobacco use among substance use disorder (SUD) treatment staff is associated with tobacco-related services received by clients

Permalink

<https://escholarship.org/uc/item/539827hw>

Authors

Guydish, Joseph
Le, Thao
Hosakote, Sindhushree
et al.

Publication Date

2022

DOI

10.1016/j.jsat.2021.108496

Peer reviewed

Tobacco use among substance use disorder (SUD) treatment staff is associated with tobacco-related services received by clients

Joseph Guydish,^a Thao Le,^a Sindhushree Hosakote,^a Elana Straus,^a Jessie Wong,^a

Cristina Martínez,^b Kevin Delucchi^c

^a Philip R. Lee Institute for Health Policy Studies
University of California San Francisco
3333 California St., Ste. 265
San Francisco, CA 94118
Emails: joseph.guydish@ucsf.edu
thao.le@ucsf.edu
sindhu.hosakote@ucsf.edu
Elana.Strasu@ucsf.edu
jessiewong314@gmail.com

^b Tobacco Control Unit, Cancer Control and Prevention Programme
Institut Català d'Oncologia-ICO, Av. Granvia de L'Hospitalet 199-203, 08908
L'Hospitalet de Llobregat, Barcelona, Spain

□ ^c Department of Psychiatry
□ University of California San Francisco
□ 401 Parnassus Ave.
□ San Francisco, CA 94143
Email: kevin.delucchi@ucsf.edu

Corresponding Author:

Joseph Guydish, PhD
Professor of Medicine and Health Policy
Philip R. Lee Institute for Health Policy Studies
University of California San Francisco
3333 California St. Ste. 265
San Francisco, CA 94118
Phone: 415-476-0954
Email: joseph.guydish@ucsf.edu

Funding: This work was supported by the California Tobacco Related Disease Research Program (TRDRP 27IR-0040, 28CP-0038), by the NCI Cancer Center Support Grant (P30 CA082103), and by the California Tobacco Control Program(CTCP 18-10025).

Disclosures: The authors have no conflicts of interest to disclose.

Citation: Guydish J, Le T, Hosakote S, Straus E, Wong J, Martínez C & Delucchi K. (2022) Tobacco use among substance use disorder (SUD) treatment staff is

associated with tobacco-related services received by clients. [Journal of Substance Abuse Treatment, 132, doi.org/10.1016/j.jsat.2021.108496](https://doi.org/10.1016/j.jsat.2021.108496)

ABSTRACT

Background: Despite disproportionately high rates of smoking among people in residential substance use disorder (SUD) treatment, few receive tobacco cessation services. Little is known about how smoking among treatment staff may impact this disparity. We explored the relationship between staff tobacco use and client tobacco use. Additionally, we examined the relationship between staff tobacco use and tobacco-related services reported by staff and clients.

Methods: Staff (n = 363) and clients (n = 639) in 24 California publicly-funded residential SUD treatment programs were surveyed in 2019-20. Staff self-reported current tobacco use, as well as their beliefs, self-efficacy, and practices regarding smoking cessation. Clients reported their tobacco use and the services received while in treatment. Regression analyses examined the adjusted and unadjusted associations between staff and client tobacco use and other outcomes.

Results: Use of any tobacco product by staff ranged from 0% to 100% by program, with an average of 32% across programs. Adjusted analyses found that higher rates of staff tobacco use were associated with higher rates of client tobacco use, and with fewer clients receiving tobacco-related counseling. In programs that had higher rates of staff tobacco use, staff were less likely to believe that clients should quit smoking in treatment and had lower self-efficacy to address smoking.

Conclusion: Higher rates of tobacco use among staff are associated with higher rates of client tobacco use and fewer clients receiving cessation counseling. Efforts to reduce tobacco use among SUD clients should be supported by efforts to reduce

tobacco use among staff. SUD treatment programs, and agencies that fund and regulate those programs, should aim to reduce the use of tobacco products among staff.

INTRODUCTION

Smoking rates among persons with substance use disorders (SUD) are higher than in the general population (Richter et al., 2002; Smith et al., 2014), and showed no decline from 2002- 2014 (Weinberger et al., 2018). This disparity suggests that population level tobacco control strategies such as education and taxation have limited effect among people with SUDs (Warner, 2006). Smokers with SUDs smoke more heavily (Ward et al., 2012), have a harder time quitting smoking (Weinberger et al., 2016), and experience both excess and premature tobacco-related mortality (Bandiera et al., 2015).

About 2.4 million Americans enter SUD specialty treatment annually (SAMHSA, 2017), and among those the smoking prevalence is about 70% (Guydish et al., 2019). Receiving smoking cessation services while in SUD treatment is associated with improvement in SUD outcomes (Baca & Yahne, 2009; Prochaska et al., 2004) and continued smoking is associated with relapse to substance use (Weinberger et al., 2017). However, in one review only 40% of SUD programs offered cessation counseling and only 26% offered cessation medication (Knudsen, 2017). Resource limitations, such as lack of staff time or training, are commonly cited to explain this discrepancy (Sheals et al., 2016). Some staff believe that quitting smoking poses a risk to sobriety, or that clients are not interested in quitting (Guydish et al., 2007), although such beliefs have been debunked (Prochaska, 2011).

One reason why tobacco services are not delivered in SUD programs may be the level of tobacco use among staff (Cookson et al., 2014; Guydish et al., 2007). Several papers report SUD staff smoking rates 10% higher than that of the U.S. general population (Bobo & Davis, 1993; Gill & Bennett, 2000; Olsen et al., 2005; Rothrauff & Eby, 2011). Other papers report staff smoking rates only slightly higher than population rates (Chisolm et al., 2010; Laschober et al., 2015; Muilenburg et al., 2016; Pagano et al., 2016). Knudsen, Studts and Studts (2012), surveying counselors from over 400 programs, reported a staff smoking rate (20%) very close to the population rate at that time. Laschober et al (2015) surveyed counselors from over 200 programs, and reported a 21% staff smoking rate. Reports from the UK and Australia found staff smoking rates 10 - 25% higher than the general population (Cookson et al., 2014; Skelton et al., 2017).

Related questions are whether staff smoking is associated with client smoking rates, and whether staff smoking rates are associated with client receipt of tobacco-related services. While we found no studies for the former question, several papers have explored the latter. Early studies of staff suggest that those who smoke, as compared to non-smokers, have more resistant views towards smoking as a treatment issue (Bobo et al., 1995; Campbell et al., 1998). Staff who smoke less often provided tobacco-related services to clients (Knudsen & Studts, 2010), and more often believed that clients are not interested in quitting smoking (Laschober et al., 2015; Skelton et al., 2017). One study found that counselor smoking status was not associated with providing cessation counseling to clients (Knudsen et al., 2012)

Studies of both staff and clients, in the same program, may illuminate any relationship between staff smoking and providing tobacco-related services to

clients. Bernstein and colleagues (1999) surveyed staff and clients after initiating a choice-based smoking program within a treatment facility. While 38% of the clients had a counselor who smoked, staff smoking was not associated with client participation in the smoking program. Olsen et al (2005) found that 97% of counselors reported providing smoking cessation counseling to clients, but only 48% of clients reported receiving such counseling. Counselor smoking status was not associated with client receipt of cessation counseling.

While prior literature has focused on smoking combustible cigarettes, it may be helpful to consider use of other tobacco products since, among SUD clients, recent use of any tobacco product is 4-5% higher than use of combustible cigarettes alone (Guydish et al., 2016; Guydish et al., 2020). Assessing the relationship between tobacco use among staff and among clients, and between staff tobacco use and client receipt of tobacco-related services, requires data collection among staff and clients in the same program. It also requires data collection in a number of programs because the program becomes the unit of analysis. For these reasons, studies using client level variables aggregated to the program level, such as the association between staff and client tobacco use, are infrequent in the SUD treatment literature.

Such studies may be important, however, with respect to tobacco use in SUD treatment. High rates of tobacco use among staff may normalize tobacco use, hinder tobacco policy development or enforcement and, where tobacco using staff are reluctant to counsel clients about tobacco use, limit the level of tobacco-related services provided. In that case, reducing tobacco use among staff may be necessary in order to reduce tobacco use among clients. To assess the association between staff tobacco use and client tobacco use we surveyed both staff and clients

in 24 California publicly-funded residential SUD treatment programs. As a secondary aim, we examined whether tobacco use among staff was associated with measures of tobacco-related services as reported by both staff and clients.

METHODS

Program Recruitment

We collected data between January 2019 and July 2020 from 24 California licensed residential SUD programs recruited in the course of three studies designed to improve tobacco-related policies and services. Program inclusion criteria were for California state-licensed residential SUD programs, where the Program Director indicated willingness to participate in a tobacco-related intervention. The 24 programs were located in 12 of California's 58 counties, ranging from Lake County in the North to San Diego County in the South. Additional information about program selection and recruitment is reported in Guydish et al. (2020).

Participants

Participants included program clients, staff, and directors. Eligible clients were all those enrolled in the program at the time of data collection. Eligible staff were all full and part time paid staff working in the program. As three program directors led more than one program, 20 directors represented the 24 programs.

Procedures

Site visits to collect client surveys were generally completed in one day. Program directors reported the number of clients enrolled in the program, for use in calculating response rates. Research staff reviewed a study information sheet with clients in small groups, and gave each client a computer tablet survey with a pre-populated ID number. The client reviewed the study information sheet on the tablet

and used a button to consent or to decline participation. The anonymous survey took about 30 minutes. After the survey, research staff assessed clients' expired carbon monoxide (CO) was using a Bedfont piCO™ hand-held monitor (Bedfont Scientific Ltd, 2018). At four programs where no site visit occurred due to COVID restrictions, clients completed the same procedures on a computer provided by their program, however CO data were not collected because research staff were not present on site. Participants received a \$20 gift card.

Directors at each program provided the research team with staff work email addresses, r, and staff were then invited by email to complete the confidential online survey. After three weekly reminders, the research team used additional strategies approved by the Director. For example, having the Director send a reminder to non-responders, or having the research team re-send the invitation to non-responders. Staff respondents received a \$25 gift card.

Program directors completed an online tobacco policy survey. Study procedures were approved by the Institutional Review Board of the University of California San Francisco.

Measures

Client Measures. In addition to demographic characteristics, participants reported their health insurance coverage. The California Medicaid program (Medi-Cal) covers residential SUD treatment (DHCS, 2020), and also covers smoking cessation counseling and medication (DHCS, 2016). All three study surveys asked participants whether they sought treatment mainly for a substance use problem, for both substance use and mental health problems, or for some other problem. However, the survey for one study also allowed respondents to say they were in treatment for mental health problems only. For the purpose of analyses, these

cases were combined into the substance use and mental health category. Current smoking status was defined by asking clients “Do you currently smoke cigarettes?” with response codes “Yes, I currently smoke,” “No, I quit smoking,” and “No, I never smoked.”

In programs where CO measures were included, we used ≤ 9 ppm to verify non-smoking status. Earlier guidelines for biochemical verification of smoking status recommended using a cutoff from 8 -10 ppm (Benowitz et al., 2002). More recent guidelines recommend a range of 5 - 6 ppm while commenting that higher cutoffs may be needed where environmental exposure is high (Benowitz et al., 2019). Indoor smoking is not permitted in public buildings in California, including residential SUD programs, however the rate of smoking among clients is very high (59.6% in the current sample), and outdoor smoking on program grounds is common. Residential SUD clients, even non-smokers, are likely to be exposed to secondhand smoke in the course of entering and exiting buildings, or during breaks where clients socialize in designated outdoor smoking areas. Given the high rate of smoking and likelihood of secondhand smoke exposure, we used the more relaxed (≤ 9 ppm) cutoff. Clients who self-reported as non-smokers, but registered CO above the cutoff were regarded as “probable smokers,” and treated as current smokers for analyses. Clients also reported whether they had used e-cigarettes, smokeless tobacco, cigars or little filtered cigars in the past month. These items were used to calculate the proportion who used at least one tobacco product in the past month. Current smokers also reported number of cigarettes per day (CPD), whether they had made a quit attempt in the past year, and whether they were thinking about quitting smoking in the next 30 days.

The survey asked clients about four tobacco-related services they may have received in the treatment program. All clients reported whether any staff member had asked if they smoke. Current smokers reported whether they had attended a smoking cessation support group (yes/no), and how often their counselor encouraged them to quit smoking or arranged an appointment to discuss quitting. The last two items were dichotomized as Never vs. Occasionally/Often/Very Often/Always. If a client received one or more of these three services, they were coded as having received any counseling. Smokers who received a referral to either a smoking cessation specialist or a telephone quitline were coded as having received any referral. Last, smokers who received any nicotine replacement therapy (NRT) or other cessation medication were coded as receiving any NRT/Pharmacotherapy.

Staff Measures. In addition to demographic characteristics, staff self-reported current smoking status, but without biochemical verification. Staff reported whether they had used e-cigarettes, smokeless tobacco, cigars or little filtered cigars in the past month. These items were used to calculate the proportion who used any tobacco product in the past month. Current smokers reported number of cigarettes per day (CPD), whether they had made a quit attempt in the past year, and whether they were thinking about quitting smoking in the next 30 days.

Staff also completed the Smoking Knowledge, Attitudes and Practices (S-KAP) survey which includes multi-item scales reflecting beliefs about addressing tobacco use in SUD treatment ($\alpha = 0.74$), self-efficacy in providing tobacco-related services ($\alpha = 0.72$), and practices ($\alpha = 0.91$) used by counselors to address smoking with clients (Delucchi et al., 2009). The Belief scale includes 7 items which ask, for example, whether quitting smoking while in treatment threatens sobriety and

whether counseling motivates clients to quit. The Self-Efficacy scale includes 9 items concerning skills in providing tobacco-related services, for example, that clients want to quit and are likely to follow the clinician's advice. The Practice scale includes 9 items which ask, for example, how often the counselor asks, advises, or assists clients with quitting smoking. All responses are scored from 1 to 5, and the mean of items comprises the scale score. Higher scale scores reflect more positive beliefs about addressing smoking, greater self-efficacy and greater use of practices to address smoking. Scale items, response codes, and mean (SD) values for the current staff sample are included in Supplemental Table 1.

Residential SUD programs include both clinical and non-clinical staff however, only clinical staff would deliver tobacco-related services. Consequently, the Beliefs scale was completed by all staff, while only clinical staff completed the Self-efficacy and Practice scales. Clinical staff were those who had an active client caseload in the past month (values ≥ 1) and/or had conducted group or individual sessions in the past week (values ≥ 1).

Program Tobacco Policy. Program directors reported on the tobacco policy at each program was measured using a survey developed for this study. The survey included 20 items drawn from prior research concerning tobacco free grounds (Mullenburg et al., 2016), smoking among staff (Cookson et al., 2014; Skelton et al., 2017), and staff and clients smoking together (Guydish et al., 2017). Each item is assigned a score of 1 if the response is aligned with strategies that discourage smoking. For example, the program receives points if clients (1 point) and staff (1 point) are not permitted to smoke outdoors on program grounds, and if the program provides tobacco-related screening (1 point), advice (1 point), counseling (1 point), referral (1 point) or education (1 point). The total possible score is 20, and in this

sample of 24 programs the tobacco policy scores ranged from 3 to 19 (median = 11). The scale items and scoring are found at <https://doi.org/10.6084/m9.figshare.14550981.v1>

Data Analysis

We report demographic and tobacco use characteristics, aggregated across programs, for clients and staff. For clients we report the proportion who received each of the four tobacco-related services and, for staff, the mean values for the three scales.

We assessed the association between staff use any tobacco product in each program as the predictor and each of 8 outcomes (client use of any tobacco product, the four client-reported tobacco service measures, and the three staff-reported measures) using general linear models. The unit of the analysis was the clinic (N=24), limiting sample size. The small sample size hindered the number of control variables included in the models. We controlled for two covariates likely to influence associations between staff smoking rates and client receipt of tobacco-related services. These are the proportion of clients in each program who were covered by Medi-Cal and for whom cessation services were covered as a healthcare benefit, and the strength of program tobacco policies which may encourage quitting and reduce tobacco-related disease in this population (Marynak, 2016). Unadjusted estimates (with 95% CI) and adjusted estimates (with 95% CI) were presented. We classified 47 cases as “probable smokers” due to discordant self-report and expired CO measures, and these cases were included in the main analyses as current smokers. As it is possible that these cases were not smokers, we repeated multivariate analyses testing the association between staff and client smoking prevalence, but excluding the 47 “probable smokers.” Last, we graphically

represent the association between staff use of any tobacco product in each program and client use of any tobacco product in its respective program using a scatterplot.

RESULTS

Demographic and tobacco use characteristics

Across all programs, 639 clients completed the survey, representing 84% of those eligible. Participation rates in individual programs ranged from 67% to 100%. Demographic, insurance, and reason for treatment variables are given in Table 1. With non-smoking status verified by expired CO \leq 9 ppm, client smoking prevalence was 67%. This prevalence includes 47 cases who self-reported as non-smokers but registered above the CO cutoff. An additional 26 cases reported no current use of combustible cigarettes, but used other tobacco products in the past month, giving a total of 454 (71%) tobacco users.

Insert Table 1 about here

Across all programs, 363 staff completed the survey, representing 80% of those eligible. Participation rates in individual programs ranged from 59% to 100%. Staff demographic characteristics are reported in Table 1. Nearly two-thirds (65.4%) had clinical responsibilities. Among staff, 21.8% self-reported as current smokers, and 32% had used any tobacco product in the past month.

Among current smokers, for clients and staff respectively, mean (SD) CPD was 9.8 (7.7) and 7.9 (5.7), proportions reporting a quit attempt in the past year were 59.8% and 65.8%, and proportions thinking of quitting smoking in the next 30 days were 32.4% and 39.2% (data not shown). As the Tobacco Policy Measure was developed for this study, we assessed the correlation between the policy measure

(reported by program directors) and the Belief, Self-Efficacy, and Practice scales (reported by program staff) in the 24 programs. The policy measure was positively and significantly correlated with the Belief ($r = 0.52$, $p < .01$) and Practice ($r = 0.42$, $p < .05$) scales, but the correlation fell short of significance for the Self-Efficacy scale ($r = 0.39$, $p = .063$). These findings suggest that tobacco policy measure reported by program directors was positively associated with tobacco-related beliefs and practices reported by staff in the same program.

Tobacco related services measures reported by clients and staff

Table 2 summarizes, in the first column, the proportion of clients who were asked about their smoking status, and who received tobacco-related counseling, medication, or referral. The denominator varies because smoking status was asked of all clients ($N = 639$), while receipt of tobacco-related services was asked only of those who reported current smoking status or quitting smoking while in the treatment program ($n = 466$). Among all clients, 64.1% had been asked their smoking status. Among tobacco users or those who quit in the program, 57.1% had received tobacco-related counseling, 26% had received cessation medication, and 32.5% had received tobacco-related referral.

Insert Table 2 about here

The rightmost column of Table 2 shows mean (SD) values for the three S-KAP scales. The denominator varies by scale, as the belief scale items were asked of all staff ($N = 363$) while the efficacy and practice scale items were asked of clinical

staff (n = 236). All mean scale scores shown occur on a scale of 1 to 5, where higher scores indicate greater belief about treating tobacco use, and greater ability to do so.

Association between staff tobacco, client tobacco use, and tobacco service measures

Results of analyses for associations between staff use of any tobacco product and both client and staff outcome measures are shown in Table 3. At the top of the table, the first unadjusted estimate shows that the rate of staff tobacco use was positively associated with the rate of client tobacco use (beta = 0.63, CI 0.13, 1.12). After adjusting for the proportion of clients who were covered by Medi-Cal and strength of the tobacco policy in each program, this association remained significant (beta = 0.59, 95% CI 0.06 - 1.12, p = 0.03). In the sensitivity analysis, excluding the 47 “probable smokers,” the finding was the same (beta = 0.64, 95% CI 0.13 - 1.15, p = 0.017, data not shown).

Insert Table 3 about here

The rightmost column of Table 3 shows the association of staff tobacco use with other outcomes, adjusting for the proportion of clients covered by Medi-Cal, and controlling for the strength of program tobacco policies. The prevalence of tobacco use among staff was significantly and inversely associated with client receipt of tobacco-cessation counseling, staff beliefs about having clients quit in drug treatment, and staff self-efficacy to assist clients with quitting.

Prevalence of past month tobacco use among both staff and clients in the 24 programs is plotted in Figure 1. The program at bottom left had 0% of staff and

20% of clients reporting recent use of tobacco products. The program at far right had 100% of staff and 86% of clients reporting use of tobacco products

Insert Figure 1 about here

DISCUSSION

In this sample of 24 publicly-funded California residential SUD treatment programs, staff smoking ranged from 0% to 57.1% by program and, collapsed across all programs, staff smoking prevalence was 21.8%. This prevalence is in contrast to the California general population smoking rate of 11.2% in 2018 (Centers for Disease Control and Prevention., 2018). We consider not only use of combustible cigarettes, but also use of other tobacco products. In the current sample, prevalence of use of any tobacco product among staff ranged from 0% to 100% and, collapsed across programs was 32%.

One explanation for elevated rates of staff smoking and use of other tobacco products may be the tradition of hiring persons who are in recovery from substance use. Doing so offers an available recruitment pool, dedicated to SUD treatment, and with lived experience relevant to the counseling mission (Guydish et al., 2017). However, if the use of tobacco products is high among clients, and if those clients enter the treatment workforce, then use of tobacco products may also be high in the workforce.

Multivariate analyses found that higher rates of staff tobacco use were associated with higher rates of client tobacco use, and with fewer clients receiving tobacco-related counseling. In programs with higher rates of staff tobacco use, staff

were less likely to believe that clients should quit smoking in the treatment program, and had lower self-efficacy to address smoking among clients.

Our analyses suggest that greater tobacco use among staff is associated with fewer tobacco services among clients, but it may depend on who is asked. When staff were asked about tobacco services they provide to clients, we see the association in unadjusted analysis but significant is lost in the adjusted analysis. This loss of significance may indicate no relationship, or a relationship that could be seen only in a larger sample. When clients were asked about services they received, the association was present and unambiguous. Reports by clients regarding tobacco-related services they received in a program may be more reliable than staff reports about tobacco services they provide, to the degree that they reflect what services clients actually received, or what services they recognized as tobacco-related.

The current findings are broadly consistent with literature showing that staff who smoke held more negative views about addressing smoking (Bobo et al., 1995; Campbell et al., 1998), more often believed that clients were not interested in quitting smoking (Skelton et al., 2017), and were less likely to implement smoking related client services (Bobo et al., 1995; Campbell et al., 1998; Knudsen & Studts, 2010; Laschober et al., 2015).

Study limitations include small sample size ($N = 24$), which restricts the number of covariates that can be controlled in analyses. The two covariates selected, whether clients had Medi-Cal insurance that covered smoking cessation services and the strength of the tobacco policy in each program, have face validity. However, other covariates could be tested and, particularly in a larger sample of programs, could result in different findings. All participating programs were

recruited in California, which has the lowest general population smoking rate (11.2%) of any US State excepting Utah, and a robust state tobacco control program (Centers for Disease Control and Prevention., 2018; Roeseler & Burns, 2010). If staff and client tobacco use rates are influenced by statewide smoking rates, then our findings concerning prevalence of staff smoking may underestimate staff smoking rates in states where population smoking rates are higher. The study sample includes adult residential SUD programs only which, in California, comprise 19% of all publicly funded treatment programs (Guydish et al., 2020). Findings may not extend to outpatient, methadone, or adolescent focused programs. All programs in this sample were publicly-funded programs. Higher rates of smoking, or of use of any tobacco product, may be more likely in publicly-funded treatment systems. Hospital-based programs, Veteran’s Affairs programs, and private healthcare systems providing their own chemical dependence services may include more professional and medical staff where smoking rates are low. However, two-thirds of current SUD treatment occurs in the public sector (Mark et al., 2007; Mechanic et al., 1995), so high rates of smoking among SUD staff may occur in the preponderance of programs. The measure of tobacco policy strength was created for this research, and has not been validated. Last, programs in this study were recruited for research on tobacco-free grounds, and had expressed interest in better addressing tobacco use among clients. Findings may be conservative if other programs not included in the study tended to have little or no interest in changing practices to address smoking.

We are aware of no other study in which staff and client tobacco-related variables were assessed in a similar or larger number of SUD treatment programs, and where staff tobacco use was assessed for relationships with both client and

staff variables. Findings show that higher rates of tobacco use among staff are associated with higher rates of tobacco use and lower rates of tobacco-related services among clients. SUD treatment programs, and agencies that fund and regulate those programs, should work to reduce use of all tobacco products among staff. This is in the interest of the treatment workforce as well as in the interest of the treatment clients. Smoke-free workplace policies, which reduce smoking (Fichtenberg & Glantz, 2002) and increase health equity (Hafez et al., 2019), should be implemented in SUD settings no less than they are implemented in virtually every other healthcare setting.

Table 1: Demographic Characteristics for Clients (N = 639) and Staff (N = 363) Across 24 SUD Treatment Programs

| | Mean (SD) or n (%) | |
|---|--------------------|---------------|
| | Clients (n=639) | Staff (n=363) |
| Age, mean | 38.5 (11.7) | 44.6 (12.5) |
| Gender, % | | |
| Male | 460 (72.2%) | 122 (33.7%) |
| Female | 166 (26.1%) | 233 (64.4%) |
| Other | 11 (1.7%) | 7 (1.9%) |
| Race/Ethnicity, % | | |
| Hispanic/Latino | 255 (39.9%) | 111 (30.8%) |
| Black or African American | 126 (19.7%) | 81 (22.4%) |
| White or Caucasian | 193 (30.2%) | 124 (34.4%) |
| Other/Multiple ¹ | 65 (10.2%) | 45 (12.5%) |
| Education, % | | |
| No HS diploma/GED | 164 (25.7%) | 7 (2.0%) |
| High school diploma or GED | 223 (34.9%) | 52 (14.7%) |
| Some college or technical/trade school ² | 186 (29.1%) | 127 (36.0%) |
| Bachelor's or Associate's | 66 (10.3%) | 104 (29.5%) |
| Graduate degree ³ | | 63 (17.9%) |
| Smoking Status, % | | |
| Current Smoker | 381 (59.6%) | 79 (21.8%) |
| Probable smoker ⁴ | 47 (7.4%) | |
| Former Smoker | 150 (23.5%) | 177 (48.8%) |
| Never Smoker | 61 (9.6%) | 107 (29.5%) |
| Any E-cigarette use past month | 47 (23.4%) | 61 (16.8%) |
| Any Tobacco Product use past month | 454 (71.1%) | 116 (32.0%) |
| Healthcare coverage, % | | |
| Medi-Cal | 453 (70.9%) | |
| Medicare | 30 (4.7%) | |
| Employer, family, or other plan ⁵ | 38 (6.0%) | |
| No healthcare coverage | 71 (11.1%) | |

| | | |
|---|-------------|--------------------|
| Don't know/not sure if covered | 47 (7.4%) | |
| In treatment for, % | | |
| Substance use | 356 (56.2%) | |
| Both substance use and mental health ⁶ | 189 (29.8%) | |
| Other reason for treatment | 89 (14.0%) | |
| Clinical Staff, n (%) | | 236 (65.4%) |

1. Client category includes American Indian or Alaska Native (2.7%), Asian/Pacific Islander (1.7%), and those reporting multiple (3.8%) and "other" (2.0%) race/ethnicity. Staff category includes American Indian or Alaska Native (0.8%), Asian (2.2%), Native Hawaiian/Pacific Islander (1.1%), those reporting multiple (4.2%), and "other" (4.2%) race/ethnicity
2. For staff, this category includes cases reporting college coursework related to counseling licensure.
3. Graduate training was not asked of clients
4. Self-reported as non-smokers but registered > 9 ppm on expired CO measure.
5. Includes cases reporting another source of health insurance (2.8%) and those reporting that source of health insurance was unknown (1.1%)
6. In one of the projects, response codes for this item included "mental health only" (with 3.2%) responses. To represent the entire sample, these cases are collapsed into "Both substance use and mental health disorders."

Table 2: Tobacco-related services reported by Clients and Staff across 24 SUD Treatment Programs

| | Mean (SD) or n (%) | |
|--|--------------------|---------------|
| | Clients | Staff |
| Client was asked about smoking ¹ | 409 (64.1%) | N/A |
| Client received any counseling ² | 265 (57.1%) | N/A |
| Client received any cessation medications ² | 121 (26.0%) | N/A |
| Client received any referral ² | 151 (32.5%) | N/A |
| Staff belief scale ³ | N/A | 3.5 (0.70) |
| Staff self- efficacy scale ⁴ | N/A | 3.2 (0.69) |
| Staff practice scale ⁴ | N/A | 2.6 (1.03) |

¹All clients (N=639)

²Self-reported current smokers and former smokers who quit in the program (N=466)

³All staff (N=363)

⁴Clinical staff (N=236)

Table 3: Association between prevalence of tobacco use among staff and selected client and staff outcomes (N = 24 programs)

| | Unadjusted Estimates (95%CI) | p-value | Adjusted Estimates (95%CI) ¹ | p-value |
|---|---------------------------------|------------------|---|------------------|
| Client tobacco use prevalence | 0.63 (0.13, 1.12) | 0.016 | 0.59 (0.06, 1.12) | 0.030 |
| Client was asked about smoking | -0.13 (-0.53, 0.27) | 0.516 | -0.11 (-0.49, 0.27) | 0.548 |
| Client received any counseling | -0.74 (-1.15, -0.33) | 0.001 | -0.69 (-1.12, -0.27) | 0.003 |
| Client received any cessation medications | -0.28 (-0.80, 0.24) | 0.272 | -0.18 (-0.69, 0.32) | 0.456 |
| Client received any referral | -0.41 (-0.84, 0.03) | 0.065 | -0.35 (-0.76, 0.07) | 0.100 |
| Staff belief scale | -0.02 (-0.02, -0.01) | <0.001 | -0.01 (-0.02, -0.01) | <0.001 |
| Staff self-efficacy scale | -0.01 (-0.02, -0.01) | <0.001 | -0.01 (-0.02, -0.01) | <0.001 |
| Staff practice scale | -0.01 (-0.02, -0.001) | 0.041 | -0.01 (-0.02, 0.001) | 0.083 |

¹Adjusted for % Med-Cal clients and Tobacco policy strength score

Figure 1: Plot of staff tobacco use prevalence v. client tobacco use prevalence (N = 24 programs)

Supplemental Table 1. Smoking Knowledge, Attitudes and Practice (S-KAP) Scale items, response codes and mean (SD) values for staff in 24 residential substance use treatment programs*

| Attitude - Beliefs Scale items | Mean (SD) |
|---|------------------|
| If a patient has been in recovery from alcoholism for less than 6 months, quitting smoking would threaten their sobriety 1 = Strongly Agree, 2 = Agree, 3 = Unsure, 4 = Disagree, 5 = Strongly Disagree | 3.2 (1.18) |
| Smoking is a personal decision which does not concern the clinician 1 = Strongly Agree, 2, 3, 4, 5 = Strongly Disagree | 3.2 (1.23) |
| Smoking cessation counseling is an important part of my agency's mission 1 = Strongly Disagree, 2 = Disagree, 3 = Unsure, 4 = Agree, 5 = Strongly Agree | 3.6 (1.24) |
| Counseling by a clinician helps motivate smokers to quit 1 = Strongly Disagree, 2, 3, 4, 5 = Strongly Agree | 3.8 (1.04) |
| Clinicians should make appointments specifically to help patients quit 1 = Strongly Disagree, 2, 3, 4, 5 = Strongly Agree | 3.8 (0.99) |
| In your opinion, what is the best point to encourage clients to stop smoking? 1 = Never, 2 = After 1 year of treatment, 3 = After 6 months of treatment, 4 = After one to three months of treatment, 5 = As soon as they begin treatment | 4.3 (1.06) |
| In your opinion, for clients who use drugs and smoke cigarettes, which should come first? 1 = Quit using drugs, 3 = Quit smoking, 5 = Quit smoking and using drugs at the same time | 2.4 (1.89) |
| <hr/> Self-Efficacy scale items <hr/> | |
| I have the required skills to help my patients quit smoking 1 = Strongly Disagree, 2, 3, 4, 5 = Strongly Agree | 3.5 (1.14) |
| My patients are concerned about smoking 1 = Strongly Disagree, 2, 3, 4, 5 = Strongly Agree | 3.2 (1.13) |

| | |
|---|------------|
| My patients follow my advice about behavior change 1 = Strongly Disagree, 2, 3, 4, 5 = Strongly Agree | 3.8 (0.76) |
| My patients want to quit smoking 1 = Strongly Disagree, 2, 3, 4, 5 = Strongly Agree | 3.1 (1.00) |
| I know where to refer patients for help with smoking cessation 1 = Strongly Disagree, 2, 3, 4, 5 = Strongly Agree | 3.6 (1.08) |
| If you counseled all your patients who smoke, what percentage do you think would try to quit smoking for more than 6 months? 1 = 0%, 1 = 1-10%, 2 = 11-20%, 3 = 21-30%, 4 = 31-40%, 5 = > 40% | 2.5 (1.46) |
| If you counseled all your patients who smoke, what percentage do you think would successfully quit smoking for more than 6 months 1 = 0%, 1, 2, 3, 4, 5 = > 40% | 2.0 (1.28) |
| How confident are you in your current ability to treat tobacco addiction? 1 = Not at all confident, 2 = Not very confident, 3 = Unsure, 4 = Confident, 5 = Very confident | 3.3 (1.17) |
| Given your clinical experience, how much emphasis do you believe should be placed on tobacco dependence treatment while in drug abuse treatment 1 = none at all, 3 = a moderate amount, 5 = a great deal | 3.7 (1.20) |

Practice scale items

| | |
|---|------------|
| Smoking cessation counseling is an important part of my job 1 = Strongly Disagree, 2, 3, 4, 5 = Strongly Agree | 3.2 (1.14) |
| In the past month, how frequently did you ask your patients whether they smoked 1 = Never, 2 = Occasionally, 3 = Often, 4 = Very Often, 5 = Always | 2.8 (1.42) |
| In the past month, how frequently did you advise patients who did smoke to quit 1 = Never, 2, 3, 4, 5 = Always | 2.4 (1.31) |

In the past month, how frequently did you assist patients who wanted to stop smoking with referrals and advice to quit 2.7 (1.50)

1 = Never, 2, 3, 4, 5 = Always

In the past month, how frequently did you arrange a follow up visit or phone call to discuss quitting? 2.1 (1.44)

1 = Never, 2, 3, 4, 5 = Always

In the past month, how frequently did you encourage patients who smoke to stop smoking completely? 2.4 (1.39)

1 = Never, 2, 3, 4, 5 = Always

In the past month, how frequently did you encourage patients who smoke to use nicotine replacement 2.4 (1.33)

1 = Never, 2, 3, 4, 5 = Always

In the past month, how frequently did you encourage patients who smoke to reduce smoking to five or fewer cigarettes per day, if patient stated they could not quit 2.3 (1.32)

1 = Never, 2, 3, 4, 5 = Always

In the past month, how frequently did you encourage clients who smoke to not smoke in the presence of infants or children 3.2 (1.72)

1 = Never, 2, 3, 4, 5 = Always

* Notes: Scale development and psychometric properties are reported in Delucchi, Tajima & Guydish, 2009. Mean scores reported are for substance use treatment staff recruited from 24 California residential treatment programs. Attitude - Belief scale items were asked of all staff (N = 363), while Self-Efficacy and Practice items were asked of clinical staff only (n = 236).

References

- Baca, C. T., & Yahne, C. E. (2009). Smoking cessation during substance abuse treatment: what you need to know. *J Subst Abuse Treat*, 36(2), 205-219. <https://doi.org/10.1016/j.jsat.2008.06.003>
- Bandiera, F. C., Anteneh, B., Le, T., Delucchi, K., & Guydish, J. (2015). Tobacco-related mortality among persons with mental health and substance abuse problems. *PLoS One*, 10(3), e0120581. <https://doi.org/10.1371/journal.pone.0120581>
- Bedfont Scientific Ltd. (2018). *piCO monitor*. <https://www.bedfont.com/shop/smokerlyzer/pico>
- Benowitz, N. L., Ahijevych, K., Jarvis, M. J., Hall, S., LeHouezec, J., Hansson, A., Lichtenstein, E., Henningfield, J., Tsoh, J., Hurt, R. D., & Velicer, W. (2002). Biochemical verification of tobacco use and cessation. *Nicotine Tob Res*, 4(2), 149-159. <https://doi.org/10.1080/14622200210123581>
- Benowitz, N. L., Bernert, J. T., Foulds, J., Hecht, S. S., Jacob, P., Jarvis, M. J., Joseph, A., Oncken, C., & Piper, M. E. (2019). Biochemical Verification of Tobacco Use and Abstinence: 2019 Update. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*. <https://doi.org/10.1093/ntr/ntz132>
- Bernstein, S. M., & Stoduto, G. (1999). Adding a choice-based program for tobacco smoking to an abstinence-based addiction treatment program. *Journal of Substance Abuse Treatment*, 17(1-2), 167-173. [https://doi.org/10.1016/s0740-5472\(98\)00071-3](https://doi.org/10.1016/s0740-5472(98)00071-3)
- Bobo, J. K., & Davis, C. M. (1993). Recovering staff and smoking in chemical dependency programs in rural Nebraska. *J Subst Abuse Treat*, 10(2), 221-227. [https://doi.org/10.1016/0740-5472\(93\)90047-6](https://doi.org/10.1016/0740-5472(93)90047-6)
- Bobo, J. K., Slade, J., & Hoffman, A. L. (1995). Nicotine addiction counseling for chemically dependent patients. *Psychiatr Serv*, 46(9), 945-947. <https://doi.org/10.1176/ps.46.9.945>
- Campbell, B. K., Krumenacker, J., & Stark, M. J. (1998). Smoking cessation for clients in chemical dependence treatment. A demonstration project. *J Subst Abuse Treat*, 15(4), 313-318. [https://doi.org/10.1016/s0740-5472\(97\)00197-9](https://doi.org/10.1016/s0740-5472(97)00197-9)
- Centers for Disease Control and Prevention. (2018). *Current Cigarette Use by Adults, Office on Smoking and Health, National Center for Chronic Disease Prevention and Health Promotion*. <https://www.cdc.gov/statesystem/cigaretteuseadult.html>
- Chisolm, M. S., Brigham, E. P., Lookatch, S. J., Tuten, M., Strain, E. C., & Jones, H. E. (2010). Cigarette smoking knowledge, attitudes, and practices of patients and staff at a perinatal substance abuse treatment center. *J Subst Abuse Treat*, 39(3), 298-305. <https://doi.org/10.1016/j.jsat.2010.06.008>
- Cookson, Strang, J., Ratschen, E., Sutherland, G., Finch, E., & McNeill, A. (2014). Smoking and its treatment in addiction services: clients' and staff behaviour and attitudes. *BMC Health Services Research*, 14, 304. <https://doi.org/10.1186/1472-6963-14-304>
- Delucchi, K. L., Tajima, B., & Guydish, J. (2009). Development of the Smoking Knowledge, Attitudes, and Practices (S-KAP) Instrument. *Journal of drug issues*, 39(2), 347-364. <https://doi.org/10.1177/002204260903900207>
- DHCS. (2016). *Comprehensive tobacco prevention and cessation services for Medi-Cal beneficiaries*. <https://www.dhcs.ca.gov/formsandpubs/Documents/MMCDAPLsandPolicyLetters/APL2016/APL16-014.pdf>
- DHCS. (2020, 2/25/2020 12:05 PM). *Essential Health Benefits*. https://www.dhcs.ca.gov/services/medical/Pages/Benefits_services.aspx
- Fichtenberg, C. M., & Glantz, S. A. (2002). Effect of smoke-free workplaces on smoking behaviour: systematic review. *BMJ*, 325(7357), 188. <https://doi.org/10.1136/bmj.325.7357.188>
- Gill, B. S., & Bennett, D. L. (2000). Addiction professionals' attitudes regarding treatment of nicotine dependence. *J Subst Abuse Treat*, 19(4), 317-318. [https://doi.org/10.1016/s0740-5472\(00\)00106-9](https://doi.org/10.1016/s0740-5472(00)00106-9)

- Guydish, Le, Campbell, Yip, Ji, & Delucchi. (2017). Drug abuse staff and clients smoking together: A shared addiction. *Journal of substance abuse treatment*, 76, 64-68. <https://doi.org/10.1016/j.jsat.2017.01.014>
- Guydish, Passalacqua, E., Tajima, B., & Manser, S. T. (2007). Staff smoking and other barriers to nicotine dependence intervention in addiction treatment settings: a review. *J Psychoactive Drugs*, 39(4), 423-433. <https://doi.org/10.1080/02791072.2007.10399881>
- Guydish, Tajima, B., Pramod, S., Le, T., Gubner, N. R., Campbell, B., & Roman, P. (2016). Use of multiple tobacco products in a national sample of persons enrolled in addiction treatment. *Drug and Alcohol Dependence*, 166, 93-99. <https://doi.org/10.1016/j.drugalcdep.2016.06.035>
- Guydish, Wahleithner, J., Williams, D., & Yip, D. (2020). Tobacco-free grounds implementation in California residential substance use disorder (SUD) treatment programs. *Journal of Addictive Diseases*, 38(1), 55-63. <https://doi.org/10.1080/10550887.2020.1713687>
- Guydish, Yip, Le, T., Gubner, N. R., Williams, D. D., & Delucchi, K. L. (2019). Tobacco Cessation Services in Addiction Treatment: What Do Clients Say? *Psychiatr Serv*, 70(3), 229-232. <https://doi.org/10.1176/appi.ps.201700565>
- Guydish, J., Kapiteni, K., Le, T., Campbell, B., Pinsker, E., & Delucchi, K. (2020). Tobacco use and tobacco services in California substance use treatment programs. *Drug and Alcohol Dependence*, 214, 108173. <https://doi.org/10.1016/j.drugalcdep.2020.108173>
- Hafez, A. Y., Gonzalez, M., Kulik, M. C., Vijayaraghavan, M., & Glantz, S. A. (2019). Uneven Access to Smoke-Free Laws and Policies and Its Effect on Health Equity in the United States: 2000-2019. *American Journal of Public Health*, 109(11), 1568-1575. <https://doi.org/10.2105/AJPH.2019.305289>
- Knudsen. (2017). Implementation of smoking cessation treatment in substance use disorder treatment settings: a review. *Am J Drug Alcohol Abuse*, 43(2), 215-225. <https://doi.org/10.1080/00952990.2016.1183019>
- Knudsen, Studts, C. R., & Studts, J. L. (2012). The implementation of smoking cessation counseling in substance abuse treatment. *J Behav Health Serv Res*, 39(1), 28-41. <https://doi.org/10.1007/s11414-011-9246-y>
- Knudsen, & Studts, J. L. (2010). The implementation of tobacco-related brief interventions in substance abuse treatment: a national study of counselors. *Journal of substance abuse treatment*, 38(3), 212-219. <https://doi.org/10.1016/j.jsat.2009.12.002>
- Laschober, T. C., Muilenburg, J. L., & Eby, L. T. (2015). Factors Linked to Substance Use Disorder Counselors' (Non)Implementation Likelihood of Tobacco Cessation 5 A's, Counseling, and Pharmacotherapy. *J Addict Behav Ther Rehabil*, 4(1). <https://doi.org/10.4172/2324-9005.1000134>
- Mark, T. L., Levit, K. R., Vandivort-Warren, R., Coffey, R. M., & Buck, J. A. (2007). Trends in spending for substance abuse treatment, 1986-2003. *Health Affairs*, 26(4), 1118-1128. <https://doi.org/10.1377/hlthaff.26.4.1118>
- Marynak, V. B., Tetlow S, et al. (2016). Tobacco Cessation Interventions and Smoke-Free Policies in Mental Health and Substance Abuse Treatment Facilities — United States. *MMWR Morb Mortal Wkly Rep* 2018;67:519–523. <http://dx.doi.org/10.15585/mmwr.mm6718a3>
- Mechanic, D., Schlesinger, M., & McAlpine, D. D. (1995). Management of mental health and substance abuse services: State of the art and early results. *Milbank Quarterly*, 73(1), 19-55. <https://doi.org/10.2307/3350312>
- Muilenburg, J. L., Laschober, T. C., Eby, L. T., & Moore, N. D. (2016). Prevalence of and Factors Related to Tobacco Ban Implementation in Substance Use Disorder Treatment Programs. *Administration and policy in mental health*, 43(2), 241-249. <https://doi.org/10.1007/s10488-015-0636-y>
- Olsen, Y., Alford, D. P., Horton, N. J., & Saitz, R. (2005). Addressing smoking cessation in methadone programs. *J Addict Dis*, 24(2), 33-48. https://doi.org/10.1300/J069v24n02_04

- Pagano, A., Guydish, J., & Le, T., et al. (2016). Smoking Behaviors and Attitudes Among Clients and Staff at New York Addiction Treatment Programs Following a Smoking Ban: Findings After 5 Years. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*, 18(5), 1274-1281. <https://doi.org/10.1093/ntr/ntv116>
- Prochaska, J. J. (2011). Smoking and mental illness--breaking the link. *The New England journal of medicine*, 365(3), 196-198. <https://doi.org/10.1056/NEJMp1105248>
- Prochaska, J. J., Delucchi, K., & Hall, S. M. (2004). A meta-analysis of smoking cessation interventions with individuals in substance abuse treatment or recovery. *J Consult Clin Psychol*, 72(6), 1144-1156. <https://doi.org/10.1037/0022-006x.72.6.1144>
- Richter, K. P., Ahluwalia, H. K., Mosier, M. C., Nazir, N., & Ahluwalia, J. S. (2002). A population-based study of cigarette smoking among illicit drug users in the United States. *Addiction (Abingdon, England)*, 97(7), 861-869. <https://doi.org/10.1046/j.1360-0443.2002.00162.x>
- Roeseler, A., & Burns, D. (2010). The quarter that changed the world. *Tobacco Control*, 19 Suppl 1(Suppl_1), i3-i15. <https://doi.org/10.1136/tc.2009.030809>
- Rothrauff, T. C., & Eby, L. T. (2011). Counselors' knowledge of the adoption of tobacco cessation medications in substance abuse treatment programs. *The American journal on addictions*, 20(1), 56-62. <https://doi.org/10.1111/j.1521-0391.2010.00095.x>
- SAMHSA. (2017). *National Survey of Substance Abuse Treatment Services (N-SSATS): 2015. Data on Substance Abuse Treatment Facilities. BHSIS Series S-88,(HHS Publication No. (SMA) 17-5031.* https://www.samhsa.gov/data/sites/default/files/2015_National_Survey_of_Substance_Abuse_Treatment_Services.pdf
- Sheals, K., Tombor, I., McNeill, A., & Shahab, L. (2016). A mixed-method systematic review and meta-analysis of mental health professionals' attitudes toward smoking and smoking cessation among people with mental illnesses. *Addiction (Abingdon, England)*, 111(9), 1536-1553. <https://doi.org/10.1111/add.13387>
- Skelton, E., Tzelepis, F., Shakeshaft, A., Guillaumier, A., Dunlop, A., McCrabb, S., Palazzi, K., & Bonevski, B. (2017). Addressing tobacco in Australian alcohol and other drug treatment settings: a cross-sectional survey of staff attitudes and perceived barriers. *Substance abuse treatment, prevention, and policy*, 12(1), 20-20. <https://doi.org/10.1186/s13011-017-0106-5>
- Smith, P. H., Mazure, C. M., & McKee, S. A. (2014). Smoking and mental illness in the U.S. population. *Tob Control*, 23(e2), e147-153. <https://doi.org/10.1136/tobaccocontrol-2013-051466>
- Ward, K. D., Kedia, S., Webb, L., & Relyea, G. E. (2012). Nicotine dependence among clients receiving publicly funded substance abuse treatment. *Drug and alcohol dependence*, 125(1-2), 95-102. <https://doi.org/10.1016/j.drugalcdep.2012.03.022>
- Warner, K. E. (2006). Tobacco Policy Research: Insights and Contributions to Public Health Policy. In *Tobacco control policy*. (pp. 3-86). Jossey-Bass/Wiley.
- Weinberger, A. H., Funk, A. P., & Goodwin, R. D. (2016). A review of epidemiologic research on smoking behavior among persons with alcohol and illicit substance use disorders. *Prev Med*, 92, 148-159. <https://doi.org/10.1016/j.ypmed.2016.05.011>
- Weinberger, A. H., Gbedemah, M., Wall, M. M., Hasin, D. S., Zvolensky, M. J., & Goodwin, R. D. (2018). Cigarette use is increasing among people with illicit substance use disorders in the United States, 2002-14: emerging disparities in vulnerable populations. *Addiction (Abingdon, England)*, 113(4), 719-728. <https://doi.org/10.1111/add.14082>
- Weinberger, A. H., Platt, J., Esan, H., Galea, S., Erlich, D., & Goodwin, R. D. (2017). Cigarette Smoking Is Associated With Increased Risk of Substance Use Disorder Relapse: A Nationally Representative, Prospective Longitudinal Investigation. *J Clin Psychiatry*, 78(2), e152-e160. <https://doi.org/10.4088/JCP.15m10062>