

UC Berkeley

UC Berkeley Previously Published Works

Title

Optimizing Trap Characteristics to Monitor the Leafhopper Bug *Leptoglossus zonatus* (Heteroptera: Coreidae) in Orchards.

Permalink

<https://escholarship.org/uc/item/53k9r31n>

Journal

Insects, 11(6)

ISSN

2075-4450

Authors

Wilson, Houston

Maccaro, Jessica

Daane, Kent

Publication Date

2020-06-09

DOI

10.3390/insects11060358

Peer reviewed

Article

Optimizing Trap Characteristics to Monitor the Leaf-footed Bug *Leptoglossus zonatus* (Heteroptera: Coreidae) in Orchards

Houston Wilson ^{1,*} , Jessica J. Maccaro ¹ and Kent M. Daane ² 

¹ Department of Entomology, University of California, Riverside, Riverside, CA 92521, USA; jessica.maccaro@ucr.edu

² Department of Environmental Science, Policy and Management, University of California, Berkeley, Berkeley, CA 94720-3114, USA; kdaane@ucanr.edu

* Correspondence: houston.wilson@ucr.edu; Tel.: +001-(559)-646-6519

Received: 6 May 2020; Accepted: 6 June 2020; Published: 9 June 2020



Abstract: The leaf-footed bug, *Leptoglossus zonatus* (Heteroptera: Coreidae), has become a key pest of almonds, pistachios, and pomegranates in California. Adults and nymphs directly feed on nuts and fruits, which reduces crop yield and quality and can facilitate pathogen infections. Current monitoring strategies require growers to actively sample the tree canopy, with no economic thresholds being developed for this pest. To improve monitoring of *L. zonatus*, a three-year study was conducted to identify an optimal trap. A hanging cross-vane panel trap was identified as the best trap type in Year 1, and subsequent work in Years 1–3 focused on refining its use by modifying surface texture and color. Results indicated that coating trap surfaces with the lubricant fluon improved trap catching ability, and adults were most frequently recovered in yellow traps. A hanging cross-vane panel trap with these features could serve as the basis for the development of a new monitoring system for this pest in orchards, which could be improved further if semiochemical lures will be developed.

Keywords: leaf-footed bug; cross-vane panel trap; trap color; *Leptoglossus*

1. Introduction

The genus *Leptoglossus* is neotropical in origin, with most species limited to Central and South America [1,2], where they are considered pests for a wide variety of crops [3–8]. In North America, the key economic species are *L. clypealis*, *L. occidentalis*, *L. phyllopus*, and *L. zonatus*. *Leptoglossus occidentalis* primarily attack coniferous trees [9,10], whereas the other three species have been recovered from a range of perennial crops that include tree nuts, citrus, peaches, and pomegranates [11–14]. The primary agricultural pest species in California are *L. clypealis* and *L. zonatus*, which are known to attack almonds, pistachios, and pomegranates when not feeding on a variety of weedy annual species [15]. Early reports first documented *L. clypealis* feeding on pistachio, which led to nut drop and epicarp lesion [16,17]. While historically *L. clypealis* has been the dominant species found on California tree nuts [18], recent surveys have noted a shift towards *L. zonatus*, which is now considered the primary species attacking these crops [15,19,20].

These species of *Leptoglossus* overwinter as adults in aggregations in sheltered areas, such as evergreen trees, shrubs, and residential structures, and in the spring disperse in search of food and reproduction sites [10,14,15,21]. In California, *L. clypealis* and *L. zonatus* complete three generations, but a fourth generation is possible if quality food sources and mild fall-winter temperatures are present [19]. While there are many plant species these leaf-footed bugs will feed on, adults typically begin to attack almonds in April–May. As almond shells harden, adults (either from the overwintering or first summer generation) move over to pistachios in May–June, which are still vulnerable during

this period. Similarly, as pistachio shells harden, the later summer generation(s) of adult leaffooted bug shift to pomegranates in August–September, and typically complete a generation while residing in this crop until late November, at which point adults move to overwintering aggregation sites. While the most extensive crop damage is due to adults, who have strong mouthparts and can disperse widely, feeding by developing nymphs can also have an impact on crop yield and quality.

Current recommendations for monitoring leaffooted bugs include beat sampling the tree canopy, visually searching trees for adults, or assessing immature nuts for signs of feeding damage [15,21]. All of these approaches are very time and labor intensive, and no economic thresholds associated with any of these sampling methods exist. Furthermore, damaging populations of *L. zonatus* adults can be sporadic and tend to arrive rapidly, which makes them difficult to predict and/or detect in a timely manner. As such, use of chemical controls is typically based on presence/absence of *Leptoglossus* spp. and largely rely on pyrethroids, like bifenthrin and lambda-cyhalothrin. Design of an effective trap and lure system for monitoring *L. zonatus* could improve sampling efficiency and facilitate the development of economic thresholds. Here, as a first step towards developing this type of system, different trap types were screened for their ability to catch *L. zonatus* under field conditions, with parallel optimization of several trap parameters.

2. Materials and Methods

A series of field studies were carried out over a three-year period in the San Joaquin Valley, California. In an initial study in Year 1, the efficacy of different trap types was compared, in order to identify the most promising one. The subsequent efforts in Years 1–3 focused on refining the use of the most efficient trap, by modifying trap color and surface coating. Working with unbaited traps required that all studies took place at sites with heavy infestations in the late summer and fall when *L. zonatus* populations are highest.

2.1. Trap Comparison Study

Five trap types and three bait treatments were evaluated at three field sites that included an olive, pistachio, and pomegranate orchard, each heavily infested with *L. zonatus*. The olive orchard was 2.8 ha (7 ac), the pistachio orchard was 0.8 ha (2 ac), and the pomegranate orchard was arranged as a hedge that was 4.6 m (15 ft) wide by 419.4 m (1376 ft) long.

Trap types included a black hanging cross-vane panel trap (Intercept Trap, Alpha Scents, West Linn, OR, USA), a 1.2 m (4 ft) tall black pyramid trap (Dead Inn, AgBio Inc., Denver, CO, USA), a 0.6 m (2 ft) tall black pyramid trap (Dead Inn, AgBio Inc., Denver, CO, USA), a 15.2 × 30.5 cm (6 × 12 in) clear sticky dual panel trap (Pherocon, Trécé Inc., Adair, OK, USA), and a green-white-yellow bucket trap (Multi-color UNI-Trap, Alpha Scents, West Linn, OR, USA). The collection buckets of the hanging cross-vane panel traps and bucket traps contained 443 mL (15 oz) of a killing solution, which consisted of 9.9 mL (2 tsp) biodegradable detergent diluted in 3.8 L (1 gal) of water.

Trap baits included split pomegranate (50 g), almond meal (40 g) mixed with crude almond oil (10 g), and a no bait control. Baits were placed in 10.2 × 15.2 cm (4 × 6 in) organandy mesh bags. These bait bags were then suspended from the center of the hanging panel trap, placed inside the collection cup at the top of the pyramid traps, attached with a binder clip to the clear sticky trap, or suspended in the collection cup inside the bucket trap. In total there were 15 unique trap × bait treatment combinations.

At each of the three sites, a randomized complete block design was used with five replicates per site. Replicates were spaced 36.6 m, 18.3 m, and 10 m apart and within replicates the different trap/bait treatments were spaced 9.1 m, 5.2 m, and 33 m apart at the olive, pistachio, and pomegranate sites, respectively. Traps were set up and monitored weekly between 7 September–18 October in Year 1. Each week the baits and collecting solution were replaced and all adult *Leptoglossus* spp. were removed, sexed, and identified to species.

2.2. Surface Coating Study

In a follow-up study, the use of fluon (Insect-a-Slip, BioQuip, Rancho Dominguez, CA, USA) was evaluated to improve recovery of *Leptoglossus* spp. in the black hanging cross-vane panel traps. There were five treatments that included a coating of undiluted fluon; dilutions of 50%, 25%, and 12.5% fluon; and a control trap with no fluon. In each fluon treatment, a single layer of solution was painted over all trap surfaces. No baits were used with any of the traps. Traps were evaluated using a randomized complete block design with five replicates at the pomegranate site described above. Traps were set up and monitored weekly from 13 November–4 December in Year 1. As before, each week, the collecting solution was replaced and all adult *Leptoglossus* spp. were removed, sexed, and identified to species.

2.3. Trap Color

The effect of the hanging cross-vane panel trap color was evaluated each fall in a heavily infested pomegranate orchard over a three-year period. Black traps were compared to red, yellow, green, blue, and white traps. Field studies utilized a randomized complete block design with five replicates. In Years 1–2, this study was located at the pomegranate site described above and took place during 4–18 December, in Year 1, and 22 August–20 November, in Year 2. In Year 3, the study took place 27 September–3 December in a 0.4 ha (1 ac) pomegranate orchard at the UC Kearney Agricultural Research and Extension Center, Parlier, California, USA. Replicates were spaced 10 m and 10.3 m apart and traps within replicates were spaced 3 m and 9.1 m apart, in Years 1–2 and 3, respectively.

2.4. Statistical Analysis

Data on insect abundance from each trial were $\log(x + 1)$ transformed and analyzed with generalized linear mixed-models using the “glmer” function in the “lme4” package in the R statistical program (<http://www.r-project.org/>). Fixed effects were evaluated through model comparison using chi-square tests via the “drop1” function. When a multilevel categorical variable was found to be significant, means were separated using post hoc Tukey contrasts (“glht” function in the “multcomp” package). Male and female *L. zonatus* were evaluated separately.

For the trap/bait study, fixed effects included “Trap Type” and “Bait,” with the random effects “Replicate Block” nested within “Site” within “Sample Week.” Replicate Block was included as a random effect, since each block contained multiple repeats of the different trap types and baits. An interaction term for “Trap Type \times Bait” was initially evaluated, but since it was found to be non-significant, the analysis presented here models these factors separately. The fluon study included fixed effect “Fluon Dilution” and the random effect “Sample Week.” Finally, the trap color study included fixed effect “Trap Color,” with random effects “Site” nested within “Sample Week” within “Year.”

3. Results

Almost all the *Leptoglossus* spp. recovered in these trials were *L. zonatus*, with *L. clypealis* rarely encountered. As such, all of the data analyses focused on *L. zonatus* alone. Analysis of the trap and bait trial data ($n = 400$, total *L. zonatus* females = 103, males = 114) indicated that capture of *L. zonatus* was influenced by trap type (males $\chi^2 = 106.8$, $p < 0.001$; females $\chi^2 = 101.0$, $p < 0.001$), but not by bait (males $\chi^2 = 0.2$, $p = 0.91$; females $\chi^2 = 0.2$, $p = 0.92$). The hanging cross-vane panel trap captured the most *L. zonatus* adults (Figure 1). In the surface treatment trial ($n = 75$, total *L. zonatus* females = 277, males = 213), *L. zonatus* catch was increased substantially by coating trap surfaces with fluon (males $\chi^2 = 33.6$, $p < 0.001$; females $\chi^2 = 20.4$, $p < 0.001$), even when highly diluted (Figure 2). Finally, data from the multi-year trap color experiment ($n = 568$, total *L. zonatus* females = 342, males = 201) indicated differences in *L. zonatus* catch across the different trap colors (males $\chi^2 = 40.6$, $p < 0.001$; females $\chi^2 = 70.0$, $p < 0.001$). Yellow traps were the most attractive, followed by blue and green traps, while white and red traps were the least attractive (Figure 3).




Figure 1. The hanging cross-vane panel trap consistently captured the most *L. zonatus*, regardless of bait type. Columns of the same color without a common letter differ significantly. Alm. = almond meal bait, Cont. = no bait control, and Pom. = pomegranate bait; Bucket Trap = green-white-yellow bucket trap, Hanging Panel = black hanging cross-vane panel trap, Pyramid - Short = 0.6 m pyramid trap, Pyramid - Tall = 1.2 m pyramid trap, and Clear Sticky Trap = clear sticky dual panel trap.




Figure 2. The addition of fluon increased trap capture of *L. zonatus*. Columns of the same color without a common letter differ significantly.




Figure 3. Yellow traps captured the most *L. zonatus*, followed by blue and green traps. X-axis defines the color of trap evaluated. Columns of the same color without a common letter differ significantly.

4. Discussion

This series of experiments demonstrated that an unbaited hanging cross-vane panel trap is attractive to *L. zonatus*, especially a yellow trap. Capture can be increased by coating trap surfaces with fluon to reduce insect ability to grip the trap surface. Inclusion of a split pomegranate or almond meal bait did not increase trap catch, indicating that these materials have low attractive ability on this insect species, and that *L. zonatus* apparently are attracted to the hanging cross-vane panel trap itself. Response of *L. zonatus* to a large dark object, such as this panel trap, is similar to observations by Panizzi [22], who reported that *L. zonatus* aggregated on unbaited plastic cylinder traps when the cylinders were first introduced into corn fields.

The cross-vane panel trap was initially developed to monitor Coleoptera in forests [23,24], as a replacement or supplement for the multiple-funnel trap [25]. Subsequent work with the cross-vane panel trap demonstrated its utility for trapping cerambycid and buprestid beetles [26–28]. Trapping efficacy for beetle species can be further enhanced, sometimes more than ten-fold, by coating trap surfaces with fluon or other lubricants [29–31]. Our results demonstrated analogous increases in trapping efficiency when traps were treated with fluon.

Cross-vane panel traps have also been used to effectively trap Hemipterans, such as triatomines (Reduviidae: Triatominae) [32], bagrada bug (Pentatomidae: *Bagrada hilaris*) [33], and brown marmorated stink bug (Pentatomidae: *Halyomorpha halys*) [34], but apparently are not well suited for spotted lanternfly (Fulgoridae: *Lycorma delicatula*) [35]. No traps specifically for *Leptoglossus* spp. have been developed. The scant literature includes a prototype bottle trap for *L. zonatus* [36], along with a study that mentions the use of multi-funnel traps for testing *L. occidentalis* attraction to caged aggregations of males and females [37]. Bycatch of non-target organisms in the cross-vane panel trap was minimal, with some honeybees (Apidae: *Apis mellifera*) recovered in blue and green traps and assassin bugs (Reduviidae: *Zelus* spp.) in blue and yellow traps (data not shown).

The effect of trap color has been widely evaluated across multiple insect orders, including Coleoptera [38–40], Lepidoptera [41–43], Hymenoptera [44–47], and Hemiptera [48–52]. The only previous study to evaluate *Leptoglossus* spp. responses to color was a laboratory study, which found that *L. zonatus* adults and nymphs were primarily attracted to blue and green [53]. These results were partially complimented in our study, in which blue and green were the second and third most attractive colors after yellow. Response of *L. zonatus* to trap color in the current study may have been skewed by differences in trap temperature, which likely varied between the lighter and darker colored traps under field conditions. This may be an important consideration, given that a recent study of *L. occidentalis*,

a relative species of *L. zonatus*, indicated that this species utilizes infrared cues to locate host plants [54]. It may be that *L. zonatus* utilizes infrared cues in a similar way. As such, subsequent field evaluations of trap color effects on *L. zonatus* could be improved by controlling for, or at least recording data on, trap temperature.

5. Conclusions

Identification of the hanging cross-vane panel trap for *L. zonatus* represents an important step forward in the development of an effective monitoring program for this pest, especially the highly mobile adult, and will allow for the future screening of candidate lures for this insect under field conditions [55]. Once paired with an attractive lure, studies will need to determine the most effective density and spatial arrangement of traps to accurately reflect orchard populations of *L. zonatus* and correlate trap captures with economic thresholds. If perfected, this type of trap and lure sampling system will reduce the sampling effort and lead to earlier and/or more accurate detection of *L. zonatus* populations in orchards. It is also possible that these traps could work well for other *Leptoglossus* pest species, such as *L. occidentalis*, which has recently invaded Europe, and may be worth further investigation in that context [56,57].

Author Contributions: Conceptualization, H.W. and K.M.D.; methodology, H.W. and J.J.M.; investigation, H.W. and J.J.M.; formal analysis, H.W.; writing—original draft preparation, H.W.; writing—review and editing, H.W., J.J.M. and K.M.D. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the California Pistachio Research Board and the Almond Board of California.

Acknowledgments: Authors thank the California Pistachio Research Board and the Almond Board of California for funding this research. Field studies and data collection were carried out with assistance from German Camacho, Jesus Ceja, Gabrielle Celaya, Tyler Colombero, Dani Evans, Javier Herrera, Garrett Morales, Joshua Reger, Austin Souza, May Yang, and Sunny Yang. Many thanks are owed as well to all collaborating growers for access to field sites to collect insects. Authors also extend thanks to Jocelyn Millar for providing useful comments and feedback on a draft version of this manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Allen, R.C. A revision of the genus *Leptoglossus* Guérin (Hemiptera: Coreidae). *Entomol. Am.* **1969**, *45*, 35–140.
- Brailovsky, H. Illustrated key for identification of the species included in the genus *Leptoglossus* (Hemiptera: Heteroptera: Coreidae: Coreinae: Anisoscelini), and descriptions of five new species and new synonyms. *Zootaxa* **2014**, *3794*, 143–178. [[CrossRef](#)] [[PubMed](#)]
- Panizzi, A.R. Desempenho de ninfas e adultos de *Leptoglossus zonatus* (Dallas, 1852) (Hemiptera: Coreidae) em diferentes alimentos. *An. da Soc. Entomológica do Bras.* **1989**, *18*, 375–389.
- Kubo, R.; Batista, A. Ocorrência e danos provocados por *Leptoglossus zonatus* (Dallas, 1852) (Hemiptera: Coreidae) em citros. *An. da Soc. Entomol. do Bras.* **1992**, *21*, 467–470.
- Rodriguez, F.C.Y.; Carmona, M.A.L.; Zuluaga, N.C.; Gamboa, J.A.Q. Plagas potenciales del cultivo de *Jatropha curcas* L., en el occidente de Antioquia, Colombia. *Rev. Fac. Nac. Agron. Medellin.* **2012**, *65*, 6823–6826.
- Raga, A.; Piza, C.; de Souza, F. Occurrence and damage of *Leptoglossus zonatus* (Dallas) (Heteroptera: Coreidae) on pomegranate, *Punica granatum* L., in Campinas, Sao Paulo. *An. Soc. Entomol. do Bras.* **1995**, *24*, 183.
- Rodrigues Netto, S.M.; Guilhem, D.J. Ocorrência de *Leptoglossus zonatus* (Dallas, 1852) (Hemiptera: Coreidae) em maracujazeiro (*Passiflora edulis* f. *flavicarpa*). *Arq. Inst. Biol. (Sao Paulo)* **1996**, *63*, 85–86.
- Pires, E.M.; Bonaldo, S.M.; Ferreira, J.A.M.; Soares, M.A.; Candan, S. New record of *Leptoglossus zonatus* (Dallas) (Heteroptera: Coreidae) attacking starfruit (*Averrhoa carambola* L.) in Sinop, Mato Grosso, Brazil. *EntomoBrasilis.* **2011**, *4*, 33–35. [[CrossRef](#)]
- Koerber, T.W. *Leptoglossus occidentalis* (Hemiptera, Coreidae), a newly discovered pest of coniferous seed. *Ann. Entomol. Soc. Am.* **1963**, *56*, 229–234. [[CrossRef](#)]
- Hedlin, A.F.; Yates III, H.O.; Tovar, D.C.; Ebel, B.H.; Koerber, T.W.; Merkel, E.P. *Cone and Seed Insects of North American Conifers*; United States Forest Service: Washington, DC, USA, 1980.

11. Burgess, P.S. *55th Annual Report for the Year Ending June 30, 1944*; University of Arizona Agricultural Experiment Station: Tuscon, AZ, USA, 1955.
12. Schaefer, C.W.; Mitchell, P.L. Foodplants of the Coreoidea (Hemiptera: Heteroptera). *Ann. Entomol. Soc. Am.* **1983**, *76*, 591–615. [[CrossRef](#)]
13. Tarango Rivera, S.H.; Banuelos, M.L.G.; Plata, M.C.C. Efecto de la alimenta de cinco especies de chinches (Hemiptera: Pentatomidae: Coreidae) en frutos de nogal pecanero. *Agric. Tec. Mex.* **2007**, *33*, 241–249.
14. Xiao, Y.; Fadamiro, H.Y. Host preference and development of *Leptoglossus zonatus* (Hemiptera: Coreidae) on Satsuma mandarin. *J. Econ. Entomol.* **2009**, *102*, 1908–1914. [[CrossRef](#)] [[PubMed](#)]
15. Daane, K.M.; Yokota, G.Y.; Bentley, W.J.; Weinberger, G.B.; Millar, J.G.; Beede, R.H. Stink bugs and leaf-footed bugs. In *Pistachio Production Manual*; Ferguson, L., Haviland, D.R., Eds.; University of California Agriculture and Natural Resources Publication 3545: Oakland, CA, USA, 2016; pp. 225–238.
16. Bolkan, H.A.; Ogawa, J.M.; Rice, R.; Bostock, R.M.; Crane, J.C. Leaf-footed bug implicated in pistachio epicarp lesion. *Calif. Agric.* **1984**, *38*, 16–17.
17. Michailides, T.J.; Rice, R.E.; Ogawa, J.M. Succession and significance of several hemipterans attacking a pistachio orchard. *J. Econ. Entomol.* **1987**, *80*, 398–406. [[CrossRef](#)]
18. Daane, K.; Yokota, G.; Krugner, R.; Steffan, S.; da Silva, P.; Beede, R.; Bentley, W.; Weinberger, G. Large bugs damage pistachio nuts most severely during midseason. *Calif. Agric.* **2005**, *59*, 95–102. [[CrossRef](#)]
19. Daane, K.M.; Yokota, G.Y.; Wilson, H. Seasonal dynamics of the leaf-footed bug *Leptoglossus zonatus* and its implications for control in almonds and pistachios. *Insects* **2019**, *10*, 255. [[CrossRef](#)]
20. Joyce, A.L.; Higbee, B.S.; Haviland, D.R.; Brailovsky, H. Genetic variability of two leaf-footed bugs, *Leptoglossus clypealis* and *Leptoglossus zonatus* (Hemiptera: Coreidae) in the Central Valley of California. *J. Econ. Entomol.* **2017**, *110*, 2576–2589. [[CrossRef](#)]
21. Zalom, F.G.; Haviland, D.R.; Symmes, E.S.; Tollerup, K.E. Leaf-footed Bug. In *UC IPM Almond Pest Management Guidelines*; University of California Agriculture and Natural Resources, Publication #3431: Oakland, CA, USA, 2017.
22. Panizzi, A.R. A possible territorial or recognition behavior of *Leptoglossus zonatus* (Dallas) (Heteroptera: Coreidae). *Rev. Bras. Entomol.* **2004**, *48*, 577–579. [[CrossRef](#)]
23. Czokajlo, D.; Ross, D.; Kirsch, P. Intercept panel trap, a novel trap for monitoring forest Coleoptera. *J. For. Sci.* **2001**, *47*, 63–65.
24. McIntosh, L.R.; Katinic, P.J.; Allison, J.D.; Borden, J.H.; Downey, D.L. Comparative efficacy of five types of trap for woodborers in the Cerambycidae, Buprestidae and Siricidae. *Agric. For. Entomol.* **2001**, *3*, 113–120. [[CrossRef](#)]
25. Lindgren, B.S. A multiple funnel trap for scolytid beetles (Coleoptera). *Can. Entomol.* **1983**, *115*, 299–302. [[CrossRef](#)]
26. De Groot, P.; Nott, R. Evaluation of traps of six different designs to capture pine sawyer beetles (Coleoptera: Cerambycidae). *Agric. For. Entomol.* **2001**, *3*, 107–111. [[CrossRef](#)]
27. Miller, R.D.; Crowe, C.M. Relative performance of Lindgren multiple-funnel, intercept panel, and colossus pipe traps in catching Cerambycidae and associated species in the Southeastern United States. *J. Econ. Entomol.* **2011**, *104*, 1934–1941. [[CrossRef](#)] [[PubMed](#)]
28. Graham, E.E.; Poland, T.M.; McCullough, D.G.; Millar, J.G. A comparison of trap type and height for capturing cerambycid beetles (Coleoptera). *J. Econ. Entomol.* **2012**, *105*, 837–846. [[CrossRef](#)] [[PubMed](#)]
29. Graham, E.E.; Mitchell, R.F.; Reagel, P.F.; Barbour, J.D.; Millar, J.G.; Hanks, L.M. Treating panel traps with a fluoropolymer enhances their efficiency in capturing cerambycid beetles. *J. Econ. Entomol.* **2010**, *103*, 641–647. [[CrossRef](#)] [[PubMed](#)]
30. Graham, E.E.; Poland, T.M. Efficacy of fluon conditioning for capturing cerambycid beetles in different trap designs and persistence on panel traps over time. *J. Econ. Entomol.* **2012**, *105*, 395–401. [[CrossRef](#)]
31. Allison, J.D.; Graham, E.E.; Poland, T.M.; Strom, B.L. Dilution of fluon before trap surface treatment has no effect on longhorned beetle (Coleoptera: Cerambycidae) captures. *J. Econ. Entomol.* **2016**, *109*, 1215–1219. [[CrossRef](#)]
32. Updyke, A.E.; Allan, B.F. An experimental evaluation of cross-vane panel traps for the collection of sylvatic Triatomines (Hemiptera: Reduviidae). *J. Med. Entomol.* **2018**, *55*, 485–489. [[CrossRef](#)]
33. Joseph, S.V. Effect of trap color on captures of bagrada bug, *Bagrada hilaris* (Hemiptera: Pentatomidae). *J. Entomol. Sci.* **2014**, *49*, 318–321. [[CrossRef](#)]

34. Chase, K.D.; Stringer, L.D.; Butler, R.C.; Liebhold, A.M.; Miller, D.R.; Shearer, P.W.; Brockerhoff, E.G. Multiple-lure surveillance trapping for *Ips* bark beetles, *Monochamus* longhorn beetles, and *Halyomorpha halys* (Hemiptera: Pentatomidae). *J. Econ. Entomol.* **2018**, *111*, 2255–2263. [[CrossRef](#)]
35. Francese, J.A.; Cooperband, M.F.; Murman, K.M.; Cannon, S.L.; Booth, E.G.; Devine, S.M.; Wallace, M.S. Developing traps for the spotted lanternfly, *Lycorma delicatula* (Hemiptera: Fulgoridae). *Environ. Entomol.* **2020**, *49*, 269–276. [[CrossRef](#)] [[PubMed](#)]
36. Barreto, M.R.; da Silva, L.G. Eficiência da armadilha “R. Bianco” para captura do percevejo *Leptoglossus zonatus* Dallas (Hemiptera: Coreidae), na cultura do milho. *EntomoBrasilis* **2016**, *9*, 84–88. [[CrossRef](#)]
37. Blatt, E.S.; Borden, J.H. Evidence for a male-produced aggregation pheromone in the western conifer seed bug, *Leptoglossus occidentalis* Heidemann (Hemiptera: Coreidae). *Can. Entomol.* **1996**, *128*, 777–778. [[CrossRef](#)]
38. Hesler, L.S.; Sutter, G.R. Effect of trap color, volatile attractants, and type of toxic bait dispenser on captures of adult corn rootworm beetles (Coleoptera: Chrysomelidae). *Environ. Entomol.* **1993**, *22*, 743–750. [[CrossRef](#)]
39. Francese, J.A.; Crook, D.J.; Fraser, I.; Lance, D.R.; Sawyer, A.J.; Mastro, V.C. Optimization of trap color for emerald ash borer (Coleoptera: Buprestidae). *J. Econ. Entomol.* **2010**, *103*, 1235–1241. [[CrossRef](#)]
40. Kim, S.H.S.; Trammel, E.C.; Lewis, B.A.; Johnson, D.T. Comparison of color attractiveness for *Agrilus ruficollis* (Coleoptera: Buprestidae): Potential for a simple trap. *J. Econ. Entomol.* **2016**, *109*, 1799–1806. [[CrossRef](#)]
41. Mitchell, E.R.; Agee, H.R.; Heath, R.R. Influence of pheromone trap color and design on capture of male velvetbean caterpillar and fall armyworm moths (Lepidoptera: Noctuidae). *J. Chem. Ecol.* **1989**, *15*, 1775–1784. [[CrossRef](#)]
42. Knight, A.L.; Fisher, A.J. Increased catch of codling moth (Lepidoptera: Tortricidae) in semiochemical-baited orange plastic delta-shaped traps. *Environ. Entomol.* **2006**, *35*, 1597–1602. [[CrossRef](#)]
43. Athanassiou, G.C.; Kavallieratos, N.G.; Mazomenos, B.E. Effect of trap type, trap color, trapping location, and pheromone dispenser on captures of male *Palpita unionalis* (Lepidoptera: Pyralidae). *J. Econ. Entomol.* **2009**, *97*, 321–329. [[CrossRef](#)]
44. Trimble, M.R.; Brach, E.J. Effect of color on sticky-trap catches of *Pholetesor ornigis* (Hymenoptera: Braconidae), a parasite of the spotted tentiform leafminer *Phyllonorycter blancardella* (Lepidoptera: Gracillariidae). *Can. Entomol.* **1985**, *117*, 1559–1564. [[CrossRef](#)]
45. Stephen, P.W.; Rao, S. Unscented color traps for non-*Apis* Bees (Hymenoptera: Apiformes). *J. Kansas Entomol. Soc.* **2005**, *78*, 373–380. [[CrossRef](#)]
46. Toler, R.T.; Evans, E.W.; Tepedino, V.J. Pan-trapping for bees (Hymenoptera: Apiformes) in Utah’s West Desert: The importance of color diversity. *Pan-Pac. Entomol.* **2005**, *81*, 103–113.
47. Beers, E.H. Effect of trap color and orientation on the capture of *Aphelinus mali* (Hymenoptera: Aphelinidae), a parasitoid of woolly apple aphid (Hemiptera: Aphididae). *J. Econ. Entomol.* **2012**, *105*, 1342–1349. [[CrossRef](#)] [[PubMed](#)]
48. Landis, J.B.; Fox, L. Lygus bugs in eastern Washington: Color preferences and winter activity. *Environ. Entomol.* **1972**, *1*, 464–465. [[CrossRef](#)]
49. Chum, -C.C.; Pinter, P.J.; Henneberry, T.J.; Umeda, K.; Natwick, E.T.; Wei, Y.-A.; Reddy, V.R.; Shrepatis, M. Use of CC traps with different trap base colors for silverleaf whiteflies (Homoptera: Aleyrodidae), thrips (Thysanoptera: Thripidae), and leafhoppers (Homoptera: Cicadellidae). *J. Econ. Entomol.* **2009**, *93*, 1329–1337.
50. Leskey, C.T.; Wright, S.E.; Short, B.D.; Khrimian, A. Development of behaviorally-based monitoring tools for the brown marmorated stink bug (Heteroptera: Pentatomidae) in commercial tree fruit orchards. *J. Entomol. Sci.* **2012**, *47*, 76–85. [[CrossRef](#)]
51. Rodriguez-Saona, R.C.; Byers, J.A.; Schiffhauer, D. Effect of trap color and height on captures of blunt-nosed and sharp-nosed leafhoppers (Hemiptera: Cicadellidae) and non-target arthropods in cranberry bogs. *Crop Prot.* **2012**, *40*, 132–144. [[CrossRef](#)]
52. Dimeglio, S.A.; Kuhar, T.P.; Weber, D.C. Color preference of harlequin bug (Heteroptera: Pentatomidae). *J. Econ. Entomol.* **2017**, *110*, 2275–2277. [[CrossRef](#)]
53. Franco-Archundia, L.S.; Gonzaga-Segura, A.J.; Jimenez-Perez, A.; Castejon-Gomez, V.R. Behavioral response of *Leptoglossus zonatus* (Heteroptera: Coreidae) to stimuli based on colors and its aggregation pheromone. *Insects* **2018**, *9*, 91. [[CrossRef](#)]
54. Takács, S.; Bottomley, H.; Andreller, I.; Zaradnik, T.; Schwarz, J.; Bennett, R.; Strong, W.; Gries, G. Infrared radiation from hot cones on cool conifers attracts seed-feeding insects. *Proc. R. Soc. B Biol. Sci.* **2009**, *276*, 649–655. [[CrossRef](#)]

55. Beck, J.J.; Gee, W.S.; Cheng, L.W.; Higbee, B.S.; Wilson, H.; Daane, K.M. Investigating host plant-based semiochemicals for attracting the leaffooted bug (Hemiptera: Coreidae), an insect pest of California agriculture. In *Roles of Natural Products and Biorational Pesticides in Agriculture*; Beck, J.J., Rering, C.C., Duke, S.O., Eds.; ACS Symposium Series; American Chemical Society: Washington, DC, USA, 2018; pp. 143–165.
56. Lesieur, V.; Lombaert, E.; Guillemaud, T.; Courtial, B.; Strong, W.; Roques, A.; Auger-Rozenburg, M.-A. The rapid spread of *Leptoglossus occidentalis* in Europe: A bridgehead invasion. *J. Pest Sci.* **2019**, *92*, 189–200. [[CrossRef](#)]
57. Lis, J.A.; Lis, B.; Gubernator, J. Will the invasive western conifer seed bug *Leptoglossus occidentalis* Heidemann (Hemiptera: Heteroptera: Coreidae) seize all of Europe? *Zootaxa* **2008**, *17410*, 66–68. [[CrossRef](#)]



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).