

UC Irvine

UC Irvine Previously Published Works

Title

Diagnosis of hemochromatosis.

Permalink

<https://escholarship.org/uc/item/5432m1wt>

Journal

Annals of Internal Medicine, 131(4)

ISSN

1056-8751

Authors

Kalantar-Zadeh, Kamyar

Luft, Friedrich C

Publication Date

1999-08-17

DOI

10.7326/0003-4819-131-4-199908170-00018

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

disease (ESRD), the relation between tissue iron stores and serum ferritin levels is altered. A relative increase in serum ferritin levels unrelated to iron stores or acute phase reactants is frequently observed (2, 3). Serum ferritin levels of 1000 $\mu\text{g/L}$ or greater in patients with ESRD are not uncommon, and iron overload is extremely rare (3, 4). In one of our dialysis units, the mean serum ferritin level is 630 $\mu\text{g/L}$, with a median value of 601 (SD, 358 $\mu\text{g/L}$). By using bone marrow iron stores, we showed that a serum ferritin level less than 200 $\mu\text{g/L}$ is highly specific for iron deficiency in patients with ESRD (2). Moreover, serum ferritin levels as high as 1200 $\mu\text{g/L}$ may still be consistent with low to normal iron stores in patients with ESRD.

In a study on nutritional assessment (5), we showed that severely malnourished patients undergoing dialysis had the highest serum ferritin levels, and well-nourished patients undergoing dialysis had lower values. We also found an inverse correlation between serum ferritin and transferrin levels ($r = -0.61$) in patients with ESRD and showed that low serum transferrin levels due to malnutrition may cause an erroneously normal to high transferrin saturation ratio, even in the presence of iron deficiency (2, 5). We suggest that the hemochromatosis criteria be modified for patients with ESRD.

Kamyar Kalantar-Zadeh, MD, MPH
University of California, San Francisco
San Francisco, CA 94143

Friedrich C. Luft, MD
Humboldt University of Berlin
13122 Berlin, Germany

References

1. Powell LW, Georg K, McDonnell SM, Kowdley KV. Diagnosis of hemochromatosis. *Ann Intern Med.* 1998;129:925-31.
2. Kalantar-Zadeh K, Wunsch PH, Fink H, Höffken B, Kleiner M, Luft FC. Diagnosis of iron deficiency anemia in renal failure patients during post erythropoietin era. *Am J Kid Dis.* 1995;26:292-9.
3. Barany P, Eriksson LC, Hultcrantz R, Pettersson E, Bergstrom J. Serum ferritin and tissue iron in anemic dialysis patient. *Min Electrolyte Metab.* 1997;23:273-6.
4. NKF-DOQI clinical practice guidelines for the treatment of anemia of chronic renal failure. National Kidney Foundation-Dialysis Outcomes Quality Initiative. *Am J Kid Dis.* 1997;30(4 Suppl 3):S192-240.
5. Kalantar-Zadeh K, Kleiner M, Dunne E, Nelson M, Ahern K, Koslowe R, et al. Total iron binding capacity-estimated transferrin concentrations in dialysis patients correlate with the subjective global assessment of nutrition. *Am J Kid Dis.* 1998;31:263-72.

Diagnosis of Hemochromatosis

To the Editor: We take issue with the hemochromatosis laboratory criteria by Powell and colleagues (1). The authors concluded that a serum transferrin saturation greater than 45%, a serum ferritin level greater than 200 $\mu\text{g/L}$ in premenopausal women, and a serum ferritin level greater than 300 $\mu\text{g/L}$ in men and postmenopausal women are indicators for primary iron overload. Although the authors explain that elevated serum ferritin levels may be due to inflammation, infection, or cancer and suggest that acute-phase reactants be obtained to exclude these entities, they ignore the fact that in patients with end-stage renal