

# UC San Diego

## UC San Diego Previously Published Works

### Title

Revisiting the Rise of Electronic Nicotine Delivery Systems Using Search Query Surveillance

### Permalink

<https://escholarship.org/uc/item/54n6j73s>

### Journal

American Journal of Preventive Medicine, 50(6)

### ISSN

0749-3797

### Authors

Ayers, John W  
Althouse, Benjamin M  
Allem, Jon-Patrick  
et al.

### Publication Date

2016-06-01

### DOI

10.1016/j.amepre.2015.12.008

Peer reviewed



Published in final edited form as:

*Am J Prev Med.* 2016 June ; 50(6): e173–e181. doi:10.1016/j.amepre.2015.12.008.

## Revisiting the Rise of Electronic Nicotine Delivery Systems Using Search Query Surveillance

John W. Ayers, PhD, MA<sup>1</sup>, Benjamin M. Althouse, PhD, ScM<sup>2,3</sup>, Jon-Patrick Allem, PhD, MA<sup>4</sup>, Eric C. Leas, MPH<sup>5</sup>, Mark Dredze, PhD<sup>6</sup>, and Rebecca Williams, PhD, MHS<sup>7,8</sup>

<sup>1</sup>Graduate School of Public Health, San Diego State University, San Diego, California <sup>2</sup>The Santa Fe Institute, Santa Fe, New Mexico <sup>3</sup>Department of Biology, New Mexico State University, Las Cruces, New Mexico <sup>4</sup>Keck School of Medicine, University of Southern California, Los Angeles, California <sup>5</sup>University of California San Diego School of Medicine, San Diego, California <sup>6</sup>Johns Hopkins Human Language Technology Center of Excellence, Baltimore, Maryland <sup>7</sup>Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, North Carolina <sup>8</sup>Center for Health Promotion and Disease Prevention, University of North Carolina, Chapel Hill, North Carolina

### Abstract

**Introduction**—Public perceptions of electronic nicotine delivery systems (ENDS) remain poorly understood because surveys are too costly to regularly implement and when implemented there are large delays between data collection and dissemination. Search query surveillance has bridged some of these gaps. Herein, ENDS' popularity in the U.S. is reassessed using Google searches.

**Methods**—ENDS searches originating in the U.S. from January 2009 through January 2015 were disaggregated by terms focused on e-cigarette (e.g., *e-cig*) versus vaping (e.g., *vapers*), their geolocation (e.g., state), the aggregate tobacco control measures corresponding to their geolocation (e.g., clean indoor air laws), and by terms that indicated the searcher's potential interest (e.g., *buy e-cigs* likely indicates shopping); all analyzed in 2015.

**Results**—ENDS searches are increasing across the entire U.S., with 8,498,180 searches during 2014. At the same time, searches shifted from e-cigarette- to vaping-focused terms, especially in coastal states and states with more anti-smoking norms. For example, nationally, e-cigarette searches declined 9% (95% CI=1%, 16%) during 2014 compared with 2013, whereas vaping searches increased 136% (95% CI=97%, 186%), surpassing e-cigarette searches. More ENDS searches were related to shopping (e.g., *vape shop*) than health concerns (e.g., *vaping risks*) or cessation (e.g., *quit smoking with e-cigs*), with shopping searches nearly doubling during 2014.

**Conclusions**—ENDS popularity is rapidly growing and evolving, and monitoring searches has provided these timely insights. These findings may inform survey questionnaire development for follow-up investigation and immediately guide policy debates about how the public perceives ENDS' health risks or cessation benefits.

## Introduction

Electronic nicotine delivery systems (ENDS) are the first tobacco product born in the online age.<sup>1,2</sup> In 2011, Ayers and colleagues found that Google searches for ENDS in the U.S. were greater than searches for smoking alternatives or cessation devices (at a time when snus was garnering more academic and media attention).<sup>3</sup>

These findings have been confirmed and expanded on by telephone surveys that find awareness and use of ENDS is increasing.<sup>4–10</sup> Yet, ENDS surveillance remains limited. Surveys focused on ENDS are often too costly to regularly implement and their results are often not revealed until long after the data are first collected. As a result, public health is unable to stay on top of potential changes in public perceptions. For example, most studies have used sampling frames designed to obtain nationally representative cross-sectional estimates of ENDS use, but little is known about how perceptions and interest around ENDS varies sub-nationally or changes over time.<sup>11</sup>

Continued analysis of Google search trends may fill some of these knowledge gaps and outline agendas for follow-up survey-based surveillance.<sup>12–15</sup> Examining the content of searches can reveal the searcher's thoughts and potential motivation for searching,<sup>16</sup> such as seeking information for purchasing ENDS, whether ENDS aid cessation, or whether using ENDS poses any health risk. This study used exploratory analyses to assess variations in ENDS searches across states and time, including comparisons of searches across state-level tobacco control policies and social norms for cigarette smoking. Identifying ENDS search trends enhances the evidence base for the continued study of ENDS and their potential regulation.

## Methods

Weekly aggregated search query trends originating in the U.S. were analyzed from January 1, 2004 through January 1, 2015 using Google Trends ([google.com/trends](http://google.com/trends)). Google Trends is a public index of search activity for specific search terms or categories of terms, measuring the fraction of searches that include the terms (or categories) in question in a user-chosen geography at a particular time relative to the total number of searches at that time (relative search volume [RSV]). The RSVs from Google Trends were supplemented by raw search volume derived from Google Adwords' search volume estimator ([adwords.google.com](http://adwords.google.com)). Hundreds of studies have used Google Trends for public health insights,<sup>17</sup> including several recent examples from this journal.<sup>18–20</sup> Herein, all searches that included *ecig/s*, *e-cig/s*, *e cig/s*, *electronic cigarette/s*, *e cigarette/s*, or *e-cigarette/s* and *vape/s*, *vaper/s*, or *vaping* were monitored, after omitting searches that also mentioned *pot* or *weed*. For instance, this would include searches like *ecig*, *best ecig*, or *what are ecigs?*

Search query trends for the composite of all ENDS search terms were described nationally. Trends were then explored by disaggregating among those that included e-cigarette (e.g., *ecig*, *electronic cigs*, and *e-cigarettes*) and vaping (e.g., *vape*, *vaper*, and *vaping*) terms. Additionally, ENDS searches were compared to searches for snus, nicotine-replacement therapies, and Chantix, replicating methods detailed elsewhere.<sup>3</sup> All relied on trend analyses,

enumerating changes in search volumes year over year and making projections through 2015 based on autoregressive integrated moving average models fit using the stepwise algorithm outlined in Hyndman and Khandakar.<sup>21</sup> Mean comparisons were made using a regression approach with years as predictors and confidence bounds estimated by using 10,000 random draws from the multivariate normal sampling distribution with mean equal to the maximum-likelihood point estimates, and variance equal to the variance–covariance matrix.<sup>22</sup>

Geographic variability in searches was explored to describe the spread of ENDS. This relied on comparing ENDS search volume across the lower 48 states by year, formally done by using the maximum likelihood to estimate the change in SD over time ( $Likelihood = \sum_x \log(N(0, a + (b - a) * x))$ , where  $a$  and  $b$  are the intercept and slope of the SD and  $N$  is the normal density). This described the year by year variability in state-specific search volume. In addition, linear and quadratic models predicting ENDS search volumes using states' longitudes were fitted, based on visual inspection of the data. Analyses were replicated across all ENDS searches and ENDS searches were disaggregated by either e-cigarette or vaping terms.

Variations in searches by tobacco control policies and social norms against smoking (“anti-smoking norms”) were explored by comparing ENDS searches across the lower 48 states according to three state-level traits: clean indoor air grades from the American Lung Association updated to 2014,<sup>23</sup> cigarette excise tax rates updated to 2014,<sup>24</sup> and the anti-smoking norms of cigarette smoking as measured by survey responses.<sup>25</sup> In addition, ENDS searches were compared across the cigarette smoking prevalences of U.S. states as derived from the 2013 Behavioral Risk Factor Surveillance System. Models were executed by fitting bivariable analyses with each of the above as a predictor variables, using a linear function for cigarette excise tax rates, anti-smoking norms, and cigarette smoking prevalence, and categorical dummy indicators for clean indoor air grades given their expression on a “A” to “F” scale. Because these analyses potentially represent multiple testing of the same general (albeit separate, and routinely treated as separate) construct, the alpha was adjusted to correspond to the four tests ( $\alpha = 0.05 \div 4 = 0.0125$ ), even though this did not change the conclusions of the results.

Finally, ENDS searches related to “shopping,” seeking information about the “health” aspects of ENDS, or seeking information about the “cessation” aspects of ENDS were quantified building on methods the authors have demonstrated elsewhere.<sup>16,26</sup> First, potential search terms that occurred within ENDS searches indicative of the searcher's motivation or interest were identified based on the authors' familiarity with ENDS searches and in consultations with ENDS experts in their respective centers. Ultimately, terms with strong face validity were selected. For example, the occurrence of the term *buy* in an ENDS query is likely indicative of shopping. For shopping-related searches, ENDS searches that included the terms *buy*, *sale/s*, *shop/s*, or *store/s* were clustered. Similarly, to aggregate health-related searches, ENDS searches that included the terms *health*, *healthy*, *risk*, *risky*, *bad*, *harmful*, *cancer*, or *lung* were clustered; for cessation-related searches, those that included *stop* or *quit* were clustered. For example, *health effects of vaping*, *are e-cigs healthy?*, or *are electronic cigarettes harmful?* would be categorized as health-related (as well as hundreds more searches with these root terms). Second, the authors monitored

shopping, health, and cessation ENDS searches as a percentage of all ENDS searches each week. The resulting trends were then analyzed using methods similar to those for analyzing other trends as detailed above.

## Results

Searches regarding ENDS continue to increase, with an estimated 8,498,180 ENDS searches during 2014. All ENDS searches during 2014 (January 1, 2014 through January 1, 2015) were 450% (95% CI=313%, 711%) higher than the authors last reported for 2010, with approximately 1,545,123 ENDS searches in 2010. ENDS continue to be more searched than other smoking alternatives or nicotine-replacement therapies. ENDS searches during 2014 were 6,606% (95% CI=3,700%, 9,800%) or 66 times, 3,899% (95% CI=2,767%, 4,850%), and 3,177% (95% CI=2,433%, 4,350%) greater than searches for snus, nicotine-replacement therapies, and Chantix, respectively.

Within the increase in ENDS searches, there was a divergence in the search terms used beginning in 2014. Searches with vaping terms (e.g., *best vapes*) were increasing alongside declines in searches that used more-traditional e-cigarette terms (e.g., *best e-cigarette*). Vaping searches first surpassed e-cigarette searches in May 2014, and by December 2014, vaping searches were 95% (95% CI=76%, 109%) more common than e-cigarette searches.

The present forecasts suggest that there will be 62% (95% CI=22%, 95%) more ENDS searches on Google in 2015 than in 2014. Searches with vaping terms are also expected to continue increasing alongside decreasing searches for e-cigarettes, such that by December 2015 vaping searches will likely be 76% (95% CI=68%, 90%) greater than e-cigarette searches.

Between 2009 and 2014, ENDS searches have gone from being concentrated in states like Florida, Nevada, and Texas with fewer searches in the Midwest to being more uniformly searched across the U.S. (Figure 2). The variation between states significantly ( $p<0.001$ ) declined from 2009 through 2014 (the mean decline in SD across states was 9 [95% CI=2, 76] RSV per year). For instance, during 2009, ENDS searches in Wisconsin were 104% (95% CI=0%, 138%) higher than the mean for the other 50 states, but by 2014, the difference was -7% (95% CI= -46%, 29%) and statistically insignificant.

Yet, two geographic variations remained by 2014. First, searches for ENDS were significantly ( $p<0.001$ ) more common in Western and Midwestern states than on the Eastern seaboard (Figure 3). Second, the shift toward vaping terms over e-cigarette terms was more common in coastal states ( $p<0.001$ ), with the exception of a few New England states. For example, for all of 2014, California had the second highest volume of ENDS searches, of which 72% (95% CI=63%, 81%) included vaping terms.

Searches regarding ENDS appeared more common in states with more cigarette smokers (Figure 4), but states with more cigarette smokers were less likely to include vaping terms in their searches, although neither trend was statistically significant ( $p=0.062$  and  $p=0.145$ , respectively). Overall, during 2014, ENDS were searched at similar rates regardless of anti-smoking norms across states ( $p=0.332$ ) or strength of clean indoor air law provisions

( $p=0.260$ ), after adjusting for multiple tests ( $\alpha=0.05 \div 4 = 0.0125$ ). Yet, more ENDS searches involved vaping terms in states where anti-smoking norms were stronger ( $p<0.004$ ). For example, an increase from the 25th to the 75th percentile for anti-smoking norms predicted a 7% (95% CI=6%, 8%) increase in the proportion of ENDS searches involving vaping terms. ENDS searches were also greater in states with lower cigarette taxes ( $p<0.001$ ), but this pattern did not favor e-cigarette or vaping terms ( $p=0.528$ ). Further inspection of these data suggests the disparity was largely driven by the highest tax states (e.g., New York or Massachusetts).

About 6% (95% CI=2%, 10%) and 11% (95% CI=9%, 14%) of all ENDS searches nationally included the terms *store/s*, *shop/s*, *sale/s*, or *buy* during 2013 and 2014, respectively (Figure 5). As these statistics suggest, the rate at which ENDS searches included shopping terms was growing over time ( $p<0.0001$  for trend). In practical terms, this suggests there were 333,092 and 934,800 shopping searches in 2013 and 2014, respectively.

By contrast, only 3% (95% CI=1%, 6%) and 2% (95% CI=1%, 4%) of all ENDS searches in 2013 and 2014, respectively, included terms indicative of a health concern (e.g., *vaping healthy* or *e-cigarette risks*). The change in ENDS searches with health terms appeared to decrease over this time period (slope, 0.8 [95% CI=0.3, 1] RSV per year;  $p<0.001$ ). Even fewer ENDS searches included cessation terms, such as *do e-cigarettes help smokers quit?*, representing 0.3% (95% CI=0.1%, 0.4%) in 2013 and 0.2% (95% CI=0.001%, 0.5%) in 2014 of all ENDS searches. This change in ENDS searches with cessation terms had a significantly negative slope ( $-0.09$  [95% CI=  $-0.13$ ,  $-0.06$ ] RSV per year,  $p<0.001$ ).

## Discussion

Thousands are searching Google for ENDS each day. ENDS searches have increased in every U.S. state, with search terms now shifting from e-cigarettes to vaping, especially in coastal states and states where anti-smoking norms are stronger. When accounting for possible search motivations, it appeared that searches indicative of shopping for ENDS were increasing, whereas searches including health or cessation topics for ENDS accounted for both a smaller proportion of searches overall and have declined over time.

These findings directly address the ENDS surveillance gaps noted in numerous policy statements and review pieces.<sup>27</sup> This study is among the first to describe the vocabulary used by the population to search for ENDS, to provide statewide estimates of ENDS interest, to estimate the volume of online shopping for ENDS, and to describe how the public seeks out information on ENDS' health and cessation implications. As such, query-based intelligence has actionable implications for the development of new research questions and further policy debate.

Even though ENDS emerged in the U.S. marketplace fewer than 10 years ago,<sup>28</sup> by 2014, these products were frequently searched in every U.S. state. Searches for ENDS appear to be falling into two broad categories: "vaping" versus ENDS products names, like e-cigarettes. "Vaping" has emerged as the equivalent of "smoking" when referring to ENDS, and the rise in vaping searches is expected to outpace all other ENDS terms. This suggests that surveys

might rely on questions that use terms like *vaping*, over less commonly used product terms. Moreover, future research might explore the cultural significance of this shift in terminology.

The largely null association between ENDS searches and existing tobacco control measures highlights how ENDS may be resilient to current tobacco control regulations. For example, ENDS were searched for even more in states with high versus low cigarette taxes<sup>29</sup> and similarly searched for in states with strong versus weak clean indoor air laws.<sup>30</sup> By contrast, there was a strong positive association between anti-smoking norms and vaping searches. This suggests ENDS are less stigmatized than combustible products and potentially are being turned to as a means of avoiding stigmatization while maintaining the sensations of cigarette smoking.<sup>31</sup>

Already, millions of Google searches have been made with the likely intention of buying ENDS. Moreover, shopping searches nearly doubled from 2013 to 2014 and are projected to increase. Unlike most other tobacco products, there are no existing federal regulations governing the online sale of ENDS.<sup>32</sup> For instance, one recent study found that 77% of children aged 14–17 years were able to successfully order ENDS online and have them delivered to their home.<sup>33</sup>

Individuals in the U.S. often endorse ENDS as smoking cessation aids and some surveys suggest that many believe using ENDS will help them quit combustible cigarettes.<sup>34–37</sup> However, only a small and declining percentage of Google searches for ENDS included terms indicative of cessation. The context of this discrepancy is critical. When primed by survey questions, individuals appear to link ENDS with cessation, but in the privacy of their own home (when no investigator is providing options), it appears that searches for ENDS and cessation are infrequent. This low level of Google searches for cessation is in line with existing evidence on the effectiveness of ENDS for cessation. For instance, a meta-analysis of population-based studies of the association between ENDS and cessation indicated that ENDS are associated with significantly lower odds of quitting combustible cigarettes.<sup>11</sup> These findings are further supported by evidence that adult smokers often use ENDS in combination with other combustible tobacco products<sup>38</sup> and that many adolescents are using ENDS without ever previously smoking cigarettes.<sup>39,40</sup>

It appeared that searches related to the potential health effects of ENDS are becoming more infrequent, whereas the evidence base for the health risks about ENDS is beginning to accumulate.<sup>41–43</sup> Additional health campaigns are needed to disseminate the scientific knowledge on ENDS use. For example, recent mass media campaigns, including Tips from Former Smokers and the California Department of Public Health's Still Blowing Smoke campaign have piloted advertisements with messages that highlight the known harms of ENDS use and the fact that many ENDS products are produced by the tobacco industry who has previously engaged in dishonest behavior ([stillblowingsmoke.org](http://stillblowingsmoke.org)). Campaigns focused on highlighting risks and encouraging potential ENDS users to understand the risks (or larger state of uncertainty) are feasible given existing infrastructure for anti-tobacco mass media campaigns. Additionally, with increasing online interest in ENDS and shifting trends toward online media consumption, it may be worthwhile to further develop infrastructure to



engage in online health communication in addition to other traditional methods that have predominantly focused on TV-based media campaigns.

## Limitations

There are several study limitations to address. First, there is a unique validation challenge with search query surveillance. Typically the validity of new measures are established by comparison with existing gold standards.<sup>44</sup> For instance, the authors have used weekly CDC influenza-like illness trends to validate Google searches for influenza,<sup>18</sup> among a handful of other search validation studies.<sup>45,46</sup> However, in most cases, as with this study, no survey-based criterion exists. Even so, searches have strong face validity and confidence in their accuracy is bolstered by the facts that Internet users are demographically similar to ENDS users,<sup>47</sup> many survey-based studies replicated the authors' earlier assessment of ENDS searches,<sup>3</sup> and aggregate searches for other tobacco products corresponded to state-level prevalences.<sup>48</sup> Second, discriminating motivation across ENDS searches is more challenging. To overcome this challenge, only highly specific search terms were used. Yet, a journalist might search *best vape store* to learn about retailers without any personal interest in shopping. Such singular scenarios probably have little impact on aggregate trends given there are thousands of ENDS searches each day. Third, because searches are analyzed at the population level, they cannot be linked to searchers' demographics like with surveys. Still, search query surveillance and big data generally have numerous strengths over traditional surveillance, especially in behavioral medicine where the thoughts and actions of the population can be passively observed in near real time.<sup>12</sup>

## Conclusions

Tobacco control has historically lagged behind online tobacco markets, leaving gaps in surveillance.<sup>49–52</sup> Nowhere is this clearer than with the rise of ENDS. ENDS have become popular during a period without strong surveillance and a slowed public health reaction. Innovative methods like search query surveillance can improve the timeliness of tobacco control surveillance, especially around ENDS. As research agendas are being outlined for ENDS in numerous commentaries and opinion pieces,<sup>53–56</sup> further consideration should be given to the potential benefits of big data streams, like Internet searches. In particular, analyses like herein can both provide critical formative feedback for more costly and labor intensive investigations, such as informing survey question wording or coverage, and provide determinative insights on questions that may not be assessable using traditional techniques.

## Acknowledgments

Research reported in this publication was supported by 5R01CA169189-02, RCA173299A, and T32CA009492 from the National Cancer Institute and U.S. Food and Drug Administration Center for Tobacco Products. The content is solely the responsibility of the authors and does not represent the official views of the funders. The funders had no role in the design, conduct, or interpretation of the study nor the preparation, review, or approval of the manuscript.

Dr. Ayers and Dr. Althouse share an equity stake in a consultancy, Directing Medicine LLC, which advises clinician scientists how to implement some of the methods embodied in this work. Their organization previously advised the larger parent project of Dr. Williams on big data analytics prior to this study, and has advised other projects based in the Lineberger Comprehensive Cancer Center in the past 5 years. Dr. Dredze has been a paid by Directing Medicine



LLC for unrelated work. Dr. Ayers and Dr. Althouse have been separately paid by the University of North Carolina for unrelated speaking engagements and travel in the past 5 years. Neither the data nor the methods described in this article are proprietary. No other financial disclosures were reported by the authors of this paper.


## References

1. Yamin CK, Bitton A, Bates DW. E-cigarettes: a rapidly growing internet phenomenon. *Ann Intern Med.* 2010; 153(9):607–609. <http://dx.doi.org/10.7326/0003-4819-153-9-201011020-00011>. [PubMed: 21041581]
2. Noel JK, Rees VW, Connolly GN. Electronic cigarettes: a new ‘tobacco’ industry? *Tob Control.* 2011; 20(1):81. <http://dx.doi.org/10.1136/tc.2010.038562>. [PubMed: 20930060]
3. Ayers JW, Ribisl KM, Brownstein JS. Tracking the rise in popularity of electronic nicotine delivery systems (electronic cigarettes) using search query surveillance. *Am J Prev Med.* 2011; 40(4):448–453. <http://dx.doi.org/10.1016/j.amepre.2010.12.007>. [PubMed: 21406279]
4. Adkison SE, O’Connor RJ, Bansal-Travers M, et al. Electronic nicotine delivery systems: international tobacco control four-country survey. *Am J Prev Med.* 2013; 44(3):207–215. <http://dx.doi.org/10.1016/j.amepre.2012.10.018>. [PubMed: 23415116]
5. Regan AK, Promoff G, Dube SR, Arrazola R. Electronic nicotine delivery systems: adult use and awareness of the ‘e-cigarette’ in the USA. *Tob Control.* 2013; 22(1):19–23. <http://dx.doi.org/10.1136/tobaccocontrol-2011-050044>. [PubMed: 22034071]
6. Vickerman KA, Carpenter KM, Altman T, Nash CM, Zbikowski SM. Use of electronic cigarettes among state tobacco cessation quitline callers. *Nicotine Tob Res.* 2013; 15(10):1787–1791. <http://dx.doi.org/10.1093/ntr/ntt061>. [PubMed: 23658395]
7. McMillen RC, Gottlieb MA, Shaefer RM, Winickoff JP, Klein JD. Trends in electronic cigarette use among U.S. adults: Use is increasing in both smokers and nonsmokers. *Nicotine Tob Res.* 2015; 17(10):1195–1202. <http://dx.doi.org/10.1093/ntr/ntu213>. [PubMed: 25381306]
8. Tan AS, Bigman CA. E-cigarette awareness and perceived harmfulness: prevalence and associations with smoking-cessation outcomes. *Am J Prev Med.* 2014; 47(2):141–149. <http://dx.doi.org/10.1016/j.amepre.2014.02.011>. [PubMed: 24794422]
9. Giovenco DP, Lewis MJ, Delnevo CD. Factors associated with e-cigarette use: a national population survey of current and former smokers. *Am J Prev Med.* 2014; 47(4):476–480. <http://dx.doi.org/10.1016/j.amepre.2014.04.009>. [PubMed: 24880986]
10. King BA, Patel R, Nguyen KH, Dube SR. Trends in awareness and use of electronic cigarettes among U.S. adults, 2010–2013. *Nicotine Tob Res.* 2015; 17(2):219–227. <http://dx.doi.org/10.1093/ntr/ntu191>. [PubMed: 25239961]
11. Grana RA, Ling PM. Smoking Revolution: A content analysis of electronic cigarette retail websites. *Am J Prev Med.* 2014; 46(4):395–403. <http://dx.doi.org/10.1016/j.amepre.2013.12.010>. [PubMed: 24650842]
12. Ayers JW, Althouse BM, Dredze M. Could behavioral medicine lead the web data revolution? *JAMA.* 2014; 311(14):1399–1400. <http://dx.doi.org/10.1001/jama.2014.1505>. [PubMed: 24577162]
13. Jo CL, Ayers JW, Althouse BM, Emery S, Huang J, Ribisl KM. U.S. consumer interest in non-cigarette tobacco products spikes around the 2009 federal tobacco tax increase. *Tob Control.* 2014; 24(4):395–399. <http://dx.doi.org/10.1136/tobaccocontrol-2013-051261>. [PubMed: 24500270]
14. Ayers JW, Althouse BM, Johnson M, Cohen JE. Circaseptan (weekly) rhythms in smoking cessation considerations. *JAMA Intern Med.* 2014; 174(1):146–148. <http://dx.doi.org/10.1001/jamainternmed.2013.11933>. [PubMed: 24166181]
15. Ayers JW, Althouse BM, Noar SM, Cohen JE. Do celebrity cancer diagnoses promote primary cancer prevention? *Prev Med.* 2014; 58:81–84. <http://dx.doi.org/10.1016/j.ypmed.2013.11.007>. [PubMed: 24252489]
16. Ayers JW, Althouse BM, Ribisl KM, Emery S. Digital detection for tobacco control: Online Reactions to the United States’ 2009 cigarette excise tax increase. *Nicotine Tob Res.* 2013; 16(5): 576–583. <http://dx.doi.org/10.1093/ntr/ntt186>. [PubMed: 24323570]

17. Nuti SV, Wayda B, Ranasinghe I, et al. The use of google trends in health care research: a systematic review. *PLoS One*. 2014; 9(10):e109583. <http://dx.doi.org/10.1371/journal.pone.0109583>. [PubMed: 25337815]
18. Santillana M, Zhang DW, Althouse BM, Ayers JW. What can digital disease detection learn from (an external revision to) google flu trends? *Am J Prev Med*. 2014; 47(3):341–347. <http://dx.doi.org/10.1016/j.amepre.2014.05.020>. [PubMed: 24997572]
19. Ayers JW, Althouse BM, Johnson M, Dredze M, Cohen JE. What’s the healthiest day?: Circaseptan (weekly) rhythms in healthy considerations. *Am J Prev Med*. 2014; 47(1):73–76. <http://dx.doi.org/10.1016/j.amepre.2014.02.003>. [PubMed: 24746375]
20. Ayers JW, Althouse BM, Emery S. Changes in internet searches associated with the “tips from former smokers” campaign. *Am J Prev Med*. 2015; 48(6):e27–e29. <http://dx.doi.org/10.1016/j.amepre.2015.03.015>. [PubMed: 25998929]
21. Hyndman, R.J., Khandakar, Y. Automatic time series forecasting: The forecast package for R 7, 2008. 2007. [www.jstatsoft.org/v27/i03](http://www.jstatsoft.org/v27/i03)
22. King G, Tomz M, Wittenberg J. Making the most of statistical analyses: Improving interpretation and presentation. *Am J Poli Sci*. 2000; 44(2):341–355. <http://dx.doi.org/10.2307/2669316>.
23. American Lung Association. American Lung Association State of Tobacco Control 2014. Washington DC: American Lung Association; 2014.
24. CDC. Federal and state cigarette excise taxes - United States, 1995–2009. *MMWR Morb Mortal Wkly Rep*. 2009; 58(19):524–527. [PubMed: 19478719]
25. Alamar B, Glantz SA. Effect of increased social unacceptability of cigarette smoking on reduction in cigarette consumption. *Am J Public Health*. 2006; 96(8):1359–1363. <http://dx.doi.org/10.2105/AJPH.2005.069617>. [PubMed: 16809588]
26. Althouse BM, Allem J-P, Childers MA, Dredze M, Ayers JW. Population health concerns during the united states great recession. *Am J Prev Med*. 2014; 46(2):166–170. <http://dx.doi.org/10.1016/j.amepre.2013.10.008>. [PubMed: 24439350]
27. Bhatnagar A, Whitsel LP, Ribisl KM, et al. Electronic cigarettes: a policy statement from the American Heart Association. *Circulation*. 2014; 130(16):1418–1436. <http://dx.doi.org/10.1161/CIR.000000000000107>. [PubMed: 25156991]
28. Pepper JK, Brewer NT. Electronic nicotine delivery system (electronic cigarette) awareness, use, reactions and beliefs: a systematic review. *Tob Control*. 2013; 23(5):375–384. <http://dx.doi.org/10.1136/tobaccocontrol-2013-051122>. [PubMed: 24259045]
29. Contreary KA, Chattopadhyay SK, Hopkins DP, et al. Economic impact of tobacco price increases through taxation: a community guide systematic review. *Am J Prev Med*. 2015; 49(5):800–808. <http://dx.doi.org/10.1016/j.amepre.2015.04.026>. [PubMed: 26188686]
30. Majeed BA, Dube SR, Sterling K, Whitney C, Eriksen MP. Opinions about electronic cigarette use in smoke-free areas among U.S. adults, 2012. *Nicotine Tob Res*. 2015; 17(6):675–681. <http://dx.doi.org/10.1093/ntr/ntu235>. [PubMed: 25358659]
31. Fairchild AL, Bayer R, Colgrove J. The renormalization of smoking? E-cigarettes and the tobacco “endgame”. *N Engl J Med*. 2014; 370(4):293–295. <http://dx.doi.org/10.1056/NEJMp1313940>. [PubMed: 24350902]
32. Henriksen L. The retail environment for tobacco: a barometer of progress towards the endgame. *Tob Control*. 2015; 24(e1):e1–e2. <http://dx.doi.org/10.1136/tobaccocontrol-2014-051884>. [PubMed: 25701879]
33. Williams RS, Derrick J, Ribisl KM. Electronic cigarette sales to minors via the internet. *JAMA Pediatrics*. 2015; 169(3):e1563. <http://dx.doi.org/10.1001/jamapediatrics.2015.63>. [PubMed: 25730697]
34. Siegel MB, Tanwar KL, Wood KS. Electronic cigarettes as a smoking-cessation: tool results from an online survey. *Am J Prev Med*. 2011; 40(4):472–475. <http://dx.doi.org/10.1016/j.amepre.2010.12.006>. [PubMed: 21406283]
35. Pearson JL, Richardson A, Niaura RS, Vallone DM, Abrams DB. E-cigarette awareness, use, and harm perceptions in U.S. adults. *Am J Public Health*. 2012; 102(9):1758–1766. <http://dx.doi.org/10.2105/AJPH.2011.300526>. [PubMed: 22813087]


36. Etter JF, Bullen C, Flouris AD, Laugesen M, Eissenberg T. Electronic nicotine delivery systems: a research agenda. *Tob Control*. 2011; 20(3):243–248. <http://dx.doi.org/10.1136/tc.2010.042168>. [PubMed: 21415064]
37. Goniewicz ML, Lingas EO, Hajek P. Patterns of electronic cigarette use and user beliefs about their safety and benefits: an internet survey. *Drug Alcohol Rev*. 2013; 32(2):133–140. <http://dx.doi.org/10.1111/j.1465-3362.2012.00512.x>. [PubMed: 22994631]
38. Sussman S, Garcia R, Cruz TB, Baezconde-Garbanati L, Pentz MA, Unger JB. Consumers' perceptions of vape shops in Southern California: an analysis of online Yelp reviews. *Tob Induc Dis*. 2014; 12(1):22. <http://dx.doi.org/10.1186/s12971-014-0022-7>. [PubMed: 25484852]
39. Wills TA, Knight R, Williams RJ, Pagano I, Sargent JD. Risk factors for exclusive e-cigarette use and dual e-cigarette use and tobacco use in adolescents. *Pediatrics*. 2015; 135(1):e43–e51. <http://dx.doi.org/10.1542/peds.2014-0760>. [PubMed: 25511118]
40. Leventhal AM, Strong DR, Kirkpatrick MG, et al. Association of electronic cigarette use with initiation of combustible tobacco product smoking in early adolescence. *JAMA*. 2015; 314(7):700–707. <http://dx.doi.org/10.1001/jama.2015.8950>. [PubMed: 26284721]
41. Hajek P, Etter JF, Benowitz N, Eissenberg T, McRobbie H. Electronic cigarettes: review of use, content, safety, effects on smokers and potential for harm and benefit. *Addiction*. 2014; 109(11):1801–1810. <http://dx.doi.org/10.1111/add.12659>. [PubMed: 25078252]
42. Maziak W. Harm reduction at the crossroads: the case of e-cigarettes. *Am J Prev Med*. 2014; 47(4):505–507. <http://dx.doi.org/10.1016/j.amepre.2014.06.022>. [PubMed: 25092121]
43. Manzoli L, Flacco ME, Fiore M, et al. Electronic cigarettes efficacy and safety at 12 months: cohort study. *PLoS One*. 2015; 10(6):e0129443. <http://dx.doi.org/10.1371/journal.pone.0129443>. [PubMed: 26061661]
44. Strauss ME, Smith GT. Construct validity: advances in theory and methodology. *Annu Rev Clin Psychol*. 2009; 5:1–25. <http://dx.doi.org/10.1146/annurev.clinpsy.032408.153639>. [PubMed: 19086835]
45. Goel S, Hofman JM, Lahaie S, Pennock DM, Watts DJ. Predicting consumer behavior with Web search. *Proc Natl Acad Sci*. 2010; 107(41):17486–17490. <http://dx.doi.org/10.1073/pnas.1005962107>. [PubMed: 20876140]
46. Althouse BM, Ng YY, Cummings DA. Prediction of dengue incidence using search query surveillance. *PLoS Negl Trop Dis*. 2011; 5(8):e1258. <http://dx.doi.org/10.1371/journal.pntd.0001258>. [PubMed: 21829744]
47. Internet & American Life Project. Pew Research Center Internet Project Survey, January 2014. Washington, DC: Pew Internet Research; 2014.
48. Cavazos-Rehg PA, Krauss MJ, Spitznagel EL, et al. Monitoring of non-cigarette tobacco use using Google Trends. *Tob Control*. 2014; 24(3):249–255. <http://dx.doi.org/10.1136/tobaccocontrol-2013-051276>. [PubMed: 24500269]
49. Cohen JE, Sarabia V, Ashley MJ. Tobacco commerce on the internet: a threat to comprehensive tobacco control. *Tob Control*. 2001; 10(4):364–367. <http://dx.doi.org/10.1136/tc.10.4.364>. [PubMed: 11740029]
50. Ribisl KM, Jo C. Tobacco control is losing ground in the Web 2.0 era: invited commentary. *Tob Control*. 2012; 21(2):145–146. <http://dx.doi.org/10.1136/tobaccocontrol-2011-050360>. [PubMed: 22345237]
51. Ayers JW, Ribisl K, Brownstein JS. Using search query surveillance to monitor tax avoidance and smoking cessation following the United States' 2009 "schip" cigarette tax increase. *PLoS One*. 2011; 6(3):e16777. <http://dx.doi.org/10.1371/journal.pone.0016777>. [PubMed: 21436883]
52. Allem, JP., Ayers, JW., Althouse, BM., Williams, R. When a ban really is not a ban: internet loopholes and Djarum flavoured cigarettes in the USA. *Tob Control*. 2015. <http://dx.doi.org/10.1136/tobaccocontrol-2015-052309>
53. Sussman, S., Baezconde-Garbanati, L., Garcia, R., et al. Forces that drive the vape shop industry and implications for the health professions. *Eval Health Prof*. 2015. <http://dx.doi.org/10.1177/0163278715586295>

54. Barrington-Trimis JL, Samet JM, McConnell R. Flavorings in electronic cigarettes: an unrecognized respiratory health hazard? *JAMA*. 2014; 312(23):2493–2494. <http://dx.doi.org/10.1001/jama.2014.14830>. [PubMed: 25383564]
55. Cobb CO, Hendricks PS, Eissenberg T. Electronic cigarettes and nicotine dependence: evolving products, evolving problems. *BMC Med*. 2015; 13:119. <http://dx.doi.org/10.1186/s12916-015-0355-y>. [PubMed: 25998379]
56. Etter JF. E-cigarettes: methodological and ideological issues and research priorities. *BMC Med*. 2015; 13:32. <http://dx.doi.org/10.1186/s12916-014-0264-5>. [PubMed: 25856794]




**Figure 1. National trends for electronic nicotine delivery systems Google searches, 2004–2014**

Both panels display the national trend for all electronic nicotine delivery systems (ENDS) searches as derived from searches originating in the U.S. that included the keywords as described in the text (e.g., “buy e-cigarettes”). Panel (a) compared ENDS searches to searches for snus, Chantix, and nicotine replacement therapies. Panel (b) compared among ENDS searches that included terms indicative of vaping (e.g., “best vaping cigarettes”) or e-cigarettes (e.g., “best e-cigarettes”). Both panels present relative search volumes (100=highest search proportion, 50=50% of the highest search proportion for all Google searches on ENDS). Forecasted values through 2015 are described in the text but not shown here.




**Figure 2. The spread of electronic nicotine delivery systems Google searches by U.S. states, 2009–2014**

Each map shows the mean annual relative search volume for all electronic nicotine delivery systems (ENDS) searches. All panels present relative search volumes (100=highest search proportion, 50=50% of the highest search proportion for all Google searches on ENDS). Years prior to 2009 were not presented because searches were near or at zero volume.




**Figure 3. Longitude predicts electronic nicotine delivery systems Google searches, 2014**  
 Panel (a) compared all ENDS searches by state to the median state longitude. Searches were measured using the mean relative search volumes (100=highest search proportion, 50=50% of the highest search proportion for all Google searches on ENDS) for all of 2014. Panel (b) compared the proportion of all ENDS searches that included terms indicative of vaping (e.g., “best vaping cigarettes”) by state to the median state longitude.





**Figure 4. Electronic nicotine delivery systems Google searches by select predictors for tobacco control and smoking social norms, 2014**

Panels (a, c, e, and g) compared all ENDS searches in 2014 by state according to the smoking prevalence, social unacceptability of smoking,(20) cigarette excise tax, and clean indoor air grade –as detailed in the text. Searches were measured using the mean relative search volumes (100=highest search proportion, 50=50% of the highest search proportion for all Google searches on ENDS) for all of 2014. Panels (b, d, f, and h) replicate the same analyses but using the proportion of all ENDS searches that included terms indicative of vaping (e.g., “best vaping cigarettes”) by state as the outcome.



**Figure 5. Electronic nicotine delivery systems Google searches including shopping, health, or cessation, 2014**

Each line shows the proportion of all ENDS searches that also included terms consistent with shopping (e.g., “buy”), health (e.g., “harmful”), or cessation (e.g., “quit”), as detailed in the text. Cessation is shown on a separate scale, given searches were rare in this category.