

Lawrence Berkeley National Laboratory

Recent Work

Title

Experimental Results on the pi-pi Cross Section

Permalink

<https://escholarship.org/uc/item/55f452j1>

Journal

Physical Review Letters, 6(7)

Authors

Anderson, Jerry A.

Bank, Vo X.

Burke, Philip G.

et al.

Publication Date

1961-02-23

UNIVERSITY OF
CALIFORNIA

Ernest O. Lawrence

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

To be submitted to Physical Review Letters

UCRL-9577

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

EXPERIMENTAL RESULTS ON THE π - η CROSS SECTION

Jerry A. Anderson, Vo X. Bang, Philip G. Burke,
D. Duane Carmony and Norbert Schmitz

February 23, 1961

EXPERIMENTAL RESULTS ON THE π - π CROSS SECTION*

Jerry A. Anderson, Vo X. Bang, Philip G. Burke,
D. Duane Carmony and Norbert Schmitz†

Lawrence Radiation Laboratory
University of California
Berkeley, California

February 23, 1961

At the Rochester Conference 1960,¹ we reported our first experimental results on the π - π cross section, which we obtained by applying the Chew-Low extrapolation method² to about 700 π -p inelastic scatterings. These events were analyzed in photographs taken with the Alvarez 72-inch hydrogen bubble chamber, which was exposed to a 1.03-Bev/c π^- beam at the Bevatron. We now have a total of 1275 inelastic events with the proton stopping in the chamber. In this paper we will use the notation, the selection and evaluation criteria, and the extrapolation procedures which have been discussed in Reference 1.

Figure 1 shows our experimental distributions,

$$F(p^2, \omega^2) = (p^2 + 1)^2 \frac{\partial^2 \sigma(p^2, \omega^2)}{\partial p^2 \partial \omega^2} ,$$

as functions of the four-momentum transfer p^2 for eight different intervals of ω^2 , the invariant mass of the di-pion. We express p^2 and ω^2 in units of the pion mass squared. Fitting polynomials in $(p^2 + 1)$ of several orders through these data we obtain results as follows.

1. Whereas at the time of the Rochester Conference a first-order polynomial adequately fitted the data for all the intervals of ω^2 , we find that a χ^2 test now reveals the necessity of a quadratic fit for the low ω^2 intervals. In the higher ω^2 intervals, although we have more events than at the lower ω^2 intervals, the linear fits are still exceedingly good. We therefore take the linear values for the extrapolation π - π cross section at the higher energy.

* This work was performed under the auspices of the U. S. Atomic Energy Commission.

† Now at the Max-Planck-Institut für Physik und Astrophysik, München, Germany.

2. Also at the time of the Rochester Conference some of the fitted curves had the unphysical feature that they went negative in the physical region. This was possible because we do not have events right up to the edge of the physical region. Since we now require quadratic fits for the lower ω^2 intervals, this difficulty has been removed in all cases except for the lowest ω^2 interval (5.5 to 7.8 m_π^2). For this interval only we had to apply a constraint forcing the fitted curve to go through the end of the physical region. We then find that a linear curve with this added constraint gives a good fit. In all other intervals the fitted curve remains positive in the physical region without this constraint.

In order to get some information about the π - π interaction from a very small number of events, Bonsignori and Selleri^{3,4,5} have made the assumption that formula (3.13) of Ref. 2 (which is exactly correct only at the pole $p^2 = -1$) approximately describes the p^2 dependence at the beginning of the physical region. With this assumption an average π - π cross section has been determined. The assumption means that all other contributions to the amplitude for single-pion production (p^2 dependence of the vertex functions of the single-pion exchange diagram, exchanges of 3, 5, or more pions, and production by collision of the pion with the nucleon core rather than with the cloud) have been neglected in comparison with the single-pion exchange at the pole. With our present statistics we are able to check the assumption of Bonsignori and Selleri. If it were justified it should be possible (as one can see from (3.12) in Ref. 2) to fit the experimental distributions (or the first part of them) of Fig. 1 by a straight line passing through zero for $p^2 = 0$. This our data shows is not possible. Thus nonpole terms are important and an extrapolation requiring a large number of events is necessary to extract the pole term.

Frazer and Fulco⁶ could explain the vector part of the electromagnetic structure of the nucleon assuming a strong π - π resonance in the $T = J = 1$ state at $\omega^2 = 11$. The $\pi^- - \pi^0$ amplitude is 50% isotopic spin 1 and 50% isotopic spin 2. We find no evidence for a π - π resonance of the width and location predicted in Reference 6.

Our data do show a rise in the π - π cross section starting at $\omega^2 = 17$, reaching a value of the order of 200 mb at $\omega^2 = 20$ to 22.


However, one must remember that it is just in this region of ω^2 that our extrapolation distance begins to get larger, making the extrapolation procedure less conclusive. We are now scanning film obtained at 1.275 Bev/c incident momentum in order to reduce the extrapolation distance in this region. Also, for ω^2 greater than 9, we have a background of events coming from the reaction $\pi^- + p \rightarrow \pi^- + p + 2\pi^0$. These events do not have a pole at $p^2 = -1$, but come from a branch cut. We are eliminating this background by means of a kinematical fit.

On the other hand, Bowcock et al.⁷ found on a later analysis of the nucleon electromagnetic structure and the low-energy pion-nucleon phase shifts that the Frazer-Fulco resonance should be shifted to about $\omega_{res}^2 = 22$. This is consistent with our present results. If we assume that our data peak at $\omega^2 = 20$ to 22 (our incident energy is insufficient to examine the high-energy side of the peak), then the height is in accord with $(2J+1) 4\pi \lambda^2 \sin^2 \delta_1$ for a p-state resonance--that is, $\sin^2 \delta_1 = 1$. Our half width (obtained from the low-energy side) is approximately $5 m_{\pi}^2$. Of course our data do not rule out a nonresonant rise in the cross section composed of s, p, d, f ... states which just happens to satisfy $12 \pi \lambda^2$ at $\omega^2 = 20$ to 22. We are currently evaluating the differential cross section in the region $\omega^2 = 20$. We hope to resolve this experimental ambiguity in the very near future.

We would like to thank Professor Luis W. Alvarez for his great interest and encouragement throughout the experiment. It is a pleasure to thank Professor Frank S. Crawford, Jr., who designed the beam, as well as Professor Arthur H. Rosenfeld and other members of the Alvarez group, for many interesting and stimulating discussions. We are indebted to Professor Geoffrey F. Chew and to Dr. James S. Ball for several discussions on the theoretical aspects and to Dr. Herbert M. Steiner for useful comments. Finally we want to thank our scanners for their help in finding and analyzing the events.

REFERENCES

1. J. A. Anderson, P. G. Burke, D. D. Carmony, and N. Schmitz, in *Proceedings of the 1960 Annual International Conference on High Energy Physics at Rochester*, p. 58.
2. G. F. Chew and F. E. Low, *Phys. Rev.* 113, 1640 (1959).
3. V. Alles-Borelli, S. Bergio, E. Perez-Ferreira, and P. Waloschek, *Nuovo cimento* 14, 211 (1959).
4. I. Dorado and N. Schmitz, *Phys. Rev.* 118, 309 (1960).
5. F. Bonsignori and F. Selleri, *Nuovo cimento* 15, 465 (1960).
6. W. R. Frazer and J. R. Fulco, *Phys. Rev.* 117, 1609 (1960).
7. F. J. Bowcock, W. N. Cottingham, and D. Lurie, *Phys. Rev. Letters* 5, 386 (1960).



MUB-604

Fig. 1. Extrapolation curves $F(p^2, \omega^2)$ at fixed ω^2

(a) $\omega^2 = 5$ to $8.2 m_\pi^2$,

(e) $\omega^2 = 8.2$ to 11 ,

(b) $\omega^2 = 11$ to 13.7 ,


(f) $\omega^2 = 13.7$ to 16.5 ,

(c) $\omega^2 = 16.5$ to 19.2 ,

(g) $\omega^2 = 19.2$ to 22 ,

(d) $\omega^2 = 22$ to 24.7 ,

(h) $\omega^2 = 24.7$ to 27.5 .



MU-22625

Fig. 2. The $\pi^- - \pi^0$ cross section as a function of the total dipion mass squared as determined by the Chew-Low method. Also shown are the maximum height of a p-state resonance and the shape of the Frazer-Fulco resonance (Phys. Rev. Letters 2, 367 (1959), Eq. (10)), assuming the parameters $\nu_r = 3.5$, $\Gamma = .3$.

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.