

UCSF

UC San Francisco Previously Published Works

Title

Structural and functional conservation of fungal MatA and human SRY sex-determining proteins

Permalink

<https://escholarship.org/uc/item/55p72852>

Journal

Nature Communications, 5(1)

ISSN

2041-1723

Authors

Czaja, Wioletta
Miller, Karen Y
Skinner, Michael K
[et al.](#)

Publication Date

2014-11-01

DOI

10.1038/ncomms6434

Peer reviewed



Published in final edited form as:

Nat Commun. ; 5: 5434. doi:10.1038/ncomms6434.

Structural and functional conservation of fungal MatA and human SRY sex determining proteins

Wioletta Czaja^{1,2,4}, Karen Y. Miller^{1,4}, Michael K. Skinner^{1,3}, and Bruce L. Miller^{1,4}

¹Center for Reproductive Biology, Washington State University Pullman, WA, 99164-4236, USA

²School of Molecular Biosciences, Washington State University Pullman, WA, 99164-4236, USA

³School of Biological Sciences, Washington State University Pullman, WA, 99164-4236, USA

⁴Department of Biological Sciences, University of Idaho Moscow, ID, 83844, USA

Abstract

Sex determination in animals and fungi is regulated by specific sex determining genes. The *Aspergillus nidulans* (*A.nidulans*) mating type gene *matA* and the human *SRY* (Sex Determining Region Y) encode proteins containing a single HMG (High Mobility Group) domain. Analysis of the amino acid sequence of MatA and SRY transcription factors revealed significant structural similarity. The human SRY protein is able to functionally replace MatA and drives the sexual cycle in the fungus *A. nidulans*. Functional studies indicate that SRY drives early fruiting body development, and hybrid MatA protein carrying the SRY HMG box is fully capable of driving both early and late stages of sexual development, including gametogenesis. Our data suggest that SRY and MatA are both structurally and functionally related and conserved in regulating sexual processes. The fundamental mechanisms driving evolution of the genetic pathways underlying sex determination, sex chromosomes and sexual reproduction in eukaryotes appear similar.

Introduction

Fungal mating type systems demonstrate remarkable structural parallels between *Mat* loci and sex chromosomes in mammals, which suggest similar mechanisms may have operated in divergent eukaryotes during the early evolution of sex determination and sex chromosomes. It has long been recognized that fungal *Mat* loci share structural and functional features with the mammalian X and Y sex determination system^{1,2,3,4,5}. The classic examples involve ascomycete *Neurospora tetrasperma* and basidiomycete *Cryptococcus neoformans*, where mating type loci encode several sex-specific genes. These genes occupy large recombination suppressed regions and similar to sex chromosomes in

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:http://www.nature.com/authors/editorial_policies/license.html#terms

Correspondence: Michael K. Skinner Center for Reproductive Biology School of Biological Sciences Washington State University Pullman, WA 99164-4236 Phone: 509-335-1524 Fax: 509-335-2176 skinner@wsu.edu.

Author Contributions

WC conceived the study. WC, KYM and BLM performed the experiments. WC, BLM and MKS designed and wrote the study. All authors edited and approved the manuscript.

The authors declare no competing financial interests.

animals are rich in repetitive elements. The functional conservation of sex determining genes between distantly related eukaryotes such as fungi and mammals remains unclear.

In this report we describe structural similarity and functional conservation between the MatA (HMG box) protein of ascomycete *Aspergillus nidulans* and human sex determination region of the Y chromosome (SRY). Sex determining HMG-domain proteins of fungal and mammalian origin belong to the SOX/MATA/TCF superfamily of transcriptional regulators containing a single HMG domain. These proteins typically bind to the minor groove of a specific DNA sequence and act as architectural factors bending the DNA and facilitating gene transcription through changes in the chromatin structure^{6,7}. The role of HMG domain proteins in sex determination have focused on mammalian models. We hypothesize that structural similarity between fungal MatA and human SRY reflects functional relatedness between both HMG box proteins in the process of sex determination.

Results

Structural comparisons

Analysis of the amino acid sequence of MatA (318aa) and SRY (204aa) transcription factors revealed a remarkable level of structural similarity (Fig. 1A), which suggests that MatA is a structural homolog of human SRY. The HMG domain region contains 33% identity and 58% positives between conserved amino acids. The SRY HMG domains from human and various mammalian species share 70% identity⁸. Interestingly, human and murine SRY/Sry factors exhibit a remarkable sequence divergence with essentially no homology outside the HMG-box⁸. Mouse Sry contains a glutamine-rich region that is completely absent in human. Sequences outside the HMG box in MatA and hSRY are less conserved, although patches of closely similar sequence can be easily identified in both N- and C-terminal protein regions. The C-terminal conservation of putative helix 3 with the His, Tyr, and Pro residues suggests potential similar DNA bending capacity. Surprisingly, there appears to be a higher level of structural conservation in the N- and C-terminal regions between fungal MatA and hSRY than between mouse Sry and human SRY. This supports the observation that mouse SRY is highly divergent based on microsatellite instability⁹.

Comparison of the HMG box of fungal MatA and human SRY reveals unexpectedly high conservation of the amino acid sequence (Fig. 1B). All 10 amino acid residues which are known to form a direct contact with the DNA helix¹⁰ are identical between fungal and human HMG boxes. Residues marked in red are known to participate in a sequence-specific binding of HMG domain to the core DNA sequence¹⁰. As shown, all residues marked in red are identical except for the isoleucine (I) in MatA and valine (V) in SRY at position 2, which are non identical but have similar physico-chemical properties. The asparagine (N), highlighted in blue is typical of HMG boxes binding to DNA in a sequence-specific manner¹¹, and as indicated these residues are identical. Furthermore, the isoleucine (I) highlighted in green is important for HMG box specific DNA bending and binding efficiency^{8,12,13}. The conservation of isoleucine residue in both fungal and human HMG boxes may reflect similar modes of DNA binding and bending. MatA also contains a cluster of basic residues (ASRISPKN) which is similar to a cluster (KAKMLPKN) found in the SRY-related proteins. These residues are located C-terminal to the HMG box and are known

to function as a kinetic clamp to prolong the lifetime of SRY-DNA complex¹⁴. These conserved residues also provide insights into the ancestral origins of SRY compared to MatA which are thought to be duplication events (e.g. SOX3, TCF-1, and LEF-1) that can impact nucleocytoplasmic trafficking¹⁵.

Functional comparison

The structural similarity between fungal MatA and human SRY may suggest a functional conservation in the sex-determining pathway or other pathways driving sexual reproduction. To further explore this possibility we used the fungal model *A.nidulans* to perform interspecies gene exchange studies and analyze the ability of the human SRY protein to drive fungal sexual cycle. Manipulations of the *matA* gene structure and associated phenotypes of the resulting transgenic fungal strains are outlined in Figure 2. The endogenous open reading frame (ORF) of the *matA* gene was replaced with the ORF sequence of the SRY, creating a transgenic *A.nidulans* strain carrying the SRY ORF under control of the *matA* promoter. A separate transgenic strain was created, where the *matA* promoter: SRY transgene was integrated ectopically into the genome of a *matA(0)* strain lacking endogenous *matA*. Therefore, two independent strains were created (e.g. endogenous and ectopic) and tested with consistent phenotypes found, although the ectopic transgene was less robust. In addition, Southern blot, PCR, and DNA sequencing methods were all used to validate the different strains for SRY and SRY-MatA hybrid strains, see Methods¹⁶ and Supplementary Fig. 1. The *matA(0)* strains have a sterile phenotype described previously¹⁶. Sexual development of the ORF swap strains was analyzed with reference to the wild type *matA* positive control strain and *matA(0)* deletion negative control strain. At the early stage of the sexual cycle, 4-5 days post-induction, the wild type strain differentiated abundant fruiting bodies (cleistothecia) (Fig. 3-1C). By contrast, *matA(0)* and ORF swap strains lacked fruiting bodies and had developed only clusters of white hülle cells, which is typically an indication of unsuccessful fertilization (Fig. 3-4C). However, at 2 weeks post-induction, the wild type strain had developed mature cleistothecia (100-200µm in diameter) with abundant meiotic progeny (ascospores) (Fig. 3-1 A,B,C). The *matA(0)* strain showed no progress in further sexual development, forming thick masses of hülle cells with occasional formation of tiny, undeveloped progenitors of cleistothecia and no ascospores (Fig. 3-4 A,B,C). The ORF swap strain at the corresponding time point differentiated small (50-100µm) cleistothecia at approximately 20-30% of the wild type fruiting body abundance (Fig. 3-3B). These cleistothecia were mostly barren with development of the internal reproductive tissue having been aborted at the early stage of ascogenous hyphae differentiation. However, a small number of viable ascospores were observed (Fig. 3-3 A,B,C). Ascospore viability was routinely assessed by plating spores on agar plates, incubating and counting growing colonies. Viability of ascospores isolated from ORF swap strains and sexual phenotypes were stable generationally and meiotically that resolves secondary mutation effects. Therefore, human SRY is able to partially support fungal sexual differentiation although formation of the meiotic progeny is inefficient. Partial function of the SRY protein in driving fungal sexual development could be a result of impaired DNA binding and/or bending due to the lack of sufficient conservation between the MatA and SRY proteins. Other possibilities may include failures in protein-protein interactions between SRY and MatA-interacting fungal proteins, altered posttranslational

modifications, or improper recognition of the downstream sex-specific target genes. Limited information is currently available regarding down-stream gene targets for MatA, but future studies may help elucidate if altered gene expression may be a factor. Any of these abnormalities could contribute to the partial function of SRY in regulating fungal sexual development.

HMG box comparison

Structural and functional analyses of the mammalian HMG box proteins suggest that the conserved DNA binding HMG motif is necessary and sufficient element of SRY protein required to initiate male development and sex determination¹⁷. Indeed, most point mutations causing human sex reversal occur *de novo* in the HMG box region and impair specific DNA binding or DNA bending affecting the assembly of sex-specific transcriptional complexes^{8,14}. Several studies in mice demonstrated that transgenic expression of mammalian SRY or SOX protein bearing an SRY-type HMG domain within the mouse embryonic gonad during the critical time frame was sufficient to trigger male sex determination and XX sex reversal^{17,18}. Exploring this idea further we sought to investigate whether the SRY HMG box can functionally replace the MatA HMG box and be sufficient to regulate the *A.nidulans* sexual cycle. The HMG box of MatA was replaced with the HMG box of SRY and a transgene carrying the HMG swap was integrated ectopically into the *matA(0)* genome. Several independently derived HMG box swap transformants were analyzed and showed consistent phenotypes. We found that the hybrid MatA protein containing an HMG box of SRY is able to support a robust sexual cycle with fruiting body and progeny formation (Fig. 3-2 A,B,C). This data reveals functional conservation of the human HMG box within the context of fungal MatA protein in driving fungal sexual reproduction.

Regulatory element comparison

Interestingly, the structural conservation between MatA and SRY extends outside the gene coding region. Analysis of the promoter region in *matA* and *SRY* genes reveals a number of conserved putative regulatory elements. The promoter region of the mammalian *SRY* has several potential binding sites for SRY protein¹⁹. The HMG domain-specific core sequences (AACAAAG) have been identified 94bp upstream of the ATG in human *SRY*²⁰, and in several other mammalian species¹⁹. Analogously, we identified an identical sequence (CAAAG) located 216bp upstream of the translational start codon ATG of *matA* gene in *A.nidulans*. Furthermore, we identified a sequence for a potential silencing element residing in the 5' *matA* flank (-931 to -960bp) which shares high similarity to the putative silencing element found in the upstream regulatory region of the *SRY* gene (Fig. 4)¹⁹. Although this region has a number of regulatory elements nearby that are similar for a number of different species, deletion of this region in studies for the Wilms' Tumor 1 (WT1), Gata4 or TFCEP2 did not show strong regulatory activity in *SRY* reporter gene analysis^{21,22,23}. Similarly, deletion of this element does not effect *matA* expression *in vivo* or sexual development¹⁶. Interestingly, we identified a functional silencer, *matA* SE, located between -512 and -248bp that bears limited similarity to *hSRY* SE¹⁶ (Fig. 4). This general region also has a large number of regulatory elements (e.g. WT1, Gata4, Cp2) that are functional in *SRY*^{19,21,22,23,24}. The structural conservation of these regulatory elements suggests similar

mechanisms might be involved in the transcriptional regulation and expression of *matA* and *SRY* (Fig. 4 and Supplementary Fig. 2). These elements²⁴ in both *matA*¹⁶ and *SRY*¹⁹ appear to have been evolutionarily conserved and may contribute to the regulation of gene expression^{16,19,24}.

Discussion

Chromosomal sex-determining regions have evolved independently in different eukaryotic species and undergone a convergence of structure and function^{1,2,3}. Consistently with this theory, it has been speculated that fungal *Mat* idiomorphs may represent primitive forms of sex chromosomes^{1,2,25,26}. The current observations highlight a remarkable structural similarity and functional conservation between the fungal *MatA* and human *SRY* sex determining factors. The HMG box of human *SRY* is able to functionally replace the HMG box of fungal *MatA* and drives robust sexual differentiation and gametogenesis in *A. nidulans*. Combined observations provide insights concerning the role and conservation of the HMG box proteins in regulation of genetic pathways underlying sex determination and sexual reproduction in eukaryotes. Clearly other transcription factors such as homeodomain proteins for fungi and mammals are required^{27,28}. For example, previous analysis of DM domain proteins, such as *DMRT* or *DMY*, have strong conservation across metazoan sexual development, from insects to mammals²⁵. Therefore, different classes (e.g. HMG and DM) of transcription factors known to be involved in sex determination are also likely to have similarities between the mammalian and fungal processes. The current study suggests HMG domain proteins such as *MatA* and *SRY* have a strong conservation in sexual development, from fungi to mammals. Therefore, a combination of HMG box proteins, and potentially other factors such as DM domain proteins, appear to be involved in the sexual development of most organisms.

Methods

Protein sequence analysis

Amino acid sequences of the *MatA* (AN4734.3) and *SRY* (Q05066) proteins were analyzed using Clustal W multiple sequence alignment using the European Bioinformatics Institute website services at <http://www.ebi.ac.uk/Tools/clustalw2/index.html>.

Vectors construction

The *SRY* coding region (612bp) was amplified from the human genomic DNA using primers *SRYF1* (5'ATGCAATCATATGCTTCTGCTATG3') and *SRYR1* (5'CTACAGCTTTGTCCAGTGGCT3'). The *SRY* fragment was cloned into the Topo cloning pcr8 vector (pcr8-*SRY*). The ORF swap destination vector pmat:*SRYORF*:mat containing the *SRY* ORF under regulation of the *matA* promoter and terminator was created as a result of recombination reaction between the pcr8-*SRY* and pmatAORF swap vector²⁹. The human *SRY* HMG box was amplified using the primers *hSRYF2* (5'AGAGTGAACCGACCCATGAAC3') and *hSRYR2* (5'CCGACGAGGTCGATACTTATAATTC3'). The *An matA* sequence 5' of the HMG box was amplified by using AnmatAF10 (5'AGTGAGGGTAGAGTCTAC3') and

AnmatHMGswapR (5'P_iTGCGGGTTTCATTGACGCTGGC3'). The *An matA* sequence downstream of the HMG box was amplified using AnmatHMGswapF (5'P_iCCTTCTGAAAAGAAGCGTCGTG3') and AnmatAR10 (5' TGGAAGCTCCAGGAGCCGCTCGA3'). The three products were then ligated and the resultant ligation product amplified using AnmatAF26 (5'GGACGAAAAGCTTCTGCATTC3') and AnmatAR11 (5'TGCCGTATGCTACCTGAGC3'). The PCR product that corresponded to the predicted size of the chimeric gene was then topo cloned into pcr8. The resultant clone was purified and validated with DNA sequencing. The *matA/SRY* HMG box fusion gene was transferred into an *An pyrG* vector (pAn7) using the Invitrogen clonase system to create pmatHMGswap.

Transformation and targeted gene replacement in *A. nidulans*

Two independent ORF swap and multiple HMG box swap strains have been created. *A. nidulans* UI461 (*yA2, pabaA1, pyroA4, argB2, matA(0)::AfangB, nku::bar*) and UI 465 (*yA2, pyrG89, pabaA1, matA(0)::AfangB, nku::bar*) were used as the recipient strains. Plasmids: pmatAORF:SRYORF and ppyro were co-transformed into the UI461 strain. About 200 individual transformants that stably integrated the ppyro plasmid were recovered. All transformants were subsequently screened on YGMTV media for a non-conidiating phenotype indicative of arginine auxotrophy. Loss of arginine prototrophy indicates homologous recombination and a gene replacement event where the *mat:SRYORF:mat* fusion construct replaced the endogenous *matA(0)::AfangB*. Out of 200 transformants one *argB*⁻ colony was isolated; transformant T52. Gene replacement and correct integration of *matAORF: SRYORF* construct was confirmed by Southern blot, PCR and DNA sequencing analysis (Supplementary Fig. 1). Arginine auxotrophy impairs sexual reproduction, therefore the transformant T52 was subsequently transformed with the *pargB* plasmid to confer arginine prototrophy. Several, independent single ascospore isolates were also derived from viable ascospores produced by the original T52 *argB*⁺ strain. The ORF swap construct was also integrated ectopically into the UI465 strain carrying *matA (0)* at the resident location. Vector pmatHMGswap was used to create the HMG box swap strain by transforming UI465 strain. Routinely with all strains Southern blots, PCR and DNA sequencing were performed to identify and validate the transformants with a single copy of the plasmid integrated at *pyrG*. Multiple independent strains on different genetic backgrounds were determined for all constructs and tested.

Analysis of the sexual phenotype.

Sexual development of wild type *matA* (A1153), *matA(0)*UI461 and *mat:hSRY:mat* (UI470) was induced under standard culture condition as previously described^{29,16}. Development of hülle cells, protocleistothecia and cleistothecia were monitored over a two week period. Viability of ascospore fruiting bodies were routinely measured by plating spores on agar plates, incubating and counting growing colonies.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.


Acknowledgements

We thank Ms Heather Johnson for assistance in preparation of the manuscript. These experiments were performed in the Department of Biological Sciences at the University of Idaho. This study was supported by NSF 0318801 and NIH NCCR P20 RR015587 grants to BLM and NIH ES012974 grant to MKS.


References

1. Fraser JA, et al. Convergent evolution of chromosomal sex-determining regions in the animal and fungal kingdoms. *PLoS Biol.* 2004; 2:e384. doi:10.1371/journal.pbio.0020384. [PubMed: 15538538]
2. Fraser JA, Heitman J. Evolution of fungal sex chromosomes. *Mol Microbiol.* 2004; 51:299–306. doi:10.1046/j.1365-2958.2003.03874.x MMI3874 [pii]. [PubMed: 14756773]
3. Fraser JA, Heitman J. Chromosomal sex-determining regions in animals, plants and fungi. *Current opinion in genetics & development.* 2005; 15:645–651. doi:10.1016/j.gde.2005.09.002. [PubMed: 16182521]
4. Idnurm A, Walton FJ, Floyd A, Heitman J. Identification of the sex genes in an early diverged fungus. *Nature.* 2008; 451:193–196. doi:10.1038/nature06453. [PubMed: 18185588]
5. Kronstad JW, Staben C. Mating type in filamentous fungi. *Annu Rev Genet.* 1997; 31:245–276. doi: 10.1146/annurev.genet.31.1.245. [PubMed: 9442896]
6. Lahtz C, Pfeifer GP. Epigenetic changes of DNA repair genes in cancer. *Journal of molecular cell biology.* 2011; 3:51–58. doi:10.1093/jmcb/mjq053. [PubMed: 21278452]
7. Odell ID, Wallace SS, Pederson DS. Rules of engagement for base excision repair in chromatin. *Journal of cellular physiology.* 2013; 228:258–266. doi:10.1002/jcp.24134. [PubMed: 22718094]
8. Harley VR, Clarkson MJ, Argentaro A. The molecular action and regulation of the testis-determining factors, SRY (sex-determining region on the Y chromosome) and SOX9 [SRY-related high-mobility group (HMG) box 9]. *Endocrine reviews.* 2003; 24:466–487. doi:10.1210/er.2002-0025. [PubMed: 12920151]
9. Chen YS, Racca JD, Sequeira PW, Phillips NB, Weiss MA. Microsatellite-encoded domain in rodent Sry functions as a genetic capacitor to enable the rapid evolution of biological novelty. *Proc Natl Acad Sci U S A.* 2013; 110:E3061–3070. doi:1300860110 [pii] 10.1073/pnas.1300860110. [PubMed: 23901118]
10. Werner MH, Huth JR, Gronenborn AM, Clore GM. Molecular basis of human 46X,Y sex reversal revealed from the three-dimensional solution structure of the human SRY-DNA complex. *Cell.* 1995; 81:705–714. [PubMed: 7774012]
11. O'Flaherty E, Kaye J. TOX defines a conserved subfamily of HMG-box proteins. *BMC genomics.* 2003; 4:13. [PubMed: 12697058]
12. King CY, Weiss MA. The SRY high-mobility-group box recognizes DNA by partial intercalation in the minor groove: a topological mechanism of sequence specificity. *Proceedings of the National Academy of Sciences of the United States of America.* 1993; 90:11990–11994. [PubMed: 8265659]
13. Klass J, et al. The role of intercalating residues in chromosomal high-mobility-group protein DNA binding, bending and specificity. *Nucleic acids research.* 2003; 31:2852–2864. [PubMed: 12771212]
14. Phillips NB, et al. SRY and human sex determination: the basic tail of the HMG box functions as a kinetic clamp to augment DNA bending. *Journal of molecular biology.* 2006; 358:172–192. doi: 10.1016/j.jmb.2006.01.060. [PubMed: 16504207]
15. Chen YS, Racca JD, Phillips NB, Weiss MA. Inherited human sex reversal due to impaired nucleocytoplasmic trafficking of SRY defines a male transcriptional threshold. *Proc Natl Acad Sci U S A.* 2013; 110:E3567–3576. doi:1300828110 [pii] 10.1073/pnas.1300828110. [PubMed: 24003159]
16. Czaja W, Miller KY, Miller BL. Complex mechanisms regulate developmental expression of the matA (HMG) mating type gene in homothallic *Aspergillus nidulans*. *Genetics.* 2011; 189:795–808. doi:10.1534/genetics.111.131458. [PubMed: 21868608]

17. Zhao L, Koopman P. SRY protein function in sex determination: thinking outside the box. *Chromosome research : an international journal on the molecular, supramolecular and evolutionary aspects of chromosome biology*. 2012; 20:153–162. doi:10.1007/s10577-011-9256-x.
18. Bergstrom DE, Young M, Albrecht KH, Eicher EM. Related function of mouse SOX3, SOX9, and SRY HMG domains assayed by male sex determination. *Genesis*. 2000; 28:111–124. [PubMed: 11105052]
19. Margarit E, et al. Identification of conserved potentially regulatory sequences of the SRY gene from 10 different species of mammals. *Biochemical and biophysical research communications*. 1998; 245:370–377. doi:10.1006/bbrc.1998.8441. [PubMed: 9571157]
20. Su H, Lau YF. Identification of the transcriptional unit, structural organization, and promoter sequence of the human sex-determining region Y (SRY) gene, using a reverse genetic approach. *American journal of human genetics*. 1993; 52:24–38. [PubMed: 8434602]
21. Hossain A, Saunders GF. The human sex-determining gene SRY is a direct target of WT1. *J Biol Chem*. 2001; 276:16817–16823. doi:10.1074/jbc.M009056200 M009056200 [pii]. [PubMed: 11278460]
22. Miyamoto Y, Taniguchi H, Hamel F, Silversides DW, Viger RS. A GATA4/WT1 cooperation regulates transcription of genes required for mammalian sex determination and differentiation. *BMC Mol Biol*. 2008; 9:44. doi:1471-2199-9-44 [pii] 10.1186/1471-2199-9-44. [PubMed: 18445271]
23. Sato Y, Shinka T, Sakamoto K, Ewis AA, Nakahori Y. The male-determining gene SRY is a hybrid of DGCR8 and SOX3, and is regulated by the transcription factor CP2. *Mol Cell Biochem*. 2010; 337:267–275. doi:10.1007/s11010-009-0308-x. [PubMed: 19902333]
24. Sekido R, Lovell-Badge R. Genetic control of testis development. *Sex Dev*. 2013; 7:21–32. doi: 000342221 [pii] 10.1159/000342221. [PubMed: 22964823]
25. Matson CK, Zarkower D. Sex and the singular DM domain: insights into sexual regulation, evolution and plasticity. *Nat Rev Genet*. 2012; 13:163–174. doi:nrg3161 [pii] 10.1038/nrg3161. [PubMed: 22310892]
26. Lee N, Bakkeren G, Wong K, Sherwood JE, Kronstad JW. The mating-type and pathogenicity locus of the fungus *Ustilago hordei* spans a 500-kb region. *Proc Natl Acad Sci U S A*. 1999; 96:15026–15031. [PubMed: 10611332]
27. Metin B, Findley K, Heitman J. The mating type locus (MAT) and sexual reproduction of *Cryptococcus heveanensis*: insights into the evolution of sex and sex-determining chromosomal regions in fungi. *PLoS Genet*. 2010; 6:e1000961. doi:10.1371/journal.pgen.1000961. [PubMed: 20502678]
28. Jameson JL, Achermann JC, Ozisik G, Meeks JJ. Battle of the sexes: new insights into genetic pathways of gonadal development. *Trans Am Clin Climatol Assoc*. 2003; 114:51–63. discussion 64-55. [PubMed: 12813911]
29. Pyrzak W, Miller KY, Miller BL. Mating type protein Mat1-2 from asexual *Aspergillus fumigatus* drives sexual reproduction in fertile *Aspergillus nidulans*. *Eukaryot Cell*. 2008; 7:1029–1040. doi:EC.00380-07 [pii] 10.1128/EC.00380-07. [PubMed: 18245277]

**Figure 1.**

Structural amino acid comparison of MatA and SRY. **(A)** Mating type protein MatA is a structural homolog of human SRY. Pairwise alignment of the amino acids residues in MatA and SRY. The HMG box is shaded in gray. Color residue reflects physical-chemical nature of amino acids. Red (small, hydrophobic), Blue (acidic), Magenta (basic), Green (hydroxyl, amine, basic). Level of conservation is indicated; (*) residue is identical, (:) conserved substitutions, (.) semi-conservative substitutions. **(B)** The HMG box of fungal MatA is highly similar to the HMG box of human SRY. Pairwise sequence alignment of the HMG box from MatA and SRY HMG. Conserved and similar (+) residues are indicated in the center line. All residues marked in color represent key amino acids making direct contact with DNA. Amino acids marked in red are involved in binding to the specific DNA core sequence. Isoleucine (I) marked in green is involved in binding the minor groove of DNA. Asparagine (N) marked in blue is typical of HMG boxes binding DNA in a sequence specific manner.

**Figure 2.**

Genetic manipulations of the *matA* locus. Schematic outline illustrating structure of *matA* in wild type *A.nidulans* and its derivative strains carrying the ORF or HMG box of SRY. The genomic, chromosomal (chr.) location of the ectopic fusion gene is indicated. 1. Structure of WT *matA* gene. 2. The HMG box of *matA* replaced with HMG box of SRY. 3. The ORF of *matA* replaced with the ORF of SRY. 4. Mutant carrying complete deletion of *matA*. Associated sexual cycle phenotypes are summarized. The elements of gene structure are distinguished by different color (legend), the distance in base pairs (bp) is shown.




Figure 3.

Human HMG box of SRY drives sexual cycle in *A. nidulans*. Column 1. Sexual differentiation in wild type *A. nidulans*. Mature ascospores (red arrow) Column 1-A. Individual, crushed fruiting body (black arrow) with released ascospores (red arrow) Column 1-B. Abundant, spherical fruiting bodies (purple-black balls) develop within the vegetative tissue among yellow conidiating conidiophores (black arrow) Column 1-C. 2. The HMG box of SRY supports a wild type sexual cycle. Abundant fruiting bodies Column 2-B,C. Representative image of wild type mature ascospores is shown for both WT and HMG box swap strains in Columns 1-A/ 2-A. Column 3. The SRY ORF partially restores sexual cycle. Scarce fruiting bodies form within clusters of hülle cells Column 3-B,C. Ascogenous tissue is mostly undeveloped Column 3-A (black arrow), with small number of mature and viable ascospores Column 3-A (red arrow). Column 4. The *matA (0)* complete knockout is sterile, does not produce fruiting bodies or ascospores Column 4-A, only clusters of hülle cells are formed Column 4-B,C (arrows).




Figure 4. Structural conservation of the promoter regions in *SRY* and *matA* genes. **A.** The putative HMG binding sequence (HB), sequence element similar to *hSRY* SE and functional *matA* SE have been identified in the proximal and distal region of *matA* promoter. **B.** Schematic representation showing the distribution and location of binding sites for transcription factors. Binding sites recognized by WT1, Gata4 and Cp2 are indicated. Hexagonal shape (silencing element, SE/"SE") indicates position of conserved but apparently non-functional sequence related to MHC1 silencer. Square shape (*matA* SE) indicates positions of the functional silencer elements in *A.nidulans*. Oval shape (HMG-box binding, HB) indicates sites for putative binding of SRY and MatA proteins. Arrows indicate transcriptional start sites. **C.** Alignment of the consensus DNA sequence for the putative regulatory elements identified in the promoter regions of *SRY* and *matA*. Level of homology and conservation is shown. Underlined bases within the silencer regions indicate binding sites for the silencer of MHC class I gene.