

UC Riverside

UC Riverside Previously Published Works

Title

Effects of single parenthood on mothers' behavior, morphology, and endocrine function in the biparental California mouse

Permalink

<https://escholarship.org/uc/item/56t5v69c>

Journal

Hormones and Behavior, 114

ISSN

0018-506X

Authors

Zhao, Meng

Harris, Breanna N

Nguyen, Catherine TY

et al.

Publication Date

2019-08-01

DOI

10.1016/j.yhbeh.2019.05.005

Peer reviewed

Highlights

1. Morphological, endocrine and behavioral effects of single motherhood on mothers are not well known.
2. We evaluated such effects in the monogamous, biparental California mouse.
3. Single mothers (SM) spent more time engaged in maternal behavior than paired mothers (PM).
4. SM had heavier livers and hearts, but smaller ovaries and uteri, than PM and non-breeding females.
5. SM had a flatter diel CORT rhythm and a smaller CORT stress response than non-breeding females.

Effects of single parenthood on mothers' behavior, morphology, and endocrine function in the
biparental California mouse

Meng Zhao^{1,a}, Breanna N. Harris², Catherine T.Y. Nguyen¹ and Wendy Saltzman^{1,*}

¹ Department of Evolution, Ecology, and Organismal Biology, University of California,
Riverside

² Department of Biological Sciences, Texas Tech University

Corresponding author:

Wendy Saltzman

Department of Evolution, Ecology, and Organismal Biology

University of California, Riverside

Riverside, CA 92521

Saltzman@ucr.edu

Phone: 951-827-6356

Fax: 951-827-4286

^a Current address: Department of Pathology, Stanford University

Abstract

Motherhood is energetically costly for mammals and is associated with pronounced changes in mothers' physiology, morphology and behavior. In ~5% of mammals, fathers assist their mates with rearing offspring and can enhance offspring survival and development. Although these beneficial consequences of paternal care can be mediated by direct effects on offspring, they might also be mediated indirectly, through beneficial effects on mothers. We tested the hypothesis that fathers in the monogamous, biparental California mouse (*Peromyscus californicus*) reduce the burden of parental care on their mates, and therefore, that females rearing offspring with and without assistance from their mates will show differences in endocrinology, morphology and behavior, as well as in the survival and development of their pups. We found that pups' survival and development in the lab did not differ between those raised by a single mother and those reared by both mother and father. Single mothers spent more time in feeding behaviors than paired mothers. Both single and paired mothers had higher lean mass and/or lower fat mass and showed more anxiety-like behavior in open-field tests and tail-suspension tests, compared to non-breeding females. Single mothers had higher body-mass-corrected liver and heart masses, but lower ovarian and uterine masses, than paired mothers and/or non-breeding females. Mass of the gastrointestinal tract did not differ between single and paired mothers, but single mothers had heavier gastrointestinal tract compared to non-breeding females. Single motherhood also induced a flattened diel corticosterone rhythm and a blunted corticosterone response to stress, compared to non-breeding conditions. These findings suggest that the absence of a mate induces morphological and endocrine changes in mothers, which might result from increased energetic demands of pup care and could potentially help maintain normal survival and development of pups.

24 Keywords: single mothers; maternal care; biparental care; morphology; corticosterone

25

26 **Highlights:**

27 1. Metabolic and behavioral effects of single motherhood on mothers have rarely been explored.

28 2. We tested for such effects in the monogamous, biparental California mouse.

29 3. Single mothers spent more time in feeding behaviors than paired mothers.

30 4. Single mothers had higher liver and heart masses, but lower ovarian and uterine masses, than
31 both paired mothers and non-breeding females.

32 5. Single motherhood induced a flattened diel corticosterone rhythm and a blunted
33 corticosterone response to stress, compared to non-breeding conditions.

34 Introduction

35 Maternal care is costly for female mammals and is associated with pronounced morphological,
36 physiological, and behavioral changes in mothers (Lonstein, 2007; Slattery and Neumann, 2008;
37 Speakman, 2008). In small mammals, for example, mothers often undergo declines in fat stores
38 during lactation. They can also experience organ remodeling, such as growth of the alimentary
39 tract and associated organs such as the liver and pancreas, which may be necessary to meet the
40 high demands of lactation (Jolicoeur et al., 1980; Kennedy et al., 1958). Consequences of the
41 costs of motherhood include reductions in thermogenesis and physical activity, loss of bone mass,
42 and disruption of sleep patterns (Speakman, 2008).

43 Mothers in some mammalian species also exhibit blunted hormonal, neural and
44 behavioral responses to stressors during late pregnancy and lactation (Brunton et al., 2008;
45 Lightman et al., 2001; Slattery and Neumann, 2008). Furthermore, pregnant and lactating rats,
46 mice and possibly humans exhibit reduced anxiety and fearfulness. After becoming mothers, for
47 example, rats (*Rattus norvegicus*) and house mice (*Mus spp.*) have reduced acoustic startle
48 responses, increased locomotion in open-field tests, increased time spent in the open arms of the
49 elevated plus maze and reduced fleeing from an intruder (Lonstein, 2007; Lonstein et al., 2014;
50 Slattery and Neumann, 2008).

51 Although most mammals are uniparental, with only mothers caring for their offspring,
52 approximately 5% are biparental, with fathers helping to rear their young (Kleiman and Malcolm,
53 1981). Depending on the species, paternal care in mammals can include a variety of behaviors,
54 including retrieving, defending, playing with, grooming and huddling offspring, as well as
55 providing them with food and a nest (Kleiman and Malcolm, 1981). In some species, such as the
56 California mouse (*Peromyscus californicus*), fathers spend as much time interacting with their

offspring as do mothers (Bester-Meredith et al., 1999; Woodroffe and Vincent, 1994). In other species, such as prairie voles (*Microtus ochrogaster*), fathers' paternal behavior may compensate for low levels of maternal care (Perkeybile et al., 2013): offspring receive more care from fathers when mothers display low care. Therefore, fathers in biparental species can reduce the energetic burden of motherhood.

Fathers in biparental species can also influence offspring survival and development. For example, in California mice, paternal care has been found to greatly enhance survival and development of pups both in the field and under energetically challenging lab conditions, such as low ambient temperature or having to work for food (Bredy et al., 2007; Cantoni and Brown, 1997; Dudley, 1974; Gubernick and Teferi, 2000; Gubernick et al., 1993; Wright and Brown, 2002). The absence of a father can also influence offspring's cognitive, emotional, and reproductive behavior. In California mice, permanent removal of the father on postnatal day 3 decreases spatial learning ability (Bredy et al., 2004), and in mandarin (*M. mandarinus*) and prairie voles, paternally deprived offspring are more anxious and less social on multiple measures (Cao et al., 2014; Jia et al., 2009; Wang et al., 2012; Yu et al., 2012).

Although the negative consequences of losing the father can be mediated by direct effects on offspring, they might also be mediated indirectly, through adverse effects on mothers. However, the effects of fathers on their mates have received very little study. Exceptions include studies in Djungarian hamsters (*Phodopus campbelli*), which found that removal of the mate might influence pup survival by altering mothers' thermoregulatory abilities (Scribner and Wynne-Edwards, 1994; Walton and Wynne-Edwards, 1997; Wynne-Edwards and Lisk, 1989), as well as a recent study in prairie voles, which showed that mothers losing their mates have altered emotionality, including increased anxiety-related behavior in the elevated plus maze and more

passive stress-coping behavior in the forced-swim test (Bosch et al., 2017).

The California mouse is monogamous and biparental in both the field and lab, and fathers engage in all the same parental behaviors (except nursing), and to a similar extent, as mothers (Gubernick and Alberts, 1987). Most research in the California mouse has focused on the biology of paternal behavior; much less has focused on maternal care (Bester-Meredith et al., 2017), and the effects of fathers on their mates have received virtually no attention.

In this study we examined the effects of parenthood, as well as the effects of mate loss, on California mouse mothers. We hypothesized that mothers rearing offspring without assistance from their mates would have poorer morphological, physiological and affective condition, as well as impaired survival and development of pups, compared to mothers housed with their mates. In addition, we hypothesized that these effects of single motherhood would be further influenced by environmental conditions. We predicted that both single mothers and mothers housed with their mates would have lower body and fat mass, and display altered profiles of corticosterone (CORT), a metabolically important, stress-responsive hormone, compared to non-breeding females; and that the effects in single mothers should be even more severe than those in mothers housed with their male mate. We further predicted that paired mothers, but not single mothers, would show less anxiety-like and depression-like behavior than non-breeding females, as seen in other species. Finally, we anticipated that these differences between single and paired mothers would be more pronounced, or might only occur, under energetically challenging conditions that increase the cost of motherhood.

Material and Methods

Animals

California mice were bred in our colony at the University of California, Riverside (UCR) and were descended from mice from the *Peromyscus* Genetic Stock Center (University of South Carolina, Columbia, SC, USA). Animals were housed in polycarbonate cages (44 × 24 × 20 cm) with aspen shavings as bedding and cotton as nesting material. Food (Purina 5001 Rodent Chow, LabDiet, Richmond, IN, USA) and tap water were provided *ad lib*. The colony was on a 14:10 light:dark cycle, with lights on at 05:00 h and lights off at 19:00 h. Room temperature was approximately 22° C, and humidity was about 57%. Cages were checked twice daily and changed weekly.

Sample sizes were based on prospective power analysis (G*Power 3; Faul et al., 2009), using the magnitude of group differences and the within-group standard errors observed in a previous study of California mice (Saltzman et al., 2015). All procedures used were in accordance with the *Guide for the Care and Use of Laboratory Animals* and were approved by the UCR Institutional Animal Care and Use Committee. UCR is fully accredited by the Association for Assessment and Accreditation of Laboratory Animal Care (AAALAC).

Experimental design

Mice were weaned at 27-31 days of age, prior to the birth of younger siblings, and housed in same-sex groups of 3-4 related and/or unrelated, age-matched individuals. At 92-111 days of age (102.6 ± 0.5 days, mean $\pm$ SE; young adulthood), 88 females were randomly assigned to two categories. Non-breeding females (NB, n=29) were housed with a male that had been vasectomized (see below) 7 days before pairing, and breeding females (n=59) were housed with an intact male. These latter males were vasectomized 16-21 days after pairing (13 to 32 days before birth of the first litter) to prevent postpartum pregnancies. Non-breeding

females were used to control for the effects of motherhood. Pair mates were no more closely related to each other than first cousins.

When its first litter was born, each breeding female was randomly assigned to one of two conditions: single mothers (SM, n=29), whose male mates were removed 1 day after the birth of the first litter, or paired mothers (PM, n=30), whose mates remained with them throughout the entire lactation period. All mothers (and NB, at a comparable time point) were then left undisturbed for 2-3 days until the beginning of the 25-day testing period (Fig. 1). At postpartum day 27 or 28, females and pups were weighed and their body composition was assessed (see below). Pups were also checked daily for eye opening (Harris et al., 2013; Zhao et al., 2018).

Approximately half of the pairs in each reproductive group (n=42: 14 NB, 14 PM, 14 SM) were housed under standard lab conditions throughout the experiment (CTRL). The remaining pairs (n=46: 15 NB, 16 PM, 15 SM) were housed from the time of pair formation in cages in which they were required to climb towers in order to gain access to food and water (CLIMB; see below; Fig. 2). Thus, we had a total of 6 experimental groups (3 reproductive groups × 2 housing conditions).

Climbing towers

A climbing-exercise paradigm modified from previous studies of house mice (Lionikas and Blizard, 2008; Mori et al., 2003), rats (Notomi et al., 2001) and California mice (Zhao et al., 2018) was used to subject the animals in the CLIMB condition to a mild physical challenge (Fig. 2). Mice were housed in cages with plastic lids connected to two wire mesh (mesh openings: ~5 mm × 5 mm) towers measuring ~8 cm in diameter and ~50 cm in height. Food was located at the top of one tower, and a water bottle was at the top of the other tower.

149

150 **Vasectomies**

151 Male mice were vasectomized using antiseptic techniques and standard surgical procedure

152 (Chauke et al., 2011). Briefly, mice were anesthetized with isoflurane gas using a vaporizer. The

153 incision region (1 cm above genital area) was shaved and sanitized, and a ventral midline

154 incision (approximately 1 cm) was made. The vas deferens was sutured and cut. All reproductive

155 structures were repositioned back in the abdominal cavity, the incision was closed with

156 absorbable sutures (Monocryl Suture 4-0 FS-2, Ethicon, San Angelo, TX) and the skin was

157 sealed using tissue glue (Vetbond Tissue Adhesive 1469SB, St. Paul, MN, USA).

158 Following surgery, males to be paired with non-breeding females were housed

159 individually in a standard cage for 7 days until pair formation. For breeding pairs, male and

160 female pair mates were housed for the first 7 days after surgery on opposite sides of a standard

161 cage divided in half by a steel mesh barrier, with food and water available in each half, to

162 prevent direct physical interaction during post-surgical recovery (Harris and Saltzman, 2013).

163 Males and females in non-breeding pairs were also housed in divided cages for 7 days at a

164 matched time point to control for any effects of separation from the pair mate. At the end of the

165 experiment, all females were euthanized and dissected to check for pregnancy. Two females

166 were found to be pregnant; therefore, their data were not used in any analyses.

167

168 **Measurements**

169 **Morphology**

170 **Body mass**

171 From the time of pair formation until the birth of the first litter in each breeding pair, or a

matched time point for NB, females were weighed to the nearest 0.01 g at 14:00-16:00 h twice weekly, at 3- to 4-day intervals, to assess overall health, habituate the animals to handling, and monitor pregnancies through patterns of mass gain. Females were also weighed at 14:00-16:00 h on test days 1, 4, 8, and 13, as well as at 11:40-12:30 h on test day 12 (Fig. 1).

Body composition (test days 1, 4, 8, 13)

Body composition was assessed with an EchoMRI-100 magnetic resonance whole-body analyzer (Echo Medical Systems, Houston, TX, USA) as previously described (Zhao et al., 2017, 2018). Each scan lasted approximately 90 s, and animals were not sedated or anesthetized. Lean and fat masses were computed both as absolute values and as percentages of total body mass (analyses using body mass as a covariate yielded similar results).

Euthanasia and organ collection (test day 14)

Females were decapitated between 13:30 and 16:00 h. Trunk blood was collected in a heparinized weighing boat, and organs [heart, liver, spleen, leg muscles (left and right triceps surae), ovaries (left and right), uteri, adrenal glands (left and right) and kidneys (left and right)] were removed and weighed as previously described (Zhao et al., 2017, 2018). At a later date, stomach, caecum, and large and small intestines were thawed, cut open, cleaned of internal contents, blotted, and weighed as previously described (Andrew et al., 2016).

Behavioral indicators of neophobia, anxiety, and depression

Novel-object test (test day 4)

As described above, mothers in some mammalian species, such as rats and mice, exhibit reduced

anxiety and fearfulness, as well as blunted responses to stressors, during late pregnancy and lactation (Brunton et al., 2008; Lightman et al., 2001; Slattery and Neumann, 2008). We evaluated whether single motherhood would further influence mothers' anxiety/fearfulness using the novel-object test, which is commonly employed to assess neophobia in small mammals (Chauke et al., 2012; Greggor et al., 2015; Sarowar et al., 2016). In this test, frequency and duration of an animal's interactions with an unfamiliar object are considered negative indices of neophobia and, correspondingly, of fear and/or anxiety (Hughes, 2007; Powell et al., 2004). Novel-object tests were administered between 14:00 h and 16:00 h. Each female's cage mate(s) (male mate and/or pups) were removed from the home cage, housed in a clean cage, and moved to a separate room. After 5 min, a golf ball (diameter: 4.8 cm) was placed in the corner of the home cage farthest from the female. Behavioral responses to the novel object were video-recorded for 5 min. Immediately after testing, the female was weighed, scanned for body composition (see above), and then reunited with its cage mate(s) in the home cage. Behavioral parameters scored included latency to approach to within 2 cm of the golf ball, bouts of approaching the ball, total duration of immobility, and total durations of time spent sniffing and touching the ball (Chauke et al., 2012).

Open-field test (test day 8)

The open-field test is a standard method for measuring anxiety-like behavior in rodents, based on findings that many pharmacological anxiolytic treatments reduce measures of anxiety-like behaviors in this paradigm (reviewed in Prut and Belzung, 2003). The amount of exploratory behavior and time spent in the center of the arena are considered to be negative measures of anxiety, while the amount of defecation is considered a positive measure (Archer, 1973; Flint et

al., 1995; Willis-Owen et al., 2007). We administered tests between 14:00 h and 16:00 h. The open-field arena was a 1.0 m × 1.0 m square with a height of 0.4 m, constructed of opaque black plastic and placed on a clean sheet of white butcher paper to enhance contrast between the arena floor and the darkly colored mice as previously described (Perea-Rodriguez et al., 2018). Tests were recorded by a GoPro HERO SESSION video camera (GoPro Inc., San Mateo, CA, USA) suspended above the arena. After each test, the arena was disinfected and the butcher paper was replaced. The open-field arena was located in an environmental chamber maintained at 1400 lux with two overhead white lights; temperature and humidity were maintained at ~23°C and ~70%, respectively. For each test, the female subject was placed in the center of the arena and video-recorded for 10 min. Immediately following testing, the female was weighed, scanned for body composition (see above), and then reunited with its cage mate(s) in the home cage.

Exploratory behaviors in the open-field arena were quantified using TopScanLite software (Clever Sys Inc., Reston, Virginia, USA). The arena was divided into two concentric regions in the software: an inner square, measuring 0.5 × 0.5 m, in the center of the arena, and an outer region extending 0.5 m from each wall to the perimeter of the inner square. Bouts of crossing the boundary between inner and outer regions, total distance moved, distance moved in each region, and duration of time spent in each region were determined. We also recorded the number of fecal boli produced during the 10 min of testing. Distance traveled, duration in the inner region of the arena, and number of crossing bouts were considered negative indicators of anxiety-like behavior, whereas number of fecal boli was considered a positive indicator (Gould et al., 2009).

Tail-suspension test (test day 12)

The tail-suspension test is widely used to evaluate depression-like behavior in rodents, with the duration of immobility considered a positive index of depression; numerous antidepressant medications reduce immobility in this paradigm (reviewed in Cryan et al., 2005, Yan et al., 2010). We used the tail-suspension test both to assess depression-like behavior and as an acute stressor for assessment of the CORT stress response.

Between 11:30 and 12:35 h, mice were suspended by their tails from a ring stand for 6 min and the duration of immobility was measured, as previously described (Zhao et al., 2017, 2018). Briefly, the ring stand was placed on an activity detector unit (MAD-1: Sable Systems International, Henderson, NV, USA) interfaced to a Macintosh computer equipped with an A-D converter and Warthog LabHelper software (<http://www.warthog.ucr.edu>). Activity was recorded every 0.004 s. Warthog LabAnalyst software (<http://www.warthog.ucr.edu>) was used for baseline correction and calculation of activity duration (Malisch et al., 2009). Duration of time spent immobile was interpreted as a positive indicator of depression-like behavior (Cryan et al., 2005). In addition, we recorded the number of fecal boli produced during the 6 min of testing as a positive indicator of anxiety.

Saccharin preference test (test day 6)

Preference for sweet solutions is frequently used as a measure of anhedonia, a common marker of depression. The relative amount of sweet solution consumed is considered a negative index of anhedonia (Moreau, 1997; Willner et al., 1992). This test has consistently demonstrated a decrease in consumption of, and preference for, highly palatable sweet solutions under chronically stressful conditions (Grippe et al., 2005; Papp et al., 1991; Strekalova et al., 2004). We characterized females' preference for saccharin (Sweet'N Low; Cumberland Packing Corp.,

264 Brooklyn, NY, USA) solution (0.2% w/v in water) vs. water as previously described (Zhao et al.,
265 2017, 2018). Briefly, each mouse's cage mate(s) were removed from the home cage at the time
266 of lights-off (19:00 h). The female remained alone in its home cage for 4 h, with access to
267 standard chow and two plastic syringes, one containing ~35 ml of water and the other containing
268 ~35 ml of 0.2% saccharin solution. Positions of the two types of liquid were randomly assigned.
269 The syringes were weighed immediately before and after the 4-h test period. The amount of
270 consumed saccharin solution divided by the overall liquid consumption was calculated as an
271 index of relative preference for saccharin solution.

272

273 **Home-cage behavioral observation (test days 5, 7, 9, 11)**

274 Females were video-recorded in their home cages under undisturbed conditions for 1 hour on test
275 days 5 (20:00-22:00 h, under red light), 7 (14:00-16:00 h), 9 (08:00-10:00 h), and 11 (02:00-
276 04:00 h, under red light). We quantified total duration of time spent in each of six mutually
277 exclusive behavior categories - feeding, drinking, nest building, sleeping (sedentary with eyes
278 closed or with face tucked into body and without pups underneath), resting (sedentary, not
279 associated with other behaviors), and maternal behaviors, including nursing (sedentary with
280 visible pups underneath), grooming pup, retrieving pup – using The Observer software (Noldus,
281 Wageningen, Netherlands).

282

283 **Blood collection (test days 2, 3, 10, 12)**

284 Blood was collected under baseline conditions on test days 2 (03:50-04:10 h), 3 (19:50-20:10 h)
285 and 10 (11:50-12:10 h) for analysis of baseline CORT concentrations across the diel cycle, and
286 immediately after tail-suspension tests (11:30-12:35 h) on test day 12 for assessment of the

CORT response to an acute stressor. Animals were anesthetized with isoflurane, and blood was collected from the retro-orbital sinus into heparinized capillary tubes within 3 min after either initial disturbance to the cage (baseline samples) or the end of the tail-suspension test (post-stress samples). Blood was centrifuged immediately for 12 min (13,300 rpm, 4 °C), and plasma was removed and stored at -80 °C.

Corticosterone radioimmunoassay

Plasma corticosterone concentrations were determined using a radioimmunoassay kit (cat. no. 07120103; MP Biomedicals, Solon, OH) previously validated for *P. californicus* (Chauke et al., 2011). The assay was run according to manufacturer's instructions, but the assay standard curve was extended down from 25 to 12.5 ng/ml (90-91% bound) and went to 1000 ng/ml (20-21% bound). Samples were diluted anywhere from 1:100 to 1:1600 in order to ensure that values were contained within the curve. All samples were run in duplicate. Intra- and inter-assay coefficients of variation, calculated using kit-provided controls, were 2.19% and 5.31% for the high control, and 9.48% and 7.82 for the low control, respectively.

Statistical analysis

Data were analyzed by analysis of covariance (ANCOVA) or repeated-measures analysis of variance (ANOVA) and Fisher's LSD for post hoc tests using SPSS. Age and/or other potentially relevant variables (see Results) were used as covariates. For each analysis, residuals were checked for skewness and visually inspected for outliers. Plasma CORT data were also tested for sphericity using Mauchly's test, and all other traits were tested for homogeneity of variance using Levene's test. Dependent variables were transformed as needed; however, data are presented as

non-transformed values in the text and figures for ease of interpretation. For paired (right and left) organs, values from the two organs were compared using a paired t-test and a Pearson correlation to gauge repeatability (Table 1), and mean values were used for subsequent analyses. For the breeding pairs, we also performed 2-way (reproductive group x housing condition) ANOVAs to determine if litter size, pups' age of eye opening, and body composition at weaning age differed between pups reared by single and paired mothers or between pups in the CTRL and CLIMB housing conditions.

Excluding such nuisance variables as age, this study generated 400 P-values, 96 of which were <0.05. These tests include a substantial amount of non-independence because the same individuals were measured for all traits, some traits were correlated, and many tests were inter-related. To compensate for non-independence in multiple related tests, we used the Adaptive False Discovery Rate procedure as implemented in PROC MULTTEST in SAS 9.4 (SAS Incorporated, Cary, NC, USA). Based on this procedure, the 71 smallest P-values would have adjusted P-values <0.05 (the highest being 0.016). For simplicity, all P-values reported in the text and figures are raw values, not adjusted for multiple comparisons. We calculated partial eta squared (η^2) effect sizes for all ANOVA and ANCOVA, and Cohen's d effect sizes for all pairwise comparisons with $P < 0.05$. We refer to P-values ≤ 0.016 and those between 0.016 and 0.05 with partial $\eta^2 > 0.06$ or Cohen's $d > 0.5$ as 'significant'. Effect size of $0.06 < \eta^2 < 0.14$ and $0.5 < d < 0.8$ is considered a medium effect, and $\eta^2 > 0.14$ $d > 0.8$ is considered a large effect (Cohen, 1988). All tests were two-tailed.

Results

Home-cage behavior

Each category of behavior (feeding, drinking, nest building, resting, sleeping, and maternal behaviors) was analyzed separately for each of the four time points / test days. Duration of feeding differed among reproductive groups ($P = 0.029$) only at 02:00-04:00 h (near the end of the active [dark] period; day 11), with SM spending more time feeding than NB ($P = 0.008$, $d = 0.70$). Feeding behavior did not differ among housing conditions and was not affected by a housing condition $\times$ reproductive group interaction at any time point. Duration of drinking, as well as nest building, did not differ reliably among reproductive groups, between housing conditions, or by a housing condition $\times$ reproductive group interaction at any time point.

Time spent resting differed among reproductive groups at all four time points (all $P < 0.005$), with NB spending more time resting than both PM (all $P < 0.020$, all $2.50 > d > 0.5$) and SM (all $P < 0.003$, all $2.00 > d > 0.5$). Time spent sleeping was also higher in NB than both PM ($P = 0.014$, $d = 0.52$) and SM ($P = 0.005$, $d = 0.65$) at 14:00-16:00 h.

No significant effects of reproductive group (SM vs. PM), housing condition or housing condition $\times$ reproductive group interaction were found on total duration of time spent in maternal behavior (nursing, grooming pup, retrieving pup) at any time point.

Behavioral indicators of neophobia, anxiety, and depression

Novel-object test

Neither housing condition, reproductive group, nor their interaction affected females' latency to approach to within 2 cm of the novel object (golf ball), number of approaches, duration of time spent sniffing and touching the novel object, or duration of immobility (Supplementary Table 1).

Open-field test

ANCOVA (with age as a covariate) revealed a significant difference among reproductive groups in number of fecal boli produced ($P = 0.014$), with both SM and PM producing more boli than NB ($P = 0.009$, $d = 0.69$; $P = 0.012$, $d = 0.62$, respectively; Fig. 3). Number of fecal boli was not influenced by a main effect of housing condition or a housing condition $\times$ reproductive group interaction.

Neither housing condition, reproductive group, nor their interaction affected females' distances traveled during the open-field test (total distance and distance within each region), duration of time spent in the inner or outer region, or number of crossing bouts (Supplementary Table 1).

Tail-suspension test

ANCOVA (with age and body mass as covariates) revealed a positive effect of body mass ($P = 0.012$) on the duration of immobility, but we found no significant differences between housing conditions or among reproductive groups, and the housing condition $\times$ reproductive group interaction was not significant.

As in the open-field test, number of fecal boli expelled differed significantly among reproductive groups, with both SM and PM producing more boli than NB ($P = 0.036$, $d = 0.76$; $P = 0.030$, $d = 0.73$, respectively). Number of fecal boli was not influenced by a main effect of housing condition or a housing condition $\times$ reproductive group interaction (Table 2; Fig. 3).

Saccharin-preference test

ANCOVA (with age as a covariate) showed that preference for saccharin solution did not differ among reproductive groups or between housing conditions, and was not affected by a housing

condition × reproductive group interaction (Supplementary Table 1).

Plasma corticosterone concentrations

Baseline corticosterone

Baseline CORT concentrations were analyzed via repeated-measures ANOVA with time of sample (04:00, 20:00, and 12:00 h) as a within-subjects factor, and housing condition and reproductive group as between-subjects factors. The data were not normally distributed, mainly because of 8 outliers (2 CTRL SM, 2 CTRL PM, 2 CTRL NB, and 2 CLIMB SM) for samples collected at 04:00 h (outlier values were ~10-fold higher than other samples, for no obvious biological reason). Therefore, data from the outliers were discarded, and the remaining data were log₁₀-transformed to meet normality assumptions; however, analysis of log₁₀-transformed data yielded similar results whether the outliers were included or not. Here we report results of analyses with the outliers omitted.

Plasma CORT concentrations (Fig. 4) varied across the three times of day (main effect of time: $F_{2,112} = 242.11$, $P < 0.001$) in a pattern that was influenced by both housing condition (time × housing condition interaction: $F_{2,112} = 3.76$, $P = 0.026$) and reproductive group (time × reproductive condition interaction: $F_{2,112} = 3.45$, $P = 0.011$). CORT levels were highest around the beginning of the active period (20:00 h, 1 h after lights-off) and lowest near the beginning of the inactive period (04:00 h, 1 h before lights-on). Neither the main effects of housing condition or reproductive group, the interaction between housing condition and reproductive group, nor the 3-way interaction between time, housing condition and reproductive group was significant.

Post-hoc tests for the interaction between time and housing condition revealed that at 04:00 h, females in standard cages (CTRL) had significantly higher CORT levels than females in

towered cages (CLIMB; $P = 0.009$, $d = 0.72$). Post-hoc tests for the interaction between time and reproductive condition revealed that at 20:00 h, both single mothers and paired mothers had lower CORT levels than non-breeding females ($P = 0.002$, $d = 0.95$; $P = 0.040$, $d = 0.52$, respectively). In addition, at 12:00 h, single mothers had higher baseline CORT levels than non-breeding females ($P = 0.030$, $d = 0.82$). No other pairwise comparisons were significant.

Corticosterone response to stress

Data were analyzed using repeated-measures ANOVA with stress condition (baseline CORT at 12:00 h, post-stress CORT [immediately after a 6-min tail-suspension test] at 12:00 h) as a within-subjects factor, and housing condition and reproductive group as between-subjects factors. The tail-suspension test significantly increased $\log_{10}$ -transformed plasma CORT levels (main effect of stress: $F_{1,66} = 345.43$, $P < 0.001$), and this effect differed among reproductive groups (stress $\times$ reproductive group interaction: $F_{66} = 4.48$, $P = 0.015$; Fig. 5). As described above, post-hoc tests indicated that baseline CORT levels at 12:00 h were significantly higher in single mothers than in non-breeding females. However, post-stress CORT levels were significantly lower in single mothers ($P = 0.011$, $d = 0.73$), but not paired mothers ($P = 0.130$), than in non-breeding females. None of the remaining main effects (housing condition, reproductive group) or interactions (stress $\times$ housing condition, stress $\times$ housing condition $\times$ reproductive group) were significant.

Morphology

Body mass

We analyzed each individual body-mass value during the test period (days 1, 4, 8, 12 and 13) and

also the mean postpartum body mass from all test days except day 12. Body mass on day 12 was excluded because it was measured immediately after the tail-suspension test; however, analyses that included test day 12 generated similar results. ANCOVA (with age as a covariate) found no main effect of housing condition, reproductive group, or interaction between housing condition and reproductive group for either mean body mass or body mass on test days 1, 4, 8 or 13. On test day 12, body mass differed among reproductive groups ($P = 0.046$); post-hoc tests revealed that SM had significantly higher body mass than NB ($P = 0.014$, $d = 0.75$; Table 2).

Body composition

Neither rank-transformed fat mass on any test day nor mean fat mass differed between housing conditions or among reproductive groups, when using age as a covariate. Similar results were found when mean or individual fat mass was expressed as percentage of total body mass, except that rank-transformed percent fat mass on test days 8 and 13 showed a main effect of reproductive group ($P = 0.014$ and $P = 0.031$ respectively), with SM having lower percent fat mass than both PM ($P = 0.043$, $d = 0.64$) and NB ($P = 0.005$, $d = 0.74$) on test day 8, and SM having lower percent fat mass than NB ($P = 0.011$, $d = 0.66$) on test day 13. Note that mean and individual fat mass, as well as percent fat mass, had significant heterogeneity of variance that was not eliminated by transforms (with NB having higher variability than SM and PM), but the differences among groups are evident.

Square root-transformed mean postpartum lean mass differed among reproductive groups ($P = 0.029$; Fig. 6) when analyzed by ANCOVA with age as a covariate, with SM having higher lean mass than NB ($P = 0.008$, $d = 0.74$). Lean mass also differed among reproductive groups for test days 4 and 8 ($P = 0.003$ and $P = 0.001$, respectively). On each of these days, both SM and

PM had significantly higher lean mass than NB ($P = 0.001$, $d = 1.02$; $P = 0.016$, $d = 0.70$, respectively). No differences were found among groups for test day 1 or 13.

When expressed as a percentage of total body mass, mean percent lean mass did not differ among reproductive groups. However, percent lean mass on test days 4 and 13 differed among reproductive groups ($P = 0.008$ and $P = 0.050$, respectively), with SM having higher percent lean mass than NB ($P = 0.002$, $d = 0.84$) on test day 4, and both SM and PM having higher percent lean mass than NB ($P = 0.042$, $d = 0.58$; $P = 0.022$, $d = 0.58$, respectively) on test day 13. Percent lean mass did not differ among groups on any other day. Neither lean mass nor percent lean mass was influenced by a main effect of housing condition or by a housing condition $\times$ reproductive group interaction (Supplementary Table 1).

Organ masses

Body mass was used as a covariate in all organ-mass analyses. For all paired organs (triceps surae, ovaries, adrenals and kidneys), the two organs showed high correlations within individual animals, similar to findings in male California mice (Andrew et al., 2017; de Jong et al., 2013; Zhao et al. 2017, 2018). Right kidneys were significantly heavier than left kidneys (Table 1), as has been seen in other mammals and with other organs (e.g. Coleman et al., 1998; Idelman, 1978), while right adrenals were significantly lighter than left adrenals. Three females from the CLIMB condition had much higher triceps surae mass than the other CLIMB females and were excluded from analysis as outliers.

ANCOVA (with age and body mass as covariates) found significant positive effects of body mass (all $P < 0.001$) and significant effects of reproductive group on mass of the heart ($P = 0.023$), liver ($P < 0.001$), stomach ($P = 0.003$), caecum ($P = 0.005$), large intestine ($P = 0.006$),

small intestine ($P = 0.014$), and uterus ($P = 0.004$), and on mean mass of the triceps surae ($P = 0.008$) and ovaries ($P = 0.002$) (Table 2; Fig. 6; Fig. 7). No main effects of housing condition or housing condition $\times$ reproductive group interactions were found.

Post-hoc tests revealed that SM had significantly higher heart mass ($P = 0.006$, $d = 1.14$) and caecum mass ($P = 0.001$, $d = 1.33$), and lower rank-transformed triceps surae mass, than NB ($P = 0.002$, $d = 0.25$). Log_{10} -transformed liver mass was higher in SM than in both PM ($P = 0.046$, $d = 0.61$) and NB ($P < 0.001$, $d = 1.32$) and was higher in PM than NB ($P = 0.023$, $d = 0.66$). NB had lower stomach mass, large intestine mass, and small intestine mass compared to both SM ($P = 0.001$, $d = 1.33$; $P = 0.037$, $d = 0.76$; $P = 0.014$, $d = 1.07$, respectively) and PM ($P = 0.014$, $d = 0.82$; $P = 0.002$, $d = 0.91$; $P = 0.009$, $d = 0.96$, respectively). Finally, SM had lower uterus mass than PM ($P = 0.001$, $d = 0.90$) and lower ovary mass than both PM ($P = 0.003$, $d = 0.66$) and NB ($P = 0.001$, $d = 0.70$). No other organ masses or lengths differed among reproductive groups or between housing conditions, and no significant or significant interactions were found (Supplementary Table 1).

Pup development

Neither litter size at weaning (range: 1–3 pups, mean $\pm$ SE = 1.9 ± 0.1) nor day of eye opening of first pup in each litter (16.4 ± 0.4 days) differed between offspring of single mothers and paired mothers or between the CTRL and CLIMB conditions. We also found no difference in pups' body mass or body composition at the time of weaning (postpartum day 27 or 28) (Table 3).

Discussion

In this study, we tested the hypothesis that mate loss would lead to poorer morphological,

physiological and affective condition in California mouse mothers, as well as impaired survival and development of pups, compared to mothers housed with their mates. We tested this hypothesis under both standard laboratory conditions and mildly challenging conditions, in which mice had to climb wire-mesh towers to obtain food and water. Because females housed in towered cages had to spend more time and energy to climb and forage for food, we expected to find more severe effects of being single under the challenging condition. We found that mothers that lost their mates shortly after giving birth showed more morphological, physiological and behavioral differences from non-breeding females than did females whose male mates were present throughout the lactational period; however, these effects were not modulated by housing condition.

Effects of housing condition

Housing under challenging conditions had a moderate effect on baseline CORT concentrations. As previously found in California mice (Harris et al., 2012) and many other species (Dallman et al., 1987; Dickmeis, 2009), CORT levels were highest around the start of the active period (20:00 h), intermediate midway through the active period (12:00 h), and lowest at the end of the active period (04:00 h). Although this pattern was seen in virtually all females in the three reproductive groups, mice in standard cages had significantly *higher* plasma CORT levels than those in towered cages at 04:00 h. A previous study in our lab showed that chronic variable stress could increase baseline CORT levels in male California mice at 09:00 h (Harris et al., 2013). In contrast, housing in towered cages in the present study did not increase CORT levels. Instead, the mice housed in standard cages had a higher CORT trough compared to the ones in towered cages, causing a more flattened pattern of the CORT rhythm. An increase in glucocorticoid

levels at the circadian trough is commonly seen under chronic stress, which, in humans, might facilitate development of the metabolic syndrome (Dallman et al., 2000) and depression (Meijer et al., 1997) in the long term. However, it is unclear what the more dynamic rhythm means in mice housed in towered cages.

However, we found no difference in the preference for saccharine between females housed in towered cages and those housed under standard conditions. Low preference for sweetened water or other rewarding substances may be an indicator of anhedonia, a component of depression-like behavior (Schrader, 1997). Stressful environmental conditions, such as restricted access to food, continuous overnight illumination, cage tilt, and intermittent white noise, can decrease responsiveness to rewards in lab rats (Willner et al., 1992) and mice (Strekalova et al., 2004). Environmental enrichment, similarly, can influence animals' preference for rewarding stimuli. For example, rats housed in enriched environments show reduced cocaine self-administration (Puhl et al., 2012). It is possible that the required amount of climbing in the towers in our study might not have been energetically demanding, in view of the California mouse's semi-arboreal lifestyle in the wild (Ribble, 1992). However, whether the towered cages in our study acted as a stressor or as enrichment is not clear. In any case, we found no other effects of towered cages, aside from changes in baseline CORT, suggesting that this housing paradigm is neither highly stressful nor particularly enriching for California mice. In a recent study of male California mice, we found other effects of being housed in these towered cages (Zhao et al., 2018), including increases in body mass and body fat, but those animals were also food-restricted every third day, which likely accounted for the effects of housing conditions.

Effects of motherhood

Compared to non-breeding females, both single mothers and paired mothers produced significantly more fecal boli in the open-field and tail-suspension tests, which is considered a positive indicator of anxiety level (Archer, 1973; Bronikowski et al., 2001; Colman et al., 2007; Flint et al., 1995). Motherhood in several uniparental mammals influences females' responses to stress: mothers exhibit blunted hormonal, neural and behavioral responses to a multitude of stressors during pregnancy and lactation (Brunton et al., 2008; Lightman et al., 2001; Slattery and Neumann, 2008), including reduced anxiety and fearfulness (Lonstein, 2007; Slattery and Neumann, 2008), potentially to protect maternal care from being inhibited by stress (Brunton et al., 2008). In contrast, we found evidence that mothers were more anxious than non-breeding females. This might indicate a difference among rodent species or perhaps between uniparental and biparental animals, but more studies on other biparental species are needed to test this hypothesis. Alternatively, since we found no other effects of motherhood on behavior in the novel-object, open-field or tail-suspension tests, the difference in production of fecal boli might reflect changes in bowel function associated with lactation (Elias and Dowling, 1976; Hammond, 1997). Indeed, mothers in our study had larger intestines, caeca, and stomachs, which could be associated with the differences in production of fecal boli.

Mothers in our study, both paired and single, had higher lean mass on test days 4, and 8 than non-breeding females. We also found lower percent fat and higher percent lean masses in single mothers and, in some cases, in paired mothers than in non-breeding females on one or more test days. Although no measures of body composition differed significantly among reproductive groups on test day 1 (3-4 days after the mothers gave birth), the differences were mostly significant beginning on test day 4, indicating that lactation increased mothers' lean mass while decreasing fat stores, but that this effect did not become apparent within the first day after

parturition.

Energy balance in female mammals is strongly influenced by reproduction; females have high energy demands during pregnancy and, especially, lactation (Boland et al., 2001; Speakman, 2008). Motherhood can induce changes in body composition by increasing utilization of fat stores, especially in capital breeders, which rely primarily on fat stores to meet the energetic demands of reproduction (Bonnet et al., 1998). Our observed effects of motherhood on body fat and lean content in California mice are similar to those in capital breeders. Whether California mice are capital or income breeders (increase food intake to meet higher energetic demands) has yet to be investigated, but one study suggested that females in the congeneric species *P. leucopus* (white-footed mouse) are likely to be income breeders (Millar, 1975).

Effects of single motherhood

In small mammals, mothers have high basal and/or resting metabolic rate and experience growth of the alimentary tract in order to meet the high energy demands of lactation (Speakman, 2008). Reproduction has been found to induce a higher mass in liver, stomach, and large and small intestine compared to nonreproducing groups (Sadowska et al., 2018). Although we did not measure metabolic rate in this study, we found that single mothers had significantly greater heart mass than non-breeding females. Heart mass is usually positively correlated with basal metabolic rate (Chappell et al., 1999; Daan et al., 1990; Meerlo et al., 1997), daily energy expenditure (Dlugosz et al., 2012), and maximal oxygen consumption (Andrew, 2017) in vertebrates. Measures of mothers' metabolic rates in future studies would be helpful to further clarify the physiological effects of single motherhood.

Liver is a key organ that governs glucose and lipid metabolism as well as numerous other

functions (Samuel and Shulman, 2016). In our study, mothers had heavier livers than non-breeding females, consistent with findings from other species that lactation can induce hepatic growth (Speakman, 2008). In addition, we found that liver mass was higher in single mothers than in paired mothers, which might indicate that single mothers have higher energy demands during lactation than paired mothers (Piersma, 1997). However, we did not find significant differences in mass of any part of the gastrointestinal tract (stomach, caecum, large intestine, or small intestine) between single and paired mothers, although both groups had significantly higher values than non-breeding females. Thus, elevated liver mass in mothers, especially single mothers, might be associated with differences in functions separate from the gastrointestinal tract, such as lipid and carbohydrate metabolism, sulfation or bile formation (Gebhardt, 1992). Alternatively, although both heart mass and liver mass can correlate positively with metabolic rate and energetic demands, thickened heart muscle and liver enlargement can also have pathological causes and/or consequences, such as hypertension and fatty liver disease, in both humans and lab rodents (Samuel and Shulman, 2017; van Nierop et al., 2013). In addition, heart and liver can influence each other's functioning, and many metabolic factors can affect both organs simultaneously (Møller and Bernardi, 2013). Thus, the proximate causes and consequences of heart and liver enlargement in single mothers are not clear.

Triceps surae is an important muscle in running behavior, and the mass of gastrocnemius and/or triceps surae typically is positively related to sprinting performance in mice (Dohm et al., 1994; Syme et al., 2005). Research in exercise physiology indicates that voluntary exercise (e.g., wheel-running) in mice leads to increases in skeletal muscle masses (Soffe et al., 2016). Single mothers in our study had significantly lower triceps surae mass than non-breeding females, possibly due to changes in their behavior, such as less locomotion and more huddling with their

pups.

We did not observe differences in triceps surae mass between females in standard and towered cages. In house mice, towers similar to the ones used in the present study have revealed a strain $\times$ exercise interaction in the forelimb biceps brachii muscle, but no effect of climbing exercise was found on hindlimb muscles (Lionikas and Blizard, 2008; Mori et al., 2003). In contrast, a similar study found that hindlimb muscles of rats housed in towered cages were heavier than those of control rats after 4 weeks of exercise, but not after 8 weeks of exercise (Notomi et al., 2001). The different findings from rats, house mice, and California mice might be accounted for by possible species differences in behavioral responses to the towers and/or by differences in the duration of housing in towered cages.

Single mothers in our study had significantly lower ovarian and uterine masses compared to paired mothers, possibly due to lack of stimulation from a (vasectomized) male mate. Masses of reproductive organs in females are commonly used as indicators of reproductive function (Kumar et al., 2000; Kumar, 2005; Lee et al., 1998) and can be influenced by several reproductive hormones (e.g. luteinizing hormone, follicle-stimulating hormone, estrogen; Dewailly et al., 2006; Dupont et al., 2000; Halpin and Charlton, 1988; Kumar et al., 1997). Secretion of these hormones, in turn, can be elicited by interactions with males, including mating behaviors (Edwards, 1970). In some rodent species, for example, mating with a sterile male can induce pseudopregnancy. Although pseudopregnancy has not been studied in California mice, it is possible that paired mothers and non-breeding females, but not single mothers, might have been pseudopregnant during the period of data collection. Additionally or alternatively, reduced ovarian and uterine masses in single mothers might have resulted more directly from increased energetic demands on these females and shunting of metabolic resources to other, more essential

organs such as heart and liver. Indeed, energetic demands can greatly influence reproductive function in female mammals at the level of the ovaries and uterus through changes in lipid metabolism (Bellefontaine and Elias, 2014; Fontana and Torre, 2016).

Single and paired mothers showed differences in duration of feeding behaviors in the home cage. Single mothers, but not paired mothers, spent more time feeding than non-breeding females at 02:00-04:00 h, indicating that single mothers may need higher food intake to meet the higher energetic demands of caring for offspring without assistance from a mate.

We found no effects of single motherhood on behavior in the novel-object, open-field or tail-suspension tests, compared to paired mothers. In contrast, a recent study in prairie voles found that mothers whose male mates were removed on post-pairing day 18, prior to the birth of their first litter, showed altered emotionality, including increased anxiety-related behavior in the elevated plus maze and more passive stress-coping in the forced-swim test, which might result from increased signaling in the brain's corticotropin-releasing factor system (Bosch et al., 2017). This might reflect a difference between California mice and prairie voles. The timeline regarding the male-female separation might also explain the different findings in the two studies – while single mothers in the present study were separated from their male mate on the first day after parturition, approximately 35 days after pair formation, the prairie vole mothers were separated from their mate prior to giving birth. The earlier and longer separation from their mate might have been more stressful for the prairie vole mothers, which might have resulted in more severe emotional effects of single motherhood.

Corticosterone

Along with many other hormonal changes during lactation, postpartum females in

uniparental rodents and humans can have elevated baseline glucocorticoid levels in order to maintain milk production and increase food intake, facilitating shifts in feeding behavior and energy utilization to meet increased metabolic demands (Tu et al., 2005; Woodside et al., 2012). In the present study, we found that single mothers had higher baseline CORT levels at 12:00 h than non-breeding females. However, at 20:00 h (around the peak of the CORT rhythm), both single and paired mothers had significantly lower CORT levels than non-breeding females, which generated a more flattened CORT rhythm in mothers. Interestingly, lactating rats also have a more flattened CORT rhythm than virgin females, although the flattening in that species occurs through an elevation of the nadir CORT level (Walker et al., 1992).

In addition to baseline CORT concentrations, CORT responses to an acute stressor differed among reproductive groups. A 6-minute tail-suspension test midway through the active period markedly elevated plasma CORT levels in all animals, as expected, but this effect was blunted in single mothers, compared to non-breeding females: single mothers had significantly higher CORT levels under baseline conditions, but significantly lower CORT levels immediately after the stressor. Neither group differed significantly from paired mothers either in baseline CORT levels at 12:00 or in stress-induced CORT levels, consistent with a previous study in our lab (Chauke et al., 2011). The effect of single motherhood on CORT responses to an acute stressor, compared to non-breeding females, contrasts markedly with the lack of effects on behavioral indices of emotionality, as described above.

Lactating females in several species (e.g., rats, sheep) are less likely than non-lactating females to exhibit glucocorticoid elevations in response to stressful stimuli (Tu et al., 2005). On the other hand, a study of a biparental primate, the common marmoset (*Callithrix jacchus*), found no differences in cortisol responses to acute stressors between paired, reproductively

experienced females that were and were not lactating (Saltzman and Abbott, 2011). Our findings of blunted CORT responses in single but not paired mothers might indicate that, with help from the male mate, motherhood in biparental species differs in fundamental ways from motherhood in uniparental species, therefore not requiring marked changes in functioning of the hypothalamic-pituitary-adrenal (HPA) axis. Removal of the mate might increase the demands on mothers in biparental species, thereby mimicking the situation in uniparental species and eliciting changes in HPA activity, similar to findings in several bird species (Wingfield and Sapolsky, 2003). For example, removal of the mate might alter mothers' food intake, activity levels, sleep patterns, thermoregulatory abilities (Walton and Wynne-Edwards, 1997), or patterning of nursing bouts, all of which could potentially affect baseline CORT levels and responses to stress.

Pups

None of the measures that we monitored in pups – survival, age of eye opening, and body mass and body composition at the age of weaning – differed between pups reared by single mothers and those reared by both mother and father. This result indicates that single mothers were able to provide a similar quality of parental care and raise a similar quality and number of offspring, compared to paired mothers, both under standard housing conditions and when housed in towered cages. However, it is possible that effects of having a single mother on the offspring could be revealed under more challenging conditions. Losing fathers in California mice has been found to impair pup survival and development in the field and in energetically challenging laboratory settings, but not under standard lab conditions (Bredy et al., 2007; Cantoni Brown, 1997; Dudley, 1974; Gubernick and Teferi, 2000; Gubernick et al., 1993; Wright and Brown,

2002). Therefore, it appears that mothers in this species can compensate for the loss of a mate by providing more maternal care, under benign living conditions but not under demanding conditions, in order to maintain offspring survival and quality. In contrast, in prairie voles, single mothers showed no evidence of strong parental compensation in response to the absence of the father, indicating a minimal effect of family structure on maternal behavior but a large effect on the total amount of pup care (Ahern et al., 2011; Bosch et al., 2017); however, effects on pups' survival, development and body composition were not reported in the same studies.

Conclusion

In conclusion, this was one of the first studies to evaluate effects of loss of a mate on mothers in a biparental mammal. Effects of single motherhood on mothers' behavior, morphology, and physiology suggest that rearing offspring alone is energetically costly and can induce organ remodeling and changes in HPA activity, even when animals are housed under benign laboratory conditions with abundant food and water, mild temperature and humidity, and absence of predators. Although single mothers in the present study were able to produce a similar number and quality of offspring and showed no differences in behavioral indicators of emotionality compared to paired mothers, it is possible that more severe effects of being a single mother on the females, and/or effects of having a single mother on the offspring, could be revealed under more challenging or ecologically relevant conditions.

Acknowledgements

We thank the staff of the Spieth Vivarium and Dr. Akiko Sato for care of the animals, and Anthony Atalla, Emma Zschunke, Duc Huynh, May Chan, Michelle Phung, Dariana Chow,

1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904

724 Oanh Nguyen, Ethan Chan, Manparbodh Kaur, Alexis Rodriguez, Phuong-Anh Do, Nathan
725 Horrell and Melina Acosta for assistance with care of the animals and data collection. We are
726 also grateful to three anonymous reviewers for helpful comments on an earlier draft of the
727 manuscript. This research was supported by NSF IOS 1256572.

728
729

1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960

References

Ahern, T. H., Hammock, E. A. & Young, L. J., 2011. Parental division of labor, coordination, and the effects of family structure on parenting in monogamous prairie voles (*Microtus ochrogaster*). *Developmental Psychobiology* 53, 118-131.

Andrew, J. R., Saltzman, W., Chappell, M. A. & Garland, Jr. T., 2016. Consequences of fatherhood in the biparental California mouse (*Peromyscus californicus*), locomotor performance, metabolic rate, and organ masses. *Physiological and Biochemical Zoology* 89, 130-140.

Andrew, J.R., 2017. The physiological, exercise performance, and morphological consequences of fatherhood in the biparental California mouse (*Peromyscus californicus*). Ph.D. dissertation, University of California, Riverside.

Archer, J., 1973. Tests for emotionality in rats and mice: a review. *Animal Behaviour* 21, 205-235.

Bellefontaine, N. & Elias, C. F., 2014. Minireview: Metabolic control of the reproductive physiology: Insights from genetic mouse models. *Hormones and Behavior* 66, 7-14.

Bester-Meredith, J. K., Burns, J. N., Conley, M. F., Mammarella, G. E., Ng, N. D. (2017) *Peromyscus* as a model system for understanding the regulation of maternal behavior. In: *Seminars in Cell & Developmental Biology*, Vol. 61. pp. 99-106. Elsevier.

Bester-Meredith, J. K., Young, L. J. & Marler, C. A., 1999. Species differences in paternal behavior and aggression in *Peromyscus* and their associations with vasopressin immunoreactivity and receptors. *Hormones and Behavior* 36, 25-38.

Boland, M., Lonergan, P. & O'callaghan, D., 2001. Effect of nutrition on endocrine parameters, ovarian physiology, and oocyte and embryo development. *Theriogenology* 55, 1323-1340.

1961
1962
1963 753 Bonnet, X., Bradshaw, D. & Shine, R., 1998. Capital versus income breeding: an ectothermic
1964 perspective. *Oikos*, 333-342.
1965 754
1966
1967 755 Bosch, O. J., Pohl, T. T., Neumann, I. D. & Young, L. J., 2017. Abandoned prairie vole mothers
1968 show normal maternal care but altered emotionality: potential influence of the brain
1969 corticotropin-releasing factor system. *Behavioural Brain Research* 341, 114-121.
1970 756
1971
1972 757
1973
1974 758 Bredy, T. W., Brown, R. E. & Meaney, M. J., 2007. Effect of resource availability on biparental
1975 care, and offspring neural and behavioral development in the California mouse
1976 759
1977 (*Peromyscus californicus*). *European Journal of Neuroscience* 25, 567-575.
1978 760
1979
1980 761 Bredy, T. W., Lee, A. W., Meaney, M. J. & Brown, R. E., 2004. Effect of neonatal handling and
1981 paternal care on offspring cognitive development in the monogamous California mouse
1982 762
1983 (*Peromyscus californicus*). *Hormones and Behavior* 46, 30-38.
1984 763
1985
1986 764 Bronikowski, A. M., Carter, P. A., Swallow, J. G., Girard, I. A., Rhodes, J. S. & Garland Jr, T.,
1987 2001. Open-field behavior of house mice selectively bred for high voluntary wheel-
1988 running. *Behavior Genetics* 31, 309-316.
1989 765
1990
1991 766
1992
1993 767 Brunton, P., Russell, J. & Douglas, A., 2008. Adaptive responses of the maternal
1994 hypothalamic-pituitary-adrenal axis during pregnancy and lactation. *Journal of*
1995 768
1996 *Neuroendocrinology* 20, 764-776.
1997 769
1998
1999 770 Cantoni, D. & Brown, R. E., 1997. Paternal investment and reproductive success in the
2000 California mouse, *Peromyscus californicus*. *Animal Behaviour* 54, 377-386.
2001 771
2002
2003 772 Cao, Y., Wu, R., Tai, F., Zhang, X., Yu, P., An, X., Qiao, X. & Hao, P., 2014. Neonatal paternal
2004 deprivation impairs social recognition and alters levels of oxytocin and estrogen receptor
2005 α mRNA expression in the MeA and NAcc, and serum oxytocin in mandarin voles.
2006 773
2007
2008 774
2009
2010 775
2011
2012
2013
2014
2015
2016

2017
2018
2019 776 Cohen, J., 1988. Statistical Power Analysis for the Behavioral Sciences. Routledge. New York,
2020
2021 777 NY.
2022
2023 778 Gebhardt R., 1992. Metabolic zonation of the liver: regulation and implications for liver function.
2024
2025 779 Pharmacology & Therapeutics 53, 275-354.
2026
2027 780 Chappell, M. A., Bech, C. & Buttemer, W. A., 1999. The relationship of central and peripheral
2028
2029 781 organ masses to aerobic performance variation in house sparrows. The Journal of
2030
2031 782 Experimental Biology 202, 2269-2279.
2032
2033 783 Chauke, M., de Jong, T. R., Garland, T. & Saltzman, W., 2012. Paternal responsiveness is
2034
2035 784 associated with, but not mediated by reduced neophobia in male California mice
2036
2037 785 (*Peromyscus californicus*). Physiology & Behavior 107, 65-75.
2038
2039 786 Chauke, M., Malisch, J. L., Robinson, C., de Jong, T. R. & Saltzman, W., 2011. Effects of
2040
2041 787 reproductive status on behavioral and endocrine responses to acute stress in a biparental
2042
2043 788 rodent, the California mouse (*Peromyscus californicus*). Hormones and Behavior 60,
2044
2045 789 128-138.
2046
2047 790 Coleman, M. A., Garland, T., Marler, C. A., Newton, S. S., Swallow, J. G. & Carter, P. A., 1998.
2048
2049 791 Glucocorticoid response to forced exercise in laboratory house mice (*Mus domesticus*).
2050
2051 792 Physiology & Behavior 63, 279-285.
2052
2053 793 Colman, R. J., Nam, G., Huchthausen, L., Mulligan, J. D. & Saupe, K. W., 2007. Energy
2054
2055 794 restriction-induced changes in body composition are age specific in mice. The Journal of
2056
2057 795 Nutrition 137, 2247-2251.
2058
2059 796 Cryan, J. F., Mombereau, C. & Vassout, A., 2005. The tail suspension test as a model for
2060
2061 797 assessing antidepressant activity: review of pharmacological and genetic studies in mice.
2062
2063 798 Neuroscience & Biobehavioral Reviews 29, 571-625.
2064
2065
2066
2067
2068
2069
2070
2071
2072

2073
 2074
 2075 799 Daan, S., Masman, D. & Groenewold, A., 1990. Avian basal metabolic rates: their association
 2076
 2077 800 with body composition and energy expenditure in nature. American Journal of
 2078
 2079 801 Physiology-Regulatory, Integrative and Comparative Physiology 259, R333-R340.
 2080
 2081 802 Dallman, M., Akana, S., Bhatnagar, S., Bell, M. & Strack, A., 2000. Bottomed out: metabolic
 2082
 2083 803 significance of the circadian trough in glucocorticoid concentrations. International
 2084
 2085 804 Journal of Obesity 24, S40.
 2086
 2087 805 Dallman, M. F., Akana, S. F., Cascio, C. S., Darlington, D. N., Jacobson, L. & Levin, N., 1987.
 2088
 2089 806 Regulation of ACTH secretion: variations on a theme of B. In: Proceedings of the 1986
 2090
 2091 807 Laurentian Hormone Conference. pp. 113-173. Elsevier.
 2092
 2093 808 De Jong, T. R., Harris, B. N., Perea-Rodriguez, J. P., & Saltzman, W., 2013. Physiological and
 2094
 2095 809 neuroendocrine responses to chronic variable stress in male California mice (*Peromyscus*
 2096
 2097 810 *californicus*): influence of social environment and paternal
 2098
 2099 811 state. Psychoneuroendocrinology, 38(10), 2023-2033.
 2100
 2101 812 Dewailly, D., Robin, G., Peigne, M., Decanter, C., Pigny, P. & Catteau-Jonard, S., 2016.
 2102
 2103 813 Interactions between androgens, FSH, anti-Müllerian hormone and estradiol during
 2104
 2105 814 folliculogenesis in the human normal and polycystic ovary. Human Reproduction Update
 2106
 2107 815 22, 709-724.
 2108
 2109 816 Dickmeis, T., 2009. Glucocorticoids and the circadian clock. Journal of Endocrinology 200, 3-22.
 2110
 2111 817 Dlugosz, E. M., Harris, B. N., Saltzman, W. & Chappell, M. A., 2012. Glucocorticoids, aerobic
 2112
 2113 818 physiology, and locomotor behavior in California mice. Physiological and Biochemical
 2114
 2115 819 Zoology 85, 671-683.
 2116
 2117
 2118
 2119
 2120
 2121
 2122
 2123
 2124
 2125
 2126
 2127
 2128

- 820 Dohm, M. R., Richardson, C. S. & Garland Jr, T., 1994. Exercise physiology of wild and
821 random-bred laboratory house mice and their reciprocal hybrids. American Journal of
822 Physiology-Regulatory, Integrative and Comparative Physiology 267, R1098-R1108.
- 823 Dudley, D., 1974. Contributions of paternal care to the growth and development of the young in
824 *Peromyscus californicus*. Behavioral Biology 11, 155-166.
- 825 Dupont, S., Krust, A., Gansmuller, A., Dierich, A., Chambon, P. & Mark, M., 2000. Effect of
826 single and compound knockouts of estrogen receptors alpha (ERalpha) and beta (ERbeta)
827 on mouse reproductive phenotypes. Development 127, 4277-4291.
- 828 Edwards, D. A., 1970. Induction of estrus in female mice: estrogen-progesterone interactions.
829 Hormones and Behavior 1, 299-304.
- 830 Elwood, R. & Broom, D., 1978. The influence of litter size and parental behaviour on the
831 development of Mongolian gerbil pups. Animal Behaviour 26, 438-454.
- 832 Flint, J., Corley, R., DeFries, J. C., Fulker, D. W., Gray, J. A., Miller, S. & Collins, A. C., 1995.
833 A simple genetic basis for a complex psychological trait in laboratory mice. Science 269.
- 834 Fontana, R. & Torre, S. D., 2016. The deep correlation between energy metabolism and
835 reproduction: a view on the effects of nutrition for women fertility. Nutrients 8, 87.
- 836 Gould, T. D., Dao, D. T., Kovacs, C. E., 2009. The open field test. In: Mood and Anxiety
837 Related Phenotypes in Mice: Characterization Using Behavioral Tests, Humana Press,
838 New York.
- 839 Greggor, A.L., Thornton, A., Clayton, N.S., 2015. Neophobia is not only avoidance: improving
840 neophobia tests by combining cognition and ecology. Current Opinion in Behavioral
841 Sciences 6, 82-89.

- 842 Grippo, A.J., Francis, J., Beltz, T.G., Felder, R.B., Johnson, A.K., 2005. Neuroendocrine and
843 cytokine profile of chronic mild stress-induced anhedonia. *Physiology & Behavior* 84,
844 697-706.
- 845 Gubernick, D. J. & Alberts, J. R., 1987. The biparental care system of the California mouse,
846 *Peromyscus californicus*. *Journal of Comparative Psychology* 101, 169.
- 847 Gubernick, D. J. & Teferi, T., 2000. Adaptive significance of male parental care in a
848 monogamous mammal. *Proceedings of the Royal Society of London. Series B: Biological*
849 *Sciences* 267, 147-150.
- 850 Gubernick, D. J., Wright, S. L. & Brown, R. E., 1993. The significance of father's presence for
851 offspring survival in the monogamous California mouse, *Peromyscus californicus*.
852 *Animal Behaviour* 46, 539-546.
- 853 Halpin, D. & Charlton, H., 1988. Effects of short-term injection of gonadotrophins on ovarian
854 follicle development in hypogonadal (hpg) mice. *Journal of Reproduction and Fertility* 82,
855 393-400.
- 856 Harris, B. N., de Jong, T. R., Yang, V. & Saltzman, W., 2013. Chronic variable stress in fathers
857 alters paternal and social behavior but not pup development in the biparental California
858 mouse (*Peromyscus californicus*). *Hormones and Behavior* 64, 799-811.
- 859 Harris, B. N. & Saltzman, W., 2013. Effect of reproductive status on hypothalamic–pituitary–
860 adrenal (HPA) activity and reactivity in male California mice (*Peromyscus californicus*).
861 *Physiology & Behavior* 112, 70-76.
- 862 Harris, B. N., Saltzman, W., de Jong, T. R. & Milnes, M. R., 2012. Hypothalamic–pituitary–
863 adrenal (HPA) axis function in the California mouse (*Peromyscus californicus*): Changes

864 in baseline activity, reactivity, and fecal excretion of glucocorticoids across the diurnal
865 cycle. *General and Comparative Endocrinology* 179, 436-450.

866 Hughes, R.N., 2007. Neotic preferences in laboratory rodents: issues, assessment and substrates.
867 *Neuroscience & Biobehavioral Reviews* 31, 441-464.

868 Idelman, S., 1978. The structure of the mammalian adrenal cortex. *General, Comparative and*
869 *Clinical*, 1-199.

870 Jia, R., Tai, F., An, S., Zhang, X. & Broders, H., 2009. Effects of neonatal paternal deprivation
871 or early deprivation on anxiety and social behaviors of the adults in mandarin voles.
872 *Behavioural Processes* 82, 271-278.

873 Jolicoeur, L., Asselin, J. & Morisset, J., 1980. Trophic effects of gestation and lactation on rat
874 pancreas. *Biomedical Research* 1, 482-488.

875 Kennedy, G., Pearce, W. M. & Parrott, D. M., 1958. Liver growth in the lactating rat. *Journal of*
876 *Endocrinology* 17, 158-160.

877 Kleiman, D. G. & Malcolm, J. R. (1981) The evolution of male parental investment in mammals.
878 In: *Parental Care in Mammals*. pp. 347-387. Springer.

879 Kumar, T. R., 2005. What have we learned about gonadotropin function from gonadotropin
880 subunit and receptor knockout mice? *Reproduction* 130, 293-302.

881 Kumar, T. R., Wang, Y., Lu, N. & Matzuk, M. M., 1997. Follicle stimulating hormone is
882 required for ovarian follicle maturation but not male fertility. *Nature Genetics* 15, 201.

883 Kumar, T. R., Wiseman, A. L., Kala, G., Kala, S. V., Matzuk, M. M. & Lieberman, M. W., 2000.
884 Reproductive defects in γ -glutamyl transpeptidase-deficient mice. *Endocrinology* 141,
885 4270-4277.

2297
2298
2299 886 Lee, H.-W., Blasco, M. A., Gottlieb, G. J., Horner II, J. W., Greider, C. W. & DePinho, R. A.,
2300
2301 887 1998. Essential role of mouse telomerase in highly proliferative organs. *Nature* 392, 569.
2302
2303 888 Lightman, S. L., Windle, R. J., Wood, S. A., Kershaw, Y. M., Shanks, N. & Ingram, C. D., 2001.
2304
2305 889 Peripartum plasticity within the hypothalamo-pituitary-adrenal axis. *Progress in Brain*
2306
2307 890 *Research* 133, 111-129.
2308
2309 891 Lionikas, A. & Blizard, D. A., 2008. Diverse effects of stanozolol in C57BL/6J and A/J mouse
2310
2311 892 strains. *European Journal of Applied Physiology* 103, 333-341.
2312
2313 893 Lonstein, J. S., 2007. Regulation of anxiety during the postpartum period. *Frontiers in*
2314
2315 894 *Neuroendocrinology* 28, 115-141.
2316
2317 895 Lonstein, J. S., Maguire, J., Meinschmidt, G. & Neumann, I. D., 2014. Emotion and mood
2318
2319 896 adaptations in the peripartum female: complementary contributions of GABA and
2320
2321 897 oxytocin. *Journal of Neuroendocrinology* 26, 649-664.
2322
2323 898 Meerlo, P., Bolle, L., Visser, G. H., Masman, D. & Daan, S., 1997. Basal metabolic rate in
2324
2325 899 relation to body composition and daily energy expenditure in the field vole, *Microtus*
2326
2327 900 *agrestis*. *Physiological Zoology* 70, 362-369.
2328
2329 901 Meijer, O., Van Oosten, R. & De Kloet, E., 1997. Elevated basal trough levels of corticosterone
2330
2331 902 suppress hippocampal 5-hydroxytryptamine_{1A} receptor expression in adrenally intact
2332
2333 903 rats: implication for the pathogenesis of depression. *Neuroscience* 80, 419-426.
2334
2335 904 Millar, J. S., 1975. Tactics of energy partitioning in breeding *Peromyscus*. *Canadian Journal of*
2336
2337 905 *Zoology* 53, 967-976.
2338
2339 906 Møller, S. & Bernardi, M., 2013. Interactions of the heart and the liver. *European Heart Journal*
2340
2341 907 34, 2804-2811.
2342
2343
2344
2345
2346
2347
2348
2349
2350
2351
2352

- 908 Moreau, J. L., 1997. Validation of an animal model of anhedonia, a major symptom of
909 depression. *Encephale* 23, 280-289.
- 910 Mori, T., Okimoto, N., Sakai, A., Okazaki, Y., Nakura, N., Notomi, T. & Nakamura, T., 2003.
911 Climbing exercise increases bone mass and trabecular bone turnover through transient
912 regulation of marrow osteogenic and osteoclastogenic potentials in mice. *Journal of Bone
913 and Mineral Research* 18, 2002-2009.
- 914 Notomi, T., Okimoto, N., Okazaki, Y., Tanaka, Y., Nakamura, T. & Suzuki, M., 2001. Effects of
915 tower climbing exercise on bone mass, strength, and turnover in growing rats. *Journal of
916 Bone and Mineral Research* 16, 166-174.
- 917 Perea-Rodriguez JP, Zhao M, Harris BN, Raqueno J, Saltzman W., 2018. Behavioral and
918 endocrine consequences of placentophagia in male California mice (*Peromyscus
919 californicus*). *Physiology & Behavior* 188:283-290.
- 920 Perkeybile, A. M., Griffin, L. & Bales, K. L., 2013. Natural variation in early parental care
921 correlates with social behaviors in adolescent prairie voles (*Microtus ochrogaster*).
922 *Frontiers in Behavioral Neuroscience* 7, 21.
- 923 Piersma T, Lindström Å., 1997. Rapid reversible changes in organ size as a component of
924 adaptive behaviour. *Trends in Ecology & Evolution* 12:134-138.
- 925 Piovanotti, M. R. A. & Vieira, M. L. s., 2004. Presence of the father and parental experience
926 have differentiated effects on pup development in Mongolian gerbils (*Meriones
927 unguiculatus*). *Behavioural Processes* 66, 107-117.
- 928 Powell, S.B., Geyer, M.A., Gallagher, D., Paulus, M.P., 2004. The balance between approach
929 and avoidance behaviors in a novel object exploration paradigm in mice. *Behavioural
930 Brain Research* 152, 341-349.

- 931 Prut, L., Belzung, C., 2003. The open field as a paradigm to measure the effects of drugs on
932 anxiety-like behaviors: a review. *European Journal of Pharmacology* 463, 3-33.
- 933 Puhl, M. D., Blum, J. S., Acosta-Torres, S. & Grigson, P. S., 2012. Environmental enrichment
934 protects against the acquisition of cocaine self-administration in adult male rats, but does
935 not eliminate avoidance of a drug-associated saccharin cue. *Behavioural Pharmacology*
936 23, 43-53.
- 937 Ribble, D. O., 1992. Lifetime reproductive success and its correlates in the monogamous rodent,
938 *Peromyscus californicus*. *Journal of Animal Ecology*, 457-468.
- 939 Sadowska J, Gębczyński AK, Konarzewski M., 2018. Long-term trait consistency in mice
940 selected for swim-induced high aerobic capacity. *Physiological and Biochemical Zoology*
941 91(4):925-932.
- 942 Saltzman, W. & Abbott, D. H., 2011. Hormonal and behavioral responses to stress in lactating
943 and non-lactating female common marmosets (*Callithrix jacchus*). *Physiology &*
944 *Behavior* 104, 446-453.
- 945 Saltzman, W., Harris, B. N., de Jong, T. R., Nguyen, P. P., Cho, J. T., Hernandez, M. & Perea-
946 Rodriguez, J. P., 2015. Effects of parental status on male body mass in the monogamous,
947 biparental California mouse *Journal of Zoology* 296, 23-29.
- 948 Samuel, V. T. & Shulman, G. I., 2016. The pathogenesis of insulin resistance: integrating
949 signaling pathways and substrate flux. *The Journal of Clinical Investigation* 126, 12-22.
- 950 Samuel, V. T. & Shulman, G. I., 2017. Nonalcoholic fatty liver disease as a nexus of metabolic
951 and hepatic diseases. *Cell Metabolism* 27, 22-41.

2465
2466
2467 952 Sarowar, T., Grabrucker, S., Föhr, K., Mangus, K., Eckert, M., Bockmann, J., Boeckers, T.M.,
2468
2469 953 Grabrucker, A.M., 2016. Enlarged dendritic spines and pronounced neophobia in mice
2470
2471
2472 954 lacking the PSD protein RICH2. *Molecular Brain* 9, 28.
2473
2474 955 Schrader, G., 1997. Does anhedonia correlate with depression severity in chronic depression?
2475
2476 956 *Comprehensive Psychiatry* 38, 260-263.
2477
2478 957 Scribner, S. J. & Wynne-Edwards, K. E., 1994. Thermal constraints on maternal behavior during
2479
2480 958 reproduction in dwarf hamsters (*Phodopus*). *Physiology & Behavior* 55, 897-903.
2481
2482 959 Slattery, D. A. & Neumann, I. D., 2008. No stress please! Mechanisms of stress
2483
2484 960 hyporesponsiveness of the maternal brain. *The Journal of Physiology* 586, 377-385.
2485
2486 961 Soffe, Z., Radley-Crabb, H., McMahon, C., Grounds, M. & Shavlakadze, T., 2016. Effects of
2487
2488 962 loaded voluntary wheel exercise on performance and muscle hypertrophy in young and
2489
2490 963 old male C57Bl/6J mice. *Scandinavian Journal of Medicine & Science in Sports* 26, 172-
2491
2492 964 188.
2493
2494 965 Speakman, J. R., 2008. The physiological costs of reproduction in small mammals. *Philosophical*
2495
2496 966 *Transactions of the Royal Society B: Biological Sciences* 363, 375-398.
2497
2498 967 Strekalova, T., Spanagel, R., Bartsch, D., Henn, F. A. & Gass, P., 2004. Stress-induced
2499
2500 968 anhedonia in mice is associated with deficits in forced swimming and exploration.
2501
2502 969 *Neuropsychopharmacology* 29, 2007-2017.
2503
2504 970 Syme, D. A., Evashuk, K., Grintuch, B., Rezende, E. L. & Garland Jr, T., 2005. Contractile
2505
2506 971 abilities of normal and “mini” triceps surae muscles from mice (*Mus domesticus*)
2507
2508 972 selectively bred for high voluntary wheel running. *Journal of Applied Physiology* 99,
2509
2510 973 1308-1316.
2511
2512
2513
2514
2515
2516
2517
2518
2519
2520

2521
2522
2523 974 Tu, M. T., Lupien, S. J. & Walker, C.-D., 2005. Measuring stress responses in postpartum
2524
2525 975 mothers: perspectives from studies in human and animal populations. *Stress* 8, 19-34.
2526
2527 976 van Nierop, B. J., van Assen, H. C., van Deel, E. D., Niesen, L. B., Duncker, D. J., Strijkers, G. J.
2528
2529 & Nicolay, K., 2013. Phenotyping of left and right ventricular function in mouse models
2530 977
2531 of compensated hypertrophy and heart failure with cardiac MRI. *PLoS One* 8, e55424.
2532 978
2533 Walker, C.-D., Lightman, S. L., Steele, M. K. & Dallman, M. F., 1992. Suckling is a persistent
2534 979
2535 stimulus to the adrenocortical system of the rat. *Endocrinology* 130, 115-125.
2536 980
2537 Walton, J. M. & Wynne-Edwards, K. E., 1997. Paternal care reduces maternal hyperthermia in
2538 981
2539 Djungarian hamsters (*Phodopus campbelli*). *Physiology & Behavior* 63, 41-47.
2540 982
2541 Wang, J., Tai, F., Yan, X. & Yu, P., 2012. Paternal deprivation alters play-fighting, serum
2542 983
2543 corticosterone and the expression of hypothalamic vasopressin and oxytocin in juvenile
2544 984
2545 male mandarin voles. *Journal of Comparative Physiology A* 198, 787-796.
2546 985
2547 Wang, Z. & Novak, M. A., 1992. Influence of the social environment on parental behavior and
2548 986
2549 pup development of meadow voles (*Microtus pennsylvanicus*) and prairie voles (*M.*
2550 987
2551 *ochrogaster*). *Journal of Comparative Psychology* 106, 163.
2552 988
2553 Willis-Owen, S.A., Flint, J., 2007. Identifying the genetic determinants of emotionality in
2554 989
2555 humans; insights from rodents. *Neuroscience & Biobehavioral Reviews* 31, 115-124.
2556 990
2557 Willner, P., Muscat, R. & Papp, M., 1992. Chronic mild stress-induced anhedonia: a realistic
2558 991
2559 animal model of depression. *Neuroscience & Biobehavioral Reviews* 16, 525-534.
2560 992
2561 Wingfield, J. & Sapolsky, R., 2003. Reproduction and resistance to stress: when and how.
2562 993
2563 *Journal of Neuroendocrinology* 15, 711-724.
2564 994
2565 Woodroffe, R. & Vincent, A., 1994. Mother's little helpers: patterns of male care in mammals.
2566 995
2567 *Trends in Ecology & Evolution* 9, 294-297.
2568 996
2569
2570
2571
2572
2573
2574
2575
2576

- Woodside, B., Budin, R., Wellman, M. K. & Abizaid, A., 2012. Many mouths to feed: The control of food intake during lactation. *Frontiers in Neuroendocrinology* 33, 301-314.
- Wright, H. W. Y., 2006. Paternal den attendance is the best predictor of offspring survival in the socially monogamous bat-eared fox. *Animal Behaviour* 71, 503-510.
- Wright, S. L. & Brown, R. E., 2002. The importance of paternal care on pup survival and pup growth in *Peromyscus californicus* when required to work for food. *Behavioural Processes* 60, 41-52.
- Wynne-Edwards, K. E. & Lisk, R. D., 1989. Differential effects of paternal presence on pup survival in two species of dwarf hamster (*Phodopus sungorus* and *Phodopus campbelli*). *Physiology & Behavior* 45, 465-469.
- Yan, H.-C., Cao, X., Das, M., Zhu, X.-H., Gao, T.-M., 2010. Behavioral animal models of depression. *Neuroscience Bulletin* 26, 327-337.
- Yu, P., An, S., Tai, F., Zhang, X., He, F., Wang, J., An, X. & Wu, R., 2012. The effects of neonatal paternal deprivation on pair bonding, NAcc dopamine receptor mRNA expression and serum corticosterone in mandarin voles. *Hormones and Behavior* 61, 669-677.
- Zhao, M., Garland, T., Chappell, M. A., Andrew, J. R., Harris, B. N. & Saltzman, W., 2018. Effects of a physical and energetic challenge on male California mice (*Peromyscus californicus*): modulation by reproductive condition. *Journal of Experimental Biology* 221, jeb168559.
- Zhao, M., Garland, T., Chappell, M. A., Andrew, J. R. & Saltzman, W., 2017. Metabolic and affective consequences of fatherhood in male California mice. *Physiology & Behavior* 177, 57-67.

Figure Captions

Fig. 1. Experimental timeline. Adult females were weighed twice weekly from pair formation to parturition, and pups were checked daily for eye-opening. NB – Non-breeding females, PM – paired mothers, SM – single mothers, CORT – corticosterone.

Fig. 2. Towered cages used in the CLIMB condition. Food was located at the top of one tower, and a water bottle was at the top of the other tower.

Fig. 3. Estimated marginal means (EMM) and associated standard errors (SE) of number of fecal boli produced during A: 10-minute open-field tests and B: 6-minute tail-suspension tests. Statistical results shown are for pairwise comparisons among reproductive groups; neither the main effect of housing condition nor the housing condition $\times$ reproductive group interaction was significant for either measure. Light gray, standard cages; dark gray, towered cages.

Fig. 4. EMM $\pm$ SE baseline plasma corticosterone concentrations across the diel cycle in female California mice. A: CORT levels of all females housed in standard cages (CTRL) or in cages with towers requiring the mice to climb to obtain food and water (CLIMB), collapsed across reproductive conditions.

Fig. 5. EMM $\pm$ SE plasma corticosterone concentrations at 12:00 h under baseline conditions and immediately after a 6-minute tail-suspension test. The tail-suspension test significantly increased plasma CORT. A: CORT levels in all females housed in standard cages (CTRL) or in

cages with towers requiring the mice to climb to obtain food and water (CLIMB), collapsed across reproductive conditions. B: CORT levels of single mothers (SM), paired mothers (PM), and non-breeding females (NB), collapsed across housing conditions.

Fig. 6. EMM $\pm$ SE of mean lean mass and organ masses of female California mice. Ovary and triceps surae masses are means of the two paired organs. Statistical results shown are for pairwise comparisons among reproductive groups; neither the main effect of housing condition nor the housing condition $\times$ reproductive group interaction was significant for any measure. Light gray, standard cages; dark gray, towered cages.

Fig. 7. EMM $\pm$ SE of gastrointestinal masses of female California mice. Statistical results shown are for pairwise comparisons among reproductive groups; neither the main effect of housing condition nor the housing condition $\times$ reproductive group interaction was significant for any measure. Light gray, standard cages; dark gray, towered cages.

Table 1. Results of Pearson correlations and paired t-tests comparing values for paired organ sizes. Positive t values indicate that left > right. Significant P values ($P \leq 0.016$ when modified for Adaptive False Discovery Rate and Cohen's $d > 0.5$ for pair-wise comparisons) are underlined.

Organ	Unit	Transform	N of paired observations	r of Pearson correlation	P of Pearson correlation	t of paired t-test	P of paired t-test	Cohen's d
Triceps surae	gram	none	80	0.86	<u>1.66E-24</u>	1.11	0.270	0.066
Ovary	gram	none	79	0.77	<u>5.77E-17</u>	0.03	0.979	0.002
Adrenal	gram	none	80	0.82	<u>1.85E-20</u>	5.82	1.22E-07	0.392
Kidney	gram	none	80	0.93	<u>1.44E-35</u>	-1.99	0.0501	0.085
Femur	gram	none	77	0.78	<u>6.66E-17</u>	-1.13	0.263	0.086
Femur	millimeter	none	77	0.62	<u>1.86E-09</u>	0.17	0.868	0.017
Foot	millimeter	none	76	0.52	<u>1.00E-06</u>	1.17	0.246	0.132

Table 2. Results of ANCOVAs comparing three reproductive groups (non-breeding females, paired mothers, single mothers) under two housing conditions (standard cages, towered cages) with significant group differences. Significant P values ($P \leq 0.016$ when modified for Adaptive False Discovery Rate and $\eta^2 > 0.06$) are underlined.

Measure	Unit	Transform	Repro. Group				Housing Condition				Repro. Group * Housing Condition			
			F	D.F.	P	Partial eta squared	F	D.F.	P	Partial eta squared	F	D.F.	P	Partial eta squared
Open field: number of fecal boli	number	none	4.50	2,80	<u>0.014</u>	0.101	0.62	1,80	0.433	0.008	0.26	2,80	0.769	0.007
Tail suspension: number of fecal boli	number	none	3.14	2,77	<u>0.049</u>	0.075	0.10	1,77	0.752	0.001	0.23	2,77	0.796	0.006
Heart mass	gram	none	3.99	2,72	<u>0.023</u>	0.100	0.07	1,72	0.790	0.001	0.46	2,72	0.635	0.013
Liver mass	gram	log ₁₀	8.55	2,72	<u>4.68E-04</u>	0.192	0.32	1,72	0.573	0.004	0.18	2,72	0.835	0.005
Triceps surae mass (mean of left and right)	gram	rank	5.25	2,69	<u>0.008</u>	0.132	0.12	1,69	0.732	0.002	0.30	2,69	0.745	0.008
Ovary mass (mean of left and right)	gram	**0.5	6.96	2,71	<u>0.002</u>	0.164	0.01	1,71	0.932	1.02E-04	0.39	2,71	0.681	0.011
Uterus mass	gram	log ₁₀	5.85	2,71	<u>0.004</u>	0.141	0.75	1,71	0.390	0.010	0.22	2,71	0.807	0.006
Stomach mass	gram	none	6.53	2,69	<u>0.003</u>	0.159	1.33	1,69	0.252	0.019	0.59	2,69	0.557	0.017
Caecum mass	gram	log ₁₀	5.78	2,69	<u>0.005</u>	0.144	0.26	1,69	0.610	0.004	1.47	2,69	0.236	0.041
Large intestine mass	gram	none	5.51	2,69	<u>0.006</u>	0.138	0.18	1,69	0.677	0.003	1.84	2,69	0.166	0.051
Small intestine mass	gram	log ₁₀	4.54	2,69	<u>0.014</u>	0.116	0.51	1,69	0.477	0.007	0.43	2,69	0.656	0.012

2843															
2844															
2845															
2846	Postpartum body mass 4	gram	none	3.20	2,78	<u>0.046</u>	0.076	1.20	1,78	0.276	0.015	0.98	2,78	0.378	0.025
2847															
2848	Postpartum lean mass														
2849	(mean of lean masses 1-4)	gram	**0.5	3.69	2,81	<u>0.029</u>	0.083	0.47	1,81	0.495	0.006	1.98	2,81	0.145	0.047
2850															
2851	Postpartum lean mass 2	gram	rank	11.20	2,80	<u>0.003</u>	0.134	0.96	1,80	0.763	0.001	0.36	2,80	0.345	0.026
2852															
2853	Postpartum lean mass 3	gram	**0.5	7.43	2,80	<u>0.001</u>	0.157	1.33	1,80	0.252	0.016	0.64	2,80	0.530	0.016
2854															
2855	Percent fat mass 2	percent	rank	4.48	2,80	<u>0.014</u>	0.101	0.01	1,80	0.947	5.60E-05	1.47	2,80	0.235	0.036
2856															
2857	Percent fat mass 3	percent	rank	3.64	2,79	<u>0.031</u>	0.084	0.09	1,79	0.771	0.001	2.25	2,79	0.112	0.054
2858															
2859	Percent lean mass 2	percent	rank	5.15	2,80	<u>0.008</u>	0.114	0.00	1,80	0.975	1.20E-05	1.45	2,80	0.242	0.035
2860															
2861	Percent lean mass 4	percent	rank	3.13	2,72	<u>0.0496</u>	0.080	0.12	1,72	0.725	0.002	2.39	2,72	0.099	0.062
2862															
2863															
2864															
2865															
2866															
2867															
2868															
2869															
2870															
2871															
2872															
2873															
2874															
2875															
2876															
2877															
2878															
2879															
2880															
2881															
2882															
2883															
2884															

Table 3. Results of ANOVAs comparing pups reared by single mothers and paired mothers in the two housing conditions (standard cages, towered cages).

Measure	Unit	Transform	Repro. Group				Housing Condition				Repro. Group * Housing Condition			
			F	D.F.	P	Partial eta squared	F	D.F.	P	Partial eta squared	F	D.F.	P	Partial eta squared
Litter size at weaning	pup	none	1.41	1,55	0.239	0.025	0.12	1,55	0.732	0.002	0.01	1,55	0.937	1.14E-04
Pups' age of eye opening (eye opening of the first pup of the litter)	day	none	0.23	1,55	0.635	0.004	0.20	1,55	0.658	0.004	0.69	1,55	0.411	0.012
Pups' fat mass (mean of all pups in litter)	gram	log ₁₀	0.15	1,51	0.704	0.003	1.28	1,51	0.263	0.024	1.09	1,51	0.302	0.021
Pups' lean mass (mean of all pups in litter)	gram	none	0.56	1,51	0.458	0.011	1.94	1,51	0.169	0.037	0.25	1,51	0.618	0.005
Pups' percent fat mass (mean of all pups in litter)	percent	log ₁₀	0.03	1,51	0.859	0.001	0.23	1,51	0.637	0.004	0.86	1,51	0.359	0.017
Pups' percent lean mass (mean of all pups in litter)	percent	rank	0.01	1,51	0.923	1.85E-04	0.94	1,51	0.336	0.018	0.46	1,51	0.503	0.009
Pups' body mass (mean of all pups in litter)	gram	none	1.14	1,53	0.290	0.021	3.11	1,53	0.084	0.055	0.28	1,53	0.600	0.005
Pups' body mass (total body mass of litter)	gram	none	2.61	1,53	0.112	0.047	3.43	1,53	0.070	0.061	0.02	1,53	0.884	4.02E-04

Supplementary Table 1. Results of ANCOVAs comparing the three reproductive groups (non-breeding females, paired mothers, single mothers) in the two housing conditions (standard cages, towered cages). Significant P values ($P \leq 0.016$ when modified for Adaptive False Discovery Rate and $\eta^2 > 0.06$) are underlined.

Measure	Unit	Transform	Repro. Group				Housing Condition				Repro. Group * Housing Condition			
			F	D.F.	P	Partial eta squared	F	D.F.	P	Partial eta squared	F	D.F.	P	Partial eta squared
Novel object: latency to approach object	second	rank	0.68	2,81	0.510	0.016	0.21	1,81	0.648	0.003	0.12	2,81	0.890	0.003
Novel object: bouts of approaching object	number	**0.5	0.04	2,81	0.966	0.001	1.08	1,81	0.303	0.013	0.54	2,81	0.584	0.013
Novel object: duration of immobility	second	log ₁₀	1.20	2,81	0.312	0.054	0.20	1,81	0.658	0.005	0.51	2,81	0.603	0.024
Novel object: durations of sniffing and touching object	second	log ₁₀	0.85	2,81	0.430	0.021	0.32	1,81	0.574	0.004	1.15	2,81	0.323	0.028
Saccharine preference (proportion of total fluid consumption)	proportion	none	0.39	2,80	0.678	0.010	4.87	1,80	<u>0.030</u>	0.057	1.44	2,80	0.243	0.035
Open field: number of crossing bouts	number	log ₁₀	0.86	2,74	0.427	0.047	0.15	1,74	0.702	0.005	0.27	2,74	0.766	0.001
Open field: duration in outer region	second	rank	0.02	2,73	0.978	0.001	0.04	1,73	0.845	0.001	1.50	2,73	0.230	0.039
Open field: duration in inner region	second	**0.5	0.27	2,74	0.762	0.007	0.29	1,74	0.592	0.004	1.52	2,74	0.225	0.039
Open field: distance traveled in outer region	meter	**0.5	2.62	2,74	0.079	0.066	0.18	1,74	0.669	0.002	0.18	2,74	0.836	0.005

2969															
2970															
2971															
2972	Open field:														
2973	distance traveled in inner	meter	**0.5	0.27	2,68	0.767	0.007	0.23	1,68	0.637	0.003	0.13	2,68	0.882	0.004
2974	region														
2975	Open field:														
2976	total distance traveled in	meter	none	1.32	2,71	0.274	0.036	0.39	1,71	0.535	0.005	0.01	2,71	0.988	3.50E-04
2977	inner and outer regions														
2978	Open field:														
2979	number of fecal boli	number	none	4.50	2,80	<u>0.014</u>	0.101	0.62	1,80	0.433	0.008	0.26	2,80	0.769	0.007
2980															
2981	Open field:														
2982	number of urine pools	number	**0.5	0.41	2,80	0.665	0.010	0.58	1,80	0.447	0.007	0.20	2,80	0.822	0.005
2983															
2984	Tail suspension:														
2985	duration of mobility	minute	none	0.51	2,81	0.602	0.013	0.07	1,81	0.793	0.001	0.14	2,81	0.869	0.004
2986															
2987	Tail suspension:														
2988	number of fecal boli	number	none	3.14	2,77	<u>0.049</u>	0.075	0.10	1,77	0.752	0.001	0.23	2,77	0.796	0.006
2989															
2990	Heart mass	gram	none	3.99	2,72	<u>0.023</u>	0.100	0.07	1,72	0.790	0.001	0.46	2,72	0.635	0.013
2991															
2992	Liver mass	gram	log ₁₀	8.55	2,72	<u>4.68E-04</u>	0.192	0.32	1,72	0.573	0.004	0.18	2,72	0.835	0.005
2993															
2994	Spleen mass	gram	log ₁₀	0.22	2,72	1.526	0.041	0.73	1,72	0.125	0.002	0.85	2,72	0.166	0.005
2995															
2996	Triceps surae mass	gram	rank	5.25	2,69	<u>0.008</u>	0.132	0.12	1,69	0.732	0.002	0.30	2,69	0.745	0.008
2997	(mean of left and right)														
2998	Ovary mass														
2999	(mean of left and right)	gram	**0.5	6.96	2,71	<u>0.002</u>	0.164	0.01	1,71	0.932	1.02E-04	0.39	2,71	0.681	0.011
3000															
3001	Uterus mass	gram	log ₁₀	5.85	2,71	<u>0.004</u>	0.141	0.75	1,71	0.390	0.010	0.22	2,71	0.807	0.006
3002															
3003	Adrenal mass														
3004	(mean of left and right)	gram	rank	1.21	2,72	0.306	0.032	0.05	1,72	0.830	0.001	0.29	2,72	0.753	0.008
3005															
3006															
3007															
3008															
3009															
3010															

3011															
3012															
3013															
3014	Kidney mass	gram	none	0.28	2,72	0.759	0.008	0.00	1,72	0.960	3.50E-05	0.53	2,72	0.590	0.015
3015	(mean of left and right)														
3016															
3017	Stomach mass	gram	none	6.53	2,69	<u>0.003</u>	0.159	1.33	1,69	0.252	0.019	0.59	2,69	0.557	0.017
3018															
3019	Caecum mass	gram	log ₁₀	5.78	2,69	<u>0.005</u>	0.144	0.26	1,69	0.610	0.004	1.47	2,69	0.236	0.041
3020															
3021	Large intestine mass	gram	none	5.51	2,69	<u>0.006</u>	0.138	0.18	1,69	0.677	0.003	1.84	2,69	0.166	0.051
3022															
3023	Small intestine mass	gram	log ₁₀	4.54	2,69	<u>0.014</u>	0.116	0.51	1,69	0.477	0.007	0.43	2,69	0.656	0.012
3024															
3025	Femur mass	gram	none	0.00	2,67	0.999	2.60E-05	0.01	1,67	0.928	1.22E-04	0.25	2,67	0.783	0.007
3026	(mean of left and right)														
3027															
3028	Femur length	millimeter	none	0.96	2,70	0.386	0.027	0.95	1,70	0.333	0.013	0.17	2,70	0.844	0.005
3029	(mean of left and right)														
3030															
3031	Foot length	millimeter	none	0.49	2,69	0.617	0.014	0.38	1,69	0.538	0.006	0.15	2,69	0.862	0.004
3032	(mean of left and right)														
3033															
3034	Tail length	millimeter	none	0.40	2,70	0.671	0.011	1.93	1,70	0.169	0.027	0.79	2,70	0.460	0.022
3035															
3036	Postpartum body mass	gram	none	0.79	2,80	0.459	0.019	0.54	1,80	0.464	0.007	1.40	2,80	0.253	0.034
3037	(mean of body masses 1-5)														
3038															
3039	Postpartum body mass 1	gram	log ₁₀	0.60	2,76	0.554	0.015	0.22	1,76	0.645	0.003	1.29	2,76	0.282	0.033
3040															
3041	Postpartum body mass 2	gram	none	0.22	2,79	0.801	0.006	0.10	1,79	0.757	0.001	1.63	2,79	0.203	0.04
3042															
3043	Postpartum body mass 3	gram	none	2.18	2,81	0.120	0.051	1.55	1,81	0.216	0.019	1.67	2,81	0.195	0.040
3044															
3045	Postpartum body mass 4	gram	none	3.20	2,78	<u>0.046</u>	0.076	1.20	1,78	0.276	0.015	0.98	2,78	0.378	0.025
3046															
3047	Postpartum body mass 5	gram	none	2.09	2,68	0.132	0.058	0.88	1,68	0.352	0.013	0.71	2,68	0.494	0.021
3048															
3049															
3050															
3051															
3052															


3053															
3054															
3055															
3056	Postpartum fat mass	gram	rank	1.06	2,81	0.353	0.025	0.09	1,81	0.766	0.001	2.20	2,81	0.117	0.052
3057	(mean of fat masses 1-4)														
3058															
3059	Postpartum fat mass 1	gram	rank	1.38	2,76	0.258	0.035	0.10	1,76	0.749	0.001	0.65	2,76	0.526	0.017
3060															
3061	Postpartum fat mass 2	gram	rank	2.82	2,80	0.066	0.066	0.00	1,80	0.996	3.39E-07	2.22	2,80	0.115	0.053
3062															
3063	Postpartum fat mass 3	gram	rank	1.89	2,80	0.158	0.045	0.03	1,80	0.874	3.20E-04	2.28	2,80	0.109	0.054
3064															
3065	Postpartum fat mass 4	gram	rank	1.74	2,73	0.183	0.046	0.05	1,73	0.830	0.001	3.02	2,73	0.055	0.077
3066															
3067	Postpartum lean mass	gram	**0.5	3.69	2,81	<u>0.029</u>	0.083	0.47	1,81	0.495	0.006	1.98	2,81	0.145	0.047
3068	(mean of lean masses 1-4)														
3069															
3070	Postpartum lean mass 1	gram	rank	2.52	2,76	0.087	0.062	0.08	1,76	0.772	0.001	1.16	2,76	0.321	0.029
3071															
3072	Postpartum lean mass 2	gram	rank	11.20	2,80	<u>0.003</u>	0.134	0.96	1,80	0.763	0.001	0.36	2,80	0.345	0.026
3073															
3074	Postpartum lean mass 3	gram	**0.5	7.43	2,80	<u>0.001</u>	0.157	1.33	1,80	0.252	0.016	0.64	2,80	0.530	0.016
3075															
3076	Postpartum lean mass 4	gram	**2	2.31	2,73	0.106	0.06	0.01	1,73	0.912	1.69E-04	1.37	2,73	0.260	0.036
3077															
3078	Percent fat mass	percent	rank	2.96	2,81	0.058	0.058	0.39	1,81	0.532	0.532	1.71	2,81	0.187	0.187
3079	(mean of percent fat masses 1-4)														
3080															
3081	Percent fat mass 1	percent	rank	1.66	2,76	0.197	0.042	0.06	1,76	0.809	0.001	0.48	2,76	0.621	0.012
3082															
3083	Percent fat mass 2	percent	rank	4.48	2,80	<u>0.014</u>	0.101	0.01	1,80	0.947	5.60E-05	1.47	2,80	0.235	0.036
3084															
3085	Percent fat mass 3	percent	rank	3.64	2,79	<u>0.031</u>	0.084	0.09	1,79	0.771	0.001	2.25	2,79	0.112	0.054
3086															
3087	Percent fat mass 4	percent	rank	1.35	2,72	0.265	0.036	0.25	1,72	0.620	0.003	2.11	2,72	0.129	0.055
3088															
3089	Percent lean mass	percent	rank	1.80	2,81	0.171	0.043	0.05	1,81	0.831	0.001	1.71	2,81	0.187	0.041
3090	(mean of percent lean masses 1-4)														
3091															
3092															
3093															
3094															

3095															
3096															
3097															
3098	Percent lean mass 1	percent	rank	0.22	2,75	0.804	0.006	0.01	1,75	0.918	1.41E-04	0.46	2,75	0.635	0.012
3099															
3100	Percent lean mass 2	percent	rank	5.15	2,80	<u>0.008</u>	0.114	0.00	1,80	0.975	1.20E-05	1.45	2,80	0.242	0.035
3101															
3102	Percent lean mass 3	percent	rank	2.79	2,79	0.068	0.066	0.12	1,79	0.733	0.001	1.77	2,79	0.177	0.043
3103															
3104	Percent lean mass 4	percent	rank	3.13	2,72	<u>0.0496</u>	0.0800	0.12	2,72	0.725	0.002	2.39	2,72	0.099	0.062
3105															
3106															
3107															
3108															
3109															
3110															
3111															
3112															
3113															
3114															
3115															
3116															
3117															
3118															
3119															
3120															
3121															
3122															
3123															
3124															
3125															
3126															
3127															
3128															
3129															
3130															
3131															
3132															
3133															
3134															
3135															
3136															


Day of Expt.	Test Day	Time	Procedure(s)
~ -59			Vasectomize NB males
~ -45			Pair formation; move half of pairs to climbing cages
~ (-29)-(-24)			Vasectomize SM & PM males; move all pairs to divided cages for 1 week
0			Birth
1		09:00-10:00 h	Remove SM fathers
3	1	14:00-16:00 h	Body composition
6	2	04:00 h	Blood sample (basal CORT)
9	3	20:00 h	Blood sample (basal CORT)
12	4	14:00-16:00 h	Novel-object test; body composition
14	5	20:00-22:00 h	Behavioral observation
15	6	19:00-23:00 h	Saccharine preference test
17	7	14:00-16:00 h	Behavioral observation
18	8	14:00-16:00 h	Open-field test; body composition
20	9	8:00-10:00 h	Behavioral observation
21	10	12:00 h	Blood sample (basal CORT)
23	11	2:00-4:00 h	Behavioral observation
24	12	11:30-12:35 h	Tail-suspension test; blood collection (post-stress CORT), body mass
26	13	14:00-16:00 h	Body composition
27	14	14:00-16:00 h	Dissection




A




B







A



B



A



B

