

UCLA

UCLA Previously Published Works

Title

Crystal Clear: Purity of consumer-level methamphetamine samples and methamphetamine-adulteration of other drugs in Los Angeles, 2023-2025

Permalink

<https://escholarship.org/uc/item/58d2h7cj>

Journal

Drug and Alcohol Dependence Reports

ISSN

2772-7246

Authors

Koncsol, Adam J
Molina, Caitlin A
Romero, Ruby
[et al.](#)

Publication Date

2026-10-01

DOI

10.1016/j.dadr.2026.100482

Copyright Information

This work is made available under the terms of a Creative Commons Attribution-NonCommercial-NoDerivatives License, available at <https://creativecommons.org/licenses/by-nc-nd/4.0/>

Peer reviewed

Crystal Clear: Purity of street-level methamphetamine samples and methamphetamine-adulteration of other drugs in Los Angeles, 2023-2025

Authors: Adam J. Koncsol, MPA^{a,b}; Caitlin A. Molina, BA, BS^a; Ruby Romero, BA^a; Megan Grabill, BA^{a,c}; Morgan E. Godvin, BA^a; Jesse Carrizal^d; Manuel Contreras^d; Oscar Arellano, BA^d; Naomi Villanueva, MA^e; Elham Jalayer, BS^g; spider davila^f; Soma Snakeoil^g; Joshua Smith, BS^h; Charles J. Santamaria, BS^h; Leslie Nuñez, BS^h; Ezinne Okonkwoⁱ; Siddarth Puri, MD^j; Joseph R. Friedman, MD, PhD, MPH^k; Chelsea L. Shover, PhD^{a,b}

- a) Division of General Internal Medicine and Health Services Research, Geffen School of Medicine, University of California, Los Angeles, Los Angeles, CA, United States
- b) Dept. of Health Policy & Management, Fielding School of Public Health, University of California, Los Angeles, Los Angeles, CA, United States
- c) Dept. of Anthropology, College of Social Sciences, University of California, Los Angeles, Los Angeles, CA, United States
- d) Homeless Outreach Program's Integrated Care System, Los Angeles, CA, United States
- e) Bienestar Human Services, Los Angeles, CA, United States
- f) Community Health Project LA, Los Angeles, CA, United States
- g) The Sidewalk Project, Los Angeles, CA, United States
- h) Geffen School of Medicine, University of California, Los Angeles, Los Angeles, CA, United States
- i) University of California, Los Angeles, Los Angeles, CA, United States
- j) San Luis Obispo County Department of Behavioral Health, San Luis Obispo, CA, United States
- k) University of California, San Diego, San Diego, CA, United States

*Correspondence to Adam J. Koncsol, koncsol@ucla.edu, 1100 Glendon Ave, Ste, 850, Los Angeles, CA 90024

Abstract

Introduction:

The United States has seen a rise in methamphetamine use disorders and methamphetamine-involved deaths. However, there is limited information on methamphetamine at the street level. We leverage data from a community-based drug checking program in Los Angeles to investigate the makeup of the retail illicit methamphetamine market and the extent methamphetamine plays a role as an adulterant in other drugs.

Methods:

Participants accessing a community-based drug checking program provided samples of drug products and completed brief interviews in Los Angeles County, California, from 2023 to 2025. Samples underwent laboratory-based qualitative and quantitative mass spectrometry. We quantitated the presence and concentration of methamphetamine as well as all other identified substances among samples expected only to be methamphetamine and samples expected to be another substance (e.g., fentanyl, heroin) where methamphetamine was found as an adulterant/contaminant.

Results:

Of N=2193 total samples, n=652 (29.7%) were methamphetamine-positive on DART-MS. In samples expected to contain methamphetamine, the mean purity was 78.2% (SD 24.0%; range <1% to ≈100%). As an unexpected component, the mean methamphetamine purity was 17.8% (SD 32.3%) and was mostly found in samples expected to be fentanyl, heroin, other amphetamines.

Conclusions:

Participants in Los Angeles seeking methamphetamine appear to frequently obtain high-purity samples that are rarely adulterated with other substances, although we observed a wide variation in concentration between samples. In contrast, drugs purchased without the expectation of containing methamphetamine had high prevalence of methamphetamine adulteration/contamination. Future research should expand methamphetamine surveillance techniques and investigate how variability in illicit markets may impact health outcomes.

Keywords: *drug checking; illicit drug supply; quantitation, harm reduction, surveillance*

Highlights:

- No other active compounds were detected for 90% of samples expected to be meth
- Samples expected to contain methamphetamine had a mean purity of 72% (SD 24%).
- Meth was found as an adulterant in opioids and stimulants in high concentrations

Introduction:

Methamphetamine is a central nervous system stimulant that carries potential health risks to multiple organ systems and is implicated in a rising number of drug-related deaths (Dickson et al., 2021; Schürer et al., 2017; Schwarzbach et al., 2020; Somma et al., 2023; Tobolski et al., 2022). Methamphetamine use appears to be increasing around the world, particularly in the Americas, Asia, and Australia (Blemings & Cunningham, 2024; Han, Compton, et al., 2021; Han, Cotto, et al., 2021; Hedegaard et al., 2020). Health consequences of methamphetamine use have also increased, with the U.S. seeing a rise in methamphetamine-related deaths from 4,500 in 2015 to approximately 30,000 in 2024 (National Center for Health Statistics, 2025). Chronic methamphetamine use increases the risk of cardiovascular harms such as myocardial toxicity, coronary artery disease, hypertension, stroke, sudden cardiac arrest, and other causes of morbidity and mortality (Dickson et al., 2021; Schürer et al., 2017; Schwarzbach et al., 2020; Somma et al., 2023; Tobolski et al., 2022). Understanding the methamphetamine that is used in real-world settings may have a number of implications for addiction, psychiatric, cardiovascular medicine more broadly. For example, the extent to which variation and adulteration in drug markets affect methamphetamine withdrawal and craving may be important for outcomes related to psychosis, mood regulation, cardiovascular and neurological wellbeing, contingency management, mitigating risk of overamping, and behavioral health interventions more broadly (Ciccarone & Shoptaw, 2022; Coffin et al., 2020; Colfax et al., 2011; Lake et al., 2020; Li & Shoptaw, 2023; Minozzi et al., 2024; Rush et al., 2025; Trivedi et al., 2021; Wang et al., 2012).

As a fully synthetic drug, methamphetamine is considerably cheaper and easier to produce than agriculturally-derived stimulants like cocaine (Jones et al., 2022; Lewis et al., 2021; Montero et al., 2022). Methamphetamine seized by law enforcement tends to have exceptionally high purity, with most recently available U.S. national averages exceeding 90% (Drug Enforcement Administration Special Testing and Research Laboratory of the Office of Forensic Science, 2024; Toske & McKibben, 2022; United States Sentencing Commission, 2024). However, drugs seized in criminal investigations are potentially interdicted higher in the supply chain prior to adulteration by lower-level sellers and may reflect purer product

than what end-buyers receive. Drug adulteration can happen at various points in the supply chain as a way to reduce costs associated with production and is typically done by bulking drug product with cheaper compounds that may mimic or intensify psychoactive effects. Such practices are driven by a combination of drug distributors saving money and people who use drugs being unable to determine the presence of any adulterants prior to sale or use. Drug checking—chemically analyzing personal samples of drugs for harm reduction and supply monitoring—can characterize methamphetamine purity and adulteration at the street level.

Los Angeles County, California has long led the United States in both number of methamphetamine-related deaths (reflecting the large population) and proportion of all drug-related deaths attributable to methamphetamine (reflecting a persistent market) (Shover et al., 2023). In recent years, an increasing number of Los Angeles County's deaths attributed to methamphetamine were also attributed to fentanyl (Shover et al., 2023). This pattern is mirrored in national data, where the fourth wave of the U.S. overdose crisis is characterized by deaths involving polysubstance combinations of fentanyl and other synthetic opioids and stimulants like cocaine and methamphetamine (Ciccarone, 2021; Friedman et al., 2026; Friedman & Shover, 2023). While public health campaigns have often focused on the threat of fentanyl adulteration of other drugs, data from testing actual drug products has been limited.

Leveraging data from a community-based drug checking program, we sought to empirically investigate the methamphetamine supply in Los Angeles. Specifically, we sought to characterize the presence and purity/concentration of methamphetamine and co-detected substances among 1) methamphetamine-positive samples expected only to be methamphetamine and 2) methamphetamine-positive samples expected to be another substance (e.g., purchased as fentanyl or heroin) where methamphetamine is functioning as an adulterant. "Purity" is used to refer specifically to the percentage by mass of methamphetamine in samples expected to be methamphetamine, whereas "concentration" is used to refer to the percentage by mass of any compound present in any drug (e.g., methamphetamine in a sample expected to be fentanyl).

Methods:

Point-of-service drug checking

Applying data sampling and methods described previously (Appley et al., 2023; Friedman et al., 2025; Shover et al., 2025; Sisco et al., 2017), anonymous participants utilizing a community-based drug checking program, *Drug Checking Los Angeles*, voluntarily provided a small sample (approximately 1-5 mg) of drug products (e.g., powders, crystals, pills, liquids, tar, etc.) for testing at four different sites in Los Angeles County, California: (1) East Los Angeles, (2) Downtown Los Angeles / Skid Row, (3) Hollywood, and (4) South-Central Los Angeles. This study uses data collected from February 2023 to December 2025. An optional anonymous survey was administered by trained staff to assess what the drug was purported to be at the time of sale and what a participant expected the sample to contain. Instances where a respondent did not know what the sample contained or declined to answer are presented as “Unknown / Declined.” Initial-point-of-service drug checking used Fourier-Transform Infrared (FTIR) spectroscopy and various BTNX-branded and lab-validated immunoassay test strips, such as fentanyl and methamphetamine test strips (Sisco et al., 2024).

Laboratory-based confirmatory analysis

Samples received through Drug Checking Los Angeles were tested in the field to provide real-time results to participants and also sent by mail to the National Institute of Standards and Technology Rapid Drug Analysis and Research (NIST RaDAR) program for laboratory-based analysis (NIST RaDAR, 2022). NIST RaDAR evaluated all samples included in the study with direct-analysis-in-real-time mass spectrometry (DART-MS) and a subset of samples with liquid-chromatography tandem mass spectrometry (LC-MS/MS).

DART-MS methods assess samples against a library of nearly 1,500 substances, including most pharmaceutical and illicitly manufactured drugs, adulterants, cutting and bulking agents, precursor chemicals, and other relevant substances (adhesives, industrial ingredients, etc.). DART-MS methods are only able to provide information on the binary presence or absence of compounds and are unable to provide information on concentration; however, they only require residual or trace amounts of a sample to be analyzed. A subset of samples analyzed with DART-MS were also analyzed with LC-MS /MS methods to quantitate

the percent mass of pre-specified compounds present in the sample. See **Supplemental Table A** for more information about the specific analytes used on the LC-MS /MS quantitation panel. More detailed descriptions of the confirmatory DART-MS and LC-MS /MS methodologies employed in this study have been previously published (Appley et al., 2023). An important limitation to all reported results is that laboratory methods may be unequipped to detect byproduct or precursor compounds related to the synthesis products for which laboratory standards do not exist.

Methamphetamine concentration values reported by NIST RaDAR were expressed as methamphetamine freebase. To account for the molecular weight of the hydrochloride salt when calculating concentration estimates, methamphetamine freebase was converted to methamphetamine HCl. Dividing the molecular weight of methamphetamine HCl (185 Da) by the molecular weight of methamphetamine freebase (149 Da) and multiplying that conversion factor (approximately 1.24) by each methamphetamine freebase value can provide a more accurate estimation of methamphetamine concentration in our sample—especially because all samples of methamphetamine tested were sold as “crystal methamphetamine” or “ice”. As a result of the freebase-to-salt concentration conversion, any samples $\geq 85\%$ methamphetamine freebase were treated as pure and were imputed to be 100% for this analysis. Similarly, samples that were considered below the limit of quantitation ($<LOQ$) were imputed to be 0.1% to calculate descriptive statistics.

Statistical analysis

The count and prevalence (%) of any co-detected compounds were calculated across methamphetamine-expected and methamphetamine-unexpected samples confirmed with DART-MS to contain methamphetamine. This highlights (1) compounds that may be present in samples expected to be methamphetamine and (2) how often methamphetamine was detected in samples expected to be other drugs, like heroin or fentanyl. For the subset of samples that underwent quantitative LC-MS /MS analysis, the concentration was assessed in terms of average, standard deviation, and range of methamphetamine, presented across participant drug expectation categories. Study protocols were approved by the UCLA IRB (IRB-22-0760 and IRB-22-1273). FTIR analyses were completed using

OPUS, version 8.7.31 (Bruker, Billerica, MA, USA). Statistical analyses were carried out using Stata, version 19.5 (StataCorp, College Station, TX, USA).

Results:

We tested N = 2,193 samples of drug product with DART-MS from February 2023 to December 2025. Among n = 652 samples found to contain methamphetamine, 67% were expected to contain methamphetamine (n = 432 methamphetamine-only, n = 2 methamphetamine + heroin mixture, n = 1 methamphetamine + fentanyl mixture, and n = 1 methamphetamine + ketamine mixture). The remaining 33% (n=220) were not expected to contain methamphetamine.

Co-prevalent compounds in methamphetamine

Table 1 shows co-detected substances among n=432 of methamphetamine-positive samples expected to only contain methamphetamine. The most commonly co-prevalent compounds were cocaine (n=9, 2.1%), ephedrine (n=8, 1.9%), lidocaine (n=8, 1.9%), and fentanyl (n=7, 1.6%). Most samples (n=387, 89.6%) expected to be methamphetamine contained only methamphetamine, with no other co-prevalent compounds detected on DART-MS.

Purity of methamphetamine

Of the n=652 samples confirmed to contain methamphetamine, a subset (n=208) underwent LC-MS /MS analysis. For quantitated samples expected to be methamphetamine (n=172), the average percent by mass of methamphetamine hydrochloride was 78.2% (SD=24.0%) and ranged from below the limit of quantitation ($\approx 0.1\%$) to virtually pure ($\approx 100\%$). **Figure 1** highlights the distribution of methamphetamine concentration among samples expected to be methamphetamine vs where methamphetamine was not expected, excluding n=11 samples where the expected drug could not be determined. For samples which were not pure methamphetamine but did not contain any additional compounds on DART-MS, the remaining percentage mass of the sample was likely to be inert precursors and byproducts related to inefficient methamphetamine synthesis.

Methamphetamine's role as an adulterant

In approximately 33% (n=220) of samples in which methamphetamine was detected, its presence was not expected by the participant. **Table 2 and Figure 2** highlight how frequently methamphetamine was detected as an adulterant in drug products expected to be other substances. Notably, methamphetamine was detected in 8.5% of expected fentanyl samples (n=62), 16.4% of expected heroin samples (n = 34), 14.0% of expected MDMA / MDA samples (n =19), and 70.4% of expected amphetamine samples (n =19). It was also detected among samples expected to be 2C-B (41.2%, n=7), benzodiazepines (11.5%, n = 6), and GHB / BDO (33.3%, n=2). In our sample, methamphetamine was occasionally detected in samples expected to be oxycodone (4.0%, n =3), powder and crack cocaine (3.8%, n=6), and was not detected in the few samples expected to be designer stimulants, cathinones, or research chemicals others brought for testing. Methamphetamine's presence as an adulterant is not limited to only crystalized or powdered drugs but was present across a wide range of various forms. For example, methamphetamine was detected in tars, typically co-prevalent with heroin, pills, primarily as a substitute to counterfeit Adderall and occasionally in expected oxycodone, liquids, and other drug forms.

Table 3 displays the concentration of methamphetamine in samples expected to be methamphetamine, as well as its concentration in other drugs (when it is present as an adulterant). When present as an adulterant in other drugs, methamphetamine was present at levels that are likely pharmacologically active and relevant for clinical outcomes. For example, among 30 samples expected to only contain fentanyl, the mean methamphetamine concentration was 17.5% by mass. Similarly, in 9 samples expected to only contain heroin, the average methamphetamine concentration was 15.2% by mass. Notably, however, was how heterogeneous methamphetamine adulteration was across samples of heroin and fentanyl, with standard deviations of methamphetamine concentration being 33.0% and 27.0%, respectively.

Discussion:

In this study of street-level methamphetamine-involved samples, we found that drugs expected to be methamphetamine overwhelmingly—90% of the time—contained only methamphetamine. In contrast, samples expected to only contain other drugs (including fentanyl, heroin, MDMA, 2C-B, and others) often

unexpectedly contained methamphetamine across a number of various forms, like crystals, powders, tars, pills, liquids, and more. Overall, this reflects the increasingly ubiquitous role of inexpensive and highly pure methamphetamine in various aspects of the illicit drug market, as well as the volatility of the drug supply in the age of synthetic drugs more generally. In our sample, the percentage mass of methamphetamine in a sample expected to be methamphetamine ranged from well below 1% to nearly 100%, with an average of 78%. These findings illustrate that the purity of methamphetamine at the street-level in Los Angeles is not as uniform or high as law enforcement seizure data suggest (95%-100%).

Variation in methamphetamine purity may have implications for the health risks of individuals exposed to the drug. Individuals expecting or accustomed to lower purity samples may be inadvertently and unwittingly exposed to a higher dose. This likely has implications for methamphetamine toxicity, including risk of psychosis, mood dysregulation, cardiovascular and pulmonary harms. Similarly, understanding variation in potency may have implications for treatment initiation and retention with medications for methamphetamine use disorder, as well as psychosocial and behavioral interventions such as contingency management, cognitive behavioral therapy, motivational interviewing, and community reinforcement approaches (Ciccarone & Shoptaw, 2022; Coffin et al., 2020; Colfax et al., 2011; Lake et al., 2020; Li & Shoptaw, 2023; Minozzi et al., 2024; Rush et al., 2025; Trivedi et al., 2021; Wang et al., 2012). Understanding methamphetamine purity at the street level can help more fully contextualize health-related harms associated with methamphetamine, along with other factors relevant to clinical care, like the dosage, frequency of use, route of administration. Additional public health interventions that include harm reduction programming for people who use methamphetamine may be able to consider that methamphetamine purity varies significantly between batches. Better understanding of purity and potency may be important to avoid or respond to 'overamping', a term used to describe a form of stimulant overdose with symptoms ranging from catatonia, anxiety, elevated heart rate, audiovisual hallucinations, and more severe outcomes like loss of consciousness, tonic-clonic seizures, and cardiac issues (Mansoor et al., 2022).

Our findings do not support the notion that fentanyl has been regularly added to drug products sold as methamphetamine in Los Angeles. Fentanyl was

detected in 1.6% of samples expected to be methamphetamine, and in the few cases where quantitative data were available, the concentrations were very low. It is unknown how many of these samples were purchased already contaminated versus being later contaminated by co-use through shared paraphernalia or storage, given that polysubstance use was commonly reported by drug checking participants. These findings may not generalize more broadly as drug markets are often highly localized and may change rapidly.

The balance between high-risk yet low probability of fentanyl adulteration in methamphetamine can inform measured public health communications. Any fentanyl cross-contamination, even trace amounts, presents a substantial risk of fatal opioid overdose to people without opioid tolerance. Thus, the recommendations to always check stimulants for fentanyl before use and to carry naloxone remain essential harm reduction strategies. However, messaging should take care not to imply or assert that methamphetamine adulteration with fentanyl is widespread. Evidence-based messaging can help promote safer practices while also maintaining credibility with people who use drugs and their communities.

Nevertheless, we did find that methamphetamine was regularly detected in samples expected to be other drugs, mostly fentanyl, heroin, and MDMA, sometimes at substantial and pharmacologically active concentrations far above trace amounts. For example, one sample of powder heroin contained 10 times more methamphetamine than heroin by weight: 81% methamphetamine to only 7% of heroin. **These findings help provide context to the role that methamphetamine often can serve in Los Angeles markets: an adulterant to other substances rather than the target of adulteration itself.**

Importantly, people who use drugs such as fentanyl, heroin, amphetamines, 2C-B, MDMA, and other phenethylamines may unknowingly be exposed to methamphetamine, which could have harmful or unpleasant consequences. When methamphetamine was detected in other drugs, it was typically detected at levels which may be pharmacologically active e.g., 17.5% of samples expected to be fentanyl contained methamphetamine. Methamphetamine was also detected across a wide range of drug forms not limited to crystals or powders, such as pills, tars, and liquids. Although methamphetamine was not detected in any designer stimulants, cathinones, or research chemicals in our sample, these compounds

may be ripe for methamphetamine adulteration or substitution given their entactogenic and stimulating effects and relatively higher market prices. Indeed, methamphetamine has been documented as an adulterant in other markets. Publicly available results from WEDINOS (U.K.) and DrugsData.org (worldwide, results through mid-2024) show that methamphetamine has been detected internationally in samples expected to be 2-CB, 3-MMC, ketamine, or benzodiazepines, among others (Erowid, 2024; Welsh Emerging Drugs and Identification of Novel Substances Project, 2026).

Unintentional methamphetamine exposure may contribute to physical health and psychiatric signs and symptoms inconsistent with the clinical presentation typical of using opioids or shorter-acting stimulants (such as amphetamine salts or MDMA) alone. Harm reduction interventions could underscore the risk of methamphetamine adulteration in other drugs and the potential harms of unintentional exposure. Test strips are one example of a validated, low-cost, and low-barrier technology that could be leveraged to help people avoid unwanted exposure to methamphetamine (Sisco et al., 2024). These findings also help to highlight the potential value of data from community-based drug checking programs. Given that illicit markets are highly volatile and constantly shifting, community-based drug checking programs empower people who use drugs to make decisions in real time while also providing aggregated epidemiological surveillance of multiple, and sometimes overlapping, drug supplies. Drug checking services in the United States face a number of unique challenges which make implementation and maintenance difficult. Lack of organizational and material resources in the context of funding-limited environments, ambiguous or contentious local and state policies, the historical not-in-my-back-yard socio-spatial stigmatization of harm reduction and addiction services are a few of the barriers to drug checking service delivery (Smith, 2020; Smith et al., 2026).

It is unclear why methamphetamine, a potent, long-acting stimulant, was so often an unexpected adulterant in opioids. These drugs, which have very different effects, are frequently used together, e.g. as a “goofball;” however, preparations in which a participant expected to contain both an opioid and methamphetamine were presented separately in the analysis. We speculate that methamphetamine

may be added because it functions as a cheap active cutting agent, producing a physiological or psychological response in a way that an inert compound, like sugar, would not. Methamphetamine is particularly inexpensive in Southern California, with some participants reporting bulk prices as low as \$5 USD per gram

The findings associated with methamphetamine in Los Angeles may further illuminate the fourth wave of the overdose crisis (Ciccarone, 2021; Friedman et al., 2026; Friedman & Shover, 2023). Largely characterized by an increase of deaths involving both fentanyl and stimulants, there is uncertainty the extent to which the fourth wave is driven by polysubstance use versus adulteration and contamination. The finding that drugs expected to be methamphetamine are nearly ubiquitously unadulterated helps to rule out the notion that fentanyl is a common adulterant in Los Angeles methamphetamine. Moreover, the finding that methamphetamine was regularly present in both fentanyl and heroin at pharmacologically relevant concentrations may help explain changing rates of stimulant-positivity in mortality records.

Limitations:

While methamphetamine samples tested through a community-based drug checking program can provide unique insights into street-level drug markets, there are several limitations to consider. Given the sensitivity of laboratory instruments, detection of adulterants at low levels could occur due to brief, accidental cross-contamination—especially in instances where a person reports that they use both substances. Cross-contamination could also occur through storing multiple drug products in the same bag, or even by touching one substance and then touching another. Although polysubstance use and co-storage information was ascertained by drug checking staff if a participant opted to share that information, it was not possible to completely distinguish between intentional adulteration and accidental cross-contamination. The use of quantitative testing, however, helps to contextualize instances wherein methamphetamine adulteration represents a large portion of the sample tested—far above levels that would be expected from accidental cross-contamination alone.

The fallibility of a participant's memory to accurately recall drug expectation may have also biased our results. Given that drug expectation was ascertained

through participant self-report, a participant may have misremembered or mixed-up bags. Despite our best efforts to sample across four geographically and culturally distinct areas of Los Angeles, this is ultimately a convenience sample that may or may not be generalizable to drugs not submitted for checking. For example, a participant may be more likely to bring a sample to check after an adverse or unusual experience or because they were already utilizing syringe service program resources. Laboratory methods are subject to various errors that may affect the precision of our estimates, including the margin of error in quantification and limitations of specific analytical methods. For example, DART-MS techniques are not generally equipped to detect some molecules, like lactose or starches, so the prevalence of some cutting agents are not reflected in these results. DART-MS methods may have also missed byproduct or precursor compounds related to the methamphetamine synthesis process for which laboratory standards may not yet exist—helping to explain the gap between samples with only methamphetamine detected yet were below 100% pure methamphetamine concentration. The LC-MS /MS quantitation panel only included a subset of all potential compounds, and substances not on this panel were unable to be quantitated. We aimed to mitigate some of these limitations via the use of multiple technologies, yet results should be interpreted in light of these potential sources of error and bias.

Conclusion:

Methamphetamine purity in Los Angeles is high on average, however, lower than the >95% values seen in wholesale seizure data. Few contaminants were detected among samples expected to be methamphetamine; however, the purity of methamphetamine in the community setting is highly heterogeneous between samples. Methamphetamine is frequently used as an adulterant in other drugs—often at pharmacologically-relevant concentrations (i.e., representing a large fraction of the sample). Understanding methamphetamine concentration among drug samples in the community has a number of potential clinical and public health implications. Public health surveillance of street-level drug markets may facilitate interventions to address population-wide increases in methamphetamine use, shifting purity, and new adulterants. Future research is needed to evaluate how these findings compare to drug markets beyond Los Angeles. Further

research is warranted to understand methamphetamine as an adulterant and the health effects of intentional and unintentional polysubstance use.

Table 1: Count and proportion of compounds detected in samples expected to contain only methamphetamine

			Methamphetamine-only expected samples	
Qualitative co-prevalent compounds	Class	Compound	<i>Count</i>	<i>Proportion</i>
	Amphetamines	Methamphetamine N, N	432	100.0%
		Dimethylamphetamine	4	0.9%
		MDAI	3	0.7%
		MDMA	1	0.2%
	Fentanyl, fentalogs, and fentanyl precursors	Fentanyl	7	1.6%
		4-ANPP	1	0.2%
		Aniline	1	0.2%
		Acetyl fentanyl	1	0.2%
	Heroin and related compounds	Heroin	2	0.5%
		6-MAM	1	0.2%
		Papaverine	1	0.2%
		Noscapine	1	0.2%
	Anesthetics	Cocaine	9	2.1%
		Lidocaine	8	1.9%
		Benzocaine	1	0.2%
	Other drugs	Ephedrine	8	1.9%
		Epinephrine / Adrenaline	2	0.5%
		Acetaminophen	1	0.2%
		Ketamine	1	0.2%
	Cutting agents & fillers	Dimethyl Sulfone (MSM)	6	1.4%
		Diethyl Phthalate	2	0.5%
		Levamisole	1	0.2%
		Triacetin	1	0.2%

	Total	N	432	100.0%
--	--------------	---	-----	--------

Note: Table shows the count and proportion (%) of compounds that were detected among n = 432 methamphetamine-positive samples which were expected to contain only methamphetamine. Samples collected February 2023 - December 2025 through a community-based drug checking program in Los Angeles.

Table 2: Methamphetamine positivity by drug expectation

Expected drug	Methamphetamine negative		Methamphetamine positive		Total samples	
	Count	Proportion	Count	Proportion	Count	Proportion
Methamphetamine	9	2.0%	432	98.0%	441	20.1%
Fentanyl	671	91.5%	62	8.5%	733	33.4%
Heroin	173	83.6%	34	16.4%	207	9.4%
Amphetamine	8	29.6%	19	70.4%	27	1.2%
MDMA / XTC	117	86.0%	19	14.0%	136	6.2%
2C-B	10	58.8%	7	41.2%	17	0.8%
Benzodiazepines	46	88.5%	6	11.5%	52	2.4%
Cocaine or Crack	152	96.2%	6	3.8%	158	7.2%
"Psilocybin" products	5	62.5%	3	37.5%	8	0.4%
Oxycodone	72	96.0%	3	4.0%	75	3.4%
Heroin and Methamphetamine mixture	0	0.0%	2	100.0%	2	0.1%
GHB or BDO	4	66.7%	2	33.3%	6	0.3%
Amoxicillin	0	0.0%	1	100.0%	1	0.0%

Cannabis	1	50.0%	1	50.0%	2	0.1%
Dexmethylphenidate	0	0.0%	1	100.0%	1	0.0%
Methamphetamine and Fentanyl mixture	1	50.0%	1	50.0%	2	0.1%
Methamphetamine and Ketamine mixture	0	0.0%	1	100.0%	1	0.0%
Fentanyl and Heroin mixture	4	80.0%	1	20.0%	5	0.2%
PCP	6	85.7%	1	14.3%	7	0.3%
Ketamine	78	98.7%	1	1.3%	79	3.6%
Unknown/declined	122	71.3%	49	28.7%	171	7.8%
Total	1541	70.3%	652	29.7%	2193	100.0%

Note: There were n = 652 samples of drug product that were confirmed positive for methamphetamine and are a subset of N = 2,193 total samples that were brought for testing at a community-based drug checking site in Los Angeles. Samples collected February 2023 - December 2025. DART-MS methods were used to qualitatively assess the presence of methamphetamine. Semi-structured ethnographic interviewing methods were used to capture what participants expected their drugs to be at the time of testing, either through what the substance was bought as or what the participant otherwise expected the sample to contain and may be subject to various limitations, such as recall bias or cross contamination. "Psilocybin" products largely included various food products, pills, or liquids, but not mushroom or fungal material.

Table 3: Methamphetamine appears in other illicit drugs, sometimes in significant concentrations

		Average methamphetamine concentration (percent by mass)	Standard Deviation	Range	N
Expected drug or substance	Methamphetamine	78.2	24.0	[<LOQ - ≈100%]	145
	Fentanyl	17.5	33.0	[<LOQ - ≈100%]	30
	Amphetamines	19.5	32.6	[<LOQ - 90.0%]	8
	Heroin	15.2	27.0	[<LOQ - 81.3%]	9
	2C-B	4.9	.	4.9%	1
	Cocaine / crack	49.3	69.6	[<LOQ - 98.6%]	2
	GHB	<LOQ	.	<LOQ	1
	Oxycodone	3.8	.	3.8%	1
	Unknown / Declined	65.9	38.3	[24.8% - ≈100%]	11
	All methamphetamine-positive samples	62.4	37.5	[<LOQ - ≈100.0%]	208

Note: Table shows the methamphetamine concentration (percent by mass), standard deviation, range, count of samples and participant's expected drug or substance among n = 208 methamphetamine-positive, LC-MS quantitated samples. Data collected February 2023 - December 2025 in Los Angeles. Semi-structured ethnographic interviewing methods were used to capture what

participants expected their sample to be at the time of testing, either through what the drug or substance was bought as or what the participant otherwise expected the sample to contain. As with other forms of self-reported data, these data may be subject to bias if a participant mixed up samples or otherwise made an error in recalling what they bought a sample as.

**Supplemental table A: LC-MS /MS
Quantitation Panel**

Class	Quantitative Analyte
Amphetamines	Methamphetamine
Fentanyl, fentalogs, and fentanyl precursors	Fentanyl Carfentanil† Fluorofentanyl 4-ANPP Ethyl 4-ANPP Phenethyl 4-ANPP
α_2- agonists	Xylazine Medetomidine
Heroin and related compounds	Heroin 6-MAM† Noscapine† Papaverine†
Anesthetics	Cocaine Lidocaine* Procaine† Tetracaine
Other drugs	Acetaminophen Caffeine† Ketamine Levamisole†

	Quetiapine (Seroquel)
Cutting agents & fillers	BTMPS*

*Added to the quantitation panel in July 2024

†Added to the quantitation panel in January 2025

Funding Acknowledgements:

This work was supported by the Centers for Disease Control and Prevention as part of Overdose Data to Action: LOCAL (CDC-RFA-CE-23-0003), National Institutes of Health (R01 DA057630), and an equipment grant from the James B. Pendleton Charitable Trust to the UCLA AIDS Institute and UCLA Center for AIDS Research. AJK was supported by the NIH/National Center for Advancing Translational Science (NCATS) UCLA CTSI (grant TL1TR001883). CLS was supported by the U.S. National Institute on Drug Abuse (grant K01DA050771). The funders had no role in designing the study, analyzing the data, or the decision to submit for publication.

Additional Acknowledgments:

We would also like to acknowledge our laboratory partners at the National Institute of Standards and Technology Rapid Drug Analysis and Research program (NIST RaDAR). We would also like to acknowledge our community partners at Bienestar Human Services, Community Health Project LA, The Sidewalk Project, and Homeless Outreach Program's Integrated Care System (HOPICS). We would also like to extend our gratitude to the anonymous participants of Drug Checking Los Angeles who made this work possible through their donation of samples for confirmatory testing.

References

- Appley, M. G., Robinson, E. L., Thomson, A., Russell, E., & Sisco, E. (2023). An analytical platform for near real-time drug landscape monitoring using paraphernalia residues. *Forensic Chemistry*, 34, 100489.
<https://doi.org/10.1016/j.forc.2023.100489>
- Blemings, B., & Cunningham, S. (2024). Temporary gains and permanent costs in methamphetamine precursor control. *International Journal of Drug Policy*, 104501. <https://doi.org/10.1016/j.drugpo.2024.104501>
- Ciccarone, D. (2021). The Rise of Illicit Fentanyls, Stimulants and the Fourth Wave of the Opioid Overdose Crisis. *Current Opinion in Psychiatry*, 34(4), 344–350.
<https://doi.org/10.1097/YCO.0000000000000717>
- Ciccarone, D., & Shoptaw, S. (2022). Understanding Stimulant Use and Use Disorders in a New Era. *Medical Clinics of North America, Substance Use Disorders*, 106(1), 81–97. <https://doi.org/10.1016/j.mcna.2021.08.010>
- Coffin, P. O., Santos, G.-M., Hern, J., Vittinghoff, E., Walker, J. E., Matheson, T., Santos, D., Colfax, G., & Batki, S. L. (2020). Effects of Mirtazapine for Methamphetamine Use Disorder Among Cisgender Men and Transgender Women Who Have Sex With Men. *JAMA Psychiatry*, 77(3), 246–255.
<https://doi.org/10.1001/jamapsychiatry.2019.3655>

Colfax, G. N., Santos, G.-M., Das, M., Santos, D. M., Matheson, T., Gasper, J.,

Shoptaw, S., & Vittinghoff, E. (2011). Mirtazapine to Reduce

Methamphetamine Use: A Randomized Controlled Trial. *Archives of General Psychiatry*, 68(11), 1168–1175.

<https://doi.org/10.1001/archgenpsychiatry.2011.124>

Dickson, S. D., Thomas, I. C., Bhatia, H. S., Nishimura, M., Mahmud, E., Tu, X. M., Lin,

T., Adler, E., Greenberg, B., & Alshawabkeh, L. (2021). Methamphetamine-

Associated Heart Failure Hospitalizations Across the United States:

Geographic and Social Disparities. *Journal of the American Heart Association*, 10(16), e018370. <https://doi.org/10.1161/JAHA.120.018370>

Drug Enforcement Administration Special Testing and Research Laboratory of the

Office of Forensic Science. (2024). *Methamphetamine Profiling Program*

Report: CY 2022 (PRB# 2024-02).

https://www.dea.gov/sites/default/files/2025-02/508_CY%202022%20MPP_LS%20Report%20PRB%202024-02.pdf

Erowid. (2024). *Test Results*. DrugsData.Org.

<https://www.drugsdata.org/results.php?>

sort=DatePublishedU+desc&start=0&a=&search_field=-&m1=-1&m2=-

1&sold_as_ecstasy=both&datefield=tested&max=2000

Friedman, J. R., Palamar, J. J., Ciccarone, D., Gaines, T. L., Borquez, A., Shover, C. L., & Strathdee,

S. A. (2026). Charting the decline of the fourth wave: US overdose deaths by race, ethnicity and substance involvement. *Addiction (Abingdon, England)*, 121(9), 2552–2558. <https://doi.org/10.1111/add.70472>

Friedman, J., Molina, C. A., Koncsol, A. J., Romero, R., Godvin, M. E., Jalayer, E., Davila, S., Arellano, O., Cowan, A., Hurley, B., & Shover, C. L. (2025). Xylazine prevalence and concentration in the Los Angeles fentanyl market, 2023–2025. *Drug and Alcohol Dependence Reports*, 16, 100364. <https://doi.org/10.1016/j.dadr.2025.100364>

Friedman, J., & Shover, C. L. (2023). Charting the fourth wave: Geographic, temporal, race/ethnicity and demographic trends in polysubstance fentanyl overdose deaths in the United States, 2010-2021. *Addiction (Abingdon, England)*, 118(12), 2477–2485. <https://doi.org/10.1111/add.16318>

Han, B., Compton, W. M., Jones, C. M., Einstein, E. B., & Volkow, N. D. (2021). Methamphetamine Use, Methamphetamine Use Disorder, and Associated Overdose Deaths Among US Adults. *JAMA Psychiatry*, 78(12), 1329. <https://doi.org/10.1001/jamapsychiatry.2021.2588>

Han, B., Cotto, J., Etz, K., Einstein, E. B., Compton, W. M., & Volkow, N. D. (2021). Methamphetamine Overdose Deaths in the US by Sex and Race and

Ethnicity. *JAMA Psychiatry*, 78(5), 564.

<https://doi.org/10.1001/jamapsychiatry.2020.4321>

Hedegaard, H., Miniño, A., & Warner, M. (2020). Drug Overdose Deaths in the United States, 1999–2018. *NCHS Data Brief*, (356).

Jones, C. M., Houry, D., Han, B., Baldwin, G., Vivolo-Kantor, A., & Compton, W. M. (2022). Methamphetamine use in the United States: Epidemiological update and implications for prevention, treatment, and harm reduction. *Annals of the New York Academy of Sciences*, 1508(1), 3–22.

<https://doi.org/10.1111/nyas.14688>

Lake, M. T., Shoptaw, S., Ipser, J. C., Takada, S., van Nunen, L. J., Lipinska, G., Stein, D. J., & London, E. D. (2020). Decision-Making by Patients With Methamphetamine Use Disorder Receiving Contingency Management Treatment: Magnitude and Frequency Effects. *Frontiers in Psychiatry*, 11.

<https://doi.org/10.3389/fpsy.2020.00022>

Lewis, D., Kenneally, M., Van denHeuvel, C., & Byard, R. W. (2021).

Methamphetamine deaths: Changing trends and diagnostic issues.

Medicine, Science and the Law, 61(2), 130–137.

<https://doi.org/10.1177/0025802420986707>

- Li, M. J., & Shoptaw, S. J. (2023). Clinical Management of Psychostimulant Withdrawal: Review of the Evidence. *Addiction (Abingdon, England)*, 118(4), 750–762. <https://doi.org/10.1111/add.16093>
- Mansoor, M., McNeil, R., Fleming, T., Barker, A., Vakharia, S., Sue, K., & Ivsins, A. (2022). Characterizing stimulant overdose: A qualitative study on perceptions and experiences of "overamping". *The International journal on drug policy*, 102, 103592. <https://doi.org/10.1016/j.drugpo.2022.103592>
- Minozzi, S., Saulle, R., Amato, L., Traccis, F., & Agabio, R. (2024). Psychosocial interventions for stimulant use disorder. *The Cochrane Database of Systematic Reviews*, 2024(2), CD011866. <https://doi.org/10.1002/14651858.CD011866.pub3>
- Montero, F., Bourgois, P., & Friedman, J. (2022). Potency-Enhancing Synthetics in the Drug Overdose Epidemic: Xylazine ("Tranq"), Fentanyl, Methamphetamine, and the Displacement of Heroin in Philadelphia and Tijuana. *Journal of Illicit Economies and Development*, 4(2), 204–222. <https://doi.org/10.31389/jied.122>
- NIST RaDAR. (2022, January 1). *RaDAR Training Materials*. NIST. <https://www.nist.gov/programs-projects/radar-training-materials>
- Rush, C. R., Santos, G.-M., McMahan, V. M., Fraser, A., Clark, J., Luna Marti, X., Walker, J. E., Shoptaw, S., & Coffin, P. O. (2025). Mirtazapine reduces

- hypothetical methamphetamine demand in humans. *Drug and Alcohol Dependence*, 274, 112769. <https://doi.org/10.1016/j.drugalcdep.2025.112769>
- Schürer, S., Klingel, K., Sandri, M., Majunke, N., Besler, C., Kandolf, R., Lurz, P., Luck, M., Hertel, P., Schuler, G., Linke, A., & Mangner, N. (2017). Clinical Characteristics, Histopathological Features, and Clinical Outcome of Methamphetamine-Associated Cardiomyopathy. *JACC: Heart Failure*, 5(6), 435–445. <https://doi.org/10.1016/j.jchf.2017.02.017>
- Schwarzbach, V., Lenk, K., & Laufs, U. (2020). Methamphetamine-related cardiovascular diseases. *ESC Heart Failure*, 7(2), 407–414. <https://doi.org/10.1002/ehf2.12572>
- Shover, C. L., Friedman, J. R., Romero, R., Buhr, R., Chu, B., Tang, A., Medina, J. A., Wisk, L., Lucas, J., & Goodman-Meza, D. (2023). Longitudinal changes in co-involved drugs, comorbidities, and demographics of methamphetamine-related deaths in Los Angeles County. *Journal of Substance Use & Addiction Treatment*, 151. <https://doi.org/10.1016/j.josat.2023.209101>
- Shover, C. L., Koncsol, A. J., Godvin, M. E., Goodman-Meza, D., Pardo, B., Poimboeuf, M., Molina, C. A., Romero, R., Feng, J., & Friedman, J. R. (2025). High variation in purity of consumer-level illicit fentanyl samples in Los Angeles, September 2023–April 2025. *International Journal of Drug Policy*, 145, 104977. <https://doi.org/10.1016/j.drugpo.2025.104977>

- Sisco, E., Pyfrom, E. M., Banta-Green, C. J., Shover, C. L., Molina, C. A., Biamont, B., & Robinson, E. L. (2024). Beyond fentanyl test strips: Investigating other urine drug test strips for drug checking applications. *Forensic Chemistry*, 40, 100594. <https://doi.org/10.1016/j.forc.2024.100594>
- Sisco, E., Verkouteren, J., Staymates, J., & Lawrence, J. (2017). Rapid detection of fentanyl, fentanyl analogues, and opioids for on-site or laboratory based drug seizure screening using thermal desorption DART-MS and ion mobility spectrometry. *Forensic Chemistry*, 4, 108–115. <https://doi.org/10.1016/j.forc.2017.04.001>
- Smith C. B. (2010). Socio-spatial stigmatization and the contested space of addiction treatment: remapping strategies of opposition to the disorder of drugs. *Social science & medicine* (1982), 70(6), 859–866. <https://doi.org/10.1016/j.socscimed.2009.10.033>
- Smith, J., Philbrick, S. M., Akiba, C. F., Wenger, L. D., Hairgrove, S., Kral, A. H., Lambdin, B. H., & Patel, S. V. (2026). Drug checking services: barriers and facilitators to implementation by U.S. syringe services programs. *Harm reduction journal*, 23(1), 55. <https://doi.org/10.1186/s12954-026-01413-1>
- Somma, V., Osekowski, M., Paratz, E., & Bonomo, Y. (2023). Methamphetamine-associated cardiomyopathy: An addiction medicine perspective. *Internal Medicine Journal*, 53(1), 21–26. <https://doi.org/10.1111/imj.15990>

- Tobolski, J., Sawyer, D. B., Song, S. J., & Afari, M. E. (2022). Cardiovascular disease associated with methamphetamine use: A review. *Heart Failure Reviews*, 27(6), 2059–2065. <https://doi.org/10.1007/s10741-022-10261-7>
- Toske, S. G., & McKibben, T. D. (2022). Monitoring methamphetamine in the United States: A two-decade review as seen by the DEA methamphetamine profiling program. *Drug Testing and Analysis*, 14(3), 416–426. <https://doi.org/10.1002/dta.3186>
- Trivedi, M. H., Walker, R., Ling, W., Cruz, A. dela, Sharma, G., Carmody, T., Ghitza, U. E., Wahle, A., Kim, M., Shores-Wilson, K., Sparenborg, S., Coffin, P., Schmitz, J., Wiest, K., Bart, G., Sonne, S. C., Wakhlu, S., Rush, A. J., Nunes, E. V., & Shoptaw, S. (2021). Bupropion and Naltrexone in Methamphetamine Use Disorder. *New England Journal of Medicine*, 384(2), 140–153. <https://doi.org/10.1056/NEJMoa2020214>
- United States Sentencing Commission. (2024, June 10). *Methamphetamine Trafficking Offenses in the Federal Criminal Justice System*. United States Sentencing Commission. https://www.ussc.gov/sites/default/files/pdf/research-and-publications/researchpublications/2024/202406_Methamphetamine.pdf
- Wang, G., Smith, L., Volkow, N., Telang, F., Logan, J., Tomasi, D., Wong, C., Hoffman, W., Jayne, M., Alia-Klein, N., Thanos, P., & Fowler, J. (2012). Decreased

dopamine activity predicts relapse in methamphetamine abusers. *Molecular Psychiatry*, 17(9), 918–925. <https://doi.org/10.1038/mp.2011.86>

Welsh Emerging Drugs and Identification of Novel Substances Project. (2026).

Sample results. WEDINOS.Wales. <https://wedinos.wales/sample/>