

# UCSF

## UC San Francisco Previously Published Works

### Title

Bridging the Gap Between Practice Guidelines and the Therapy Room: Community-Derived Practice Adaptations for Psychological Services with Transgender and Gender Diverse Adults in the Central United States.

### Permalink

<https://escholarship.org/uc/item/58d8418q>

### Journal

Professional Psychology: Research and Practice, 53(4)

### ISSN

0735-7028

### Authors

Hope, Debra  
Holt, Natalie  
Woodruff, Nathan  
[et al.](#)

### Publication Date

2022-08-01

### DOI

10.1037/pro0000448

Peer reviewed



# HHS Public Access

Author manuscript

*Prof Psychol Res Pr.* Author manuscript; available in PMC 2023 November 22.

Published in final edited form as:

*Prof Psychol Res Pr.* 2022 August ; 53(4): 351–361. doi:10.1037/pro0000448.

## **Bridging the Gap Between Practice Guidelines and the Therapy Room: Community-Derived Practice Adaptations for Psychological Services with Transgender and Gender Diverse Adults in the Central United States**

**Debra A. Hope,**

University of Nebraska-Lincoln

**Natalie R. Holt,**

University of Nebraska-Lincoln

**Nathan Woodruff,**

Trans Collaborations Local Community Board Richard Mocarski

**Heather Meyer,**

University of Nebraska at Kearney

**Jae A. Puckett,**

Michigan State University

**Joshua Eyer,**

University of Alabama

**Shelley Craig<sup>\*</sup>,**

University of Toronto

**Jamie Feldman<sup>\*</sup>,**

University of Minnesota

**Jay Irwin<sup>\*</sup>,**

University of Nebraska - Omaha

**John Pachankis<sup>\*</sup>,**

Yale University

**K. J. Rawson<sup>\*</sup>,**

Northeastern University

**Jae Sevelius<sup>\*</sup>,**

University of California, San Francisco

**Sim Butler**

---

Address correspondence to: Debra A. Hope, Ph.D., Department of Psychology, University of Nebraska-Lincoln, Lincoln, NE 68588-0308. dhope1@unl.edu.

<sup>\*</sup>Shelley Craig, Jamie, Feldman, Jay Irwin, John Pachankis, K. J. Rawson, and Jae Sevelius are members of the Trans Collaborations National Advisory Board. They are listed in alphabetical order.

The authors report no conflicts of interest.

University of Alabama

## Abstract

Individuals who identify as transgender or gender diverse (TGD) are presenting at mental health clinicians' offices with increasing frequency. Many TGD clients are seeking care related to affirming their gender identity but also may present with anxiety, depression, trauma, substance abuse, or other problems for which a clinician may commonly provide services. Some clinicians may hesitate to accept TGD clients into their practice if they have little specialized training to work with this population in an affirming manner, especially in more underserved areas where a generalist practice is the norm. Numerous professional associations and experts have developed guidelines for affirmative behavioral health care for TGD people. However, what is needed are community informed recommendations to bridge from the official guidelines to clinicians' in-session activities. The Trans Collaborations Practice Adaptations for Psychological Interventions for Transgender and Gender Diverse Adults are derived from iterative interviews with TGD community members and affirming mental health clinicians in the Central United States. The 12 practice adaptations are intended to guide clinicians to adapt their usual treatment approach to be TGD affirming, especially in underserved and rural areas. The practice adaptations cover numerous aspects of practice including the office setting and paperwork, understanding gender identity and incorporating it into the case conceptualization, therapist's self-awareness, and referrals. The Trans Collaborations Practice Adaptations will help clinicians work confidently and competently with adult TGD clients, regardless of the presenting problem, to ensure TGD communities receive the best interventions for their behavioral health concerns.

---

A client seeking treatment for depression crosses out male/female and writes *genderqueer* on the intake paperwork. The mental health clinician is familiar with non-binary gender identities but is unsure about whether to ask about it in session and wonders if it is relevant for the cognitive therapy that is their typical intervention for depression. In another office, a client who initially presented as a cisgender man seeking help for stress following the dissolution of a romantic relationship has indicated they would like to explore their gender identity and the gender dysphoria they have experienced for many years. The mental health clinician is immediately worried they might ask for a letter for gender affirmation surgery and wonders if they should let the client know right away that is outside of their scope of practice. Unless a clinician specializes in gender therapy or has had many similar cases, it is unlikely they know how to provide culturally-informed care for these clients. Yet, if they are outside of a large urban center, referral options may be limited. This paper is for the many mental health clinicians who have noticed an increase in similar clients in their practices but find many of the materials on how to provide culturally appropriate care either too general to be useful or based on well-resourced urban areas that do not represent their community.

## Transgender and Gender Diverse Clients and the Need for Psychological Services

National surveys indicate that individuals who identify as transgender or gender diverse (TGD<sup>1</sup>) are more likely than not to seek psychological services at some point in

their lives (James et al., 2016). Reasons for seeking services vary by individual and often reflect the same reasons cisgender people seek services, but two factors probably increase the likelihood of seeking services as well. First, structural oppression resulting in marginalization<sup>2</sup> stress experienced by TGD people (Hendricks & Testa, 2012) increases risk for a variety of mental health problems including anxiety (e.g., Bouman et al., 2017), depression (e.g., Nuttbrock et al., 2010), substance abuse (e.g., Gonzalez et al., 2017), and suicidality (e.g., Goldblum et al., 2012). Second, even when there are no mental health concerns, under current Standards of Care set by the World Professional Association for Transgender Health (WPATH; Coleman et al., 2012), access to services for medical gender affirmation, including surgeries and hormones, may require a psychosocial assessment or mental health evaluation. Despite this high need, few mental health clinicians have received training for working with TGD clients and/or the training they received has been based on outdated models that pathologized TGD identities (e.g., Holt et al., 2020). Not surprisingly, when TGD people seek mental health care, they may be refused care or receive inadequate care (Holt et al., 2020; James et al., 2016).

### Current Resources for Mental Health Clinicians Serving TGD Clients

In order to help more mental health clinicians be well-prepared to work with TGD clients, various professional organizations have established TGD-specific practice guidelines. Two prominent sets of guidelines come from the American Psychological Association (APA) and the American Counseling Association (ACA). The APA's *Guidelines for Psychological Practice with Transgender and Gender Nonconforming People* (APA Guidelines; American Psychological Association, 2015) consist of 16 guidelines across five areas, one of which is Assessment, Therapy, and Intervention. The APA Guidelines provide a succinct summary of research and basic information about TGD people followed by examples of how to apply the information. The Association for Lesbian, Gay, Bisexual, and Transgender Issues in Counseling (ALGBTIC; 2009) developed the ALGBTIC Competencies for Counseling Transgender Clients published by the ACA. Similar to the APA Guidelines, the ACA competencies are grounded in empirical research whenever possible. There are 11 to 17 specific competencies under each of eight domains (e.g. Human Growth and Development; Social and Cultural Foundations). The ACA document does not review the literature in the same way as the APA Guidelines as the format is a list of specific competencies, many of which reflect summaries of the literature (e.g., the role of marginalization stress on TGD health). Another potential resource for behavioral health clinicians who work with TGD clients is the WPATH Standards of Care (Coleman et al., 2012). The mental health section of the Standards and related materials (e.g. Fraser, 2009; Lev, 2009) emphasize clinicians' role in gender affirmation care and guidelines for writing letters for hormones or gender

<sup>1</sup>Individuals may use a variety of labels to describe their gender identity including transgender, non-binary, genderqueer, queer, transgender man, transgender woman, gender non-conforming and many others. For the purposes of this paper, the term *transgender and gender diverse* (TGD) will be used as an umbrella term for everyone whose gender identity does not match the gender/sex they were assigned at birth. Individuals whose gender identity matches their gender/sex assigned at birth will be referred to as *cisgender*. We acknowledge that terminology varies across time and across subgroups in communities and any choice excludes the label used by some people, including some of our research participants.

<sup>2</sup>We use the term *marginalization stress* instead of *minority stress* to properly locate the problem with the experience of marginalization itself, not the marginalized person, as requested by our Local Community Board that guides, advises, and engages in research efforts as part of our community-academic partnership.

affirmation surgeries, with limited guidance for general mental health care. Any general information in the WPATH Standards of Care on behavioral health care with TGD clients is less specific than either the APA Guidelines or the ACA Competencies.

These are strong professional guidelines for behavioral health care for TGD people but the guidelines are often too general to help a Clinician understand what to do moment by moment in a session. Some excellent resources that may aid clinicians including recent books by Chang et al. (2018), Kauth and Shipherd (2018), and Singh and dickey (2017), as well as a clinical article by Puckett and colleagues (Puckett et al., 2018). Additionally, Austin and Craig (2015) were among the first to address TGD behavioral health from a cognitive-behavioral (CBT) perspective. They have developed an affirmative CBT for anxiety and depression for gender diverse adolescents (Austin et al, 2018), with discussion of extending it to adults (Austin et al., 2017).

The published guidelines, competencies, and new books and research are invaluable for clinicians seeking to provide TGD-affirming care. These resources are particularly strengthened by perspectives from TGD-identified clinicians. The nascent nature of this area necessitates recommendations that are grounded in the needs of diverse TGD communities. Especially given the history of marginalization of TGD people by mental health communities (Ciszek et al., 2021; Riggs et al., 2019), it is essential that the voices of TGD individuals are central in identifying what constitutes TGD-affirmative behavioral health care.

## **Bridging the Gap between Practice Guidelines and Real-World Practice**

*The goal of the Trans Collaborations Practice Adaptations for Psychological Services for TGD Adults described here is to inform psychological services, especially psychotherapy, for adult TGD clients through the lens of TGD communities and clinicians in highly underserved regions of the US. Our primary audience is clinicians with less experience working with TGD communities (a large majority of clinicians in underserved areas), but some Practice Adaptations also will help a more experienced clinician adapt to a rural context. The Practice Adaptations are meant to be a data-driven bridge between professional statements such as APA's Guidelines (APA, 2015) and the daily in-session work of mental health clinicians. Most resources emphasize gender affirming-related care and treatment for gender dysphoria. The Practice Adaptations may be helpful for such care but the emphasis is on adapting a clinician's general approach to make interventions more appropriate for TGD clients seeking care for a variety of problems. Given the evidence in the literature (e.g., Mizock & Lundquist, 2016) and our own work (e.g., Meyer et al., 2020) that marginalization within the therapy context is common, we placed particular emphasis on the many ways stigma can be enacted in the therapeutic context. The Practice Adaptations were informed by the voices of TGD people living in the U.S. Central Great Plains with rural and urban environments that differ in terms of available services and cultural context from the urban coastal areas where most research and professional publications originate. Thus, they are meant to be culturally sensitive to these cultures in the middle of the U.S. in particular. However, overlap with other published recommendations suggest they may be universally good practice for psychological services with TGD adults. The Practice Adaptations also*

were informed by voices of clinicians working in a variety of outpatient mental health settings who understand the nuances and demands of clinical practice.

## Community Participation to Inform the Practice Adaptations

The primary methodology and results for the study underlying the development of the Practice Adaptations are detailed elsewhere (Holt et al., 2020; Meyer et al., 2020; Mocarski et al., 2019) and available on our website ([go.unl.edu/transcollaborations](https://go.unl.edu/transcollaborations)). The Practice Adaptations were developed via data collection and feedback from our Community Board comprised of six TGD individuals, TGD adults in the community, and behavioral health clinicians (e.g., psychologists, social workers, counselors) they identified as TGD-affirmative. Given the intention to develop materials useful to clinicians in non-coastal, underserved areas, all community members and clinicians resided in the Central Great Plains. Our National Advisory Board of scholars from a variety of disciplines provided input on the various drafts. All procedures for the underlying studies to support development of the Practice Adaptations were approved by the Institutional Review Board at the University of Nebraska-Lincoln.

To be as broadly applicable as possible, the Practice Adaptations focus on aspects of psychological services that are common across therapeutic approaches. These so-called *common factors* include the therapeutic alliance, a credible rationale and intervention, and a confidential setting that reinforces the credibility of the clinician and process (e.g. Frank & Frank, 2004; Kazantzis et al., 2017; Wampold, 2015). Although the role of common factors on clinical outcomes has been debated (e.g. Hofmann & Barlow, 2014), there is substantial evidence that these common factors are important in reducing symptoms (Lambert, 2009). Common factors are a necessary, if not always a sufficient part, of successful outcomes (Hofmann & Barlow, 2014; Laska et al., 2014). Given the Practice Adaptations are focused more on commonalities across therapeutic approaches, recommendations to employ a clinician's usual approach is noted at several places where a more specific intervention is indicated. This broad application also ensures that many clinicians can utilize these Practice Adaptations as opposed to those specializing in a particular approach (e.g., CBT).

## Trans Collaborations Practice Adaptations for Psychological Services for TGD Adults

The following 12 Practice Adaptations were drawn directly from the community and clinician interviews and advisory board feedback. A list of the Practice Adaptations are presented in Table 1. Reviews of the relevant literature were considered where available and appropriate, with greater emphasis placed on empirical studies than case reports. Ordering of the Practice Adaptations does not reflect order of importance and the Practice Adaptations are of varying complexity. In certain places the gender identity of clinicians is mentioned explicitly but it should be noted that cisgender and TGD clinicians will use these Practice Adaptations according to how their own identities show up in the therapy room. The Practice Adaptations assume that clinicians have basic knowledge of gender identity and terminology relevant to the experiences of TGD people. Many clinicians do not have this basic knowledge and should seek continuing education training on working with TGD

clients and consult relevant scientific literature (Whitman & Han, 2017). This information is available from many sources, including recent books (Kauth & Shipherd, 2018; Singh & dickey, 2017;), and APA guidelines and ACA competencies, so it is not provided here.

**Practice Adaptation 1: Setting. *Provide a TGD affirmative public face to services, including the website, office features, paperwork, referral process, and office staff.***

The therapeutic setting (e.g. Frank & Frank, 2004) predominantly refers to the physical space where therapy occurs. However, contact with clients begins before they walk into the therapy office, such as through online materials, calling to set an appointment, and completing intake materials. The client's progression into the actual therapy office can communicate messages of inclusion or marginalization.

In our interviews and focus groups, the first step in seeking a clinician for many TGD individuals was to seek recommendations from their online or in-person community, if they have one. From there, many potential clients seek online information and described scouring websites for signals a clinician may be competent and affirmative for TGD clients. Elsewhere we have described ways clinicians can signal they are affirmative (Holt et al., 2020). The messages can be subtle if needed. In some rural communities, clinicians we interviewed reported there can be hesitancy to be too explicit in public materials for fear of backlash that impairs their ability to make a living.

Once a TGD individual has selected a clinician for an initial consultation, they encounter the physical space of the therapeutic setting. Cochran and colleagues (2018) and Puckett et al. (2018) provided excellent recommendations for making one's office space and procedures TGD-affirming, many of which are similar to what our participants reported. Community participants in our interviews identified that markers of stigma and lack of support in the waiting room, paperwork, and initial encounters with a clinician or support staff often act as barriers to care or cause stress and anxiety. For example, a binary option for gender on intake paperwork conveys a stigmatizing message, whereas a blank space for specifying gender and the name you wish to be called allows maximum flexibility for a wide variety of gender identities and individual situations (Holt et al., 2019; Puckett et al., 2020). (Another option is to defer discussion of these identity markers until in the privacy of the therapy room.) Stigma can be enacted via interactions with all staff members, necessitating training staff on use of correct names and pronouns and avoiding comments on changes in clients' appearances.

In certain settings, clinicians have less control over the messages conveyed by their setting such as an affiliation with a conservative religious health organization, group practice settings where the clinician is not the owner, or a geographic location where privacy may be limited. For example, one participant described being distressed by visual reminders that their clinician worked within a Catholic health organization and that they would likely terminate care prematurely, despite having positive experiences with that individual clinician. Such situations are compounded in rural spaces where TGD individuals may be

bound to clinicians associated with unwelcoming institutions or influenced by non-affirming state and local policies because there are no other options. The onus is on clinicians to help TGD clients negotiate stigmatizing spaces and understand what limitations, if any, the clinician may have in helping them. The clinician should also be taking steps to decrease the sources of stigma within their workplace so that clients can be more affirmed in seeking care.

**Practice Adaptation 2: Perceptions of Cultural Messages. *Explore the meaning of cultural events and messages individually with each client because cisgender people and TGD people may interpret such messages differently.***

Consuming TGD-related media from a variety of sources is helpful for staying up-to-date on terminology and current issues that are important to the community. However, even if cisgender clinicians and their TGD clients hear the same cultural messages about TGD issues and see the same TGD public role models, the perception of those messages and role models may vary. For example, our interviews indicated cisgender clinicians might view TGD celebrities as providing useful role models or having a pro-TGD educational message for society. The messages may be more complex or even negative for a TGD client. Public TGD figures often have a great deal of privilege to finance their gender affirmation that is not available to (or even desired by) other TGD people. Also, some TGD communities, such as transgender men of color, are rarely represented in the media (Capuzza & Spencer, 2017; Rood et al., 2017). Furthermore, many TGD media portrayals show TGD people as a form of trans normativity, where the TGD character is upper-middle class, white, post-gender affirmation procedures, heterosexual, and, paradoxically, without sexuality—all while being played by a cisgender actor (Mocarski et al., 2019).

**Practice Adaptation 3: Clinician's Experience of Gender. *Be aware of the experience of your own gender and how you may automatically enact gender roles that may convey either a marginalizing or an affirmative message to TGD clients.***

At the heart of the therapeutic alliance is a relationship between two human beings, albeit a relationship with different roles and constraints than typical day-to-day interactions (Newhill et al., 2003). However, both the clinician and client bring in perspectives, skills, habits and expectations from their other relationships that have gendered overtones. For cisgender clinicians in particular, heightened awareness of how they automatically enact gender roles based on their perception of the other person's gender is important. For example, a clinician who is a cisgender man holding the door for a TGD client to precede them into the therapy space is enacting dominant cultural messages about men holding doors. A transgender woman who enacts traditional gender norms may perceive this act as neutral or perhaps affirming. A transgender man may have the opposite reaction. A transgender man may step aside to allow a cisgender woman to enter first, as an act of claiming an expression of



their gender identity. The purpose here is not to give guidance on who should enter a room first but to heighten awareness of how we often convey gendered messages automatically. TGD clinicians similarly must be aware of their experiences of gender given that we are all subject to the internalization of gendered messages and expectations (Butler, 2011).

**Practice Adaptation 4: Comfort with Various Gender Identities/Expressions.**  
***Recognize and be comfortable with variability and fluidity of gender identity and expression that may vary in unpredictable ways over time for a given client. Your role is to support and affirm each individual gender journey, not diagnose their gender identity or impose a particular journey on the client.***

One theme that emerged in our interviews was the importance of clinicians being affirming of, and often even encourage, a client's experience of ambiguity and/or fluidity of their gender identity as well as recognize that no two gender journeys are identical. Through our interviews, it became clear that more experienced clinicians understand that for some clients this self-discovery and exploration of identity terms can be a circuitous process. The gate-keeping role for access to medical gender affirmation services can create a sense the client must "prove" their gender identity to the clinician. Clinicians can address this directly with clients by telling them whatever they say about their gender identity is acceptable and it is not for the clinician to decide what is "right" nor impose standards based on the clinician's own gender identity journey. Sometimes gender identity fluidity may itself be the resolution (e.g. Fiani & Han, 2019). Therapeutic approaches (e.g. many cognitive behavioral therapies) that include setting and monitoring explicit treatment goals can overtly or covertly convey messages that there is a correct choice or a specific timeline, and once a choice is made, it must not be changed. Witnessing and supporting the process without demanding a resolution is experienced as helpful for TGD clients. Thus an intervention for anxiety or depression, for example, can proceed in the context of gender identity fluidity rather than requiring that it be resolved before addressing presenting problems or expecting the anxiety or depression to remit only when a person experiences no shifts in their gender experience.

Although treatment of gender dysphoria and gender affirmation care are not the primary focus of these Practice Adaptations, clients may seek treatment for a behavioral health or substance abuse problem at the same time they are taking steps on their gender journey. Clinicians who do not specialize in gender affirmation services may have a prototypical view of the process. An important theme across the interviews was that each journey of gender affirmation is unique. Although some clients went to a clinician to see if they qualified for a diagnosis of gender dysphoria, most clinicians described themselves more as a witness and sounding board rather than a diagnostician. The role of clinicians seems to be supporting the self-discovery process rather than directing it. Clinicians can encourage this process by facilitating exploration of emotions and cognitions such as celebrating achievement of desired outcomes around gender affirmation. Clinicians can also help with problem solving and planning (e.g., developing contingencies for handling negative outcomes when coming out to family members). This should be incorporated as needed into an overall treatment plan.

**Practice Adaptation 5: Managing Marginalization. *Recognize the pervasive stigma and marginalization that many TGD people face in large and small ways in their daily lives. Trust the client's perceptions of their experiences as you help the client find appropriate coping responses, drawing on the client's strengths, resilience, and support system.***

Experiences of bias and discrimination are, unfortunately, an inevitable part of a TGD individual's life experience. In both large and small ways from family, friends, strangers, institutions, and cultural messages of cisgender hegemony, TGD people encounter oppression and marginalization (Puckett et al., 2021). This includes experiencing job loss, misgendering, rejection, public outing, systemic barriers, and verbal and physical aggression (James et al., 2016; Klein & Golub, 2016; Testa et al., 2012), which can be exacerbated for rural TGD communities compared to urban counterparts. Many of the participants in our interviews displayed resiliency when facing these challenges and, for some, learning to manage stress from these adversities was part of their gender journey. Regardless of their own gender identity, clinicians may not immediately see bias when it occurs and must listen carefully to the client's experiences. Then they must balance helping their TGD clients cope with this adversity while checking their own outrage at what occurred. The ability to be angry at unfair treatment can be a privilege not shared by the TGD client who is often trying to cope with the reality in which they live. On the other hand, the client may feel validated and seen knowing that their clinician shares their views on this marginalization. As such, this is nuanced and there is no one-size fits all approach that will work for all clients.

Two aspects of experiences with marginalization tend to be addressed in therapy. First, TGD clients may benefit from interventions to improve coping with the stress of marginalization itself. This could include specific stress management interventions such as relaxation, making life changes to live in a more TGD-affirmative space, and/or seeking online or in-person social support. Second, and perhaps the greater challenge, is disentangling the various aspects of the marginalization experience including the circumstances, other people's motivations and behaviors, and the client's reactions. A common strategy for coping with negative experiences with a cisgender client experiencing anxiety or depression includes exploration of whether their perceptions of the situation may be distorted (e.g. overestimation of the likelihood or cost of negative outcomes; Beck, 2011). As noted by Chapman et al. (2013), such an approach is problematic for experiences of marginalization, especially if the clinician has a majority identity. Challenging a client's experience of biased treatment or discrimination extends the marginalization into the therapy session by suggesting they do not know what happened to them. For example, a client who is going to a family reunion for the first time after a social transition may be anxious about the reactions of family members and potential disruption of the occasion. The clinician can help the client decide whether it is important to them to attend, whether they can safely attend, and how they will manage any unsupportive or hostile reactions. Another aspect that can be explored is that the TGD client is not responsible for how others react to them. Identifying the true source of the problem (other people or social institutions) is key to empowerment rather than continued marginalization in therapy.

**Practice Adaptation 6: Non-Defensive Stance. *Take responsibility for minimizing marginalization experiences within the therapy room and be open and non-defensive when you make errors or misunderstand a client.***

Despite best intentions, clinicians are likely to make occasional stigmatizing errors in session such as using the wrong pronoun or wrong name. Interviewees reported that repeated instances of microaggressions (e.g. Nadal et al., 2012) were highly problematic in therapy. An occasional error can be handled within the ongoing work, based on the clinician's own style and case conceptualization of the client. In our interviews, it was clear that the reactions of others are particularly important to clients who are in the midst of an active gender affirmation process and positive reactions can increase self-efficacy and mood. This suggests that how a clinician handles a rupture in the therapeutic alliance (e.g., Safran & Muran, 2000) when they stigmatize a client should include careful consideration of the meaning of the event to the client, as this will be highly variable across time and across clients. When clinicians make mistakes, it is important to acknowledge that these have occurred and make a repair while not monopolizing the space or launching into extended discussion of the clinician's knowledge, discomfort, or other feelings as this shifts the focus away from the client's care and onto the clinician, which can recapitulate marginalization.

**Practice Adaptation 7: Clinicians Educate Themselves. *Take responsibility for educating yourself about the local cultural context, social and legal climate, and local available services for TGD communities. Be knowledgeable about clinical guidelines, mental health literature, and needs of TGD clients. Clients are not responsible for educating clinicians and deserve the highest quality, culturally responsive services.***

One of the most frequent themes in the interviews with TGD community members was their frustration at having to educate clinicians about what it means to be TGD and the care they need. Participants recount bringing stacks of books or papers to clinicians that may not have ever been read or returned. Clinicians can glean pertinent information on their own from sources such as the ACA Competencies (ALGBTIC, 2009). However, the clinician must also delve into the context in which clients live including TGD resources, affirmative and discriminatory employers, clinicians, and religious communities, availability of appropriate healthcare referrals, relevant local, state, and national legal protections or lack thereof, and the cultures found in the local TGD communities. Both clinicians and community members emphasized that it is preferable for clinicians to be involved in the community by attending public events such as pride celebrations or taking on advocacy roles. On the other hand, joining TGD-related organizations or using events as a marketing opportunity was seen as exploitive by our interviewees. Increased likelihood of encountering clients outside of the office (whether by the clinician being visible in the local TGD community or simply by living in a small town) necessitates an early discussion of the likelihood that could happen and how both parties will handle it. Furthermore, community involvement by the clinician

means that discussion of such activities could become part of the therapy hour. The clinician has the responsibility to maintain a therapeutic focus.

**Practice Adaptation 8: Share Power. *Take responsibility for sharing power with TGD clients as much as possible, especially given the history of marginalization in the healthcare system. Recognize the impact of the gatekeeper role (clinicians providing letters for medical gender affirmation) on exacerbating the power differences that are always part of therapy.***

TGD clients come to mental health services in the context of a system that has historically marginalized TGD people and many clients have had personal negative experiences (e.g. Mizock & Lundquist, 2016). Mental health clinicians' gatekeeper role for access to medical gender affirmation services can exaggerate the power difference inherent in the therapeutic relationship (shuster, 2016). This difference can be heightened in underserved locales with few options for care and increased opportunities for multiple relationships. Even for clients who were not actively seeking a letter, the specter of the gatekeeper role made power differentials between clients and clinicians more salient.

Our data revealed a wide variety of responses about the clinicians serving as evaluators for medical gender affirmation. Some TGD clients experienced the letter requirement as demeaning while other participants were more comfortable with the requirement. TGD clients seeking a letter may resist disclosing doubts about affirming their gender, mental health problems, or other important information because they fear it will prevent them from getting a desired letter. Clinicians in this dual role of evaluator and clinician may or may not be aware of what a client is holding back and this can interfere with care. Awareness of the power differential can help clinicians thoughtfully negotiate it within their own theoretical orientation and style, just as they would negotiate other dual role relationships (Zur, 2016). For example, in our interviews, many clinicians chose to state up front what their policies were regarding provision of letters which can provide transparency to this process.

**Practice Adaptation 9: Incorporate Gender into Case Conceptualization. *Include the entire identity and experience of TGD clients in case conceptualization including past, current, and future gender identities and expression, experiences of marginalization, the affirmative or stigmatizing context of their lives (family, work, legal system), gender roles, sexual orientation, gain/loss of gender-related privilege, and intersectionality with other identities (e.g., race/ethnicity, religion, socioeconomic status, age, ability).***

Case conceptualization has been described as the "heart of evidence-based practice" (Bieling & Kuyken, 2003) and a road map to guide treatment (Aston, 2009), integrating relevant

scientific findings and theory and the client's presenting problem, specific characteristics and context through the clinician's expertise and judgement (APA, 2006). Incorporating gender into the case conceptualization for TGD clients means consideration of their gender, associated gender roles, *and* their status as an oppressed group. Additionally, all of this can change over time if the client is actively engaged in medical or social affirmation of their gender during the course of therapy. Of course, intersectionality means gender is occurring in the context of other sociocultural identities which must be considered within case conceptualization.

The experiences and gender presentations of TGD communities are far too varied to have a generic "transgender, gender diverse" case conceptualization. As a clinician comes to understand a client's gender at a given point in time, the clinician must also understand the meaning of the gender the client holds, how gender will be enacted (e.g., gender roles), how others are likely to perceive and react to the client's gender and gendered behaviors, and the client's experience of those reactions. This is not the entire case conceptualization for most clients but is a crucial context for understanding any presenting problems such as anxiety, depression, or substance abuse, or decisions about gender affirmation. We would argue that while this aspect of case conceptualization may appear more complex for TGD clients due to our default assumptions and experience with a traditional gender binary, a similar process could be used to incorporate a deep understanding of gender in all case conceptualizations.

### Specifics of TGD Case Conceptualization

Several aspects of TGD peoples' experiences of gender are especially crucial to consider when formulating a case. First, although a case conceptualization for a cisgender individual may include the experience of marginalization or privilege for women and men (in the context of other identities such as race/ethnicity), marginalization stress for a TGD person warrants special attention as the experience of being a transgender woman, even if one is rarely misgendered, differs from being a cisgender woman for example. Even our research participants who reported rarely being misgendered, described times when their TGD identity became salient or was disclosed. Second, the case conceptualization should incorporate the TGD client's experience of change in their gender, if applicable to them. This could include both positive and negative aspects of their own sense of self and self-identity, positive and negative reactions from others, and the experience of gaining or losing cultural privilege related to perceived gender. Third, written documentation of the case conceptualization should utilize identity terms and descriptors that the client uses themselves to help situate the client's experience and ensure respectful and affirming communication. Fourth, regardless of whether a clinician tends to include the environmental context in their case conceptualizations, inclusion is essential for TGD clients. Consistent with research (e.g., Tebbe & Moradi, 2016), many of our clinicians highlighted the importance of a good support network and emphasized it in their clinical work with TGD clients. The environment can offer support that helps buffer stressors, or it may create proximal stressors through lack of acceptance, overt threats, or biased treatment. The larger environmental context also affects TGD clients, such as anti-TGD legislation (e.g., Hatzenbuehler & McLaughlin, 2014). An understanding of the client's experience in their environment and extent to which they engage with media messages beyond their direct experience can help inform the case

conceptualization. This will allow the clinician to have a better understanding of whether particular cognitive or emotional reactions should best be understood as originating from within the client or as a reaction to environmental context, or a combination of the two.

**Practice Adaptation 10: Identify Focus of Services. *Recognize that a TGD identity may not be the primary organizing identity for an individual and TGD-related concerns may not be the primary focus of therapy.***

The TGD individuals who participated in our study identified as part of the community to the extent they were willing to participate. However, many also did not see their TGD identity as an organizing part of their daily lives. In the interviews, participants reported feeling stigmatized when clinicians assumed that any psychological or medical problems were associated with being TGD, and sometimes they were unable to access the same care a cisgender individual would receive for similar presenting complaints.

**It is not always about being TGD.**

When working with TGD clients, clinicians may assume that the goals of therapy will be related to gender affirmation or gender dysphoria. However, TGD clients seek therapy for anxiety, depression, substance abuse, or a host of other problems for which any person seek services. Identifying as TGD should always be part of the context, but medical or social affirmation and gender dysphoria may not be the focus of intervention. In fact, gender dysphoria is not always be an appropriate diagnosis for clients living in their affirmed gender without related distress. Case conceptualization can guide treatment selection for the presenting problem and help clinicians identify any needed adaptations for a TGD client. For example, incorporating an understanding of current and past gender identity will add important nuance to a treatment plan that includes a general intervention such as increasing assertiveness skills to improve social relationships as part of a treatment plan for depression.

**Practice Adaptation 11: TGD-Affirmative Referrals. *Consider the implications of referrals to needed services that may not be TGD-affirming and may exacerbate the problem such as a referral for hospitalization in a non-affirming facility. Problem-solve to get the best possible option, even if it is not a traditional solution.***

Given many TGD clients have a personal or vicarious experience of marginalization in healthcare settings, they may need support to follow through on a referral, even if the new clinician is likely to be affirming. Making referrals for additional services, such as drug and alcohol detoxification or hospitalization for suicidality is a routine part of most clinical practice. Especially in rural settings with few options, these referrals can present additional challenges if there are no TGD-affirming options. Hospitalization for a transgender woman with acute suicidal risk who will be placed in a hospital known to be non-affirming with a male roommate and called by the wrong name can exacerbate the crisis. Clinicians in our

interviews described the importance of thinking through the implications of a referral with a client and creatively problem solving to get the best possible option.

**Practice Adaptation 12: Advocacy. *Be prepared to advocate for TGD clients and plan ahead about how such advocacy fits within the overall approach and case conceptualization for a given client. A clinician and client may decide together that the clinician's privilege that comes from being well educated and/or cisgender can be used to a client's benefit.***

Many of the interviews included descriptions of clinicians as advocates for their TGD clients, in line with previous literature (Hope et al., 2020; Singh & Burnes, 2010). This included making telephone calls, writing letters, identifying support groups, and similar activities. A clinician who calls a physician's office to gauge whether they are TGD-affirming may get greater detail and buffer the client from ignorant or inappropriate responses. Although many clinicians are more accustomed to helping clients learn to advocate for themselves and that may be a long-term therapeutic goal, in our interviews the clinician using privilege they may have (e.g., education, cisgender) to help mitigate the client's marginalization was an important component of advocacy.

## Discussion and Implications

There is a growing array of high-quality sources for mental health clinicians who serve TGD communities, including professional guidelines, lists of competencies and rich clinical writing from clinicians with extensive experience. Although some of those sources offer more general guidance (Austin & Craig, 2015; Puckett et al., 2018; Richmond, et al., 2017), much of it is focused on treating gender dysphoria and social, medical, and legal gender affirmation. In contrast, the Practice Adaptations are meant to be appropriate for TGD people seeking care for a variety of presenting problems. Using the lens of TGD communities and clinicians in rural and underserved locales, the purpose of these Practice Adaptations is to fill a gap between current resources and what happens with specific clients in the therapy room in a concise document. The Practice Adaptations focus mostly on common factors or more general aspects of psychological interventions to be most broadly applicable. A key strength of these Practice Adaptations is that they evolved from an iterative research process (Holt et al., 2020; Meyer et al., 2020) involving TGD community voices, community mental health clinicians, and academic scholars from a variety of disciplines.

Although culturally adapted approaches improve outcomes in other minority groups (e.g., Griner & Smith, 2006), there is little information about the efficacy of psychosocial interventions, culturally adapted or not, for TGD clients. Other than interventions focused on HIV risk (e.g., Garofalo et al., 2016), there is a paucity of affirming research on clinical interventions with TGD adults (Budge & Moradi, 2018; Budge et al., 2020). Evidence-based information on how to adapt the most effective psychological interventions to meet the needs of TGD communities can spur clinical trials to identify best practices and help reduce

mental health disparities. We envision these Practice Adaptations would allow a clinical scientist to reformulate a particular treatment protocol for a disorder or group of disorders to make it TGD-affirmative and then test the efficacy of the resulting protocol with TGD participants. We are currently undertaking such an effort.

### Limitations

Several limitations to the Practice Adaptations should be noted. First, the Practice Adaptations are limited by the sources of data from which they are drawn. We purposefully collected data from TGD communities in the Central Great Plains as the experiences of these TGD communities are underrepresented in the research and clinical literature. It is possible additional Practice Adaptations might have emerged if the data were collected in a different part of the U.S. or in a different health system. However, overlap with clinical recommendations such as those found in Kauth and Shipherd (2018), Puckett et al. (2018), and Singh and dickey (2017) suggest the Practice Adaptations are more broadly applicable. Second, all of our community participants were adults and interviews with clinicians focused on care for adults. It is unknown whether the adaptations are appropriate for behavioral health care for children and youth. There are other excellent resources on evidence-based interventions for adolescents (e.g., Austin & Craig, 2015). Finally, the Practice Adaptations do not stand on their own. We assumed that clinicians would have background on TGD communities including resiliency, marginalization stress, HIV risk and other health disparities, aspects of social, legal, and medical gender affirmation, and the local context. This knowledge is crucial for TGD affirmative services and represents basic competency. This knowledge also will continue to evolve, necessitating clinicians engage in continuing education and life-long learning about issues impacting TGD communities. The focus of the Practice Adaptations is to bridge the gap between that background and psychological services from the perspective of TGD clients in underserved areas, especially for interventions for general mental health concerns.

### Conclusion

Professional understanding of what constitutes high quality health care for individuals who identify as TGD has evolved substantially since the first Harry Benjamin Standards of Care in 1979. Similarly, the TGD communities have changed as more people have access to the knowledge and language to recognize themselves as having a gender that differs from the sex they were assigned at birth and there is increased visibility of TGD experiences. However, structural oppression and resulting mental health disparities remain pervasive. Reducing barriers to behavioral health care means that serving TGD clients must move from a highly specialized service available only in resource-rich areas to the scope of practice for the average clinician. Eventually, this work must be supported by clinical research that identifies how to provide the most effective care, which, by definition, also must be affirming. These Practice Adaptations will help achieve those dual goals.

### Acknowledgments

DEBRA HOPE received her PhD in clinical psychology from the State University of New York – University at Albany. She is current Aaron Douglas Professor of Psychology and Associate Vice Chancellor and Dean of Graduate Education at the University of Nebraska-Lincoln. She is co-founder of Trans Collaborations ([go.unl.edu/](http://go.unl.edu/))



[transcollaborations](#)). Her research interests include reducing health and mental health disparities for sexual and gender minorities through community-based participatory research.

NATALIE HOLT received her PhD in clinical psychology from the University of Nebraska-Lincoln. She is currently a VA Quality Scholars fellow at the VA Tennessee Valley Healthcare System. Her research interests focus on improving access and delivery of high quality, evidence based behavioral healthcare for transgender and gender diverse communities and reducing marginalization and stigma in healthcare settings for LGBTQ+ communities.

NATHAN WOODRUFF is co-founder of Trans Collaborations ([go.unl.edu/transcollaborations](http://go.unl.edu/transcollaborations)) and Chair of the Trans Collaborations Local Community Board. He is an experienced workshop leader for diversity and inclusion, especially in health and mental health settings.

RICHARD MOCARSKI received his PhD in communication and information sciences from the University of Alabama. He currently serves as the Associate Vice President for Research at San José State University. He is a co-founder of Trans Collaborations ([go.unl.edu/transcollaborations](http://go.unl.edu/transcollaborations)). His research interests include better understanding the sociocultural impacts on narrative exclusion that leads to marginalization and how to combat these impacts through community-based interventions with the transgender and gender diverse communities.

HEATHER MEYER received her PhD in advertising from the University of Texas at Austin. She is currently Professor of Marketing at the University of Nebraska at Kearney. Her research focuses on qualitative research methods, branding, and consumer culture. In particular, she studies how consumers utilize brands in order to signal aspects of their identity to others.

JAE PUCKETT received their PhD in clinical psychology from the University of Massachusetts Boston. They are an Assistant Professor in the Ecological-Community Psychology program at Michigan State University. Their research focuses on minority stress, coping, and resilience in transgender and gender diverse individuals.

JOSHUA EYER obtained a PhD in Clinical (Health) Psychology from the University of North Carolina at Charlotte and postdoctoral specialization in clinical health psychology research. He is an associate research professional in the Alabama Life Research Institute at the University of Alabama. His research interests include community-engaged research into sexual and gender minority health, rural integrated behavioral health, and other health projects, including an implementation project responding to the opioid crisis in rural Alabama.

SHELLEY CRAIG received her PhD in social welfare from Florida International University. She is the Canada Research Chair in Sexual and Gender Minority Youth (SGMY) and is Professor at the Factor Inwentash Faculty of Social Work at the University of Toronto. Her research interests include exploring the role of technology and developing clinical models of care to enhance the resilience of sexual and gender minorities.

JAMIE FELDMAN received her MD and PhD in anthropology from the University of Illinois. She is currently is an Associate Professor in the Department of Family Medicine and Community Health at the University of Minnesota, and faculty in the Institute for Sexual and Gender Health. Dr. Feldman is a co-author of World Professional Association of Transgender Health's (WPATH) Standards of Care. Her research interests include the intersection of gender identity, use of gender affirming hormone therapy, and health care systems.

JAY IRWIN is an Associate Professor of Sociology at the University of Nebraska at Omaha. He received his PhD in Medical Sociology (with a Graduate Certificate in Gerontology) at University of Alabama at Birmingham in 2009. Dr. Irwin's primary research focuses on LGBTQ health, particularly the health and well-being of transgender individuals. Of key interest to Dr. Irwin is the mental health of these communities, as well as access to appropriate medical care. JOHN PACHANKIS received his PhD in clinical psychology from the State University of New York – Stony Brook. He is the Susan Dwight Bliss Associate Professor at the Yale School of Public Health and director of the Yale Initiative for LGBTQ Mental Health. His research focuses on the psychological experiences of sexual and gender minority individuals, including processes of identity formation and identity concealment; the cognitive, emotional, and behavioral consequences of stigma-based rejection and exclusion; and affirmative mental health treatments for sexual and gender minority populations.

K.J. RAWSON works at the intersections of the Digital Humanities and Rhetoric, LGBTQ+, and Feminist Studies. Focusing on archives as key sites of cultural power, he studies the rhetorical work of queer and transgender archival collections in both brick-and-mortar and digital spaces. He is founder and director of the Digital Transgender Archive, an award-winning online repository of trans-related historical materials, and he is the co-chair of the editorial board of the Homosaurus, an international LGBTQ linked data vocabulary.

JAE SEVELIUS is a licensed clinical psychologist and Professor of Medicine at the University of California, San Francisco, Director of the UCSF Center for AIDS Prevention Studies (CAPS), and co-founder of the UCSF Center of Excellence for Transgender Health. Dr. Sevelius' research on gender affirmation and health focuses on the development and testing of peer-led, intersectionality-informed interventions to promote sexual health and resilience among transgender women of color affected by HIV in California and São Paulo, Brazil.

SIM BUTLER received his PhD in communication and information sciences from the University of Alabama. He currently an assistant professor, the Director of the Alabama Debate Society, and the Trans Collaboration representative at the University of Alabama. His research interests include cultural narratives, forensic speaking, and community based, participatory research and engagement.

This work was support in part by grant number R21MH108897 from the National Institute of Mental Health, a Great Plains IDeA-CTR (U54GM115458) pilot award, a University of Nebraska-Lincoln Office of Research and Economic Development Revision Award, a University of Nebraska-Lincoln College of Arts and Science ENHANCE award, and a University of Nebraska Systems Science Team Building Award.

This material is based upon work supported by the Office of Academic Affiliations, Department of Veterans Affairs. VA National Quality Scholars Program and with use of facilities at VA Tennessee Valley Healthcare System, Nashville Tennessee

The authors wish to thank the members of the local community board, TGD community members and clinicians who graciously gave their time to guide development of these Adaptations. Jamie Longwell and Heather Barnes provided transcription services and we are indebted to them for their careful work.

All procedures were approved by the University of Nebraska-Lincoln Institutional Review Board.

## References

- American Psychological Association. (2015). Guidelines for psychological practice with transgender and gender nonconforming people. *American Psychologist*, 70(9), 832–864. 10.1037/a0039906 [PubMed: 26653312]
- APA Presidential Task Force on Evidence-Based Practice (2006). Evidence-based practice in psychology. *American Psychologist*, 61(4), 271–285. 10.1037/0003-066X.61.4.271 [PubMed: 16719673]
- Association of Lesbian, Gay, Bisexual, and Transgender Issues in Counseling. (2009). Competencies for counseling with transgender clients. Author.
- Aston R. (2009). A literature review exploring the efficacy of case formulations in clinical practice. What are the themes and pertinent issues? *The Cognitive Behaviour Therapist*, 2(2), 63–74. 10.1017/S1754470X09000178
- Austin A, & Craig SL (2015). Transgender affirmative cognitive behavioral therapy: Clinical considerations and applications. *Professional Psychology: Research and Practice*, 46(1), 21–29. 10.1037/a0038642
- Austin A, Craig SL, & Alessi E. (2017). Affirmative cognitive behavioral therapy with transgender and gender nonconforming adults. *Psychiatric Clinics of North America*, 40(1), 141–156. 10.1016/j.psc.2016.10.003 [PubMed: 28159140]
- Austin A, Craig SL & D'Souza S. (2018). An AFFIRMative cognitive behavioral intervention for transgender youth: Preliminary effectiveness. *Professional Psychology: Research and Practice*, 49(1), 1–8. 10.1037/pro0000154
- Beck JS (2011) *Cognitive Therapy: Basics and Beyond*. 2nd Edition. Guilford Press.
- Bieling PJ, & Kuyken W. (2003). Is cognitive case formulation science or science fiction. *Clinical Psychology: Science and Practice*, 10(1), 52–69. 10.1093/clipsy/10.1.52
- Bouman WP, Claes L, Brewin N, Crawford JR Millet N, Fernandez-Aranda F, Arcelus J. (2017). Transgender and anxiety: A comparative study between transgender people and the general population. *International Journal of Transgenderism*, 18(1), 16–26. 10.1080/15532739.2016.1258352
- Budge SL, & Moradi B. (2018). Attending to gender in psychotherapy: Understanding and incorporating systems of power. *Journal of Clinical Psychology*, 74(11), 2014–2027. 10.1002/jclp.22686 [PubMed: 30238453]
- Budge SL, Sinnard MT, & Hoyt WT (2020). Longitudinal effects of psychotherapy with transgender and nonbinary clients: A randomized control pilot trail. *Psychotherapy*, 58(1), 1–11. 10.1037/pst0000310 [PubMed: 32567869]
- Butler J. (2011). *Gender trouble: Feminism and the subversion of identity*. New York, NY: Routledge.

- Capuzza JC, & Spencer LG (2017). Regressing, progressing, or transgressing on the small screen? Transgender characters on US scripted television series. *Communication Quarterly*, 65(2), 214–230. 10.1080/01463373.2016.1221438
- Chang SC, Singh AA, & dickey I. M. (2018). *A clinician's guide to gender-affirming care: Working with transgender and gender nonconforming clients*. New Harbinger Publications.
- Chapman LK, DeLapp CTR, & Williams MT (2013). Cognitive-behavioral treatment of social anxiety among ethnic minority parents, part 2: Bridging the gap in treatment. *Directions in Psychiatry*, 33(3), 163–176.
- Ciszek E, Mocarski R, Price S, & Almeida E. (2021). Discursive stickiness: Affective institutional texts and activist resistance. *Public Relations Inquiry*, 10(3), 295–310. 10.1177/2046147X211008388
- Cochran BN, Reed OM, & Gleason HA (2018). Providing a welcoming clinic environment. (p. 44–60). In Kauth MR, & Shipherd JC (Eds.), *Adult transgender care: An interdisciplinary approach for training mental health professionals*. Routledge.
- Coleman E, Bockting W, Botzer M, Cohen-Kettenis P, DeCuypere G, Feldman J, Fraser L, Green J, Knudson G, Meyer WJ, Monstrey S, Adler RK, Brown GR, Devor AH, Ehrbar R, Ettner R, Eyler E, Garofalo R, Karasic DH, ... Zucker K (2012). Standards of Care for the Health of Transsexual, Transgender, and Gender-Nonconforming People, Version 7. *International Journal of Transgenderism*, 13(4), 165–232. 10.1080/15532739.2011.700873
- Fiani CN, & Han HJ (2019). Navigating identity: Experiences of binary and non-binary transgender and gender non-conforming (TGNC) adults. *International Journal of Transgenderism*, 20(2–3), 181–194. 10.1080/15532739.2018.1426074 [PubMed: 32999605]
- Fraser L. (2009). Psychotherapy in the World Professional Association for Transgender Health's Standards of Care: Background and recommendations. *International Journal of Transgenderism*, 11(2), 110–126. 10.1080/15532730903008057
- Frank JD, & Frank J. (2004). Therapeutic components shared by all psychotherapies. In Freeman A, Mahoney MJ, DeVito P, & Martin D. (Eds.), *Cognition and psychotherapy*, 2nd ed (pp. 45–78). Springer.
- Garofalo R, Kuhns LM, Reisner SL, & Mimiaga MJ, (2016). Behavioral interventions to prevent HIV transmission and acquisition for transgender women: A critical review. *Journal of Acquired Immune Deficiency Syndromes*, 72(3), S220–S225. 10.1097/QAI.0000000000001084 [PubMed: 27429186]
- Goldblum P, Testa RJ, Pflum S, Hendricks ML, Bradford J, & Bongar B. (2012). The relationship between gender-based victimization and suicide attempts in transgender people. *Professional Psychology: Research and Practice*, 43(5), 468–475. 10.1037/a0029605
- Gonzalez CA, Gallego JD & Bockting WO (2017). Demographic characteristics, components of sexuality and gender, and minority stress and their associations to excessive alcohol, cannabis, and illicit (noncannabis) drug use among a large sample of transgender people in the United States. *Journal of Primary Prevention*, 38(4), 419–445. 10.1007/s10935-017-0469-4 [PubMed: 28405831]
- Griner D, & Smith TB (2006). Culturally adapted mental health intervention: A meta-analytic review. *Psychotherapy: Theory, Research, Practice, Training*, 43(4), 531–548. 10.1037/0033-3204.43.4.531
- Hatzenbuehler ML & McLaughlin KA (2014). Structural stigma and hypothalamic-pituitary-adrenocortical axis reactivity in lesbian, gay, and bisexual young adults. *Annals of Behavioral Medicine*, 47(1), 39–47. 10.1007/s12160-013-9556-9 [PubMed: 24154988]
- Hendricks ML, & Testa RJ (2012). A conceptual framework for clinical work with transgender and gender nonconforming clients: An adaptation of the minority stress model. *Professional Psychology: Research and Practice*, 43(5), 460–467. 10.1037/a0029597
- Hofmann SG, & Barlow DH (2014). Evidence-based psychological interventions and the common factors approach: The beginnings of a rapprochement. *Psychotherapy*, 51(4), 510–513. 10.1037/a0037045 [PubMed: 25111379]
- Holt NR, Hope DA, Mocarski R, Meyer H, King R, & Woodruff N. (2020). The provider perspective on behavioral health care for transgender and gender nonconforming individuals in the Central

- Great Plains: A qualitative study of approaches and needs. *American Journal of Orthopsychiatry*, 90(1), 136–146. 10.1037/ort0000406 [PubMed: 30920242]
- Holt NR, Hope DA, Mocarski R, & Woodruff N. (2019). First impressions online: The inclusion of transgender and gender nonconforming identities and services in mental healthcare providers' online materials in the USA. *International Journal of Transgenderism*, 20(1), 49–62. 10.1080/15532739.2018.1428842 [PubMed: 31217753]
- Hope DA, Woodruff N, & Mocarski R (2020). Advocacy opportunities from academic-community partnerships: Three examples from Trans Collaborations. *the Behavior Therapist*, 43(7), 247–250.
- James SE, Herman JL, Rankin S, Keisling M, Mottet L, & Anafi M. (2016). The Report of the 2015 US Transgender Survey, Washington, DC: National Center for Transgender Equality. <https://transequality.org/sites/default/files/docs/usts/USTS-Full-Report-Dec17.pdf>
- Kazantzis N, Dattilio FM, & Dobson KS, (2017). *The therapeutic relationship in cognitive behavioral therapy: A clinician's guide*. Guilford Press.
- Kauth MR, & Shipherd JC (Eds.). (2018). *Adult transgender care: An interdisciplinary approach for training mental health professionals*. Routledge.
- Klein A, & Golub SA (2016). Family rejection as a predictor of suicide attempts and substance misuse among transgender and gender nonconforming adults. *LGBT Health*, 3(3), 193–199. 10.1089/lgbt.2015.0111 [PubMed: 27046450]
- Lambert MJ, (2005). Early response to psychotherapy: Further evidence for the importance of common factors rather than “placebo effects.” *Journal of Clinical Psychology*, 61(7), 855–869. 10.1002/jclp.20130 [PubMed: 15827996]
- Laska KM, Gurman AS, & Wampold BE (2014). Expanding the lens of evidence-based practice in psychotherapy: A common factors perspective. *Psychotherapy*, 51(4), 467–481. 10.1037/a0034332 [PubMed: 24377408]
- Lev AI (2009). The ten tasks of the mental health provider: Recommendations for revision of the World Professional Association for Transgender Health's Standards of Care. *International Journal of Transgenderism*, 11(2), 74–99. 10.1080/15532730903008032
- Meyer HM, Mocarski R, Holt NR, Hope DA, King RE, & Woodruff N. (2020). Unmet expectations in health care settings: Experiences of transgender and gender diverse adults in the central great plains. *Qualitative health research*, 30(3), 409–422. 10.1177/1049732319860265 [PubMed: 31328642]
- Mizock L, & Lundquist C. (2016). Missteps in psychotherapy with transgender clients: Promoting gender sensitivity in counseling and psychological practice. *Psychology of Sexual Orientation and Gender Diversity*, 3(2), 148–155. 10.1037/sgd0000177
- Mocarski R, King R, Butler S, Holt NR, Huit TZ, Hope DA, Meyer HM, & Woodruff N. (2019). The rise of transgender and gender diverse representation in the media: Impacts on the population. *Communication, Culture & Critique*, 12(3), 416–433. 10.1093/ccc/tcz031
- Nadal KL, Skolnik A, & Wong Y. 2012). Interpersonal and systemic microaggressions toward transgender people: Implications for counseling. *Journal of LGBT Issues in Counseling*, 6(1), 55–82. 10.1080/15538605.2012.648583
- Newhill CE, Safran JD, & Muran JC (2003). *Negotiating the therapeutic alliance: A relational treatment guide*. Guilford Press.
- Nuttbrock L, Hwahng S, Bockting W, Rosenblum A, Mason M, Macri M, & Becker J. (2010). Psychiatric impact of gender-related abuse across the life course of male-to-female transgender persons. *Journal of Sex Research*, 47(1), 12–23. 10.1080/00224490903062258 [PubMed: 19568976]
- Puckett JA, Aboussouan AB, Ralston AL, Mustanski B, & Newcomb ME (2021). Systems of cissexism and the daily production of stress for transgender and gender diverse people. *International Journal of Transgender Health*. Advance online publication. 10.1080/26895269.2021.1937437
- Puckett JA, Barr SM, Wadsworth LP, Thai J. (2018). Considerations for clinical work and research with transgender and gender diverse individuals. *the Behavior Therapist*, 41, 253–262.

- Puckett JA, Brown N, Dunn T, Mustanski B, & Newcomb ME (2020). Perspectives from transgender and gender diverse people on how to ask about gender. *LGBT Health*, 7(6), 305–311. 10.1089/lgbt.2019.0295 [PubMed: 32598211]
- Richmond K, Burnes TR, Singh AA, & Ferrara M. (2017). Assessment and treatment of trauma with TGNC clients: A feminist approach. (p. 191–212). In Singh AA, & dickey l. m. (Eds.). *Affirmative counseling and psychological practice with transgender and gender nonconforming clients*. American Psychological Association.
- Riggs DW, Pearce R, Pfeffer CA, Hines S, White F, & Ruspini E. (2019). Transnormativity in the psy disciplines: Constructing pathology in the Diagnostic and Statistical Manual of Mental Disorders and Standards of Care. *American Psychologist*, 74(8), 912–924. 10.1037/amp0000545 [PubMed: 31697127]
- Rood BA, Reisner SL, Puckett JA, Surace FI, Berman AK, & Pantalone DW (2017). Internalized transphobia: Exploring perceptions of social messages in transgender and gender-nonconforming adults. *International Journal of Transgenderism*, 18(4), 411–426. 10.1080/15532739.2017.1329048
- Safran JD, & Muran JC (2000). *Negotiating the therapeutic alliance: A relational treatment guide*. Guilford Press.
- shuster s. m. (2016). Uncertain expertise and the limitations of clinical guidelines in transgender healthcare. *Journal of Health and Social Behavior*, 57(3), 319–332. 10.1177/0022146516660343 [PubMed: 27601408]
- Singh AA, & Burnes TR (2010). Shifting the counselor role from gatekeeping to advocacy: Ten strategies for using the competencies for counseling with transgender clients for individual and social change. *Journal of LGBT Issues in Counseling*, 4(3–4), 241–255. 10.1080/15538605.2010.525455
- Singh AA, & dickey l. m. (2017). *Affirmative counseling and psychological practice with transgender and gender nonconforming clients*. American Psychological Association.
- Tebbe EA, & Moradi B. (2016). Suicide risk in trans populations: An application of minority stress theory. *Journal of Counseling Psychology*, 63(5), 520–533. 10.1037/cou0000152 [PubMed: 27089059]
- Testa RJ, Sciacca LM, Wang F, Hendricks ML, Goldblum P, Bradford J, & Bongar B. (2012). Effects of violence on transgender people. *Professional Psychology: Research and Practice*, 43(5), 452–459. 10.1037/a0029604
- Wampold BE (2015). How important are the common factors in psychotherapy? An update. *World Psychiatry*, 14(3), 270–277. 10.1002/wps.20238 [PubMed: 26407772]
- Whitman CN, & Han H. (2017). Clinician competencies: Strengths and limitations for work with transgender and gender non-conforming (TGNC) clients. *International Journal of Transgenderism*, 18(2), 154–171. 10.1080/15532739.2016.1249818
- Zur O. (Ed.). (2016). *Multiple relationships in psychotherapy and counseling: Unavoidable, common, and mandatory dual relations in therapy*. Taylor & Francis.

### Public Significance Statement

This paper provides several recommendations for mental health clinicians to adapt their therapeutic approach for affirming services with transgender and gender diverse (TGD) adult clients. The adaptations are accompanied by practical guidance for clinicians about how to work with TGD clients in the therapy room. The recommendations are particularly applicable for clinicians working in geographic locations with limited resources for TGD communities.

**Table 1.**

**Practice Adaptations**

<b>Trans Collaborations Practice Adaptations for Psychological Services for TGD Adults</b>
Practice Adaptation 1: Setting. <i>Provide a TGD affirmative public face to services, including the website, office features, paperwork, referral process, and office staff.</i>
Practice Adaptation 2: Perceptions of Cultural Messages. <i>Explore the meaning of cultural events and messages individually with each client because cisgender people and TGD people may interpret such messages differently.</i>
Practice Adaptation 3: Clinician’s Experience of Gender. <i>Be aware of the experience of your own gender and how you may automatically enact gender roles that may convey either a marginalizing or an affirmative message to TGD clients.</i>
Practice Adaptation 4: Comfort with Various Gender Identities/Expressions. <i>Recognize and be comfortable with variability and fluidity of gender identity and expression that may vary in unpredictable ways over time for a given client. Your role is to support and affirm each individual gender journey, not diagnose their gender identity or impose a particular journey on the client.</i>
Practice Adaptation 5: Managing Marginalization. <i>Recognize the pervasive stigma and marginalization that many TGD people face in large and small ways in their daily lives. Trust the client’s perceptions of their experiences as you help the client find appropriate coping responses, drawing on the client’s strengths, resilience, and support system.</i>
Practice Adaptation 6: Non-Defensive Stance. <i>Take responsibility for minimizing marginalization experiences within the therapy room and be open and non-defensive when you make errors or misunderstand a client.</i>
Practice Adaptation 7: Clinicians Educate Themselves. <i>Take responsibility for educating yourself about the local cultural context, social and legal climate, and local available services for TGD communities. Be knowledgeable about clinical guidelines, mental health literature, and needs of TGD clients. Clients are not responsible for educating clinicians and deserve the highest quality, culturally responsive services.</i>
Practice Adaptation 8: Share Power. <i>Take responsibility for sharing power with TGD clients as much as possible, especially given the history of marginalization in the healthcare system. Recognize the impact of the gatekeeper role (clinicians providing letters for medical gender affirmation) on exacerbating the power differences that are always part of therapy.</i>
Practice Adaptation 9: Incorporate Gender into Case Conceptualization. <i>Include the entire identity and experience of TGD clients in case conceptualization including past, current, and future gender identities and expression, experiences of marginalization, the affirmative or stigmatizing context of their lives (family, work, legal system), gender roles, sexual orientation, gain/loss of gender-related privilege, and intersectionality with other identities (e.g., race/ethnicity, religion, socioeconomic status, age, ability).</i>
Practice Adaptation 10: Identify Focus of Services. <i>Recognize that a TGD identity may not be the primary organizing identity for an individual and TGD-related concerns may not be the primary focus of therapy.</i>
Practice Adaptation 11: TGD-Affirmative Referrals. <i>Consider the implications of referrals to needed services that may not be TGD-affirming and may exacerbate the problem such as a referral for hospitalization in a non-affirming facility. Problem-solve to get the best possible option, even if it is not a traditional solution.</i>
Practice Adaptation 12: Advocacy. <i>Be prepared to advocate for TGD clients and plan ahead about how such advocacy fits within the overall approach and case conceptualization for a given client. A clinician and client may decide together that the clinician’s privilege that comes from being well educated and/or cisgender can be used to a client’s benefit.</i>

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript