

UC Irvine

UC Irvine Previously Published Works

Title

Postoperative Pain Management in Children of Hispanic Origin: A Descriptive Cohort Study.

Permalink

<https://escholarship.org/uc/item/58r8k95h>

Journal

Anesthesia and Analgesia, 122(2)

Authors

Brown, Rebecca
Zolghadr, Sheeva
Gulur, Padma
et al.

Publication Date

2016-02-01

DOI

10.1213/ANE.0000000000001042

Peer reviewed



HHS Public Access

Author manuscript

Anesth Analg. Author manuscript; available in PMC 2017 February 01.

Published in final edited form as:

Anesth Analg. 2016 February ; 122(2): 497–502. doi:10.1213/ANE.0000000000001042.

Postoperative Pain Management in Children of Hispanic Origin: A Descriptive Cohort Study

Rebecca Brown,

Department of Anesthesiology and Perioperative Care, University of California Irvine, California

Michelle A. Fortier, PhD,

Department of Anesthesiology and Perioperative Care, University of California Irvine, California;

Department of Pediatric Psychology, CHOC Children's Hospital, Orange, California

Sheeva Zolghadr,

Department of Anesthesiology and Perioperative Care, University of California Irvine, California

Padma Gulur, MD,

Department of Anesthesiology and Perioperative Care, University of California Irvine, California

Brooke N. Jenkins, and

Department of Anesthesiology and Perioperative Care, University of California Irvine, California

Zeev N. Kain, MD, MBA

Corresponding Author: Dr. Michelle A. Fortier, UCI Center on Stress and Health, 505 S. Main St., Suite 940, Orange, CA 92868, Tel: 714-456-2833, mfortier@uci.edu.

The authors declare no conflict of interests.

Reprints will not be available from the authors.

Disclosures

Name: Rebecca Brown

Contribution: Ms. Brown carried out the initial analyses, drafted the initial manuscript, and revised the manuscript.

Attestation: Ms. Brown approved the final manuscript as submitted. Ms. Brown attests to the integrity of the original data and the analysis reported in this manuscript.

Name: Michelle A. Fortier, PhD

Contribution: Dr. Fortier conceptualized and designed the study, reviewed and revised the manuscript.

Attestation: Dr. Fortier approved the final manuscript as submitted. Drs. Fortier and Kain attest to the integrity of the original data and the analysis reported in this manuscript. Dr. Fortier is the archival author.

Name: Sheeva Zolghadr

Contribution: Ms. Zolghadr carried out the initial analyses.

Attestation: Ms. Zolghadr approved the final manuscript as submitted.

Name: Padma Gulur, MD

Contribution: Dr. Gulur contributed to revisions of the manuscript.

Attestation: Dr. Gulur approved the final manuscript as submitted.

Name: Brooke N. Jenkins

Contribution: Mrs. Jenkins contributed to revisions of the manuscript.

Attestation: Mrs. Jenkins approved the final manuscript as submitted.

Name: Zeev N. Kain, MD, MBA

Contribution: Dr. Kain conceptualized and designed the study, reviewed and revised the manuscript, and approved the final manuscript as submitted.

Attestation: Dr. Kain approved the final manuscript as submitted. Dr. Kain attests to the integrity of the original data and the analysis reported in this manuscript.

This manuscript was handled by: James A DiNardo, MD

Department of Anesthesiology and Perioperative Care, University of California Irvine, California; Department of Pediatrics, CHOC Children's Hospital Orange and University of California Irvine, California; Child Study Center, Yale University School of Medicine, New Haven, Connecticut

Abstract

Background—It has been established that pain is frequently undertreated in children following outpatient surgery. Very few studies, however, have investigated this phenomenon in ethnically diverse populations.

Methods—This study included 105 families of children age 2–15 years of Hispanic origin and low income undergoing outpatient tonsillectomy and adenoidectomy surgery. Participating parents completed baseline and demographic packets. Recorded postoperative pain ratings and administration of analgesics at home for one week were collected during home visits.

Results—Despite the high (70%, 99% CI[57%, 82%]) incidence of significant pain in the first 24 hours home, 32% (95% CI[20%, 45%]) of the children received 0–1 dose of analgesia. Overall, 21% children (99% CI[11%, 35%]) received 4 or less total doses of pain medication over the entire week following surgery. Of the total analgesic doses administered to children in the week following surgery, only 44% (99% CI[40%, 47%]) were in accepted ranges.

Conclusions—Despite experiencing significant postoperative pain, Hispanic children assessed in this study received suboptimal analgesic therapy at home.

Introduction

Medical advances in recent years have allowed an increasing proportion of pediatric surgery to be performed in the outpatient setting, transitioning responsibility of postoperative pain management to parents in the home setting.¹ To adequately treat pain associated with these procedures, the American Society of Anesthesiologists (ASA) recommends that caregivers employ proactive, around-the-clock treatment regimens, inclusive of developmentally appropriate analgesic therapy.² Unfortunately, there is a significant gap in the translation of these practice guidelines to pain management in the home setting. Specifically, in a previous study published in *Pediatrics* in 2009, we examined children's pain levels and parental pain management practices at home following surgery. In this sample, we identified a significant discrepancy between parent perception of high postoperative pain and low dosing of analgesics.³ As is the case with other pediatric pain studies, however, this previous study included a minimally diverse population (77% non-Hispanic white) and thus has restricted external validity of the study to populations of diverse ethnicities and races.

Although a substantial body of literature has recognized ethnic and racial disparities in pain in adults, few studies have assessed the impact of cultural factors on the pain experience of children.^{4–8} This gap in current literature necessitates further investigation, especially considering the growing Hispanic population in the United States.⁹ It has been suggested that Hispanic cultural values, including a tendency toward stoicism and subsequent inhibition of behaviors that may burden others may put Hispanic patients at risk for under treatment of pain.¹⁰ In addition, Hispanic families may be more likely to experience systems-based barriers to optimal pain management such as lower socioeconomic status

(SES) or lower health literacy relative to their non-Hispanic White counterparts.^{7,11} Indeed, a previous study by our research program found that Hispanic parents were significantly more likely to demonstrate misconceptions regarding analgesics (e.g., issues of addiction, fear of side-effects) and pain expression in children as compared to non-Hispanic White parents.^{12,13} Given our findings that these parental misconceptions are related to under treatment of children's postoperative pain,¹⁴ we suggest that Hispanic children of lower SES may be highly vulnerable to inadequate pain management at home. Therefore, the purpose of this present investigation was to systematically examine postoperative pain management patterns within a large population of Hispanic children of low SES.

Methods

IRB/Consent

The CHOC Children's institutional review board approved the study. Written informed consent was obtained from all participants or the parents or legal guardians for minor subjects in their primary language.

Clinical Trial Registration

This study did not involve assigning patients to treatment groups.

Participants—This National Institute of Health supported study was conducted Children's Hospital of Orange County (CHOC Children's) during the years 2011–2013. The study included a cohort of 197 children ages 2–15 years who were scheduled to undergo outpatient ENT, general, orthopedic, and urologic surgery. Only Hispanic children whose parent(s) were English- or Spanish-speaking and of lower SES as defined by federal poverty guidelines were eligible for participation in the study. Exclusion criteria included American Society of Anesthesiology (ASA) classification III or IV and history of developmental delay, chronic illness, and premature birth. The exclusion criteria were employed in order to standardize the sample as physical and emotional impairments can contribute to confounding variations in pain sensitivity and emotional response to surgery, and to challenges in pain evaluation.

In order to control for variations in surgical severity and postoperative pain, the present report only includes data of children from the larger cohort undergoing outpatient T&A. Of the original sample of 197 children, 70 had a surgical procedure that was not a tonsillectomy or adenoidectomy (T&A) and were therefore not included in the analyses, resulting in a sample size of 126. Of the 126 children undergoing T&A and their parents who agreed to participate in the study, 22 dropped out (were lost to follow-up or did not return an adequate number of home surveys), yielding 105 who completed postoperative pain and analgesia follow-up monitoring.

Measures

Patient Characteristics: Acculturation Rating Scale for Mexican Americans-II: The Acculturation Rating Scale for Mexican Americans-II (ARSMA-II) is used to assess multidimensional orientation toward Mexican and Anglican culture.¹⁵ Through a self-report

questionnaire on aspects of daily living (e.g. "I think in Spanish/English," "I enjoy English/Spanish language TV," etc.), ARSMA-II generates scores along two linear subscales: an Anglo-Orientation Subscale and Mexican-Orientation Subscale. Respondents are then categorically scaled into 5 acculturation groups: very Mexican oriented, Mexican oriented to bicultural, Anglo oriented to bicultural, strongly Anglo oriented, and very assimilated. These groups note degrees of assimilation, or loss of Mexican cultural identity as Anglican culture is acquired, and development of bicultural orientation. Generational status is also noted in this survey, with "first generation" representing individuals born in Mexico or another country; "second generation" representing individuals born in the United States with at least one parent born in another country; and so on. Although the questionnaire is specific to Mexican-American culture, it has demonstrated functionality as a measure of acculturation in other Hispanic cultural groups and is frequently used as such.^{16,17} In addition, the majority of the cohort in the present study was of Mexican background.

Child Pain Report: Faces Pain Scale-Revised: The Faces Pain Scale-Revised (FPS-R) is a self-report measure used to assess pain in children.¹⁸ Children are provided a schematic with five facial expressions ranging from neutral ("no pain") to one representative of the "worst pain ever," and prompted to indicate that which best represents how they feel. A cutoff score of 3 (out of 10) is considered representative of clinically significant pain. This measure has demonstrated excellent convergent validity with other pain scales.^{19,20} FPS-R is widely validated for children aged 4 and up.²⁰

Analgesic Administration: Parents recorded the type (i.e., acetaminophen, ibuprofen, hydrocodone, codeine), strength, and quantity of pain medication given to their children on postoperative days 1 through 7. In order to provide real-time assessment of pain management, parents also recorded the time each dose was administered.

Procedure—Potential participants were identified through surgery schedules, and eligibility was assessed by review of electronic medical records for exclusionary criteria. Families were approached in the preoperative holding area by a bilingual research associate and were provided an information sheet and verbal description of the study. After parents provided informed consent and children gave appropriate assent, participants indicated their preferred language for written information and were provided demographic and baseline surveys.

All T&A procedures were performed by a single surgeon. Following surgery, children were taken to the Post Anesthesia Care Unit (PACU) where pain assessment and analgesia administration was completed by nurses consistent with hospital standard of care. Surgical technique and pain management was not standardized in order to maintain external validity. Prior to discharge, research associates provided parents with a home packet containing record forms for assessing children's pain severity and analgesic administration on each of the postoperative days. Research associates reviewed record forms with parents and verbalized how to administer FPS-R and demonstrated with the actual measure. All patients were discharged home directly from the PACU within 3 hours of surgery. Upon discharge, caretakers received verbal and standard written analgesic instructions for management of postoperative pain at home by the PACU nurse. The instructions advised around the clock

administration of acetaminophen by mouth or rectum as directed on medication packaging. Additionally, caregivers were instructed to give codeine or hydrocodone every 3–4 hours as needed if patients actively complained of pain.

At home, parents completed daily pain medication record forms for one week following surgery describing all analgesics given within each 24 hour period. Parents were also asked to obtain their child's self-report of average pain intensity over the course of each day through administration of FPS-R. Parents recorded this single daily FPS-R pain score in the provided pain record packets. If no pain medication was given on any of the postoperative days, parents were asked to indicate the reason (e.g., child not in pain, child refused medication, no medication prescribed, etc.).

Data Analysis Strategy—Descriptive statistics were used to analyze demographics, baseline characteristics, and the two primary outcomes of this study: postoperative pain intensity and analgesic administration at home. Normally distributed statistics are reported as means and standard deviations (SD), and skewed data are reported as medians and interquartile ranges (IQR). Pain intensity was analyzed such that FPS-R scores ≥ 3 defined clinically significant pain. One sample t-tests were used to compare FPS-R scores to this cut off value. This allowed us to determine whether the sample on average experienced clinically significant pain. In order to determine whether children received accepted doses of analgesics, all individual medication doses reported by parents were converted to milligrams, and divided by the child's weight obtained from hospital records. The normalized doses were then categorized as accepted or sub-accepted using the following cutoff values: acetaminophen at 10 mg/kg/dose, ibuprofen at 5 mg/kg/dose, codeine at 0.5 mg/kg/dose, and hydrocodone at 0.1 mg/kg/dose. The doses were then further subgrouped as markedly sub-accepted and moderately sub-accepted. Given the patterns of pain recovery found in previous literature,^{3,21} analysis of pain assessment and frequency of analgesic administration was focused on postoperative days 1 – 3 and week 1. Using the Clopper-Pearson method, 99% confidence intervals are provided on all point estimates.^{22,23} All study analysis was performed using SPSS 20.0 (IBM, Armonk, NY).

Results

Descriptive data

Table 1 presents the demographic and baseline characteristics of 105 participating patients and parents who completed the survey packets and pain and analgesic records. Strong Hispanic cultural identity was prevalent in this sample. Most parents were born in Mexico and spoke primarily Spanish (74%). Just 3 parents who completed the survey and pain and analgesic records were not of Hispanic background (all children included in the study were Hispanic). Of the Spanish speakers, 64% indicated English as a second language. ARSMA-II responses defined 48% of parents as “very Mexican oriented” and 33% of parents as “Mexican oriented to bicultural,” and 19% as “slightly Anglo-oriented.” One-way analyses of variance were performed to examine acculturation group differences in pain-report and analgesic dosing. No significant patterns emerged amongst these outcome variables within this cohort.

Pain Assessment at Home

Figure 1 presents the percentage of children experiencing clinically significant pain on postoperative days 1 - 3 and week 1, as assessed by child self-report (FPS-R). FPS-R scores classified the peak percentage (70%, 99% CI[57%, 82%]) of children in significant pain in the first 24 hours following surgery (IQR: (2,6)). Parents recorded children as reporting clinically significant pain on the first 2 post-operative days (day 1: 70% experienced clinically significant pain: 99% CI[57%, 82%], $t(90) = 6.05$, $p < .001$, 95% CI[1.20, 2.38]; day 2: 64% experienced clinically significant pain, 99% CI[53%, 80%]; $t(85) = 5.393$, $p < .001$, 95% CI[1.03, 2.23], with incremental improvements in pain though the week. By one week after surgery, 28% (99% CI[16%, 42%]) of children remained in clinically significant pain (IQR: (0,4)).

Frequency of Analgesic Administration by Parents

Figure 2 displays the frequency of analgesic doses administered to children on postoperative days 1 - 3 and week 1. Overall, 21% of children (99% CI[11%, 35%]) received 4 or less total doses of pain medication over the entire week following surgery. The median total analgesic dose frequency delivered to children over the first 3 days was 7 (3, 9). 32% of children (99% CI[20%, 45%]) received 0 or just 1 dose of analgesia on the first postoperative day at home. On day 3 following surgery, a large portion of children (39%, 99% CI[26%, 52%]) received 0 or 1 dose of analgesia despite 56% of children (99% CI[41%, 70%]) self-reporting significant pain to parents through FPS-R. On days 1, 2, 3 and week 1 following surgery, 18, 20, 16, and 12 children received at least one dose of opioid pain medication, respectively.

Accepted Dosing

Table 2 presents the median doses of analgesia given by parents in aggregate over postoperative week 1, as well as the distribution of accepted, moderately sub-accepted, and severely sub-accepted doses administered over that time. Overall, the majority of normalized (mg/kg) analgesic doses were sub-accepted: only 44% (99% CI[40%, 47%]) were in accepted ranges. Acetaminophen was the most common drug administered by parents, comprising 78.4% (99% CI[75%, 81%]) of the total doses given. However, only 40.7% (99% CI[36.7%, 44.8%]) of acetaminophen doses were in the accepted range, and 9.6% (99% CI[7.3%, 12.2%]) of doses were markedly low (less than 5.0 mg/kg). Ibuprofen was the only drug for which the slight majority of doses administered (52.9%, 99% CI[21.9%, 82.4%]) was in the accepted range, and for which the median dose given was of acceptable (7.6 mg/kg). Sub-accepted dosing was particularly prevalent with codeine, as 76.7% (99% CI[67.6%, 84.4%]) were moderately sub-accepted and only only 17.0% (99% CI[10.4%, 25.5%]) of doses in the accepted range.

Reasons for Not Administering Analgesia

Of the caregivers who provided reasons for not administering analgesia over the first 3 postoperative days, 63% (99% CI[38.4%, 82.8%]) indicated that their child was not in pain, 19% (99% CI[5.1%, 41.9%]) indicated that their child refused to take medication, and the remainder (19%; 99% CI[5.1%, 41.9%]) selected "other" reasons or declined to answer.

Discussion

Under the conditions of this study, we found that many of the participating low income children of Hispanic origin experienced significant pain in the week following outpatient T&A surgery and received suboptimal pain management in the home setting. These findings are consistent with a previous study examining postoperative pain management in a non-Hispanic White population of children undergoing T&A.³

Around-the-clock analgesic therapy is the current ASA recommended practice for postoperative pain management and prevention.² In the home setting, accepted treatment can be achieved through maintenance of a regular analgesia dosing schedule, typically one dose every 4–6 hours.^{2,24} Accordingly, dosing every 4–6 hours should result in 12 - 18 total doses of analgesia in the first 3 postoperative days alone. Despite parents being told to follow package instructions of administering medication every 4–6 hours, we found that parents in this sample administered only 7 (3, 9) total doses over the entire first 3 postoperative days. Parents often believe that it is best to give as little medication as possible in order to avoid risks and side effects.¹⁴ As such, they may choose to wait until their child is experiencing significant pain before administering analgesia. Unfortunately, this practice subjects the child to unnecessary discomfort in the time leading to the parent perceiving the pain and administering treatment, and in the time it takes for the drug to become efficacious. Less than half of the doses administered by parents in this study were of accepted value. Given the lack of evidence supporting the efficacy of sub-accepted dosing,²⁴ it is unclear whether or not the few doses provided by parents actually provided symptom relief.

Cultural barriers may contribute to the suboptimal pain management practices seen in this study. The majority of the participants in the present study were strongly oriented toward traditional Hispanic culture, were first generation Americans, and selected Spanish as their preferred language. Hispanic culture may be associated with increased stoicism, and subsequent inhibition of behaviors that may burden others.¹⁰ The cultural emphasis on stoicism may result contribute to decreased expression of pain by children and subsequent lower pain scores and medication dosing relative to their non-Hispanic White counterparts. This is of particular relevance in this study, given that pain intensity was assessed by child self-report. A prior study by our group found that child temperament and emotionality played a role in how children expressed pain following surgery, and was subsequently predictive of parental analgesic administration.²⁵ Though not directly assessed in this study, it is likely that such cultural factors as stoicism play a role in development of child temperament and emotionality and in the manner in which they report pain. A greater understanding of the relationship between cultural norms and pain assessment would aid providers in delivering culturally sensitive care for Hispanic children and improving the translation of care guidelines to the home setting.

Additionally, factors related to health systems may pose further barriers to adequate pain management following surgery. For example, inadequate health literacy and numeracy is common amongst Hispanic families of lower acculturation in the United States,²⁶ and profoundly impacts the ability to understand discharge instructions, medication administration, and symptomatology.^{27,28} A study by Ciampa et al found that low

acculturation amongst Hispanic caretakers is independently correlated with a decreased ability to perform child health care tasks, such as dosing liquid medication.²⁷ Therefore, both acculturation and low literacy can impact the translation of pain management instructions to the home setting. Other factors that can impact pain management include such systemic issues as access to care and lack of insurance. In the context of post-operative pain management, these factors can limit parents' ability to pay for and obtain medication. Hispanic families of lower SES have been shown to experience more such health care systems barriers than their non-Hispanic White counterparts.⁴ Both providers and researchers would benefit from a greater understanding of the impact of such health systems barriers on care for this vulnerable population.

This study has several methodological limitations that should be noted. First, because parents received pain assessment and analgesia surveys at discharge, they may have been increasingly alerted to behaviors associated with pain, and been more conscious of treating pain. Second, the data used to assess analgesic dosing and pain scores was collected by parent report, and thus subject to reporting and collecting (i.e.: coaching) bias. However, because of evidence suggesting that Hispanic patients may underreport pain, the occurrence of suboptimal pain management may be more profound than what was seen in this investigation.¹⁰ It is important to note that the purpose of this study was solely to focus on the management of postoperative pain in children of Hispanic origin and as such the study did not enroll a control group of non-Hispanic children. Such a study could be carried out in the future. Lastly, our study focused exclusively on pain management in children undergoing T&A. Because this procedure can be associated with postoperative dysphagia, some children may be unwilling to consume oral analgesics. Hence the analgesic patterns identified in this study may differ from those of other procedures.

We conclude that Hispanic children of lower SES experienced significant pain following outpatient surgery and were inadequately treated with analgesic therapy in the home setting. These findings lend evidentiary support to the need for further investigation into attitudinal- and systems-based barriers to postoperative pain management in this population. We anticipate that such findings will aid in the development of culturally- and demographically-sensitive interventions that will aid in the translation of clinical practice guidelines to the home setting. Such efforts are vital for achieving critical reductions in pediatric pain disparities.

Acknowledgments

Funding: Supported by Grant Number R21HD071798 from the Eunice Kennedy Shriver National Institute of Child Health & Human Development. The content is solely the responsibility of the authors and does not necessarily represent the official views of the Eunice Kennedy Shriver National Institute Of Child Health & Human Development or the National Institutes of Health.

References

1. Cullen KA, Hall MJ, Golosinskiy A. Ambulatory surgery in the United States, 2006. Natl Health Stat Report. 2009;1–25. [PubMed: 19294964]

2. American Society of Anesthesiologists Task Force on Acute Pain Management. Practice guidelines for acute pain management in the perioperative setting: an updated report. *Anesthesiology*. 2004; 100:1573–81. [PubMed: 15166580]
3. Fortier MA, MacLaren JE, Martin SR, Perret D, Kain ZN. Pediatric pain after ambulatory surgery: Where's the medication? *Pediatrics*. 2009; 124:588–95.
4. Anderson KO, Green CR, Payne R. Racial and ethnic disparities in pain: causes and consequences of unequal care. *J Pain*. 2009; 10:1187–204. [PubMed: 19944378]
5. Campbell CM, Edwards RR, Fillingim RB. Ethnic differences in responses to multiple experimental pain stimuli. *Pain*. 2005; 113:20–6. [PubMed: 15621360]
6. Edwards CL, Fillingim RB, Keefe F. Race, ethnicity and pain. *Pain*. 2001; 94:133–7. [PubMed: 11690726]
7. Green CR, Anderson KO, Baker TA, et al. The Unequal Burden of Pain: Confronting Racial and Ethnic Disparities in Pain. *Pain Medicine*. 2003; 4:277–94. [PubMed: 12974827]
8. Portenoy RK, Ugarte C, Fuller I, Haas G. Population-based survey of pain in the United States: Differences among White, African American, and Hispanic subjects. *The Journal of Pain*. 2004; 5:317–28. [PubMed: 15336636]
9. U.S. Census Bureau. Projections Show a Slower Growing, Older, More Diverse Nation a Half Century from Now. 2012. at <http://www.census.gov/newsroom/releases/archives/population/cb12-243.html>
10. Duggleby W. Helping Hispanic/Latino home health patients manage their pain. *Home Healthcare Nurse*. 2003; 21:174–9. [PubMed: 12637823]
11. Wallace SP, Gutierrez VF, Castaneda X. Access to preventive services for adults of Mexican origin. *J Immigr Minor Health*. 2008; 10:363–71. [PubMed: 17939052]
12. Fortier, MA.; Tan, ET.; Maurer, EL.; Gomez, SH.; Kain, DI. Parental Attitudes Regarding Analgesic Use for Children: Differences in Ethnicity and Language. American Society of Anesthesiologists; Chicago, IL: 2011.
13. Batista ML, Fortier MA, Maurer EL, Tan E, Huszti HC, Kain ZN. Exploring the impact of cultural background on parental perceptions of children's pain. *Children's Health Care*. 2012; 41:97–110.
14. Zisk Rony RY, Fortier MA, Chorney JM, Perret D, Kain ZN. Parental postoperative pain management: attitudes, assessment, and management. *Pediatrics*. 2010; 125:e1372–8. [PubMed: 20498177]
15. Cuéllar I, Arnold B, Maldonado R. Acculturation Rating Scale for Mexican Americans-II: A revision of the original ARSMA scale. *Hispanic Journal of Behavioral Sciences*. 1995; 17:275–304.
16. Johnson KL, Carroll JF, Fulda KG, Cardarelli K, Cardarelli R. Acculturation and self-reported health among Hispanics using a socio-behavioral model: the North Texas Healthy Heart Study. *BMC public health*. 2010; 10:53. [PubMed: 20122263]
17. Wiley JF, Cloutier MM, Wakefield DB, et al. Acculturation determines BMI percentile and noncore food intake in Hispanic children. *J Nutr*. 2014; 144:305–10. [PubMed: 24453127]
18. Hicks CL, von Baeyer CL, Spafford PA, van Korlaar I, Goodenough B. The Faces Pain Scale-Revised: toward a common metric in pediatric pain measurement. *Pain*. 2001; 93:173–83. [PubMed: 11427329]
19. Perrott DA, Goodenough B, Champion GD. Children's ratings of the intensity and unpleasantness of post-operative pain using facial expression scales. *European Journal of Pain*. 2004; 8:119–27. [PubMed: 14987621]
20. Stinson JN, Kavanagh T, Yamada J, Gill N, Stevens B. Systematic review of the psychometric properties, interpretability and feasibility of self-report pain intensity measures for use in clinical trials in children and adolescents. *Pain*. 2006; 125:143–57. [PubMed: 16777328]
21. Huth MM, Broome ME. A snapshot of children's postoperative tonsillectomy outcomes at home. *Journal for Specialists in Pediatric Nursing: JSPN*. 2007; 12:186–95. [PubMed: 17594298]
22. Newcombe, Robert G. Two-Sided Confidence Intervals for the Single Proportion: Comparison of Seven Methods. *Statistics in Medicine*. 1998; 17:857–72. [PubMed: 9595616]
23. Wilson EB. Probable Inference, the Law of Succession, and Statistical Inference. *Journal of the American Statistical Association*. 1927; 22:209–12.

24. Finley GA, McGrath PJ, Forward SP, McNeill G, Fitzgerald P. Parents' management of children's pain following 'minor' surgery. *Pain*. 1996; 64:83–7. [PubMed: 8867249]
25. Zolghadr S, Fortier MA, Winckler B, Kain ZN. An examination of child temperament as a predictor of parental post-operative analgesic administration. *J Ped Surg*. 2014; 49:1315–8.
26. Kutner MGE, Jin Y, Paulson C. *The Health Literacy of America's Adults: Results from the 2003 National Assessment of Adult Literacy*. 2006
27. Ciampa PJ, White RO, Perrin EM, et al. The association of acculturation and health literacy, numeracy and health-related skills in Spanish-speaking caregivers of young children. *J Immigr Minor Health*. 2013; 15:492–8. [PubMed: 22481307]
28. Yin HS, Dreyer BP, Foltin G, van Schaick L, Mendelsohn AL. Association of low caregiver health literacy with reported use of nonstandardized dosing instruments and lack of knowledge of weight-based dosing. *Ambulatory pediatrics : the official journal of the Ambulatory Pediatric Association*. 2007; 7:292–8. [PubMed: 17660100]




Figure 1. Percentage of children in significant pain as reported by child FPS-R rating.




Figure 2. Number of analgesic doses administered on postoperative days 1 (A), 2 (B), 3 (C) and week 1 (D)

Table 1

Demographic and Baseline Characteristics of the Study Sample

Parameter	Sample (N=105)
Children	
Age in yrs, mean (SD)	6.5 (3.0)
Gender, % Male (n)	58 (61)
Ethnicity, % Hispanic (n)	100 (105)
Parent who completed packets	
Age in yrs, median (IQR)	33 (29 – 39)
Years of education, mean (SD)	10.5 (3.5)
Annual household income, median (IQR)	\$18,660 (\$14,400 – 24,970)
Primary Language, % (n)	
English	26 (21)
Spanish	74 (81)
Generational status, % (n)	
1 st generation American	77 (76)
2 nd generation or greater	23 (23)
Acculturation (ARSMA-II) ^a , % (n)	
Very Mexican oriented	48 (47)
Mexican oriented to bicultural	33 (33)
Anglo oriented to bicultural	13 (13)
Strongly Anglo oriented	5 (5)
Very assimilated	1 (1)
Ethnicity	
Mexican	55 (54)
Other Latino/Hispanic/Spanish	42 (41)
Non-Hispanic	3 (3)

^a Acculturation Rating System for Mexican Americans-II

Table 2

Median dose of analgesia administered (mg/kg/dose) over post-operative days 1–7, and distribution of accepted and sub-accepted dosing

	Acetaminophen n = 1005	Ibuprofen n = 17	Codeine n = 176	Hydrocodone n = 74
Median (IQR)	9.318* (7.548 – 12.28)	7.614 3.810 – 36.06)	0.423* (0.390 – 0.497)	0.0883* (0.0759 – 0.135)
% of doses falling within mg/kg/dose range	< 5: 9.6	< 2.5: 11.8	< 0.25: 6.3	< 0.05: 5.4
	5–10: 49.8	2.5–5: 35.3	0.25–0.5: 76.7	0.05–0.1: 45.9
	10–15: 40.7	5–7.5: 52.9	0.5–0.75: 17.0	0.1–0.15: 48.6

* mg/kg/dose falls below standard accepted dose: Acetaminophen: 10 mg/kg/dose; Codeine: 0.5 mg/kg/dose; Ibuprofen: 5 mg/kg/dose; Hydrocodone: 0.1 mg/kg/dose

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript