

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

BUBBLE TRACKS IN A HYDROGEN-FILLED GLASER CHAMBER

### Permalink

<https://escholarship.org/uc/item/58w277z7>

### Author

Wood, John G.

### Publication Date

1953-02-15

UCRL 2494

UNIVERSITY OF CALIFORNIA

UNIVERSITY OF  
CALIFORNIA

*Radiation  
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UCRL-2494  
Unclassified -  
No Standard Distribution

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

BUBBLE TRACKS IN A HYDROGEN-FILLED  
GLASER CHAMBER

John G. Wood

February 15, 1954

Berkeley, California

BUBBLE TRACKS IN A HYDROGEN-FILLED  
GLASER CHAMBER

John G. Wood

Radiation Laboratory, Department of Physics  
University of California, Berkeley, California

February 15, 1954

Glaser<sup>1</sup> first showed that the boiling of superheated ethyl ether was sensitive to ionizing radiation, and later photographed the bubble tracks in his apparatus. A liquid hydrogen chamber would have a number of attractive features, and it is therefore not surprising that several groups have been working to produce such a device. At Chicago, Hildebrand and Nagle<sup>2</sup>, working in cooperation with Glaser, have produced superheated liquid hydrogen, and have shown that it boils sooner in the presence of a  $\gamma$ -ray source. We have taken the next step, but with somewhat different technique, and can report the photography of bubble tracks in hydrogen.

The bulk of the Chicago apparatus was immersed in a bath of hydrogen which was boiling at atmospheric pressure, but the bubble chamber was in the vapor space, and superheated by a coil of resistance wire. This arrangement involves problems of heat transfer and temperature measurement, in addition to some serious problems if photography were to be attempted. We have therefore immersed our bubble chamber in a bath of hydrogen which boils at high pressure; this provides a convenient heat reservoir, whose temperature is easily controlled by a pressure regulator. First, hydrogen is condensed into the bubble chamber at a pressure somewhat higher than that over the bath. After temperature equilibrium has been established, the following cycle is initiated: (1) The pressure in the bubble chamber is suddenly reduced to one atmosphere; (2) At a variable time after the pressure release, an electronic circuit triggers

a stroboscopic lamp, which takes a photograph of the chamber, and (3) pressure is reapplied to the chamber to condense any bubbles that may have formed.

We have been unable to duplicate the long times of superheat reported by the Chicago group, but we have been using considerably higher degrees of superheat. We were discouraged by our inability to attain the long times of superheat, until the track photographs showed that it was not important in the successful operation of a large bubble chamber. Tracks have even been observed in cases where the liquid hydrogen was not completely condensed in the chamber prior to expansion.

I wish to thank Drs. Luis W. Alvarez, Frank S. Crawford, Jr., and M. Lynn Stevenson for their advice and help in these experiments. I am indebted to Mr. A. J. Schwemin for help with the electronic circuits, and to Messrs. H. Powell and T. Robbins for their cooperation with the glass work.

---

This work has been performed under the auspices of the Atomic Energy Commission.

<sup>1</sup>D. A. Glaser, Phys. Rev. 87, 665 (1952)

<sup>2</sup>R. H. Hildebrand and D. E. Nagle, Phys. Rev. 92, 517 (1953)




Fig. 1.  
1-1/2 Inch Diameter Chamber  
Irradiated with a Po-Be Source.