

UC Davis

UC Davis Previously Published Works

Title

Factorial combinations of protein interactions generate a multiplicity of florigen activation complexes in wheat and barley

Permalink

<https://escholarship.org/uc/item/5975c7pn>

Journal

The Plant Journal, 84(1)

ISSN

0960-7412

Authors

Li, Chengxia
Lin, Huiqiong
Dubcovsky, Jorge

Publication Date

2015-10-01

DOI

10.1111/tpj.12960

Peer reviewed

Factorial combinations of protein interactions generate a multiplicity of florigen activation complexes in wheat and barley

Chengxia Li^{1,2}, Huiqiong Lin¹ and Jorge Dubcovsky^{1,2,3,*}

¹Department Plant Sciences, University of California, Davis, CA, USA,

²Howard Hughes Medical Institute, Chevy Chase, MD, USA, and

³Gordon and Betty Moore Foundation, Palo Alto, CA, USA

Received 9 December 2014; revised 22 July 2015; accepted 27 July 2015; published online 7 August 2015.

*For correspondence (e-mail jdubcovsky@ucdavis.edu).

SUMMARY

The FLOWERING LOCUS T (FT) protein is a central component of a mobile flowering signal (florigen) that is transported from leaves to the shoot apical meristem (SAM). Two FT monomers and two DNA-binding bZIP transcription factors interact with a dimeric 14-3-3 protein bridge to form a hexameric protein complex. This complex, designated as the 'florigen activation complex' (FAC), plays a critical role in flowering. The wheat homologue of FT, designated FT1 (= VRN3), activates expression of *VRN1* in the leaves and the SAM, promoting flowering under inductive long days. In this study, we show that FT1, other FT-like proteins, and different FD-like proteins, can interact with multiple wheat and barley 14-3-3 proteins. We also identify the critical amino acid residues in FT1 and FD-like proteins required for their interactions, and demonstrate that 14-3-3 proteins are necessary bridges to mediate the FT1–TaFDL2 interaction. Using *in vivo* bimolecular fluorescent complementation (BiFC) assays, we demonstrate that the interaction between FT1 and 14-3-3 occurs in the cytoplasm, and that this complex is then translocated to the nucleus, where it interacts with TaFDL2 to form a FAC. We also demonstrate that a FAC including FT1, TaFDL2 and Ta14-3-3C can bind to the *VRN1* promoter *in vitro*. Finally, we show that relative transcript levels of *FD-like* and *14-3-3* genes vary among tissues and developmental stages. Since FD-like proteins determine the DNA specificity of the FACs, variation in *FD-like* gene expression can result in spatial and temporal modulation of the effects of mobile FT-like signals.

Keywords: *Triticum aestivum*, *Hordeum vulgare*, flowering, florigen activation complex, FLOWERING LOCUS T, FDL, 14-3-3, VRN1.

INTRODUCTION

Florigen is a systemic signal that is produced in leaves in response to the stimulus of inductive day length and is transported to the shoot apex to induce flowering (Zeevaert, 2006). Recent studies in model plant species *Arabidopsis* and rice have revealed that FLOWERING LOCUS T (FT) (= HEADING DATE 3 (HD3A) in rice) is the main component of florigen (Corbesier *et al.*, 2007; Tamaki *et al.*, 2007). In both species, FT transcription is activated in the leaf vascular tissue and the encoded protein is transported to the shoot apical meristem (SAM) through the phloem (Corbesier *et al.*, 2007; Tamaki *et al.*, 2007).

FT encodes a RAF-kinase-inhibitor-like protein with a unique phosphatidylethanolamine-binding protein (PEBP) domain. Crystal structure studies in mammals suggest that

PEBP domain proteins contain an anion-binding pocket composed of highly conserved residues, which can bind to anions, phosphate groups and phospholipids (Banfield *et al.*, 1998; Serre *et al.*, 1998; reviewed in Granovsky and Rosner, 2008). Both FT and HD3A proteins have a conserved anion-binding pocket. Although it is unclear whether the anion-binding pocket binds to anions or phospholipids in plants, a tyrosine (Y85 in FT and Y87 in HD3A) in the pocket region has been shown to be critical for flowering regulation (Hanzawa *et al.*, 2005; Ahn *et al.*, 2006; Taoka *et al.*, 2013).

Wheat FT1 (= VRN3) is the closest homologue to *Arabidopsis* FT (Yan *et al.*, 2006). FT1 is transcribed in the leaves under long days, and its expression is closely

correlated with the transcription of its target gene, the meristem identity gene *VRN1* in the apex, suggesting that *FT1* also encodes a mobile signal in wheat (Yan *et al.*, 2006; Li and Dubcovsky, 2008; Pearce *et al.*, 2013).

In Arabidopsis, FD and FD PARALOG (FDP) are basic leucine zipper (bZIP) transcription factors that interact with FT and activate the expression of the floral meristem identity gene *APETALA1* (*AP1*) (Abe *et al.*, 2005; Wigge *et al.*, 2005). The C-terminus of FD has a short motif that is the potential target of calcium-dependent protein kinases (CDPKs), and a single amino acid substitution (a serine/threonine substituted by an alanine) in this motif disrupts FD protein function (Abe *et al.*, 2005). Other members of the bZIP protein family, including OsFD1 from rice (Taoka *et al.*, 2011), TaFDL2 and TaFDL6 from wheat (Li and Dubcovsky, 2008) and DLF1 from maize (Muszynski *et al.*, 2006) also have been shown to interact with FT orthologues (HD3A in rice, FT1 in wheat and ZCN8 in maize).

Many of the FT-interacting bZIP proteins were identified by yeast two-hybrid assays. However, a recent study in rice suggests that these yeast two-hybrid interactions are likely mediated by endogenous yeast 14-3-3 proteins, since no direct interaction between OsFD1 and HD3A could be detected by NMR titration assays, isothermal titration calorimetry or pull-down assays (Taoka *et al.*, 2011). Indeed, 14-3-3 proteins have been identified as FT-interacting proteins in a previous study in tomato (Pnueli *et al.*, 2001). Taoka *et al.* (2011) demonstrated that HD3A first interacts with 14-3-3 proteins in the cytoplasm, and that the resulting protein complex then translocates to the nucleus where it binds to OsFD1. The resultant protein complex, henceforth the florigen activation complex (FAC), then interacts with the promoter of the meristem identity gene *OsMADS15* and induces its transcription (Taoka *et al.*, 2011).

The crystal structure of the rice FAC is predicted to be a deep W-shaped structure with two 14-3-3 molecules forming a W-shaped dimer via their N-terminal regions. One HD3A monomer binds to the C-terminal region of each 14-3-3 monomer, extending the W-shape structure. Two positively charged pockets formed at the corners at the inner base of the 'W' then interact with the phosphorylated C-terminus of OsFD1 proteins (Taoka *et al.*, 2011). The resulting hexameric complex is essential for HD3A-mediated flowering promotion in rice and provides a mechanistic basis for florigen function in flowering (Taoka *et al.*, 2011).

14-3-3 proteins are present in all eukaryotes and regulate diverse cellular functions through protein–protein interactions. Most plant species have multiple 14-3-3 paralogs that provide sequence and functional diversity (Paul *et al.*, 2012). Plant 14-3-3 proteins can interact with a wide range of proteins including kinases, transcription factors, structural proteins, ion channels and pathogen defense-related proteins. Through these interactions, plant 14-3-3 proteins

can modulate enzyme activity, protein structure, stability, and subcellular localization, affecting multiple physiological and developmental processes (reviewed in Denison *et al.*, 2011).

To understand better how different FT, FD and 14-3-3 proteins regulate flowering in the temperate cereals, we characterized their interactions by both yeast two-hybrid and *in vivo* bimolecular fluorescent complementation (BiFC) assays. We identified the critical amino acids in FT1 and FD-like proteins required for their interactions with 14-3-3 proteins, and demonstrated that 14-3-3 proteins are necessary to mediate FT1–TaFDL2 interaction. BiFC assays showed that FT1–14-3-3 interactions occur in the cytoplasm, and that the tetrameric complex is then translocated to the nucleus where it interacts with TaFDL2 to form the hexameric FAC. Finally, we established that the FAC including FT1, TaFDL2 and Ta14-3-3C can bind to the *VRN1* promoter.

RESULTS

Wheat FT1 protein interacts with multiple 14-3-3 proteins

Sequences of five previously identified barley 14-3-3 paralogs A to E (Schoonheim *et al.*, 2007) were used to search and clone their wheat orthologues, which were assigned the same letters as the corresponding barley genes. In addition, an unclassified 14-3-3 protein (GenBank accession number BAK01024) with high sequence similarity to rice Os14-3-3H (also known as OsGF14h or UniProt Q2R1D5.1) was identified and cloned from barley, and was designated Hv14-3-3H. We did not identify a sequence for this gene in *T. aestivum*, but found a copy in a *T. urartu* transcriptome database (Krasileva *et al.*, 2013). Tu14-3-3H, Hv14-3-3H and Os14-3-3H are placed in a separate cluster from the other five 14-3-3 proteins in a phylogenetic analysis (Figure S1), indicating that they diverged from the other 14-3-3 early in the evolution of this gene family.

Full-length coding regions of these six 14-3-3 proteins were cloned into yeast vector pGADT7 as preys, and FT1 was cloned into pGBKT7 as bait. Yeast two-hybrid assays revealed that FT1 interacts with 14-3-3 proteins A to E, but not with 14-3-3H (Figure 1a), suggesting some functional differentiation between the two 14-3-3 clades (Figure S1).

Identification of the critical amino acid residues in FT1 protein responsible for interactions with 14-3-3 proteins

Two amino acid residues Y85 and R132 of the Arabidopsis FT protein have been previously shown to be critical for flowering (Hanzawa *et al.*, 2005; Ahn *et al.*, 2006) and for interaction with 14-3-3 proteins (Taoka *et al.*, 2011). We created point mutations in wheat FT1 by site-directed mutagenesis at these two positions (Y85 and R130), at four positions predicted as surface-exposed residues (R62, T66, P94 and F101, Taoka *et al.*, 2011), and at two additional




Figure 1. Interactions between FT1 and 14-3-3 proteins.

(a) Yeast two-hybrid assays between FT1 and six 14-3-3 proteins.

(b) Mutations of critical amino acid residues in FT1 protein abolish protein interactions with 14-3-3 proteins in yeast two-hybrid assays. FT1 and mutants were used as baits and 14-3-3 proteins were used as preys. SD medium lacking leucine and tryptophan (-L-W) was used to select for yeast transformants containing both bait and prey vectors. Interaction strength was tested on SD media lacking leucine, tryptophan, histidine and adenine (-L-W-H-A).

sites (P77S and G33E) that have been recently shown to delay flowering when mutated in wheat (Lv *et al.*, 2014). Yeast two-hybrid assays showed that FT1 mutations R62K, T66I, P94L, F101A, and R130K abolished the interactions with 14-3-3 proteins. Y85H abolished interaction with Ta14-3-3E and reduced interactions with Ta14-3-3A-D proteins (Figure 1b). The G33E and P77S mutations did not affect interactions with the 14-3-3 proteins, suggesting that their effect on wheat flowering is not mediated by the disruption of these protein-protein interactions (Figure 1b).

Interactions between FD-like and 14-3-3 proteins in wheat and barley

Wheat *TaFDL2*, *TaFDL6*, *TaFDL13* and *TaFDL15* were identified from the EST database TGI (The Gene Indices, <http://compbio.dfci.harvard.edu/tgi/>) and have been characterized in a previous study (Li and Dubcovsky, 2008). We show in this study that the previously cloned *TaFDL2* and *TaFDL6* proteins can interact as prey with all six 14-3-3 proteins in yeast two-hybrid assays (Figure 2). Previously cloned *TaFDL15* interacted strongly with Ta14-3-3A, B and C, weakly with Ta14-3-3E, and not at all with Ta14-3-3D and Hv14-3-3H. Previously cloned *TaFDL13* did not interact with any 14-3-3 proteins (Figure 2).

Re-analyses of these four *FDL* genes using the wheat genomic sequences released by the International Wheat

Genome Sequencing Consortium (IWGSC 2014; Appendix S1) showed that the previously cloned *TaFDL2* cDNA was a partial transcript missing the 5' terminal end and that *TaFDL13* is an alternative splicing form of *TaFDL15*. We cloned the full-length coding region of *TaFDL2* and confirmed by yeast two-hybrid assays that the full-length *TaFDL2* protein also interacts with all six 14-3-3 proteins (Figure S2). Comparison of the *TaFDL13* transcript and genomic sequences revealed that the *TaFDL13* transcript is an alternative splicing form generated by the retention of the single intron present in *TaFDL15*. The alternative splice variant includes a premature stop codon that results in the truncation of the last 30 amino acids (Appendix S1), which may explain the inability of the encoded *TaFDL13* protein to interact with any of the 14-3-3 proteins.

In addition, we identified two barley FD-like proteins from the GenBank sequence database, which are not orthologous to any of the previously characterized wheat FD-like proteins, and designated them as HvFDL4 (GenBank BAK05116) and HvFDL5 (GenBank BAJ92629). Yeast two-hybrid assays showed that HvFDL4 interacts with all six 14-3-3 proteins (Figure 2), whereas HvFDL5 interacts only with Ta14-3-3A, Ta14-3-3C and Hv14-3-3H (Figure 2). The wheat homologues of *HvFDL4* and *HvFDL5* (*TaFDL4* and *TaFDL5*) were also obtained from the IWGSC and their sequences are described in Appendix S1. The relationships




Figure 2. Interactions between FD-like proteins and 14-3-3 proteins.

14-3-3 proteins were used as baits, wild type and mutant TaFDL2, TaFDL6, TaFDL15, HvFDL4, and HvFDL5 were used as preys. SD-L-W-H-A medium was used to test interactions. Tri-M = triple mutant S310A–T311A–S312A.

among the different FDL proteins described in this study are summarized in a phylogenetic analysis generated from the conserved bZIP and CDPK protein regions (Figure S3b).

The C-terminal regions of the proteins encoded by the canonical *FD-like* transcripts all contain a short motif targeted by CDPKs (Figure S3a), and an alanine substitution of a serine/threonine residue within this motif disrupts FD function in *Arabidopsis* (Abe *et al.*, 2005). We generated substitution mutations for the five FD-like proteins and confirmed by yeast two-hybrid assays that this phosphorylation site was required for interactions between FD-like proteins and 14-3-3 proteins in wheat and barley (Figure 2). An alanine substitution of the serine residue was enough to disrupt all interactions between FD-like and 14-3-3 proteins (Figure 2) except TaFDL6, for which a S310A/T311A/S312A triple mutation was required to abolish TaFDL6 interactions with Ta14-3-3A to E (Figure 2). The interaction between TaFDL6 and Hv14-3-3H was not affected by any of the mutations, presented alone or in combination (Figure 2).

Dimerization of 14-3-3 proteins

We then investigated the dimer formation ability of wheat and barley 14-3-3 proteins by yeast two-hybrid assays and tested the position effect of the GAL4 DNA-binding domain (BD) and activation domain (AD) on 14-3-3 dimerization. When the GAL4-BD and GAL4-AD were placed at the N-terminus of the fusion proteins (pGBKT7 and pGADT7 vectors), only Ta14-3-3A and Hv14-3-3H were able to form homodimers, whereas the other four 14-3-3 proteins (B to

E) did not (Figure 3a). By contrast, when the GAL4-BD and GAL4-AD were placed at the C-terminus of the fusion proteins (pGBKCg and pGADCg vectors), all 14-3-3 proteins were able to form homodimers (Figure 3a). These results suggest that the addition of the GAL4-BD or AD to the N-terminal regions of the 14-3-3 proteins interferes with the formation of homodimers.

Among all the heterodimer combinations, only six were able to form heterodimers when the GAL4-BD and AD were placed at the N-terminus (Figure S4). Ta14-3-3A and Ta14-3-3C, which did not form heterodimers when the GAL4 domains were placed at the N-terminus, were able to heterodimerize when the GAL4 domains were placed at the C-terminus (Figure S4). Taken together these results suggest that the N-terminus GAL4 interfered with the formation of 14-3-3 homodimers and heterodimers in yeast. We did not test the position effect of the C-terminus GAL4 on all possible 14-3-3 heterodimer pairs in yeast because those interactions were confirmed by BiFC assays in rice protoplasts, as described below.

BiFC assays were conducted using a split yellow fluorescent protein (YFP) system (Bracha-Drori *et al.*, 2004) in which 14-3-3 proteins were fused to the N-terminal half of YFP (N-YFP) and the C-terminal half of YFP (C-YFP), respectively. Construct pairs containing one N-YFP fusion and one C-YFP fusion were co-transformed into rice protoplasts. All six 14-3-3 proteins were able to form homodimers (Figure 3b) and heterodimers with each other (Figure S4) in rice protoplasts, suggesting limited specificity in 14-3-3 dimerization. Although both N-YFP and C-YFP were placed




Figure 3. Dimerization of 14-3-3 proteins.

(a) Homodimer formation tests by yeast two-hybrid assays. Bait is listed first in each pair combination.

(b) Homodimer formation of 14-3-3 proteins by BiFC in rice protoplasts. The split YFP system was used in BiFC assays.

at the N-terminus of all fusion proteins, we did not observe any interference on 14-3-3 dimerization in protoplasts.

Subcellular localization and *in vivo* interaction of FT1, 14-3-3 and TaFDL2

We fused FT1, TaFDL2 and Ta14-3-3C proteins with the green fluorescent protein (GFP) and determined the subcellular localization of these proteins in wheat protoplasts. Microscopy analysis detected fluorescence from FT1-GFP


in both the cytoplasm and nucleus; and from Ta14-3-3C-GFP predominantly in the cytoplasm and weakly in the nucleus (Figure 4a), suggesting that the interactions between FT1 and 14-3-3 proteins are most likely to occur in the cytoplasm. Fluorescence from TaFDL2-GFP was only detected in the nucleus (Figure 4a).

To validate the protein interactions among FT1, TaFDL2 and 14-3-3 proteins, we conducted BiFC assays using wheat protoplasts. Among the six 14-3-3 proteins, Ta14-3-3A

Figure 4. Subcellular localization and *in vivo* interaction of FT1, 14-3-3 and TaFDL2.

(a) Subcellular localization images of FT1-GFP, TaFDL2-GFP and Ta14-3-3C-GFP in wheat protoplasts.

(b) BiFC assays confirming interactions of FT1–Ta14-3-3A, FT1–14-3-3C, TaFDL2–Ta14-3-3A and TaFDL2–Ta14-3-3C in wheat protoplasts. The split YFP system was used in BiFC assays.



and Ta14-3-3C showed stronger interactions in yeast with both FT1 and FD-like proteins, and therefore were selected for the BiFC assays. FT1 and TaFDL2 were fused to N-YFP and Ta14-3-3A and Ta14-3-3C were fused to C-YFP. The

reconstituted YFP signal by FT1-14-3-3 interaction was detected in the cytoplasm, where 14-3-3 proteins were mainly localized (Figure 4b). The TaFDL2–14-3-3 BiFC signal was mainly detected in the nucleus (Figure 4b),

indicating that the FT1–14-3-3 complex needs to be translocated from the cytoplasm to the nucleus to interact with TaFDL2.

14-3-3 proteins are required for the FT–FDL2 interaction

We showed previously that the TaFDL2 protein can interact with FT1 in a yeast two-hybrid assay and can bind to the *VRN1* promoter *in vitro* (Li and Dubcovsky, 2008). Transgenic wheat plants overexpressing *FT1* showed parallel increases in *VRN1* transcripts both in leaves and shoot apical meristem (SAM), suggesting that FT1 provides transcriptional activation of *VRN1* through its protein interaction with TaFDL2 (Li and Dubcovsky, 2008; Pearce *et al.*, 2013). However, we were not able to detect a direct interaction between an FT1–TaFDL2 protein complex and the *VRN1* promoter (Li and Dubcovsky, 2008). Since both FT1 and TaFDL2 were able to interact with multiple 14-3-3 proteins (Figures 1 and 2), we decided to test if the FT1–TaFDL2 interaction detected before in yeast two-hybrid assays was mediated by endogenous yeast 14-3-3 proteins. We cloned one of the yeast 14-3-3 proteins (Sc14-3-3, GenBank EDN60443) and tested its interaction with FT1 and TaFDL2. Indeed, both FT1 and TaFDL2 were able to interact with the cloned yeast 14-3-3 protein (Figure S5).

We then examined whether FT1 mutants FT1-R62K and FT1-P94L were still able to interact with TaFDL2 by yeast two-hybrid assays. These two mutant proteins lost their ability to interact with 14-3-3 proteins (Figure 1), and were also unable to interact with TaFDL2 (Figure 5). Similarly, when TaFDL2–14-3-3 interaction was disrupted by the TaFDL2-S323A mutation, TaFDL2 lost its ability to interact with FT1 (Figure 5). These data suggest that a 14-3-3 bridge is essential to mediate FT1–TaFDL2 interactions and, therefore, for the formation of the FACs.

The FAC including FT1, Ta14-3-3C and TaFDL2 can bind to the *VRN1* promoter

Using electrophoretic mobility shift assays (EMSAs), we have shown before that the TaFDL2 protein binds to the *VRN1* promoter *in vitro* (Li and Dubcovsky, 2008). However, no super-shift was observed in the presence of both FT1 and TaFDL2 proteins (Li and Dubcovsky, 2008). Here, we performed additional EMSAs with Ta14-3-3C protein included in the binding reactions. TaFDL2 protein alone was sufficient to bind to *VRN1* promoter, whereas FT1 or Ta14-3-3C alone did not bind to DNA (Figure 6). When both TaFDL2 and Ta14-3-3C were added to the binding reaction, a super-shift was detected, and the presence of all three proteins (TaFDL2, Ta14-3-3C and FT1) was able to further delay the mobility of the protein–DNA complex (Figure 6). These data indicate that the FAC including FT1, Ta14-3-3C, and TaFDL2 has the ability to bind the *VRN1* promoter *in vitro*.

In the absence of Ta14-3-3C, the binding reaction including both FT1 and TaFDL2 resulted in a DNA shift similar to the one observed when only the TaFDL2 protein was incubated with the DNA probe (Figure 6). This result provides additional evidence that FT1 and TaFDL2 cannot interact in the absence of the 14-3-3 protein bridge.

Different FT-like proteins interact with different subsets of 14-3-3 proteins

In addition to *FT1*, five additional *FT-like* genes were described before in barley and wheat, and their phylogenetic relationship were also characterized (Lv *et al.*, 2014). To test the interactions between the different FT-like and 14-3-3 proteins, we cloned TaFT2 from wheat, and HvFT3 (DQ411319), HvFT4 (DQ411320), HvFT5 (EF012202) from




Figure 5. Yeast two-hybrid interactions between FT1 and TaFDL2 and the effect of mutations on their interaction.

SD-L-W was used to select for yeast transformants and SD-L-W-H-A was used to test interactions. Dilution factors from left to right are 1, 1:10, and 1:100.

VRN1-DNA	+	+	+	+	+	+	+	+
TaFDL2	-	+	+	+	-	-	-	+
Ta14-3-3C	-	-	+	+	+	-	+	-
FT1	-	-	-	+	-	+	+	+




Figure 6. Interactions between FAC component proteins and *VRN1* promoter by electrophoretic mobility shift assays.

The DNA probe is a fragment of the *VRN1* promoter containing bZIP binding sites CACGTA and CACGTC (Li and Dubcovsky, 2008). The proteins used in the DNA-protein binding reactions include purified FT1-GST, TaFDL2-GST and Ta14-3-3C-GST. A delay in the electrophoretic mobility of the DNA fragment indicates binding to the protein.

barley. We also cloned TaTFL1 from wheat and *HvCEN1* from barley, which are orthologues of the related *Arabidopsis TERMINAL FLOWER 1 (TFL1)* and *Antirrhinum CENTRORADIALIS (CEN)*, respectively. The results from these yeast two-hybrid assays are summarized in Table 1. The different FT-like proteins showed differences in their interactions with the wheat 14-3-3 proteins. HvFT3 and HvFT5 interacted only with Ta14-3-3C, whereas HvFT4 interacted with Ta14-3-3A, B, C and D proteins. Surprisingly, wheat TaFT2, which is the closest homologue of FT1, did not interact with any of the tested 14-3-3 proteins (Table 1). TaTFL1 showed a very weak interaction with Ta14-3-3C and no interactions with the other 14-3-3 proteins. *HvCEN1* did not interact with any of the wheat 14-3-3 proteins (Table 1).

FDL and 14-3-3 transcript levels in different tissues and developmental stages

Previous studies have shown that *FT-like* genes (*FT1* to *FT5*) in barley and wheat are expressed predominantly in the leaves, and have limited expression in the SAM (Faure *et al.*, 2007; Lv *et al.*, 2014). We show by qRT-PCR that *FD-like* and *14-3-3* genes are expressed at relatively high levels in both leaves and SAM under both long-day (LD, 16 h light) and short-day (SD, 8 h light) photoperiod (Figure S6a,b), and also at different stages of SAM development (Figure S6c,d). Analyses of previously published RNA-seq datasets (Choulet *et al.*, 2014) show expression of

FDL and *14-3-3* genes in roots, leaves, stems, spikes, and grains (Figure S6e,f).

Interestingly, relative abundances among *FDL* genes varied across tissues and developmental stages (Figure S6a,c,e). For example, transcript levels of *TaFDL6* were higher than those of *TaFDL2* in leaves (LD and SD), stems, roots and grains, but not in the SAM and spikes, where *TaFDL2* transcript levels were higher than those of *TaFDL6* (Figure S6a,c,e). *TaFDL15* also showed marked differences in transcript levels, varying from being the most highly expressed *FDL* gene in developing spikes, stems, and leaves (RNA-seq, Zadok sage 23), to having relative low transcript levels in young leaves and SAM (Figure S6a), roots, and grain (Figure S6e).

The relative transcript levels of the wheat *14-3-3* genes also showed some variability across tissues and developmental stages but the differences were less evident than for the *FDL* genes (Figure S6b,d,f). *Ta14-3-3A* was consistently the most highly expressed *14-3-3* gene in all the qRT-PCR and RNA-seq studies, and *Ta14-3-3B* was the second highest, except for roots and leaves (Zadok 23) in the RNA-seq study (Figure S6b,d,f). The other three wheat *14-3-3* genes were expressed at relatively lower levels and showed limited variability across tissues and developmental stages (Figure 6b,d).

DISCUSSION

Differential functions of *FT-like* proteins

In addition to *FT*, the *FT-like* family in *Arabidopsis* includes five other members: *TWIN SISTER OF FT (TSF)*, *TERMINAL FLOWER 1 (TFL1)*, *BROTHER OF FT AND TFL1 (BFT)*, *MOTHER OF FT AND TFL1 (MFT)*, and *Arabidopsis thaliana CENTRORADIALIS homologue (ATC)*. *TSF* functions as a florigenic activator (Kardailsky *et al.*, 1999; Kobayashi *et al.*, 1999; Yamaguchi *et al.*, 2005; Andres and Coupland, 2012), whereas *TFL1*, *ATC* and *BFT* function as floral repressors (Bradley *et al.*, 1997; Mimida *et al.*, 2001; Yoo *et al.*, 2010). *MFT* is involved in seed germination (Xi *et al.*, 2010). Some of these *FT-like* genes have been shown to function through interactions with FD-like proteins. For examples, *TFL1* has been shown to modulate the transcriptional repression of its target genes in an FD-dependent manner (Hanano and Goto, 2011), and *Arabidopsis BFT* and *Chrysanthemum Anti-florigenic protein (CsAFT)* have been shown to compete with *FT* for FD binding and to function as floral repressors (Higuchi *et al.*, 2013; Ryu *et al.*, 2013).

In rice, both *HD3A* and *RICE FLOWERING LOCUS T1 (RFT1)* encode mobile flowering signals that function as flowering promoters (Komiya *et al.*, 2009; Tsuji *et al.*, 2011). Overexpression of barley *HvFT1*, *HvFT2* and *HvFT3* in rice led to early flowering, indicating that these three *FT-like* genes can act as flowering promoters (Kikuchi *et al.*, 2009). Similarly, overexpression of *FT1* and *FT2* *Brachy-*

Table 1 Yeast two-hybrid interactions of different 14-3-3 with FT1 and FT-like proteins

Bait	Prey						
	FT1	TaFT2	HvFT3	HvFT4	HvFT5	TaTFL1	HvCEN1
14-3-3A	Yes ^a	No	No	Yes	No	No	No
14-3-3B	Yes	No	No	Yes	No	No	No
14-3-3C	Yes ^a	No	Yes	Yes	Yes	Weak	No
14-3-3D	Yes	No	No	Yes	No	No	No
14-3-3E	Yes	No	No	No	No	No	No
14-3-3H	No	No	No	No	No	No	No

^aThese interactions were validated by BiFC (Figure 4).

podium orthologues (*Bradi2g07070* and *Bradi1g48830*) resulted in extremely early flowering in transgenic *Brachypodium* plants (Wu *et al.*, 2013). Transgenic wheat plants overexpressing *FT1* show an upregulation of other *FT-like* genes suggesting that *FT1* acts as a positive regulator of other *FT-like* genes (Lv *et al.*, 2014). In spite of its importance, *FT1* does not seem to be essential for flowering in wheat. Combined mutations in *FT-A1* and *FT-B1* in tetraploid wheat or downregulation of all *FT1* homoeologs by RNA interference in hexaploid wheat, delay flowering only 2–4 weeks under LD (Lv *et al.*, 2014). Therefore, it is likely that other *FT-like* genes have overlapping roles in the promotion of wheat flowering.

Potential competitive interactions among FT-like, FD-like and 14-3-3 proteins

The interactions of some FT-like proteins with the same 14-3-3 proteins (Figure S7) can result in competitive interactions. For example, FT1, HvFT3, HvFT4, HvFT5 and TaTFL1 all interact with Ta14-3-3C (Table 1 and Figure S7) and may compete for this interaction. Interestingly, Ta14-3-3C is also able to interact with all the FDL proteins tested so far creating the opportunity for additional competitive interactions (Figures 2 and S7). Proteins Ta14-3-3A, Ta14-3-3B and Ta14-3-3D also interact with more than one FT-like proteins (FT1 and HvFT4) and may also compete for these interactions (Table 1 and Figure S7). FT1 and HvFT4 may be less affected by these competitive interactions because they can interact with a large number of different 14-3-3 proteins (Table 1 and Figure S7). By contrast, the related HvFT3 and HvFT5 proteins (80–84% identical in sequence) interact only with 14-3-3C suggesting that they can be very sensitive to competition for interactions with this particular protein.

FT1 and TaFT2 share high protein identity (~78%), but they do not seem to compete for common interactors. These two proteins have similar functions as strong flowering activators (Wu *et al.*, 2013; Lv *et al.*, 2014) but show no overlaps in their 14-3-3 protein interaction profiles. FT1 interacted with five out of the six 14-3-3 proteins tested so far, whereas TaFT2 showed no interactions with any of

these 14-3-3 proteins. This result suggests that some functional differentiation may exist between these two proteins.

The functions of the other FT-like proteins in wheat and barley are still not well characterized, so it is currently not possible to predict the effect of the different 14-3-3 interactions. A detailed functional characterization of FT-like proteins and their protein partners including 14-3-3 and FD-like proteins will help determine whether they act redundantly or antagonistically to FT1 in wheat flowering regulation.

Dimerization among 14-3-3 proteins

The selectivity of the interactions between FT-like and 14-3-3 proteins contrasts with the less selective interactions between 14-3-3 monomers. Our BiFC results in rice protoplasts showed no selectivity in the interactions among wheat 14-3-3 proteins. Therefore, dimerization between different 14-3-3 proteins is likely to depend on their relative affinities and abundances. In mammalian and yeast systems it has been demonstrated that the dimerization of 14-3-3 proteins is not completely random, and that there are preferences for certain dimer combinations (Paul *et al.*, 2012). For example, the mammalian 14-3-3 σ shows a preference to form homodimers, whereas 14-3-3 β is present primarily as heterodimers (Wilker *et al.*, 2005). The two yeast 14-3-3 proteins also show an inclination to form heterodimers rather than homodimers (Chaudhri *et al.*, 2003).

In plants, 14-3-3 dimerization is complex and still not well understood. Among the eight Arabidopsis 14-3-3 proteins tested so far, all can form homodimers as well as heterodimers with each other in yeast two-hybrid assays and native gel western blot analysis (Wu *et al.*, 1997). Proteomic profiling results indicate that 14-3-3 ω can dimerize with at least 10 of the 12 14-3-3 proteins expressed in Arabidopsis (Chang *et al.*, 2009). In cotton, none of the six cotton 14-3-3 proteins were able to form homodimers in yeast two-hybrid, and only some combinations of heterodimers were detected (Zhang *et al.*, 2010). However, the yeast two-hybrid vectors used in the cotton study (pGBKT7 and pGADT7) have both GAL4-BD and AD at the N-terminus of the fusion proteins, which could have interfered with the

interactions between 14-3-3 proteins, as demonstrated in this study (Figure 3a). Additional yeast two-hybrid experiments with the GAL4-BD and AD at the C-terminus of the fusion proteins will be required to interpret the cotton results correctly.

Even though the interactions among 14-3-3 proteins were not very selective, their interactions with different FT and FD-like proteins showed some specificity (Table 1 and Figure S7) and can play a role in the modulation of the effects of different FT-like mobile proteins. To understand these effects better we characterized the spatial and temporal expression of the different components of the FACs.

Differential expression profiles of *FT-like*, *FD-like* and *14-3-3* genes

Previous studies have shown differences in the relative levels of *FT-like* genes expression at different photoperiods (Lv *et al.*, 2014). When plants of the winter wheat variety Jagger were grown under LD for 5 weeks without vernalization, *FT3* transcript levels were significantly lower than those of the other *FT-like* genes. By contrast, when plants from the same variety were grown under SD, transcript levels of *FT3* (and also *FT4* and *FT5*) were significantly higher than those of *FT1* and *FT2* (Lv *et al.*, 2014). These results suggest a specific regulatory role of *FT3* (and possibly *FT4* and *FT5*) under short days.

Mobile FT-like proteins can reach multiple plant tissues via the phloem. Therefore, tissue specific responses likely depend on the other two members of the FAC. The FDL members are particularly important since they determine the potential DNA targets of the FAC. For example, the ectopic expression of FD in the leaves of transgenic Arabidopsis plants, results in high expression levels of its *AP1* target in the leaves, where these two genes are not normally expressed (Wigge *et al.*, 2005). The wheat *FDL2* gene is normally expressed in both apices and leaves (Figure S6), and its *VRN1* target is also expressed in both tissues (when FT1 is present).

In this study we show that the relative abundances of *FDL* transcripts vary significantly across tissues and developmental stages (Figure S6a,c,e). These differences are likely to be reflected in the relative abundance of FAC complexes and can contribute to the spatial and temporal modulation of the effects of the mobile FT-like signals. The relative transcript levels of the different wheat *14-3-3* genes also show some temporal and spatial variability (Figure S6b,d,e), which may also contribute to the modulation of the effects of the FT-like mobile proteins across tissues and developmental stages.

FD-like transcription factors determine the DNA target of florigen activation complexes

We show in this study that among the three components of the particular FAC that interacts with the *VRN1* pro-

motor, only TaFDL2 is essential for DNA binding (Figure 6). This agrees with previous results in rice, which indicated that the FDL recruited to the FAC is critical in the determination of the DNA specificity of the complex (Taoka *et al.*, 2011). This has been demonstrated for rice proteins OsFD1 and OsFD2, both of which can interact with HD3A and 14-3-3 proteins to form FACs (Tsuji *et al.*, 2013). If OsFD1 is recruited, the resulting FAC acts to promote flowering; whereas if OsFD2 is recruited, the resulting FAC functions in rice leaf development (Tsuji *et al.*, 2013). This result demonstrates that the presence of different FD-like genes has the potential to determine which genes are activated or repressed by their interacting FT-like proteins. It also shows that some of the FAC can be involved in the regulation of processes other than flowering. The presence of different FT-like proteins in the complex may determine the promoting or repressing role of the FAC or affect the intensity of their responses.

In a previous study, we showed that both TaFDL2 and TaFDL6 proteins interact with FT1, but only TaFDL2 was able to bind to the *VRN1* promoter (Li and Dubcovsky, 2008). These results suggested that TaFDL6 may target other genes, a possibility reinforced here by the observation that TaFDL6 interacts with all six 14-3-3 proteins (Figure 2) and that therefore, can be part of different FACs.

Additional FD-like proteins TaFDL15, HvFDL4 and HvFDL5 also interact with multiple 14-3-3 proteins (Figure 2). Therefore, the combinations of different FT-like, FD-like and 14-3-3 proteins can potentially generate a large number of different FACs that plants can use to decode the information carried by FT-like proteins. We are currently screening for mutations of individual FD-like and FT-like proteins to understand better the specific and overlapping functions of the different FACs.

Potential downstream targets of florigen activation complexes

The wheat *VRN1* gene encodes a MADS-box transcription factor that is important but not essential for wheat flowering (Chen and Dubcovsky, 2012). This gene is closely related to Arabidopsis paralogous genes *APETALA 1* (*AP1*), *CAULIFLOWER* (*CAL*) and *FRUITFULL* (*FUL*) (Trevaskis *et al.*, 2003; Yan *et al.*, 2003), which have overlapping functions in the induction of flowering (Ferrández *et al.*, 2000). In the temperate cereals, there are two additional MADS-box genes similar to *VRN1*, designated as *FUL2* (= *HvMADS8* = *OsMADS15*) and *FUL3* (= *HvMADS3* = *OsMADS18*) (Preston and Kellogg, 2006), but their role in the regulation of flowering is currently unknown.

In this study we show that a FAC including FT1, Ta14-3-3C and TaFDL2 can bind to the *VRN1* promoter *in vitro* (Figure 6). In rice, the FAC including HD3A, OsFD1 and GF14c (= 14-3-3C) proteins binds to the promoter of *OsMADS15*, which is the rice orthologue of *FUL2* (Taoka

et al., 2011). This result suggests that wheat *FUL2* (and may be *FUL3*) promoters could also be potential downstream targets of some wheat FACs.

A recent study has demonstrated that the induction of *FT1* is associated with the upregulation of GA biosynthetic genes and the downregulation of GA catabolic genes in the SAMs of wheat (Pearce *et al.*, 2013). Therefore, it would be interesting to investigate if the promoters of any of the biosynthetic GA20ox or catabolic GA2ox genes are targets of FACs including FT1.

In summary, our study shows that the multiple FT-like, FDL and 14-3-3 wheat proteins have the potential to generate a very large number of different FACs. These multiple factorial protein combinations provide the complexity and flexibility required to modulate differential responses to mobile FT-like signals in different tissue and developmental stages.

EXPERIMENTAL PROCEDURES

Plant materials and growth conditions

Rice japonica variety Kitaake and wheat variety Chinese Spring were used for protoplast extraction. Rice plants were grown in the dark at 28°C for 2 weeks before leaves and stems were harvested for protoplast extraction following procedures described before (Bart *et al.*, 2006). We adapted the rice protoplast extraction protocol to wheat protoplast extraction, but modified the growth conditions as following: wheat plants were grown under LD (16 h light/8 h dark) at temperatures of 22°C during the day and 18°C during the night for 2 weeks before protoplast extraction.

For cloning, wheat variety Chinese Spring and barley Golden Promise plants were grown under LD at temperatures of 22°C during the day and 18°C during the night for 4 weeks before leaves were collected for total RNA extraction and cDNA synthesis.

Phylogenetic analysis

Phylogenetic analysis of the 14-3-3 proteins was performed using full-length proteins. A neighbor-joining tree was constructed using the program MEGA 5.0 (Tamura *et al.*, 2011). Bootstrap confidence values were calculated based on 1000 iterations and only values larger than 50 are shown in the respective nodes.

GenBank accession numbers used in phylogenetic analysis include AtGF14chi = GRF1 (At4g09000), AtGF14omega = GRF2 (At1g78300), AtGF14psi = GRF3 (At5g38480), AtGF14phi = GRF4 (At1g35160), AtGF14upsilon = GRF5 (At5g16050), AtGF14lambda = GRF6 (At5g10450), AtGF14nu = GRF7 (At3g02520), AtGF14kappa = GRF8 (At5g65430), AtGF14mu = GRF9 (At2g42590), AtGF14epsilon = GRF10 (At1g22300), OsGF14a (Os08g0480800), OsGF14b (Os04g0462500), OsGF14c (Os08g0430500), OsGF14d (Os11g0546900), OsGF14e (Os02g0580300), OsGF14f (Os03g0710800), OsGF14g (Os01g0209200), OsGF14h (Q2R1D5), Ta14-3-3A (AAR89812), Ta14-3-3B (AGA84102), Ta14-3-3C (AEQ53933), Ta14-3-3D (BAB11740), Ta14-3-3E (BAB11739), Hv14-3-3A (CAA44259), Hv14-3-3B (CAA63658), Hv14-3-3C (CAA74592), HvGF14D (DQ295785), HvGF14E (DQ295786) and Hv14-3-3H (BAK01024). The two letters in front of gene or protein names indicate the organism from which the sequence or clone was obtained (At = *Arabidopsis thaliana*, Ta = *Triticum aestivum*, Hv = *Hordeum vulgare*, Os = *Oryza sativa*).

The phylogenetic analysis of the FDL proteins was performed using only the conserved region of the proteins including the bZIP domain and the CDPK target region (Figure S3a). The resulting neighbour-joining tree is presented in Figure S3(b) and the complete FDL sequences (DNA and protein) are presented in Appendix S1. The rice FD-like proteins (and their corresponding genes) included in the phylogeny analysis are: OsFD2 (Os06g0720900), OsFD3 (Os02g0833600), OsFD4 (Os08g0549600), Os05g048970, Os01g0813100 and OsJ_10665 (CM000140).

Yeast two-hybrid assays

Yeast vectors pGBKT7 (GAL4 DNA-BD), pGADT7 (GAL4 activation domain, AD) and yeast strain Y2HGold (Clontech, Mountain View, CA, USA) were used in the yeast two-hybrid assays. Two additional pairs of modified Gateway (Invitrogen) bait/prey vectors were also used in this study: (i) N-terminal GAL4 fusion vectors pLAW10 and pLAW11, which are the Gateway versions of pGBKT7 and pGADT7, respectively (Cantu *et al.*, 2013); and (ii) C-terminal GAL4 fusion vectors pGBKCG and pGADCg (Stellberger *et al.*, 2010). For all Gateway compatible clonings, pDONR/Zeo (Life Technologies, Grand Island, NY, USA), was used to generate the entry vectors. Primer sequences and cloning strategies are listed in Table S1. All constructs were verified by digests and sequencing. Yeast two-hybrid assays were performed according to the manufacturer's instructions (Clontech). Transformants were selected on SD medium lacking leucine (–L) and tryptophan (–W) plates and re-plated on SD medium lacking –L, –W, histidine (–H) and adenine (–A) to test the interactions.

Site-directed mutagenesis

QuikChange II XL site-directed mutagenesis kit (Agilent, Technologies, <http://www.agilent.com/>) was used to generate single amino acid substitution mutants for FT and FD-like proteins. All the mutant constructs were verified by sequencing. Primers used to generate the mutated genes are listed in Table S2.

Protein subcellular localization

To reduce plasmid size and increase the protoplast transfection efficiency, we engineered a smaller Gateway destination vector containing the GFP reporter. The 2 × 35S promoter-GFP6-Gateway cassette from pMDC43 (Arabidopsis Biological Resource Center, stock # CD3-741) was inserted in pUC57-KAN between restriction sites *EcoRI* and *HindIII* to generate Gateway destination vector pUC57-GFP. Entry vectors generated for yeast two-hybrid assays using pDONR/Zeo system (Table S1) were used to transfer *FT1*, *TaFDL2* or *Ta14-3-3C* into pUC57-GFP by site-specific recombination to generate GFP-FT1, GFP-TaFDL2 and GFP-Ta14-3-3C fusions. The resultant constructs were used in the transfection assays to determine the subcellular localization of these proteins in wheat protoplasts. Wheat protoplast isolation and transfection were conducted as described before (Bart *et al.*, 2006).

Bimolecular fluorescence complementation assays

Bimolecular fluorescence complementation assays using split YFP system were conducted using protocols described before (Brachardori *et al.*, 2004). The pDONR entry vectors generated for yeast two-hybrid (Table S1) were used to generate BiFC constructs: each protein pair was recombined into the Gateway destination vectors pY736 and pY735 to generate N-terminal YFP and C-terminal YFP fusion proteins. All constructs were verified by sequencing. Rice and wheat protoplast isolation and transfection were conducted as described (Bart *et al.*, 2006).

Electrophoretic mobility shift assays

The generation of FT1-GST and TaFDL2-GST fusion constructs was described previously (Li and Dubcovsky, 2008). Entry vector pDONR/Zeo-Ta14-3-3C generated for yeast two-hybrid was used to transfer Ta14-3-3C into pDEST15 (Gateway compatible, Invitrogen, <http://www.thermofisher.com/us/en/home/brands/invitrogen.html>) to generate the Ta14-3-3C-GST fusion construct. Protocols for GST-tagged protein expression and purification and for PCR procedures to amplify the *VRN1* promoter fragment containing two hybrid bZIP binding sites were described before (Li and Dubcovsky, 2008). DNA-protein binding reactions were carried out as described by Taoka *et al.* (2011) and separated on 5% polyacrylamide gels in $0.5 \times$ TBE buffer and stained with ethidium bromide.

Gene expression analysis by qRT-PCR

For the qRT-PCR studies, plants from diploid wheat (*T. monococcum* L., genome A^m) accessions DV92 and PI 349049 were grown for 8 weeks under short-day (SD, 8 h of light) or for 6 weeks under SD followed by 2 weeks under long-day (LD, 16 h of light). Both accessions have a spring growth habit and carry the same *vrn2* allele, but differ in their *VRN1* alleles. DV92 carries the *vrn1* allele that is not expressed under SD, and PI34049 carries the natural mutant allele *Vrn1g* that has a 34-bp deletion in the promoter region and is expressed under SD resulting in faster SAM development (Dubcovsky *et al.*, 2006 and Pearce *et al.*, 2013).

RNA samples extracted from leaves and apices were used to compare the relative transcript levels of five *FDL* and five *14-3-3* genes. Comparisons between leaves and apices were performed only in DV92, whereas comparisons across SAM developmental stages included samples from both DV92 and PI 349049. Plant growth conditions, apex dissection, RNA extraction, cDNA synthesis and qRT-PCR were conducted as described before (Pearce *et al.*, 2013). Briefly, qRT-PCR was performed using SYBR Green[®] and a 7500 Fast Real-Time PCR system (Applied Biosystems, <http://www.thermofisher.com/us/en/home/brands/applied-biosystems.html>), and transcript levels were expressed as linearized fold-*ACTIN* levels calculated by the formula $2^{(ACTIN\ CT - TARGET\ CT)}$. Primer sequences used in the qRT-PCR studies are listed in Table S3.

Published RNA-seq datasets (Choulet *et al.*, 2014) including different tissues (spike, leaf, grain, stem, and roots) and developmental stages (Zadoks *et al.*, 1974) were used to generate expression profiles of *FDL* and *14-3-3* genes among different tissues and developmental stages. Fragments per kilobase of transcript per million mapped reads (FPKM) values and their respective standard errors were obtained from one of the developmental stages for each of the five tissues and were used to generate the graphs presented in Figure S6(e,f).

ACKNOWLEDGEMENTS

This project was supported by the Howard Hughes Medical Institute and the Gordon and Betty Moore Foundation and by NRI grant numbers 2011-67013-30077 and 2011-68002-30029 from the USDA National Institute of Food and Agriculture. The authors thank Dr. Stephen Pearce and Mr. Hans Vasquez-Gross (University of California, Davis) for their help with expression analyses of the *FDL* and *14-3-3* genes.

CONFLICT OF INTEREST

The authors have no conflict of interest to declare.

SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article.

Figure S1. Neighbor-Joining tree of wheat, barley, rice and *Arabidopsis* 14-3-3 proteins.

Figure S2. Interactions between full-length TaFDL2 and 14-3-3 proteins.

Figure S3. Phylogenetic analysis of FDL proteins.

Figure S4. Dimerization analysis of 14-3-3 proteins by yeast two-hybrid and BiFC in rice protoplasts.

Figure S5. Yeast two-hybrid interactions among FT1, FDL2 and yeast 14-3-3 (Sc14-3-3).

Figure S6. Transcript levels of *FDL* and *14-3-3* genes in different tissues and developmental stages.

Figure S7. An interaction map summarizing the identified interactions among FT-like, FD-like and 14-3-3 proteins by yeast two-hybrid assays.

Table S1. Primers used for yeast two-hybrid constructs (restriction sites are underlined).

Table S2. Primers used in mutant generation by site-directed mutagenesis (mutations underlined).

Table S3. Primers used in the Q-RT-PCR analysis

Appendix S1. Sequence annotations of the wheat (A and B genomes) and barley FD-like genes used in phylogenetic analysis.

REFERENCES

- Abe, M., Kobayashi, Y., Yamamoto, S., Daimon, Y., Yamaguchi, A., Ikeda, Y., Ichinoki, H., Notaguchi, M., Goto, K. and Araki, T. (2005) FD, a bZIP protein mediating signals from the floral pathway integrator FT at the shoot apex. *Science*, **309**, 1052–1056.
- Ahn, J.H., Miller, D., Winter, V.J., Banfield, M.J., Lee, J.H., Yoo, S.Y., Henz, S.R., Brady, R.L. and Weigel, D. (2006) A divergent external loop confers antagonistic activity on floral regulators FT and TFL1. *EMBO J.* **25**, 605–614.
- Andres, F. and Coupland, G. (2012) The genetic basis of flowering responses to seasonal cues. *Nat. Genet.* **13**, 628–639.
- Banfield, M.J., Barker, J.J., Perry, A.C. and Brady, R.L. (1998) Function from structure? The crystal structure of human phosphatidylethanolamine-binding protein suggests a role in membrane signal transduction. *Structure*, **6**, 1245–1254.
- Bart, R., Chern, M., Park, C., Bartley, L. and Ronald, P.C. (2006) A novel system for gene silencing using siRNAs in rice leaf and stem-derived protoplasts. *Plant Methods*, **2**, 13.
- Bracha-Drori, K., Shichrur, K., Katz, A., Oliva, M., Angelovici, R., Yalovsky, S. and Ohad, N. (2004) Detection of protein-protein interactions in plants using biomolecular fluorescence complementation. *Plant J.* **40**, 419–427.
- Bradley, D., Ratcliffe, O., Vincent, C., Carpenter, R. and Coen, E. (1997) Inflorescence commitment and architecture in *Arabidopsis*. *Science*, **275**, 80–83.
- Cantu, D., Yang, B., Ruan, R., Li, K., Menzo, V., Fu, D., Chern, M., Ronald, P.C. and Dubcovsky, J. (2013) Comparative analysis of protein-protein interactions in the defense response of rice and wheat. *BMC Genom.* **14**, 166.
- Chang, I.F., Curran, A., Woolsey, R., Quilici, D., Cushman, J.C., Mittler, R., Harmon, A. and Harper, J.F. (2009) Proteomic profiling of tandem affinity purified 14-3-3 protein complexes in *Arabidopsis thaliana*. *Proteomics*, **9**, 2967–2985.
- Chaudhri, M., Scarabel, M. and Aitken, A. (2003) Mammalian and yeast 14-3-3 isoforms form distinct patterns of dimers *in vivo*. *Biochem. Biophys. Res. Commun.* **300**, 679–685.
- Chen, A. and Dubcovsky, J. (2012) Wheat TILLING mutants show that the vernalization gene *VRN1* down-regulates the flowering repressor *VRN2* in leaves but is not essential for flowering. *PLoS Genet.* **8**, e1003134.
- Choulet, F., Alberti, A., Theil, S. *et al.* (2014) Structural and functional partitioning of bread wheat chromosome 3B. *Science*, **345**, 1249721.

- Corbesier, L., Vincent, C., Jang, S. et al. (2007) FT protein movement contributes to long-distance signaling in floral induction of Arabidopsis. *Science*, **316**, 1030–1033.
- Denison, F.C., Paul, A., Zupanska, A.K. and Ferl, R.J. (2011) 14-3-3 proteins in plant physiology. *Semin. Cell Dev. Biol.* **22**, 720–727.
- Dubcovsky, J., Loukoianov, A., Fu, D., Valarik, M., Sanchez, A. and Yan, L. (2006) Effect of photoperiod on the regulation of wheat vernalization genes *VRN1* and *VRN2*. *Plant Mol. Biol.* **60**, 469–480.
- Faure, S., Higgins, J., Turner, A. and Laurie, D.A. (2007) The *FLOWERING LOCUS T*-like gene family in barley (*Hordeum vulgare*). *Genetics*, **176**, 599–609.
- Ferrández, C., Gu, Q., Martienssen, R. and Yanofsky, M.F. (2000) Redundant regulation of meristem identity and plant architecture by *FRUITFULL*, *APETALA1* and *CAULIFLOWER*. *Development*, **127**, 725–734.
- Granovsky, A.E. and Rosner, M.R. (2008) Raf kinase inhibitory protein: a signal transduction modulator and metastasis suppressor. *Cell Res.* **18**, 452–457.
- Hanano, S. and Goto, K. (2011) Arabidopsis TERMINAL FLOWER1 is involved in the regulation of flowering time and inflorescence development through transcriptional repression. *Plant Cell*, **23**, 3172–3184.
- Hanzawa, Y., Money, T. and Bradley, D. (2005) A single amino acid converts a repressor to an activator of flowering. *Proc. Natl Acad. Sci. USA*, **102**, 7748–7753.
- Higuchi, Y., Nayumi, T., Oda, A., Nakano, Y., Sumitomo, K., Fukai, S. and Hisamatsu, T. (2013) The gated induction system of a systemic floral inhibitor, antiflorigen, determines obligate short-day flowering in chrysanthemums. *Proc. Natl Acad. Sci. USA*, **110**, 17137–17142.
- International Wheat Genome Sequencing Consortium (2014) A chromosome-based draft sequence of the hexaploid bread wheat (*Triticum aestivum*) genome. *Science*, **345**, 1251788.
- Kardailsky, I., Shukla, V.K., Ahn, J.H., Dagenais, N., Christensen, S.K., Nguyen, J.T., Chory, J., Harrison, M.J. and Weigel, D. (1999) Activation tagging of the floral inducer *FT*. *Science*, **286**, 1962–1965.
- Kikuchi, R., Kawahigashi, H., Ando, T., Tonooka, T. and Handa, H. (2009) Molecular and functional characterization of PEBP genes in barley reveal the diversification of their roles in flowering. *Plant Physiol.* **149**, 1341–1353.
- Kobayashi, Y., Kaya, H., Goto, K., Iwabuchi, M. and Araki, T. (1999) A pair of related genes with antagonistic roles in mediating flowering signals. *Science*, **286**, 1960–1962.
- Komiya, R., Ikegami, A., Tamaki, S., Yokoi, S. and Shimanoto, K. (2009) A gene network for long-day flowering activates *RFT1* encoding a mobile flowering signal in rice. *Development*, **135**, 767–774.
- Krasileva, K.V., Buffalo, V., Bailey, P. et al. (2013) Separating homeologs by phasing in the tetraploid wheat transcriptome. *Genome Biol.* **14**, R66.
- Li, C. and Dubcovsky, J. (2008) Wheat FT protein regulates *VRN1* transcription through interactions with FDL2. *Plant J.* **55**, 543–554.
- Lv, B., Nitcher, R., Han, X., Wang, S., Ni, F., Li, K., Pearce, S., Wu, J., Dubcovsky, J. and Fu, D. (2014) Characterization of *FLOWERING LOCUS T1* (*FT1*) gene in *Brachypodium* and wheat. *PLoS ONE*, **9**, e94171.
- Mimida, N., Goto, K., Kobayashi, Y., Araki, T., Ahn, J.H., Weigel, D., Murata, M., Motoyoshi, F. and Sakamoto, W. (2001) Functional divergence of the *TFL1*-like gene family in Arabidopsis revealed by characterization of a novel homologue. *Genes Cells*, **6**, 327–336.
- Muszynski, M.G., Dam, T., Li, B., Shirbroun, D.M., Hou, Z., Bruggemann, E., Archibald, R., Ananiev, E.V. and Danilevskaya, O.N. (2006) *delayed flowering1* encodes a basic leucine zipper protein that mediates floral inductive signals at the shoot apex in maize. *Plant Physiol.* **142**, 1523–1536.
- Paul, A., Denison, F.C., Schultz, E., Zupanska, A.K. and Ferl, R.J. (2012) 14-3-3 phosphoprotein interaction networks—does isoform diversity present functional interaction specification? *Front. Plant Sci.* **3**, 190.
- Pearce, S., Vanzetti, L.S. and Dubcovsky, J. (2013) Exogenous gibberellins induce wheat spike development under short days only in the presence of *VERNALIZATION 1*. *Plant Physiol.* **163**, 1433–1445.
- Pnueli, L., Gutfinger, T., Hareven, D., Ben-Naim, O., Ron, N., Adir, N. and Lifschitz, E. (2001) Tomato SP-interacting proteins define a conserved signaling system that regulates shoot architecture and flowering. *Plant Cell*, **13**, 2687–2702.
- Preston, J.C. and Kellogg, E.A. (2006) Reconstructing the evolutionary history of paralogous *APETALA1/FRUITFULL*-like genes in grasses (Poaceae). *Genetics*, **174**, 421–437.
- Ryu, J.Y., Lee, H.J., Seo, P.J., Jung, J.H., Ahn, J.H. and Park, C.M. (2013) The Arabidopsis floral repressor BFT delays flowering by competing with FT for FD binding under high salinity. *Mol. Plant*, **7**, 377–387.
- Schoonheim, P.J., Veiga, H., Pereira Dda, C., Friso, G., van Wijk, K.J. and de Boer, A.H. (2007) A comprehensive analysis of the 14-3-3 interactome in barley leaves using a complementary proteomics and two-hybrid approach. *Plant Physiol.* **143**, 670–683.
- Serre, L., Vallée, B., Bureaud, N., Schoentgen, F. and Zelwer, C. (1998) Crystal structure of the phosphatidylethanolamine-binding protein from bovine brain: a novel structural class of phospholipid-binding proteins. *Structure*, **6**, 1255–1265.
- Stellberger, T., Häuser, R., Baiker, A., Pothineni, V.R., Haas, J. and Uetz, P. (2010) Improving the yeast two-hybrid system with permuted fusions proteins: the Varicella Zoster Virus interactome. *Proteome Sci.* **8**, 8.
- Tamaki, S., Matsuo, S., Wong, H.L., Yokoi, S. and Shimamoto, K. (2007) Hd3a protein is a mobile flowering signal in rice. *Science*, **316**, 1033–1036.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. and Kumar, S. (2011) MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* **28**, 2731–2739.
- Taoka, K., Ohki, I., Tsuji, H. et al. (2011) 14-3-3 proteins act as intracellular receptors for rice Hd3a florigen. *Nature*, **476**, 332–335.
- Taoka, K., Ohki, I., Tsuji, H., Kojima, C. and Shimamoto, K. (2013) Structure and function of florigen and the receptor complex. *Trends Plant Sci.* **18**, 287–294.
- Trevaskis, B., Bagnall, D.J., Ellis, M.H., Peacock, W.J. and Dennis, E.S. (2003) MADS box genes control vernalization-induced flowering in cereals. *Proc. Natl Acad. Sci. USA*, **100**, 13099–13104.
- Tsuji, H., Taoka, K. and Shimamoto, K. (2011) Regulation of flowering in rice: two florigen genes, a complex gene network, and natural variation. *Curr. Opin. Plant Biol.* **14**, 45–52.
- Tsuji, H., Nakamura, H., Taoka, K. and Shimamoto, K. (2013) Functional diversification of FD transcription factors in rice, components of florigen activation complexes. *Plant Cell Physiol.* **54**, 385–397.
- Wigge, P.A., Kim, M.C., Jaeger, K.E., Busch, W., Schmid, M., Lohmann, J.U. and Weigel, D. (2005) Integration of spatial and temporal information during floral induction in Arabidopsis. *Science*, **309**, 1056–1059.
- Wilker, E.W., Grant, R.A., Artim, S.C. and Yaffe, M.B. (2005) A structural basis for 14-3-3 sigma functional specificity. *J. Biol. Chem.* **280**, 18891–18898.
- Wu, K., Lu, G., Sehnke, P. and Ferl, R.J. (1997) The heterologous interactions among plant 14-3-3 proteins and identification of regions that are important for dimerization. *Arch. Biochem. Biophys.* **339**, 2–8.
- Wu, L., Liu, D., Wu, J., Zhang, R., Qin, Z., Liu, D., Li, A., Fu, D., Zhai, W. and Mao, L. (2013) Regulation of *Flowering Locus T* by a microRNA in *Brachypodium distachyon*. *Plant Cell*, **25**, 4363–4377.
- Xi, W., Liu, C., Hou, X. and Yu, H. (2010) *MOTHER OF FT AND TFL1* regulates seed germination through a negative feedback loop modulating ABA signaling in Arabidopsis. *Plant Cell*, **22**, 1733–1748.
- Yamaguchi, A., Kobayashi, Y., Goto, K., Abe, M. and Araki, T. (2005) *TWIN SISTER OF FT (TSF)* acts as a floral pathway integrator redundantly with *FT*. *Plant Cell Physiol.* **46**, 1175–1189.
- Yan, L., Loukoianov, A., Tranquilli, G., Helguera, M., Fahima, T. and Dubcovsky, J. (2003) Positional cloning of wheat vernalization gene *VRN1*. *Proc. Natl Acad. Sci. USA*, **100**, 6263–6268.
- Yan, L., Fu, D., Li, C., Blechl, A., Tranquilli, G., Bonafede, M., Sanchez, A., Valarik, M., Yasuda, S. and Dubcovsky, J. (2006) The wheat and barley vernalization gene *VRN3* is an orthologue of *FT*. *Proc. Natl Acad. Sci. USA*, **103**, 19581–19586.
- Yoo, S.J., Chung, K.S., Jung, S.H., Yoo, S.Y., Lee, J.S. and Ahn, J.H. (2010) *BROTHER OF FT AND TFL1 (BFT)* has *TFL1*-like activity and functions redundantly with *TFL1* in inflorescence meristem development in Arabidopsis. *Plant J.* **63**, 241–253.
- Zadoks, J.C., Chang, T.T. and Konzak, C.F. (1974) A decimal code for the growth stages of cereals. *Weed Res.* **14**, 415–421.
- Zeevaert, J.A.D. (2006) Florigen coming of age after 70 years. *Plant Cell*, **18**, 1783–1789.
- Zhang, Z.T., Zhou, Y., Li, Y., Shao, S.Q., Li, B.Y., Shi, H.Y. and Li, X.B. (2010) Interactome analysis of the six cotton 14-3-3s that are preferentially expressed in fibers and involved in cell elongation. *J. Exp. Bot.* **61**, 3331–3344.