UC Irvine

UC Irvine Previously Published Works

Title

Retinal sensitivity measured by the pupillary light reflex in RCS and albino rats.

Permalink

<https://escholarship.org/uc/item/59s181w5>

Journal

Vision research, 22(9)

ISSN

0042-6989

Authors

Trejo, LJ
Cicerone, CM

Publication Date

1982

DOI

10.1016/0042-6989(82)90081-5

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

RETINAL SENSITIVITY MEASURED BY THE PUPILLARY LIGHT REFLEX IN RCS AND ALBINO RATS

LEONARD J. TREJO* and CAROL M. CICERONE

Department of Psychology, C-009, University of California, San Diego, La Jolla, CA 92093, U.S.A.

(Received 14 July 1981; in revised form 17 February 1982)

Abstract—The effects of retinal degeneration on the sensitivity of the retina were studied in the Royal College of Surgeons (RCS) rat by measuring the light reflex of the pupil in response to ganzfeld (full field) flashes. Light reflex thresholds were measured for animals from 32 to 683 days of age, and an age-related decrease in sensitivity of 5.2 log units (maximum) was measured. In contrast, thresholds for non-dystrophic albino controls increased only slightly during a comparable period. RCS rat thresholds increased more for short wavelength light than for long wavelength light. The end result was an altered action spectrum of the light reflex which largely, but not exclusively, reflected cone function. Even in cases of advanced degeneration the light reflex thresholds we measured showed significant input from rods. Pupillary dark adaptation measured following ganzfeld bleaches (10%) with test stimuli of two different wavelengths revealed two mechanisms; a photopic mechanism ($\lambda_{\max} = 520$) determined thresholds early in dark adaptation, but later a scotopic mechanism ($\lambda_{\max} = 500$) participated in the light reflex.

INTRODUCTION

The Royal College of Surgeons (RCS) rat suffers from hereditary retinal degeneration (Dowling and Sidman, 1962) and has served as an animal model of the class of inherited human diseases, retinitis pigmentosa (Young, 1977). Retinitis pigmentosa (RP) is a retinal degeneration syndrome with various modes of inheritance. The most commonly reported initial symptom is night blindness which may initially spare the central region. However, photopic vision and central vision eventually fail in most cases where the patient reaches a sufficiently advanced age (for a review see Krill, 1972).

In comparison with what is known about the etiology of retinitis pigmentosa there exists a relative wealth of data concerning the nature and the time course of the physical changes which occur in the retina of the RCS rat (e.g. Bourne *et al.*, 1938; Cicerone *et al.*, 1979; Dowling and Sidman, 1962; LaVail *et al.*, 1974; Lucas *et al.*, 1955; Noell, 1977). From the work of Cicerone *et al.* and of LaVail *et al.*, it is known that both rods and cones degenerate, but the rods do so more extensively, and in a shorter period of time. The result is a drastic change from mostly rods in the retinas of young RCS rats to mostly cones in rats with advanced degeneration. Accordingly, Cicerone *et al.* have shown that all ganglion cells with measureable sensitivity in RCS rats with advanced degeneration appear to be driven by cone inputs. On the other hand, the spectral sensitivity of single ganglion cells in normal rats invariably matches a rhodopsin action spectrum when measured under the same

(dark adapted) conditions. Although the exclusive presence of units with photopic sensitivity in the RCS retina provides evidence for the selective survival of cones, it is conceivable that many scotopic units remain undetected. In fact, ganglion cells with low sensitivity are routinely encountered in the optic tract of older RCS rats (Cicerone *et al.*, 1979).

In mammals, the pupil responds to a flash of light with a transient constriction known as the pupillary light reflex. The relative ease of observing the light reflex and its high sensitivity motivated our use of this response to estimate retinal sensitivity. In humans, the evidence that the receptors which mediate vision are the same receptors responsible for the light reflex is compelling (Lowenstein and Loewenfeld, 1969). The direct light reflex of the rat is similar to that of humans, but unlike humans, the existence of a consensual pupillary light reflex in rats has not been demonstrated (Lowenstein and Loewenfeld, 1969). As a control for possible extraretinal influences we determined at the outset of this study that cutting the optic nerve of the rat's eye abolished the pupillary light reflex. This confirmed that in the rat, as in humans, the light reflex depends on stimulation of photoreceptors in the retina. The pupillary system has a considerable gain and a unitary response, presumed to reflect activity integrated over large retinal areas (Bouma, 1965; Schweitzer, 1955). These characteristics made the pupillary light reflex an ideal response with which to study the absolute sensitivity of the retina to light and the spectral sensitivity throughout the lifespan of the RCS rat. The pupillary light reflex measurements we report here provide evidence for a rod contribution persisting in advanced degeneration. This input, although reduced by over 4 log units as

*Present address: Department of Ophthalmology RJ-10, University of Washington, Seattle, WA 98195, U.S.A.

compared to the normal, is reflected in the pupillary light reflex of animals as old as 297 days postnatal.

A preliminary report of some of these findings has been presented elsewhere (Trejo and Cicerone, 1978).

METHODS

Animals

Eleven pink-eyed beige and tan hooded RCS rats were selected as subjects from a colony maintained at the University of California, San Diego. Our colony was derived by mating of siblings from RCS rats descended from the colony originally maintained at Harvard University. Six Sprague-Dawley albino rats served as controls.

All rats were reared from birth in a darkened room where light exposure was limited to brief periods of illumination from red overhead lights during cage cleaning and during preparation of the animals for experiments. Ophthalmoscopic examinations revealed no cataracts or other gross ocular anomalies in any of the animals used in this study.

Apparatus

A uniform, full-field illumination of variable wavelength and intensity was presented to the eye of the rat (see Fig. 1). The pupil of the eye was viewed continuously with the aid of an infrared sensitive television camera (Panasonic WV-1350) and a video

monitor. Responses of the pupil to ganzfeld flashes were also videotaped for later, more detailed analyses.

Procedure

Animal preparation. Each animal was dark adapted for a minimum of 12 hours before each experimental session. Under dim red illumination and mild ether narcosis the animal was placed in a restraining jacket and a restraining box. The box was placed on a platform with the animal's head protruding into the integrating sphere and 15–30 min was allowed to ensure complete recovery from the ether exposure.

Threshold measurements. The threshold criterion was a transient, 0.2 mm constriction in the diameter of the pupil in response to a 500 msec flash. The observer waited until the pupil diameter was within about 1 mm of its dark adapted diameter and the animal was still before delivering a test flash. Since the latency of the light reflex was stable, the observer could discount other pupillary movements which were randomly related to stimulus onset.

Thresholds were determined for calibrated combinations of interference filters (Ditric Optics, Marlboro, MA, 3 cavity, 8–12 nm bandwidth at half height) and neutral density filters (Oriel) in the beam. The sequence of wavelengths was quasirandomized with the restriction that the 500 nm filter was always used at the beginning, in the middle, and at the end as a check for changes in the level of adaptation during the session.

Time-course and intensity-response functions. The amplitude of pupillary constriction was measured as a function of the intensity of the stimulus to yield intensity-response functions. The luminance of the stimulus was systematically increased in 0.5 log unit steps, and the corresponding pupillary movements were videotaped. During analysis the sequences were viewed on a video monitor at an overall magnification of $19.7\times$ and measurements were made on stopped images at 0.2 sec steps. These measurements were used to track the time course of the light reflex (see Fig. 2). The maximum pupillary constriction for each stimulus obtained from these measurements was used to plot the intensity-response functions. Our measurements showed that the slopes of these functions did not vary significantly with wavelength (e.g. Fig. 6).

Dark adaptation. Pupillary sensitivity to light of 500 and of 600 nm was measured during recovery from exposure to a white ganzfeld bleach produced by a 30 sec exposure to the full output of a 150 W xenon arc lamp in the integrating sphere. In this condition the sphere produced an irradiance of $12.36 \log \text{ quanta mm}^{-2}$ at the corneal surface. Using Cone's (1963) analysis of Dowling's (1963) 5 min bleaches we estimated that this light bleached about 10% of the pigment of the retina. (The time-weighted average pupil diameter during the bleach was 0.8 mm, and this value was used in the calculation.)

Fig. 1. Diagram of the apparatus used to measure pupillary responses in a ganzfeld. An infrared light source, IL, which was a tungsten filament bulb housed in a tube covered with a Wratten 87C infrared filter, irradiated the left eye of the rat, R, held in the restraining box, B. The infrared sensitive TV camera and lens assembly produced a magnified ($13\times$) image of the pupil on a video monitor, TV. Pupil images could also be recorded on a videocassette recorder, VCR. When the observer activated the shutter, S, a 500 msec pulse of light from the xenon arc lamp, L, passed through interference and neutral density filters, F, and an aperture in the integrating sphere, IS, to produce on the surface of the sphere an arc image, i, which diffusely illuminated the rest of the sphere and the left eye of the rat by multiple reflections from the interior surface.

An experienced observer decided when a stimulus produced a threshold constriction. As a check on this method three entire dark adaptation sequences were videotaped and subsequently analyzed as described above. This more objective method produced dark adaptation curves which did not differ systematically from those produced by the observer's judgments. Due to the speed of recovery and to the necessity of changing filters manually the number of points obtained for each run was not large.

Calibration. The irradiance at the cornea with each interference or neutral density filter interposed in the light beam was measured with an electronic radiometer (Model 450, E. G. & G. Electro-optics Division; Salem, MA) positioned at the location occupied by the rat's eye.

Since we lacked comparative data concerning the relative size and light gathering properties of the eyes of RCS rats and of the control strain, all stimulus values are reported in units of corneal irradiance. To estimate the retinal irradiance produced by our threshold stimulus for the control animals, Cone's (1963) estimates of 12% absorption by the ocular media, 23% absorption of 500 nm light by rods, and 40×10^4 rods mm^{-2} , and Block's (1969) estimates of 2.97 mm for the posterior nodal distance and 1.34 for the index of refraction at the cornea of the albino rat eye were used. Our measured values of corneal irradiance of our 500 nm light ($5.38 \log$ quanta $\text{sec}^{-1} \text{mm}^{-2}$) and pupil diameter (3 mm) were used to compute a value of 1.75×10^{-1} quantum absorbed rod $^{-1} \text{sec}^{-1}$ at threshold.

RESULTS

The pupillary light reflex

Plots of net pupillary constriction as a function of post-stimulus time at six levels of intensity, over a four log unit range, for a 361-day old albino control appear in Fig. 2. At threshold stimulation (0.2 mm decrease in pupil diameter) the pupil begins to con-

Fig. 2. Plots of net pupillary constriction as a function of poststimulus time at six levels of intensity for a 361-day old albino control. The 500 msec test flash was set at 2.21 (circles), 2.69 (triangles), 3.13 (slashed circles), 4.22 (inverted triangles), 5.13 (diamonds), and 6.12 (squares) $\log$ quanta $\text{sec}^{-1} \text{mm}^{-2}$ measured at the cornea.

strict approx. 400 msec after stimulus onset and reaches a minimum diameter near 1 sec. The redilation following threshold constriction requires between 800 msec and 1 sec. For the highest levels of our stimuli, constriction time rarely exceeds 3 sec, whereas dilatation time to the pre-stimulus diameter may exceed 30 sec. These are general characteristics of mammalian pupillary light reflexes which have been previously described (Lowenstein and Loewenfeld, 1969).

As the RCS rat ages, the time-course of the pupillary constriction to suprathreshold light levels becomes more prolonged. Figure 3 shows plots of net pupillary constriction as a function of post-stimulus time at different levels of intensity for RCS rats of ages 45, 135, 363, 432 and 524 days of age. These can be compared to the plots of pupillary constriction for the 361-day old albino rat shown in Fig. 2. In addition to the prolonged time course, the pupillary

Fig. 3. Plots of net pupillary constriction as a function of poststimulus time for RCS rats of ages 45, 135, 363, 432 and 524 days of age. The intensity levels (in units of $\log$ quanta $\text{sec}^{-1} \text{mm}^{-2}$ of corneal irradiance) of the 500 msec test stimuli are noted to the right of the plots. The error bars for the plots of the responses of the 524-day old RCS indicate 2 SEM for the most variable point in each plot.

light reflex for RCS animals older than 135 days of age exhibits two phases of constriction at high intensity levels. Furthermore, the older the RCS rat, the more light is required to attain a specified maximal constriction as compared to the 361-day old control. An indication of variability in these plots is given by the error bars for the 524-day old RCS rat. These indicate two standard errors of the mean (SEM) for the most variable data points of the set of plotted averages.

Pupillary sensitivity vs age

Pupillary ganzfeld thresholds for four albino control rats are plotted in Fig. 4. The corneal irradiance at threshold of our 500 nm light stimulus is plotted as a function of postnatal age in days. Measurements were made between 53 and 705 days of age with multiple measurements made on certain individuals.

In the same figure thresholds are plotted for 11 RCS rats ranging between 32 and 683 days of age. For the RCS rats, thresholds increase steadily between 32 and 683 days of age, resulting in progressively larger threshold elevation relative to controls. Exponential functions were fit to each set of data, and these are shown as solid lines in Fig. 4. The equation for the function describing the RCS rat data is given by

$$\log I(t) = 7.5 - 5.8 e^{-(t/303)} \quad (1)$$

and that for the albino control data by

$$\log I(t) = 3.5 - 1.9 e^{-(t/500)} \quad (2)$$

Fig. 4. Pupillary light reflex thresholds for albino control rats and for RCS rats of different ages. The ordinate gives the intensity of a 500 nm, ganzfeld flash (500 ms) in log quanta $\text{sec}^{-1} \text{mm}^{-2}$ measured at the cornea. The abscissa indicates postnatal age in days. Thresholds of the four albino control rats were measured at 53, 78, 166, 203, 208, 281, 361, 427, 433 and 705 days of age. Different open symbols denote different control animals. The line through the control thresholds is described by an exponential function, $\log I(t) = 3.5 - 1.9 e^{-(t/500)}$, where I is corneal irradiance and t is age in days. Thresholds of 11 RCS rats at various ages are marked by closed symbols. Measurements for RCS rats were made at 27, 32, 45, 52, 81, 82, 84, 86, 92, 112, 132, 157, 244, 315, 389, 432, 494, 524, 541, 589, 663, 666 and 683 days of age. Different animals are indicated with different symbols. The function drawn through the RCS rat data has an equation of the form $\log I(t) = 7.5 - 5.8 e^{-(t/303)}$, where I is corneal irradiance and t is age in days.

where I is the threshold irradiance in quanta $\text{sec}^{-1} \text{mm}^{-2}$ at the cornea and t is age. The error bars shown at the right of each curve represent two standard deviations of the discrepancy between data and prediction for each model. The time constant of the fit to the RCS data is significant ($F_{(1,23)} = 191.74$), predicting 4.62 log units of threshold elevation between 31 and 683 days. The time constant of the albino fit is also significant ($F_{(1,10)} = 11.56$), predicting 1.25 log units of threshold elevation between 53 and 705 days.

Intensity-response functions

The maximum pupillary constriction for different intensities of a 60°, 500 nm flash of 500 msec duration is plotted in Fig. 5 for a control rat and for RCS rats of 31, 165 and 298 days of age. (In preliminary experiments, pupillary responses were measured in response to a 60° field flashed on a diffusing screen which the animal viewed from the restraining box.) The function for the 31-day old RCS resembles that of the control closely enough to permit superposition if it is displaced horizontally by 1.7 log units, implying that the same mechanisms mediate the light reflex in the normal and in the 1-month old RCS rat.

Superposition of the intensity-response functions of the 165- and 298-day old RCS rats upon that of the control is not possible. Thus it is apparent that in addition to overall losses of sensitivity there are changes in the relative sensitivity of the mechanisms which mediate the light reflex at different intensities.

Fig. 5. Intensity-response functions for an albino control and three RCS rats of different ages. The abscissa gives the corneal irradiance of a 60° , 500 nm flash (500 msec) in $\log \text{quanta sec}^{-1} \text{mm}^{-2}$. The ordinate gives the maximum amplitude of pupillary constriction measured as the net change in the diameter of the pupil in mm. Symbols are used as follows: circles, a control rat; squares, a 31-day old RCS rat; hexagons, a 165-day old RCS rat; triangles, a 298-day old RCS rat. The dashed line intercepts each function at threshold, defined as an 0.2 mm constriction.

Intensity-response functions measured with a 500 and 600 nm test are plotted for the control rat and the 31- and 298-day old RCS rats in Fig. 6. Each set of points has been scotopically equated by shifting the set generated by the 600 nm light to the left 1.38 log units, the difference required for a Dartnall nomo-

gram curve peaking at 500 nm (Ebrey and Honig, 1977). This results in a superposition of the lower branches of the curves for the two wavelengths and shows that the action spectrum of the pupillary light reflex at scotopic levels is likely to be that of rhodopsin. Even for the 298-day old RCS rat, at lower

Fig. 6. Intensity response functions for 500 and 600 nm light. The ordinate gives the net pupillary constriction measured as the change in diameter in mm. The abscissa gives the corneal irradiance of a 60° , 500 nm flash in $\log \text{quanta sec}^{-1} \text{mm}^{-2}$. Open symbols are responses to 500 nm wavelength flashes, closed symbols to 600 nm wavelength flashes. Points for the 600 nm flash have been displaced to the left by 1.38 log units, the difference required for scotopic equivalence. Symbols are used as follows: circles, a control rat; squares, a 31-day old RCS rat; triangles, a 298-day old RCS rat. The dashed line intercepts each function at threshold, defined as an 0.2 mm constriction. The smooth curves were fit by eye and have no theoretical significance.

Fig. 7. Spectral sensitivity of the pupillary light reflex for a 205-day old albino control rat measured in a ganzfeld. Thresholds at each wavelength were determined from observations of the light reflex on the video monitor. Each point corresponds to the relative intensity of the stimulus required to produce an 0.2 mm pupillary constriction. The ordinate gives the log relative sensitivity of the light reflex with the value for 500 nm light arbitrarily set equal to zero. The abscissa gives the wavelength of the stimulus. The dashed curve is a Dartnall nomogram curve with a peak of 500 nm.

intensities the action spectrum of the pupillary light reflex appears to be that of rhodopsin.

The separation in the upper branches of the curves corresponds to a gain of 0.7 log unit for the 600 nm stimulus at high intensities.

This difference is consistent with a nomogram curve for a pigment with a peak absorption near 520 nm, and thus the data suggest that at higher intensities the light reflex is driven by a mechanism whose action spectrum has a peak wavelength near 520 nm.

Spectral sensitivity of the pupillary light reflex

The dark adapted pupillary spectral sensitivity of a control rat measured in the ganzfeld at 205 days of age is plotted in Fig. 7. A Dartnall nomogram curve with a peak of 500 nm was eye-fit by vertical sliding. In no case was the pupillary spectral sensitivity of any of the control animals fit better by a nomogram curve of a significantly different peak wavelength.

In Fig. 8, the upper set of points is the mean spectral sensitivity derived from two separate sessions in the ganzfeld for a single RCS rat at 35 and 37 days of age. The agreement with the 500 nm nomogram suggests that, as for the control animals, rhodopsin-based mechanisms mediated threshold responses in this RCS rat. The middle set of points, determined under the same conditions for a 135-day old RCS rat still agree well with the 500 nm nomogram curves, but the

Fig. 8. Spectral sensitivity of the pupillary light reflex of RCS rats measured in a ganzfeld. (Conventions are as in Fig. 6.) Symbols are used as follows: circles, mean thresholds for an RCS rat from measurements made at 35 and 37 days of age; triangles, points determined from threshold intercepts of intensity response functions for a 135-day old RCS rat; squares, points determined from threshold intercepts of intensity response functions for a 363-day old RCS rat. The dashed curve is a 500 nm Dartnall nomogram; the solid curve is a 520 nm Dartnall nomogram curve.

lowest set of points, for a 363-day old RCS rat, are fit better by a nomogram curve with its peak at 520 nm, as shown by the solid line.

Dark adaptation

The time-course of dark adaptation in a 78-day old albino control rat was measured after exposure to a white ganzfeld which bleached an estimated 10% of the photopigment in the retina. At the top of Fig. 9, thresholds measured with 500 msec ganzfeld flashes of lights of wavelengths 500 and 600 nm are plotted as a function of time after the bleach. The data points for the 600 nm light have been displaced vertically by 1.38 log units to equate them to those for the 500 nm light according to a Dartnall nomogram curve which peaks at 500 nm. For times greater than about 30 min after the bleach, the data points for 500 nm light fall close to those for 600 nm light, which is consistent with rods determining threshold at these times. There is a clear and consistent separation of the 500 nm data points from those for 600 nm for earlier times, suggesting that thresholds soon after the bleach are determined by a photopic mechanism.

The same kind of dark adaptation experiments were conducted for a 157-day old RCS rat and a 297-day old RCS rat, the results of which are also

Fig. 9. Dark adaptation in a 78-day old control rat and two RCS rats of ages 157 and 297 days. Thresholds for a flash (500 msec) of 500 nm light (open circles) or 600 nm light (closed circles) in the ganzfeld following the extinction of a 30 sec exposure to the unfiltered output of the xenon arc lamp which bleached approximately 10% of the visual pigment in the retina. The ordinate gives the corneal irradiance at threshold in $\log \text{quanta sec}^{-1} \text{mm}^{-2}$. The abscissa gives the time after extinction of the bleaching light in minutes. The data for the 600 nm test flashes have been displaced by 1.38 log units to equate them to the 500 nm data points according to a Dartnall nomogram curve peaking at 500 nm.

shown in Fig. 9. As with the albino control, thresholds appear to be photopically determined early in dark adaptation, while later they appear to be scotopically determined. For these RCS animals, thresholds are generally higher than those for the control at comparable times after the bleach. Also, the change from photopic to scotopic determination of thresholds occurs later, near 40 min after the bleach.

We measured dark adaptation in three other RCS rats at ages of 76, 85, 122 and 297 days and in three other albino controls at ages of 66, 114, 196, 279 and 433 days. In every case we found clear evidence for scotopic as well as photopic function in dark adaptation. The RCS animals consistently showed higher thresholds and a change from photopic to scotopic determination of thresholds which occurred later than in the controls. Thus, all our data is consistent with the examples we show.

DISCUSSION

Pupillary sensitivity with age

The intensity-response functions in Figs 5 and 6 show that there is an overall loss of sensitivity and a change in the relative sensitivity of the low intensity and high intensity components of the light reflex of the pupil with age in the RCS rat. As the retina degenerates, progressively more light is required to produce a given constriction.

We find that the increase in RCS thresholds (Fig. 4) can be modeled as an exponential loss of sensitivity. The RCS thresholds rise steadily after about 30 days postnatal, where they approximate control thresholds. Using a value of corneal irradiance of $7.5 \log \text{quanta sec}^{-1} \text{mm}^{-2}$ as the asymptotic value, k_1 , we fit the RCS thresholds of Fig. 4 to the model

$$\log I(t) = k_1 - k_2 e^{-(t/\tau)}$$

from which equations (1) and (2) were derived (see Fig. 4). The least squares estimates were $\tau = 303.03$ days and $k_2 = 5.83 \log \text{ quanta sec}^{-1} \text{ mm}^{-2}$. This equation predicts a threshold of $2.22 \log \text{ quanta sec}^{-1} \text{ mm}^{-2}$ at 30 days postnatal which agrees closely with the control thresholds. The agreement of the curve plotted from this equation and the data of Fig. 4 is quite good, but we cannot deny that this data would also be consistent with other models. We did attempt linear fits to the data, which accounted equally well for the variability of data. However, the nature of the linear fit was unsatisfactory, for the residuals were generally negative for high ages, positive for middle ages, and negative for early ages. The same cannot be said for the albino data where exponential and linear fits were equally satisfactory.

The electroretinogram also diminishes with age in RCS rats, but it is impossible to measure beyond about 90 days postnatal (Dowling and Sidman, 1962; Perlman, 1978). Our pupillary light reflex thresholds extend the range of retinal sensitivity measurements to about 700 days and suggest prolonged persistence of functional photoreceptor elements in the RCS retina.

The decline in sensitivity of the RCS rats was probably not significantly altered by repeated exposures to the testing situation. Close inspection of Fig. 4 reveals that the first tests for several animals were performed at various advanced ages (each animal in Fig. 4 is represented by a unique symbol). The thresholds in these first tests fit well within the overall pattern of age-related sensitivity loss. There is no consistent trend towards lower thresholds in these first exposures as we might expect if the testing situation had significantly accelerated the loss of sensitivity.

Spectral sensitivity

The dark adapted spectral sensitivity of the pupillary light reflex measured in control rats is similar to that in the RCS rat at 1 month of age (Figs 7 and 8). At intermediate stages of the disease there is a relative increase in the effectiveness of long wavelength light (Fig. 8, 216-day old animal) and in later stages (Fig. 8, 363 day-old animal) pupillary sensitivity may be photopically driven. When a photopic action spectrum is obtained, it appears to be well fit by a Dartnall nomogram curve for a single visual pigment with peak absorption near 520 nm. In parallel with this finding, the peak of the pupillary spectral sensitivity curve is displaced toward longer wavelengths in RP patients with severe sensitivity loss (Alexandridis and Weddigen, 1971). In both cases the wavelength of peak sensitivity is displaced, presumably owing to the change in the composition of the pupillomotor receptor population.

The change in the spectral sensitivity of the light reflex (Figs 6 and 8) is probably the result of the changing proportion of rods to cones in the RCS retina (LaVail *et al.*, 1974, Cicerone *et al.*, 1979). Our measurements show that the pupillary light reflex

exhibits input from a scotopic mechanism even in cases of advanced degeneration (298 days of age). To our knowledge this is the first study to show the functional involvement of a rhodopsin-based mechanism at such advanced age in the RCS rat. It is likely that the pathways which mediate the light reflex can integrate activity over large retinal areas (Schweitzer, 1955; Bouma, 1965). The gain introduced by pooling over such large areas could suffice to provide a scotopic input to the pupillary light reflex even with very few rod photoreceptors remaining.

Dark adaptation

During dark adaptation in the control rats a change in the relative sensitivity of the pupillary system to 500 and 600 nm lights occurs in the functions defining threshold. This feature is consistent with photopic determination of thresholds early in dark adaptation and scotopic determination subsequently. Although it may be difficult to distinguish more than one branch in the dark adaptation curve of an RCS rat with advanced degeneration there is a change in the relative sensitivity to light of 500 and of 600 nm. Cicerone *et al.* (1979) found that the Dartnall nomogram curve which matched the spectral sensitivity of ganglion cells in older RCS rats had its peak at 520 nm. The relative sensitivity difference to lights of 500 and 600 nm predicted by the 520 nm nomogram is a factor of 0.79 log unit (Ebrey and Honig, 1977). This value is close to the relative sensitivity difference for 500 and 600 nm lights measured early in dark adaptation in both the RCS and albino rats we examined. Birch and Jacobs (1975) who used a behavioral method, and Cicerone (1976), who used the electroretinogram, have also reported a visual mechanism in the rat with peak sensitivity at 520 nm. This suggests that the same receptor pigment(s) which underlies ganglion cell response in advanced retinal degeneration also determines the photopic spectral sensitivity of the pupillary system of the rat.

In the late phase of dark adaptation in both RCS and control rats, the mean difference in sensitivity to 500 and 600 nm lights approximately matches the absorption spectrum of rat rhodopsin (Bridges, 1959) or a 500 nm Dartnall nomogram curve (Ebrey and Honig, 1977). Thus it appears that the scotopic inputs to the pupillary system in both groups of rats also depend on the same receptor pigment, rhodopsin.

The changes in the two functions describing dark adaptation and in the time to branching between these functions shown by the RCS rat data are consistent with a model of the disease in which receptor damage and subsequent sensitivity loss progress at different rates for rods and cones. In dark adaptation, the sensitivity of the mechanisms driven by rods or cones is inversely related to the vertical position of the functions which describe threshold in the dark adaptation plot. With age, both mechanisms "rise" in the plot but the scotopic mechanism rises faster. Eventually only the tail of the scotopic mechanism

may be detectable as the rest is hidden above the photopic one. As a result, the time at which the scotopic mechanism begins to determine thresholds will occur progressively later in dark adaptation. It should be noted here that the qualitative changes in the shape of the pupillary dark adaptation function which correlate with the severity of visual sensitivity loss in RP patients reported by Alexandridis and Weddigen (1971) and by Berson *et al.* (1969) are similar to the pattern of age related changes we see in this study for RCS rat dark adaptation.

Time-course of response

At suprathreshold levels of stimulation we noticed an anomaly in the time-course of pupillary constriction in some RCS rats over 5 months of age (Fig. 3). Specifically, constriction occurred in two distinct phases separated by an inflection. The time required for the pupil to redilate appeared to increase with age in RCS rats. Such responses have not been observed in unaffected controls and have not been described in any of the literature known to us. Near threshold, the responses of RCS rats and controls are similar, though the RCS rat responses appear to be slightly slower. Since we measured only the peak constriction, this should not affect any of the results which depend on threshold data. For the intensity-response functions, the first peak was chosen as the response value whenever multiple peaks occurred in the response. These time-course differences may prove important in characterizing the temporal properties of the visual system in RCS rats.

Acknowledgements—This research was supported by grants EY02055 and EY03015 to C.M.C. We thank Eureka Harami for help with the experiments. We thank two anonymous reviewers for helpful comments and suggestions.

REFERENCES

- Alexandridis E. and Weddigen A. (1971) Pupillenlichtreflexe bei Heredodegeneratio pigmentosa retinae. *Albrecht v. Graefes Arch. klin. exp. Ophthalm.* **182**, 250–260.
- Berson E. L., Gouras P., Gunkel R. D. and Myrianthopoulos N. C. (1969) Rod and cone responses in sex-linked retinitis pigmentosa. *Archs Ophthalm.* **81**, 215–225.
- Birch D. and Jacob G. (1975) Behavioral measurements of rat spectral sensitivity. *Vision Res.* **15**, 687–691.
- Block M. T. (1969) A note on the refraction and image formation of the rat's eye. *Vision Res.* **9**, 705–711.
- Bouma H. (1965) Receptive systems mediating certain light reactions of the pupil of the human eye. *Philips Res. Repts., Suppl.* No. 5.
- Bourne M. C., Campbell D. A. and Tansley L. (1938) Hereditary degeneration of the rat retina. *Br. J. Ophthalm.* **22**, 613–623.
- Bridges C. D. B. (1959) Visual pigments of some common laboratory mammals. *Nature* **184**, 1727–1728.
- Cicerone C. M. (1976) Cones survive rods in the light-damaged eye of the albino rat. *Science* **194**, 1183–1185.
- Cicerone C. M., Green D. G. and Fisher L. J. (1979) Cone inputs to ganglion cells in hereditary retinal degeneration. *Science* **203**, 1113–1115.
- Cone R. A. (1963) Quantum relations of the rat electroretinogram. *J. Physiol., Lond.* **198**, 17–38.
- Dowling J. E. (1963) Neural and photochemical mechanisms of visual adaptation in the rat. *J. gen. Physiol.* **46**, 1287.
- Dowling J. E. and Sidman R. L. (1962) Inherited retinal dystrophy in the rat. *J. cell. Biol.* **14**, 73–109.
- Ebrey T. G. and Honig B. (1977) New wavelength dependent visual pigment nomograms. *Vision Res.* **17**, 147–151.
- Krill A. E. (1972) Retinitis pigmentosa: a review. *The Sight Saving Review*.
- LaVail M. M., Sidman M., Rausin R. and Sidman R. L. (1974) Discrimination of light intensity by rats with inherited retinal degeneration: a behavioral and cytological study. *Vision Res.* **14**, 693–702.
- Lowenstein O. and Loewenfeld I. E. (1969) The pupil. In *The Eye* (Edited by Davson H.), Vol. 3. Academic Press, New York.
- Lucas D. R., Attfield M. and Davey J. B. (1955) Retinal dystrophy in the rat. *J. Pathol. Bacteriol.* **70**, 469–474.
- Noell W. K. (1977) Cortical representation of retinal degeneration. In *Retinitis pigmentosa: Clinical Implications of Current Research* (Edited by Landers III M. B., Wolbarsht M. L., Dowling J. E. and Latties A.), Plenum Press, New York.
- Perlman I. (1978) Dark adaptation in abnormal (RCS) rats studied electroretinographically. *J. Physiol.* **278**, 161–175.
- Schweitzer N. M. J. (1955) Threshold measurements on the light reflex of the pupil in the dark adapted eye. *Documenta ophthalm.* **10**, 1–78.
- Trejo L. J. and Cicerone C. M. (1978) Pupillary light reflex shows shift from scotopic to photopic spectral sensitivity with age in rats (RCS) with inherited retinal dystrophy. *Invest. Ophthalm. Visual Sci., Suppl.* 114.
- Young R. W. (1977) Visual cell renewal and the problem of retinitis pigmentosa. In *Retinitis Pigmentosa: Clinical Implications of Current Research* (Vol. 77 in *Advances in Experimental Medicine and Biology*). (Edited by Landers III M. B., Wolbarsht M. L., Dowling J. E. and Latties A.), Plenum Press, New York.