

UCSF

UC San Francisco Previously Published Works

Title

Treatment Efficacy of MEDI3902 in Pseudomonas aeruginosa Bloodstream Infection and Acute Pneumonia Rabbit Models

Permalink

<https://escholarship.org/uc/item/5bs3h3mv>

Journal

Antimicrobial Agents and Chemotherapy, 63(8)

ISSN

0066-4804

Authors

Le, Hoan N
Tran, Vuvi G
Vu, Trang TT
[et al.](#)

Publication Date

2019-08-01

DOI

10.1128/aac.00710-19

Peer reviewed



Treatment Efficacy of MEDI3902 in *Pseudomonas aeruginosa* Bloodstream Infection and Acute Pneumonia Rabbit Models

Hoan N. Le,^a Vuvi G. Tran,^a Trang T. T. Vu,^a Emmanuelle Gras,^{a,b} Vien T. M. Le,^a Marcos Gabriel Pinheiro,^{a,c} Fábio Aguiar-Alves,^{a,c} Erika Schneider-Smith,^a Henry Clay Carter,^a Bret R. Sellman,^d C. Kendall Stover,^d Antonio DiGiandomenico,^d Binh An Diep^a

^aDivision of HIV, Infectious Diseases, and Global Medicine, Department of Medicine, University of California, San Francisco, San Francisco, California, USA

^bFrançois Rabelais University, Tours, France

^cPathology Program, Fluminense Federal University, Rio de Janeiro, Brazil

^dMicrobial Sciences, AstraZeneca, Gaithersburg, Maryland, USA

ABSTRACT *Pseudomonas aeruginosa* is a challenge for clinicians due to increasing drug resistance and dwindling treatment options. We report on the activity of MEDI3902, an antibody targeting type 3 secretion protein PcrV and Psl exopolysaccharide, in rabbit bloodstream and lung infection models. MEDI3902 prophylaxis or treatment was protective in both acute models and exhibited enhanced activity when combined with a sub-therapeutic dose of meropenem. These findings further support MEDI3902 for the prevention or treatment of serious *P. aeruginosa* infections.

KEYWORDS *Pseudomonas aeruginosa*, bloodstream infections, immunotherapy, pneumonia

Pseudomonas aeruginosa is the second most common cause of nosocomial pneumonia, health care-associated pneumonia, and ventilator-associated pneumonia and the seventh most common cause of nosocomial bloodstream infections (1). These infections are associated with significant mortality and morbidity, increased intensive care unit and hospital length of stay, and substantial economic burden (2). Extensively drug-resistant (XDR) and pandrug-resistant (PDR) *P. aeruginosa* infections have been reported on multiple continents and are associated with higher mortality rates (3–6). Therefore, novel therapeutics targeting this problematic pathogen are urgently needed to address this significant unmet medical need.

MEDI3902 is a bispecific monoclonal antibody (MAb) targeting the type 3 secretion (T3S) protein PcrV and the Psl exopolysaccharide and is currently under clinical evaluation for the prevention of pneumonia in ventilated patients (Effort to Prevent Nosocomial Pneumonia Caused by *Pseudomonas aeruginosa* in Mechanically Ventilated Subjects [EVADE] study, ClinicalTrials.gov NCT02696902). Both virulence determinants have been implicated in immune evasion (7–9) MEDI3902 mediates three distinct mechanisms of action, as follows: (i) binding to PcrV prevents T3S injectisome-mediated cytotoxicity and damage to host cells (10), (ii) binding to Psl promotes opsonophagocytic killing of *P. aeruginosa* by host effector cells, and (iii) binding to Psl inhibits the bacterium's ability to attach to host epithelial cells (11, 12). Mouse models have been used extensively to demonstrate preclinical protective efficacy, although it should be noted that this animal species is known to be extremely resistant to the effects of Gram-negative lipopolysaccharide (LPS) (13). In contrast to rodents and monkeys, rabbits are exquisitely similar to humans and chimpanzees in their susceptibilities to LPS (13).

Therefore, we evaluated MEDI3902 efficacy in a rabbit model of bloodstream infection. Rabbits were randomized for intravenous administration with (i) 15 mg/kg of

Citation Le HN, Tran VG, Vu TTT, Gras E, Le VTM, Pinheiro MG, Aguiar-Alves F, Schneider-Smith E, Carter HC, Sellman BR, Stover CK, DiGiandomenico A, Diep BA. 2019. Treatment efficacy of MEDI3902 in *Pseudomonas aeruginosa* bloodstream infection and acute pneumonia rabbit models. *Antimicrob Agents Chemother* 63:e00710-19. <https://doi.org/10.1128/AAC.00710-19>.

Copyright © 2019 Le et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Antonio DiGiandomenico, DiGiandomenicoA@MedImmune.com, or Binh An Diep, binh.diep@ucsf.edu. H.N.L. and V.G.T. contributed equally to this work.

Received 3 April 2019

Returned for modification 3 May 2019

Accepted 19 May 2019

Accepted manuscript posted online 3 June 2019

Published 25 July 2019




FIG 1 MEDI3902 prophylaxis and treatment improves survival outcome in a rabbit bloodstream infection model. (A to D) Comparisons of Kaplan-Meier survival curves (A), \log_{10} CFU/lung (B), \log_{10} CFU/spleen (C), and \log_{10} CFU/kidneys (D) for rabbits administered intravenously with 15 mg/kg MEDI3902 at 24 h before infection (hbi) ($n = 6$), 15 mg/kg MEDI3902 at 1 h postinfection ($n = 6$), and 15 mg/kg control IgG at 24 h before infection ($n = 6$) with strain 6077. A one-sided log rank (Mantel-Cox) test was used to evaluate survival of control IgG- versus MEDI390-treated animals. Bacterial densities for animals pretreated with 15 mg/kg control IgG were compared to those pretreated or treated with 15 mg/kg MEDI3902 by nonparametric one-way analysis of variance (ANOVA) with Kruskal-Wallis test, followed by Dunn's multiple-comparison test. Filled symbols represent data from dead animals, and open symbols represent data from surviving animals that were euthanized at 96 h postinfection.

body weight control IgG at 24 h before infection, (ii) 15 mg/kg MEDI3902 at 24 h before infection, and (iii) 15 mg/kg MEDI3902 at 1 h postinfection and then challenged by intravenous injection of the cytotoxic *P. aeruginosa* strain 6077 contained within 1.5 ml lactated Ringer's solution at 4.4e8 CFU/ml. Rabbits were monitored every 2 h for the first 36 h postinfection and then at least three times daily thereafter. All survivors were euthanized at 96 h postinfection. Overall survival rates were 0% (0/6) for rabbits administered control IgG, 100% (6/6) for animals receiving MEDI3902 24 h before infection ($P < 0.001$ versus control IgG group by log rank test), and 67% (4/6) for those administered MEDI3902 at 1 h postinfection ($P < 0.001$ versus control IgG by log rank test) (Fig. 1A). Bacterial counts from homogenized organs were significantly reduced in vital organs of rabbits administered with MEDI3902 at 24 h before infection (Kruskal-Wallis test with Dunn's multiple-comparison test for each MEDI3902 dosage versus control IgG), but some loss in protection was observed when delivered at 1 h postinfection (Fig. 1B to D). The lower limit of bacterial detection in our organ burden model was 1.72 log CFU/organ.

For treatment studies in the rabbit model of pneumonia, rabbits were randomized for intravenous treatment with MEDI3902 at 1, 5, and 15 mg/kg or control IgG (15 mg/kg) at 1 h postinfection. To induce pneumonia, a 1.5-ml inoculum containing 9e7 CFU/ml of strain 6077 was delivered directly into the lungs of anesthetized rabbits through a pediatric endotracheal tube (outside diameter of 2.5 mm) and then immediately removed after instillation of bacterial inoculum. Rabbits were monitored every 2 h for the first 36 h after infection and three times daily thereafter. Rabbits that became severely ill as defined by signs of pulmonary dysfunction (respiration rate of >75 breaths/minute, cyanosis, and cough) were euthanized for humane reasons and scored as nonsurvivors. All survivors were euthanized at 96 h postinfection. Overall survival




FIG 2 Treatment with MEDI3902 improves survival outcomes in a dose-dependent manner in a rabbit acute pneumonia model. (A to E) Comparisons of Kaplan-Meier survival curves (A), lung weight-to-body weight (LW/BW $\times 10^3$) ratio (B), \log_{10} CFU/lung (C), \log_{10} CFU/spleen (D), and \log_{10} CFU/kidneys (E) for rabbits intravenously administered 15 mg/kg MEDI3902 ($n = 6$), 5 mg/kg MEDI3902 ($n = 6$), 1 mg/kg MEDI3902 ($n = 6$), or 15 mg/kg control IgG ($n = 6$) at 1 h postinfection (hpi). A one-sided log rank (Mantel-Cox) test was used to evaluate survival. LW/BW and bacterial densities for animals treated at 1 h postinfection with control IgG were compared to those treated with different concentrations of MEDI3902 by nonparametric one-way ANOVA with Kruskal-Wallis test, followed by Dunn’s multiple-comparison test. Filled symbols represent data from dead animals, and open symbols represent data from surviving animals that were euthanized at 96 h postinfection.

rates were 0% (0/6) for rabbits treated with control IgG, compared to 100% (6/6) for those treated with MEDI3902 at 15 or 5 mg/kg and 66% (4/6) for those treated with 1 mg/kg ($P < 0.001$ for treatment with various doses of MEDI3902 compared to control IgG group by log rank test; Fig. 2A). The lung weight-to-body weight (LW/BW) ratio, a quantitative measurement of pulmonary edema and a surrogate marker of the severity of acute lung injury, was significantly reduced for all rabbits treated with MEDI3902 compared to control IgG-treated rabbits (Kruskal-Wallis test with Dunn’s multiple-comparison test for each MEDI3902 dosage versus control IgG; Fig. 2B). Bacterial counts in the lungs, spleen, and kidneys were significantly reduced in rabbits treated with MEDI3902 at 15 mg/kg and at 5 mg/kg compared to those in control IgG-treated rabbits (Kruskal-Wallis test with Dunn’s multiple-comparison test for each MEDI3902 dosage versus control IgG) (Fig. 2C to E).

We next sought to evaluate the activity of MEDI3902 used adjunctively with meropenem in the rabbit model of pneumonia. For these studies, subprotective dosages of meropenem to simulate drug resistance and MEDI3902 were identified. Intravenous (i.v.) treatment with meropenem starting at 3 h postinfection with 75 mg/kg and 10 mg/kg meropenem every 4 hours (q4h) protected virtually all treated rabbits against death in the rabbit acute pneumonia model (data not shown), whereas 1 mg/kg of meropenem i.v. q4h was nonprotective. We also found in preliminary studies that intravenous treatment of 15 mg/kg MEDI3902 at 3 h postinfection was also nonprotective. We next sought to determine whether adjunctive therapy with MEDI3902 (15 mg/kg) and meropenem (1 mg/kg q4h) conferred protection against acute pneumonia. Compared to control IgG-treated rabbits that all rapidly succumbed to infection, intravenous treatment with 15 mg/kg MEDI3902 or 1 mg/kg meropenem q4h alone significantly delayed the time to death, but ultimately, the vast majority of these animals died of profound acute respiratory failure, as evidenced from the LW/BW ($\times 10^3$) ratio of > 10 (survival analysis performed by log rank test; LW/BW analysis performed by Kruskal-




FIG 3 Adjunctive therapy of MEDI3902 with a subtherapeutic dose of meropenem in a rabbit acute pneumonia model. (A to E) Comparisons of Kaplan-Meier survival curves (A), lung weight-to-body weight (LW/BW $\times 10^3$) ratio (B), \log_{10} CFU/lung (C), \log_{10} CFU/spleen (D), and \log_{10} CFU/kidneys (E) for rabbits administered intravenously with (i) 15 mg/kg MEDI3902 ($n = 7$), (ii) 1 mg/kg meropenem i.v. q4h ($n = 8$), (iii) a 15 mg/kg MEDI3902 plus 1 mg/kg meropenem i.v. q4h ($n = 8$) combination, or (iv) 15 mg/kg of control IgG ($n = 8$) at 3 h postinfection with strain 6077. A one-sided log rank (Mantel-Cox) test was used to evaluate survival. LW/BW and bacterial densities for animals pretreated with 15 mg/kg control IgG were compared to those treated with MEDI3902 alone, meropenem alone, or the combination by nonparametric one-way ANOVA with Kruskal-Wallis test, followed by Dunn's multiple-comparison test. Filled symbols represent data from dead animals, and open symbols represent data from surviving animals that were euthanized at 96 h postinfection.

Wallis test with Dunn's multiple-comparison test for each MEDI3902 dosage versus control IgG (Fig. 3A and B). In contrast, the treatment combination of 15 mg/kg MEDI3902 and 1 mg/kg meropenem q4h significantly protected against death (log rank test) (Fig. 3A) by protecting against acute lung injury (Fig. 3B) and reducing bacterial counts in the lungs, spleen, and kidneys (Kruskal-Wallis test with Dunn's multiple-comparison test for each MEDI3902 dosage versus control IgG) (Fig. 3C to E).

In conclusion, our results confirm the protective activity of MEDI3902 used preclinically in prophylaxis and treatment regimens, as well as when used adjunctively with antibiotic.

ACKNOWLEDGMENTS

Bret R. Sellman, C. Kendall Stover, and Antonio DiGiandomenico are employed by AstraZeneca.

Patent 2014/0072,577 describing MEDI3902* has been filed by AstraZeneca.

This work was funded by a grant from AstraZeneca.

REFERENCES

1. Gaynes R, Edwards JR, National Nosocomial Infections Surveillance System. 2005. Overview of nosocomial infections caused by Gram-negative bacilli. *Clin Infect Dis* 41:848–854. <https://doi.org/10.1086/432803>.
2. Kyaw MH, Kern DM, Zhou S, Tunceli O, Jafri HS, Falloon J. 2015. Healthcare utilization and costs associated with *S. aureus* and *P. aeruginosa* pneumonia in the intensive care unit: a retrospective observational cohort study in a US claims database. *BMC Health Serv Res* 15:241. <https://doi.org/10.1186/s12913-015-0917-x>.
3. Bodro M, Sabé N, Tubau F, Lladó L, Baliellas C, González-Costello J, Cruzado JM, Carratalà J. 2015. Extensively drug-resistant *Pseudomonas aeruginosa* bacteremia in solid organ transplant recipients. *Transplantation* 99:616–622. <https://doi.org/10.1097/TP.0000000000000366>.
4. Buhl M, Peter S, Willmann M. 2015. Prevalence and risk factors associated with colonization and infection of extensively drug-resistant *Pseudomonas aeruginosa*: a systematic review. *Expert Rev Anti Infect Ther* 13: 1159–1170. <https://doi.org/10.1586/14787210.2015.1064310>.
5. Fernandes M, Vira D, Medikonda R, Kumar N. 2016. Extensively and pan-drug resistant *Pseudomonas aeruginosa* keratitis: clinical features, risk factors, and outcome. *Graefes Arch Clin Exp Ophthalmol* 254: 315–322. <https://doi.org/10.1007/s00417-015-3208-7>.
6. Mayr A, Hinterberger G, Lorenz IH, Kreidl P, Mutschlechner W, Lass-Flörl C. 2017. Nosocomial outbreak of extensively drug-resistant *Pseudomonas aeruginosa* associated with aromatherapy. *Am J Infect Control* 45: 453–455. <https://doi.org/10.1016/j.ajic.2016.11.002>.

7. Mishra M, Byrd MS, Sergeant S, Azad AK, Parsek MR, McPhail L, Schlesinger LS, Wozniak DJ. 2012. *Pseudomonas aeruginosa* Psl polysaccharide reduces neutrophil phagocytosis and the oxidative response by limiting complement-mediated opsonization. *Cell Microbiol* 14:95–106. <https://doi.org/10.1111/j.1462-5822.2011.01704.x>.
8. Rangel SM, Diaz MH, Knoten CA, Zhang A, Hauser AR. 2015. The Role of ExoS in dissemination of *Pseudomonas aeruginosa* during pneumonia. *PLoS Pathog* 11:e1004945. <https://doi.org/10.1371/journal.ppat.1004945>.
9. Thanabalasuriar A, Sureward BG, Willson ME, Neupane AS, Stover CK, Warren P, Wilson G, Keller AE, Sellman BR, DiGiandomenico A, Kubes P. 2017. Bispecific antibody targets multiple *Pseudomonas aeruginosa* evasion mechanisms in the lung vasculature. *J Clin Invest* 127: 2249–2261. <https://doi.org/10.1172/JCI89652>.
10. Warren P, Varkey R, Bonnell JC, DiGiandomenico A, Camara M, Cook K, Peng L, Zha J, Chowdury P, Sellman B, Stover CK. 2014. A novel anti-PcrV antibody providing enhanced protection against *Pseudomonas aeruginosa* in multiple animal infection models. *Antimicrob Agents Chemother* 58:4384–4391. <https://doi.org/10.1128/AAC.02643-14>.
11. DiGiandomenico A, Warren P, Hamilton M, Guillard S, Ravn P, Minter R, Camara MM, Venkatraman V, Macgill RS, Lin J, Wang Q, Keller AE, Bonnell JC, Tomich M, Jermutus L, McCarthy MP, Melnick DA, Suzich JA, Stover CK. 2012. Identification of broadly protective human antibodies to *Pseudomonas aeruginosa* exopolysaccharide Psl by phenotypic screening. *J Exp Med* 209:1273–1287. <https://doi.org/10.1084/jem.20120033>.
12. DiGiandomenico A, Keller AE, Gao C, Rainey GJ, Warren P, Camara MM, Bonnell J, Fleming R, Bezabeh B, Dimasi N, Sellman BR, Hilliard J, Guenther CM, Datta V, Zhao W, Gao C, Yu XQ, Suzich JA, Stover CK. 2014. A multifunctional bispecific antibody protects against *Pseudomonas aeruginosa*. *Sci Transl Med* 6:262ra155. <https://doi.org/10.1126/scitranslmed.3009655>.
13. Warren HS, Fitting C, Hoff E, Adib-Conquy M, Beasley-Topliffe L, Tesini B, Liang X, Valentine C, Hellman J, Hayden D, Cavaillon JM. 2010. Resilience to bacterial infection: difference between species could be due to proteins in serum. *J Infect Dis* 201:223–232. <https://doi.org/10.1086/649557>.