

# UC Davis

## UC Davis Previously Published Works

### Title

Lifestyle and Clinical Correlates of Hepatocellular Carcinoma in South Texas: A Matched Case-control Study.

### Permalink

<https://escholarship.org/uc/item/5ck1b0b3>

### Journal

Clinical gastroenterology and hepatology : the official clinical practice journal of the American Gastroenterological Association, 15(8)

### ISSN

1542-3565

### Authors

Ramirez, Amelie G  
Muñoz, Edgar  
Parma, Dorothy Long  
et al.

### Publication Date

2017-08-01

### DOI


10.1016/j.cgh.2017.03.022

Peer reviewed

# Accepted Manuscript

Lifestyle and clinical correlates of hepatocellular carcinoma in South Texas: a matched case-control study

Amelie G. Ramirez, Edgar Muñoz, Dorothy Long Parma, Joel E. Michalek, Alan E.C. Holden, Timothy D. Phillips, Bradley H. Pollock



PII: S1542-3565(17)30322-1  
DOI: [10.1016/j.cgh.2017.03.022](https://doi.org/10.1016/j.cgh.2017.03.022)  
Reference: YJCGH 55166

To appear in: *Clinical Gastroenterology and Hepatology*  
Accepted Date: 18 March 2017

Please cite this article as: Ramirez AG, Muñoz E, Parma DL, Michalek JE, Holden AEC, Phillips TD, Pollock BH, Lifestyle and clinical correlates of hepatocellular carcinoma in South Texas: a matched case-control study, *Clinical Gastroenterology and Hepatology* (2017), doi: 10.1016/j.cgh.2017.03.022.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 *Title (max 120 char) -- Lifestyle and clinical correlates of hepatocellular carcinoma in South*  
2 **Texas: a matched case-control study**

3  
4 *Short title (max 45 char) -- Liver cancer in Texas: a case-control study*

5  
6 *Authors – Amelie G. Ramirez,<sup>1,2</sup> Edgar Muñoz,<sup>1</sup> Dorothy Long Parma,<sup>1</sup> Joel E Michalek,<sup>2</sup> Alan*  
7 *E.C. Holden,<sup>1</sup> Timothy D. Phillips,<sup>3</sup> Bradley H. Pollock<sup>4</sup>*

8  
9 <sup>1</sup>*Institute for Health Promotion Research, University of Texas Health Science Center at San*  
10 *Antonio, 7411 John Smith Drive, Suite 1000, San Antonio, TX 78229*

11  
12 <sup>2</sup>*Dept. of Epidemiology and Biostatistics, University of Texas Health Science Center at San*  
13 *Antonio, 7703 Floyd Curl Drive, Mail Code 7933, San Antonio, TX 78229*

14  
15 <sup>3</sup>*Dept. of Veterinary Integrative Biosciences, Texas A&M University, 4458 TAMU, College*  
16 *Station, TX 77843*

17  
18 <sup>4</sup>*Dept. of Public Health Sciences, School of Medicine, The University of California, Davis, One*  
19 *Shields Avenue, Med-Sci 1C, Davis, CA 95616*

20  
21 *Grant Support -- This project was supported by the Cancer Prevention Research Institute of*  
22 *Texas (CPRIT) RP120462; the Institute for Health Promotion Research (IHPR) U01 CA86117*  
23 *and U54 CA153511; and the Cancer Therapy and Research Center (CTRC) at The University of*  
24 *Texas Health Science Center at San Antonio, an NCI-designated Cancer Center (P30*  
25 *CA054174).*

26  
27 *Correspondence -- Amelie G. Ramirez*  
28 *Institute for Health Promotion Research, Dept. of Epidemiology and Biostatistics,*  
29 *University of Texas Health Science Center at San Antonio*  
30 *7411 John Smith Drive, Suite 1000, San Antonio, TX 78229, U.S.A.*  
31 *[ramirezag@uthscsa.edu](mailto:ramirezag@uthscsa.edu)*  
32 *(210) 562-6500; FAX (210) 562-6545*

33  
34 *Disclosures -- The authors declare that there are no financial or personal conflicts of interest.*

35  
36 *Author Contributions – AGR and BHP contributed to study concept and design; obtained*  
37 *funding; study supervision; and critical revision of the manuscript for important intellectual*  
38 *content. EM contributed to study concept and design; data acquisition, analysis and*  
39 *interpretation; statistical analysis; drafting and critical revision of the manuscript. DLP and TDP*  
40 *contributed to study concept and design; study supervision; data acquisition, analysis and*  
41 *interpretation; drafting and critical revision of the manuscript. JEM and AECH contributed to*  
42 *study concept and design; statistical analysis; and drafting and critical revision of the manuscript.*

43  
44 *Acknowledgments – The authors thank Amanda Sintes for her assistance with interviews of*  
45 *participants, and Allen Trevino, Holiday Harris, and Norma Ketchum for their assistance with*  
46 *data management and their comments on an earlier version of the manuscript.*

1 The incidence of hepatocellular carcinoma (HCC) in the United States is rising even as overall  
2 cancer rates are declining, and Latinos are disproportionately affected, especially in Texas. Our  
3 case-control study sought to determine relative etiologic contributions of lifestyle-related risk  
4 factors in South Texas.

5

## 6 **Methods**

7 Between October 2012 and July 2014, we identified, consented and interviewed in person 51  
8 HCC adult cases (diagnosed within the past 12 months and residing in Bexar or any of the 7  
9 surrounding counties) from clinics at University Hospital and the Cancer Therapy and Research  
10 Center at UT Health San Antonio; and 104 adult controls, 32 from a related study<sup>1,2</sup> and 72  
11 randomly selected residents in the study counties. From these 51 cases and 104 controls, we  
12 created 42 matched pairs, with exact matching on sex, ethnicity (Hispanic, Non-Hispanic), and  
13 age category (18–57 years, >57 years). Urine samples (10 mL each) and serum samples (150 µL  
14 each) were assessed for biomarkers of previous aflatoxin exposure using procedures previously  
15 described.<sup>3</sup> The primary aflatoxin exposure serum biomarker was an aflatoxin B<sub>1</sub> (AFB<sub>1</sub>)-lysine  
16 DNA adduct. We also assessed urinary aflatoxin M<sub>1</sub>. We used matched logistic regression to  
17 estimate odds ratios (OR) and 95% confidence intervals. The protocol was approved by a local  
18 Institutional Review Board.

19

## 20 **Results**

21 Most matched cases and controls were Latino (67%). Compared to controls, HCC cases were  
22 more likely to report Medicare or Medicaid, lower income, less education, more lifetime alcohol  
23 use, and more smoking (Figure 1). Cases were less likely to report hypercholesterolemia [0.11

1 (0.02–0.51)], more likely to report hepatitis C [183.74 (27.37–∞), cirrhosis 2.17 (33.3–∞), and  
2 transfusions [4.35 (1.60–11.84)], and less likely to be taking aspirin [0.31 (0.11–0.85)], statins  
3 [0.03 (0–0.20)], and omega-3/fish oil [0.10 (0.01–0.78)]. Cases did not differ significantly from  
4 controls with regard to reported consumption of products made from corn (data not shown).

5 Relative to controls, cases were more likely to have HCV antibodies [174.3 (26.2–∞)] and have  
6 detectable aflatoxin levels in blood [6.09 (1.10–33.71)] and urine [3.42 (1.07–10.91)].

## 8 **Discussion**

9 To our knowledge, this is the first epidemiologic study relating HCC and aflatoxin exposure in a  
10 U.S. population, particularly a Latino-majority study population. A recent review of non-U.S.  
11 studies<sup>4</sup> reported that aflatoxin may play a causative role in 4.6% to 28.2% of all global HCC  
12 cases, and that the combined effect of AFB<sub>1</sub> exposure and HBV infection appears additive rather  
13 than multiplicative. In the current study, serum (AFB<sub>1</sub>) and urine (AFM1) aflatoxin levels and  
14 HCV infection were significantly higher in cases than controls. Our results also support the  
15 observations of others that HCV infection is a major risk factor for HCC incidence in the U.S.  
16 Although HCV infection is a strong risk factor for development of HCC, rising trends in HCC  
17 incidence among Latinos cannot be attributed solely to HCV infection. Our data on a majority-  
18 Latino cohort suggest that HCC cases are poor and controls were more affluent, and smoked  
19 more with a higher prevalence of cirrhosis than controls. The lack of case-control differences for  
20 corn-based foods, a suspected risk factor for HCC, may be due to information bias from use of a  
21 12-month dietary recall assessment. We hypothesized that contaminated corn products  
22 contributed to the increased risk of HCC and expected to observe increased consumption of corn-  
23 based foods in cases relative to controls; a possible reason we did not may be that cases and

1 controls are consuming corn products differentially contaminated with aflatoxin (e.g.,  
2 commercial vs. home-grown). Random assays of corn products purchased by cases and controls  
3 should be done in a future study to verify this. Study strengths include case-control matching on  
4 sex, ethnicity, and age; and in-person interviewing of subjects in their language of choice. A  
5 possible weakness is that cases and controls were not randomly selected with the same inclusion  
6 criteria; therefore, selection bias may exist. Cases were selected as a convenience sample from  
7 local medical facilities, whereas controls were selected from an existing prevention study and  
8 randomly from the surrounding county populations. A larger study will be needed to address  
9 sources of aflatoxin exposure as a risk factor for HCC.

10

- 1 Figure 1
- 2 Association between lifestyle and clinical factors and hepatocellular carcinoma risk in South
- 3 Texas. Odds ratios (95% CI) derived from matched logistic regression.
- 4

ACCEPTED MANUSCRIPT

## 1 **References**

- 2 1. ClinicalTrials.gov. Phase 2 Reduction of Dietary Mycotoxin Exposure by ACCS100  
3 (RDMEACCS100). Available from URL:  
4 <https://clinicaltrials.gov/ct2/show/NCT01677195?term=NCT01677195&rank=1> [accessed 03-  
5 24-2015].
- 6 2. Pollock BH, Elmore S, Romoser A, et al. Intervention trial with calcium montmorillonite clay  
7 in a south Texas population exposed to aflatoxin. *Food Addit Contam Part A Chem Anal Control*  
8 *Expo Risk Assess* 2016;33(8): 1346-54.
- 9 3. Johnson NM, Qian G, Xu L, et al. Aflatoxin and PAH exposure biomarkers in a U.S.  
10 population with a high incidence of hepatocellular carcinoma. *Sci Total Environ* 2010;408(23):  
11 6027-31.
- 12 4. Wu HC, Santella R. The role of aflatoxins in hepatocellular carcinoma. *Hepat Mon*  
13 2012;12(10 HCC): e7238.



