

UC Irvine

UC Irvine Previously Published Works

Title

Combinatorial congruences and ψ -operators

Permalink

<https://escholarship.org/uc/item/5cv4b5q6>

Journal

Finite Fields and Their Applications, 12(4)

ISSN

1071-5797

Author

Wan, Daqing

Publication Date

2006-11-01

DOI

10.1016/j.ffa.2005.08.006

Peer reviewed

Combinatorial Congruences and ψ -Operators

Daqing Wan*

dwan@math.uci.edu

Department of Mathematics
University of California, Irvine
CA 92697-3875

December 23, 2013

Abstract

The ψ -operator for (φ, Γ) -modules plays an important role in the study of Iwasawa theory via Fontaine's big rings. In this note, we prove several sharp estimates for the ψ -operator in the cyclotomic case. These estimates immediately imply a number of sharp p -adic combinatorial congruences, one of which extends the classical congruences of Fleck (1913) and Weisman (1977).

1 Combinatorial Congruences

Let p be a prime, $n \in \mathbb{Z}_{>0}$. Throughout this paper, let $[x]$ denote the integer part of x if $x \geq 0$ and $[x] = 0$ if $x < 0$. In the author's course lectures [4] on Fontaine's theory and p -adic L-functions given at UC Irvine (spring 2005) and at the Morningside Center of Mathematics (summer 2005), the following two congruences were discovered.

Theorem 1.1. *For integers $r \in \mathbb{Z}$, $j \geq 0$, we have*

$$\sum_{k \equiv r \pmod{p}} (-1)^{n-k} \binom{n}{k} \binom{\frac{k-r}{p}}{j} \equiv 0 \pmod{p^{\lfloor \frac{n-1-jp}{p-1} \rfloor}}.$$

We shall see that the theorem comes from a simple estimate of $\psi(\pi^n)$ for the cyclotomic φ -module.

*Partially supported by NSF. The author thanks Z.W. Sun for helpful discussions on combinatorial congruences

Theorem 1.2. *For integer $j \geq 0$, we have*

$$\sum_{\substack{i_0 + \dots + i_{p-1} = n \\ i_1 + 2i_2 + \dots \equiv r \pmod{p}}} \binom{n}{i_0 i_1 \dots i_{p-1}} \binom{\frac{i_1 + 2i_2 + \dots - r}{p}}{j} \equiv 0 \pmod{p^{\lfloor \frac{n(p-1) - jp - 1}{p-1} \rfloor}}.$$

As we shall see, this theorem comes from a simple estimate of $\psi(\pi^{-n})$ for the cyclotomic φ -module. Note that when $p = 2$, Theorem 1.2 is equivalent to Theorem 1.1.

The above two congruences can be extended from p to $q = p^a$, where a is a positive integer. To do so, it suffices to estimate the a -th iterate $\psi^a(\pi^n)$. This can be done by induction. The estimate of $\psi^a(\pi^n)$ for $n > 0$ leads to

Theorem 1.3. *For integers $r \in \mathbb{Z}$, $j \geq 0$ and $a > 0$, we have*

$$\sum_{k \equiv r \pmod{p^a}} (-1)^{n-k} \binom{n}{k} \binom{\frac{k-r}{p^a}}{j} \equiv 0 \pmod{p^{\lfloor \frac{n - p^{a-1} - jp^a}{p^{a-1}(p-1)} \rfloor}}.$$

The estimate of $\psi^a(\pi^n)$ for $n < 0$ leads to

Theorem 1.4. *Let*

$$S_j(n, r, p^a) = \sum_{\substack{i_0 + \dots + i_{p^a-1} = n \\ i_1 + 2i_2 + \dots \equiv r \pmod{p^a}}} \binom{n}{i_0 \dots i_{p^a-1}} \binom{(i_1 + 2i_2 + \dots - r)/p^a}{j}.$$

Then for integer $j \geq 0$, we have

$$S_j(n, r, p^a) \equiv 0 \pmod{p^{\lfloor \frac{(an - a + 1)(p-1) - j(ap - a + 1) - 1}{p-1} \rfloor}}.$$

As Z.W. Sun informed me, the special case $j = 0$ of Theorem 1.1.1 was first proved by Fleck [1] in 1913, and the special case of Theorem 1.1.3 for $j = 0$ was first proved by Weisman [5] in 1977. A different extension of Theorem 1.1.1 and Weisman's congruence has been obtained by Z.W. Sun [2] using different combinatorial arguments. Motivated by applications in algebraic topology, Sun-Davis [3] proved yet another extension:

$$\sum_{k \equiv r \pmod{p^a}} (-1)^{n-k} \binom{n}{k} \binom{\frac{k-r}{p^a}}{j} \equiv 0 \pmod{p^{\left(\text{ord}_p([n/p^{a-1}]!) - j - \text{ord}_p(j!)\right)}}.$$

2 The operator ψ

Let p be a fixed prime. Let π be a formal variable. Let

$$A^+ = \mathbb{Z}_p[[\pi]]$$

be the formal power series ring over the ring of p -adic integers. Let A be the p -adic completion of $A^+[\frac{1}{\pi}]$, and let $B = A[\frac{1}{p}]$ be the fraction field of A . The rings A^+ , A and B correspond to $A_{\mathbb{Q}_p}^+$, $A_{\mathbb{Q}_p}$ and $B_{\mathbb{Q}_p}$ in Fontaine's theory.

We shall not discuss the Galois action on A , which is not needed for our present purpose. The Frobenius map φ acts on the above rings by

$$\varphi(\pi) = (1 + \pi)^p - 1.$$

If we let $[\varepsilon] = 1 + \pi$, then $\varphi([\varepsilon]) = [\varepsilon]^p$. The map φ is injective of degree p . This gives

Proposition 2.1. $\{1, \pi, \dots, \pi^{p-1}\}$ (and $\{1, [\varepsilon], \dots, [\varepsilon]^{p-1}\}$) is a basis of A over the subring $\varphi(A)$.

Definition 2.2. The operator $\psi : A \rightarrow A$ is defined by

$$\psi(x) = \psi \left(\sum_{i=0}^{p-1} [\varepsilon]^i \varphi(x_i) \right) = x_0 = \frac{1}{p} \varphi^{-1}(\text{Tr}_{A/\varphi(A)}(x)),$$

where $x : A \rightarrow A$ denotes the multiplication by x as $\varphi(A)$ -linear map.

Example 2.3.

$$\psi([\varepsilon]^n) = \begin{cases} [\varepsilon]^{n/p}, & \text{if } p \mid n; \\ 0, & \text{if } p \nmid n. \end{cases}$$

It is clear that ψ is φ^{-1} -linear:

$$\psi(\varphi(a)x) = a\psi(x) \quad \forall a, x \in A.$$

Example 2.4. Let a be a positive integer relatively prime to p . Then

$$\psi\left(\frac{1}{(1 + \pi)^a - 1}\right) = \frac{1}{(1 + \pi)^a - 1}.$$

In fact,

$$\begin{aligned} \psi\left(\frac{1}{[\varepsilon]^a - 1}\right) &= \psi\left(\frac{1}{[\varepsilon]^{ap} - 1} \cdot \frac{[\varepsilon]^{ap} - 1}{[\varepsilon] - 1}\right) \\ &= \frac{1}{[\varepsilon]^a - 1} \psi\left(1 + [\varepsilon]^a + \dots + [\varepsilon]^{(p-1)a}\right) \\ &= \frac{1}{[\varepsilon]^a - 1} = \frac{1}{(1 + \pi)^a - 1}. \end{aligned}$$

By p -adic continuity, the above example holds for any p -adic unit $a \in \mathbb{Z}_p^*$. In the general theory of (φ, Γ) -modules, it is important to find the fix points of ψ for applications to p -adic L-functions and Iwasawa theory. In the simplest cyclotomic case, we have the following description for the fixed points (see [4]).

Proposition 2.5.

$$A^{\psi=1} = \frac{1}{\pi} \mathbb{Z}_p \oplus \mathbb{Z}_p \oplus \left\{ \sum_{k=0}^{\infty} \varphi^k(x) \mid x \in \bigoplus_{i=1}^{p-1} [\varepsilon]^i \varphi(a_i + \pi \mathbb{Z}_p[[\pi]]), \sum_{i=1}^{p-1} a_i = 0 \right\},$$

where $a_i \in \mathbb{Z}_p$.

For example, if a is a positive integer relatively prime to p , then the element

$$\frac{a}{(1+\pi)^a - 1} - \frac{1}{\pi} \in (A^+)^{\psi=1}$$

gives the cyclotomic units and the Euler system. This element is the Amice transform of a p -adic measure which produces the p -adic zeta function of \mathbb{Q} . This type of connections is conjectured to be a general phenomenon for (φ, Γ) -modules coming from global p -adic Galois representations.

3 Sharp estimates for ψ

The ring A is a topological ring with respect to the (p, π) -topology. A basis of neighborhoods of 0 is the sets $p^k A + \pi^n A^+$, where $k \in \mathbb{Z}$ and $n \in \mathbb{N}$. The operator ψ is uniformly continuous. This continuity will give rise to combinatorial congruences.

For $s \in A^+$, one checks that

$$\begin{aligned} \psi(\pi^p s) &= \psi([\varepsilon] - 1)^p s \\ &= \psi([\varepsilon]^p - 1)s + pss_1 \\ &= \pi\psi(s) + p\psi(ss_1) \in (p, \pi)\psi(sA^+). \end{aligned}$$

In particular,

$$\psi(\pi^p A^+) \subset (p, \pi)A^+.$$

Thus, by iteration, we get

Proposition 3.1 (Weak Estimate). *Let $n \geq 0$. Then*

$$\psi(\pi^n A^+) \subset (p, \pi)^{[n/p]} A^+ = \sum_{j=0}^{[n/p]} \pi^j p^{[n/p]-j} A^+.$$

Since the exponent $[(n - jp)/p]$ is decreasing in j , this proposition implies that for $x \in \pi^n A^+$, we have

$$\psi(x) = \sum_{j=0}^{\infty} a_j \pi^j, \quad a_j \in \mathbb{Z}_p, \quad \text{ord}_p(a_j) \geq [(n - jp)/p].$$

This already gives a non-trivial combinatorial congruence. Let r be an integer. Let us calculate $\psi(\pi^n [\varepsilon]^{-r})$ in a different way.

Lemma 3.2.

$$\psi(\pi^n [\varepsilon]^{-r}) = \sum_{j \geq 0} \pi^j \sum_{k \equiv r \pmod{p}} (-1)^{n-k} \binom{n}{k} \binom{(k-r)/p}{j}.$$

Proof. Since $\pi = [\varepsilon] - 1$ and $[\varepsilon] = 1 + \pi$, we have

$$\begin{aligned} \psi(\pi^n [\varepsilon]^{-r}) &= \psi(([\varepsilon] - 1)^n [\varepsilon]^{-r}) \\ &= \psi \left(\sum_{k=0}^n (-1)^{n-k} \binom{n}{k} [\varepsilon]^{k-r} \right) \\ &= \sum_{k \equiv r \pmod{p}} (-1)^{n-k} \binom{n}{k} [\varepsilon]^{(k-r)/p} \\ &= \sum_{k \equiv r \pmod{p}} (-1)^{n-k} \binom{n}{k} \sum_{j \geq 0} \binom{(k-r)/p}{j} \pi^j \\ &= \sum_{j \geq 0} \pi^j \sum_{k \equiv r \pmod{p}} (-1)^{n-k} \binom{n}{k} \binom{(k-r)/p}{j}. \end{aligned}$$

□

Comparing the coefficients of π^j in this equation and the weak estimate, we get

Corollary 3.3 (Weak Congruence). *Let $n \geq 0$. We have*

$$\sum_{k \equiv r \pmod{p}} (-1)^{n-k} \binom{n}{k} \binom{(k-r)/p}{j} \equiv 0 \pmod{p^{[(n-jp)/p]}}.$$

The above simple estimate is crude and certainly not optimal since we ignored a factor of π . We now improve on it.

Theorem 3.4 (Sharp Estimate I). *For $n \geq 0$, we have*

$$\psi(\pi^n A^+) \subset \sum_{j=0}^{\lfloor n/p \rfloor} \pi^j p^{\lfloor \frac{n-1-jp}{p-1} \rfloor} A^+.$$

Proof. We prove the theorem by induction. The theorem is trivial if $n \leq p-1$. Write

$$\varphi(\pi) = (1 + \pi)^p - 1 = \pi^p - p\pi s_1, \quad s_1 \in A^+.$$

Then,

$$\psi(\pi^p s) = \psi((\varphi(\pi) + p\pi s_1)s) = \pi\psi(s) + p\psi(\pi s_1 s).$$

This proves that the theorem is true for $n = p$. Let $n > p$. Assume the theorem holds for $\leq n-1$. It follows that

$$\psi(\pi^n A^+) = \psi(\pi^p \pi^{n-p} A^+) \subseteq \pi\psi(\pi^{n-p} A^+) + p\psi(\pi^{n+1-p} A^+).$$

By the induction hypothesis, the right side is contained in

$$\begin{aligned} & \pi \sum_{j=0}^{\lfloor (n-p)/p \rfloor} \pi^j p^{\lfloor \frac{n-p-1-jp}{p-1} \rfloor} A^+ + p \sum_{j=0}^{\lfloor (n+1-p)/p \rfloor} \pi^j p^{\lfloor \frac{n-p-jp}{p-1} \rfloor} A^+ \\ &= \sum_{j=1}^{\lfloor n/p \rfloor} \pi^j p^{\lfloor \frac{n-1-jp}{p-1} \rfloor} A^+ + \sum_{j=0}^{\lfloor (n+1-p)/p \rfloor} \pi^j p^{\lfloor \frac{n-1-jp}{p-1} \rfloor} A^+. \end{aligned}$$

□

The function $\lfloor (n-1-jp)/(p-1) \rfloor$ is decreasing in j and vanishes for $j \geq \lfloor n/p \rfloor$. Comparing the coefficients of π^j in the lemma and the above sharp estimate, we deduce

Corollary 3.5 (Sharp Congruence I). *Let $r \in \mathbb{Z}$.*

$$\sum_{k \equiv r \pmod{p}} (-1)^{n-k} \binom{n}{k} \binom{(k-r)/p}{j} \equiv 0 \pmod{p^{\lfloor \frac{n-1-jp}{p-1} \rfloor}},$$

where $j \geq 0$ is a non-negative integer.

Theorem 3.6 (Sharp Estimate II). *For $n > 0$, we have*

$$\psi\left(\frac{1}{\pi^n}A^+\right) \subseteq \sum_{j=0}^{\lfloor n(p-1)/p \rfloor} \frac{1}{\pi^{n-j}} p^{\lfloor \frac{n(p-1)-jp-1}{p-1} \rfloor} A^+.$$

Proof. Note that

$$\varphi(\pi)/\pi = \pi^{p-1} + \binom{p}{1}\pi^{p-2} + \cdots + \binom{p}{p-1} \in (\pi^{p-1}, p),$$

so $(\varphi(\pi)/\pi)^n \in (\pi^{p-1}, p)^n$. Then

$$\begin{aligned} \psi\left(\frac{1}{\pi^n}A^+\right) &= \psi\left(\frac{1}{\varphi(\pi)^n} \left(\frac{\varphi(\pi)}{\pi}\right)^n A^+\right) \\ &= \frac{1}{\pi^n} \psi\left(\left(\frac{\varphi(\pi)}{\pi}\right)^n A^+\right) \\ &\subseteq \frac{1}{\pi^n} \sum_{i=0}^n p^{n-i} \psi(\pi^{i(p-1)} A^+). \end{aligned}$$

By Sharp Estimate I, we have

$$\psi(\pi^{i(p-1)} A^+) \subseteq \sum_{j=0}^{\lfloor i(p-1)/p \rfloor} \pi^j p^{\lfloor \frac{i(p-1)-1-jp}{p-1} \rfloor} A^+.$$

Then,

$$\begin{aligned} \psi\left(\frac{1}{\pi^n}A^+\right) &\subseteq \sum_{j=0}^{\lfloor n(p-1)/p \rfloor} \frac{1}{\pi^{n-j}} \sum_{\lfloor jp/(p-1) \rfloor \leq i \leq n} p^{n-i+\lfloor \frac{i(p-1)-jp-1}{p-1} \rfloor} A^+ \\ &\subseteq \sum_{j=0}^{\lfloor n(p-1)/p \rfloor} \frac{1}{\pi^{n-j}} p^{\lfloor \frac{n(p-1)-jp-1}{p-1} \rfloor} A^+. \end{aligned}$$

□

Corollary 3.7 (Sharp Congruence II). *Let*

$$S_j(n, r, p) = \sum_{\substack{i_0 + \cdots + i_{p-1} = n \\ i_1 + 2i_2 + \cdots \equiv r \pmod{p}}} \binom{n}{i_0 \cdots i_{p-1}} \binom{(i_1 + 2i_2 + \cdots - r)/p}{j}.$$

Then integer $j \geq 0$, we have

$$S_j(n, r, p) \equiv 0 \pmod{p^{\lfloor \frac{n(p-1)-1-jp}{p-1} \rfloor}}.$$

Proof.

$$\begin{aligned}
& \psi \left(\frac{1}{\pi^n} [\varepsilon]^{-r} \right) \\
&= \frac{1}{\pi^n} \psi \left(\left(\frac{[\varepsilon]^p - 1}{[\varepsilon] - 1} \right)^n [\varepsilon]^{-r} \right) \\
&= \frac{1}{\pi^n} \psi \left((1 + [\varepsilon] + \cdots + [\varepsilon]^{p-1})^n \cdot [\varepsilon]^{-r} \right) \\
&= \frac{1}{\pi^n} \sum_{\substack{i_0 + \cdots + i_{p-1} = n \\ i_1 + 2i_2 + \cdots \equiv r \pmod{p}}} [\varepsilon]^{(i_1 + 2i_2 + \cdots - r)/p} \binom{n}{i_0 \cdots i_{p-1}} \\
&= \frac{1}{\pi^n} \sum_{\substack{i_0 + \cdots + i_{p-1} = n \\ i_1 + 2i_2 + \cdots \equiv r \pmod{p}}} \sum_{j \geq 0} \pi^j \binom{n}{i_0 \cdots i_{p-1}} \binom{(i_1 + 2i_2 + \cdots - r)/p}{j} \\
&= \sum_{j=0}^{\infty} \frac{1}{\pi^{n-j}} S_j(n, r, p).
\end{aligned}$$

The function $[(n(p-1) - jp - 1)/(p-1)]$ is decreasing in j and vanishes for $j \geq [n(p-1)/p]$. Comparing the coefficients of $\frac{1}{\pi^{n-j}}$, the congruence follows. \square

4 Sharp estimates for ψ^a

Let a be a positive integer. In this section, we extend the sharp estimates for ψ to ψ^a .

Theorem 4.1 (Sharp Estimate I). *For $n \geq 0$, we have*

$$\psi^a(\pi^n A^+) \subseteq \sum_{j=0}^{[n/p^a]} \pi^j p^{\lfloor \frac{n-p^{a-1}-jp^a}{p^a-1(p-1)} \rfloor} A^+.$$

Proof. We prove the theorem by induction on a . The theorem is true if $a = 1$. Assume now $a \geq 2$ and assume that the theorem holds for $a - 1$.

Then,

$$\begin{aligned}
\psi^a(\pi^n A^+) &= \psi(\psi^{a-1} \pi^n A^+) \\
&\subseteq \psi\left(\sum_{i=0}^{\lfloor n/p^{a-1} \rfloor} \pi^i p^{\lfloor \frac{n-p^{a-2}-ip^{a-1}}{p^{a-2}(p-1)} \rfloor} A^+\right) \\
&\subseteq \sum_{i=0}^{\lfloor n/p^{a-1} \rfloor} \sum_{j=0}^{\lfloor i/p \rfloor} \pi^j p^{\lfloor \frac{n-p^{a-2}-ip^{a-1}}{p^{a-2}(p-1)} \rfloor + \lfloor \frac{i-1-jp}{p-1} \rfloor} A^+ \\
&\subseteq \sum_{j=0}^{\lfloor n/p^a \rfloor} \pi^j \sum_{pj \leq i \leq \lfloor n/p^{a-1} \rfloor} p^{\lfloor \frac{n-p^{a-2}-ip^{a-1}}{p^{a-2}(p-1)} \rfloor + \lfloor \frac{i-1-jp}{p-1} \rfloor} A^+.
\end{aligned}$$

The exponent of p for a fixed j is decreasing in i and hence the minimum exponent of p is attained when $i = \lfloor n/p^{a-1} \rfloor$. The minimum exponent is computed to be

$$\left\lfloor \frac{n-p^{a-2}-\lfloor n/p^{a-1} \rfloor p^{a-1}}{p^{a-1}-p^{a-2}} \right\rfloor + \left\lfloor \frac{\lfloor n/p^{a-1} \rfloor - 1 - jp}{p-1} \right\rfloor = \left\lfloor \frac{n-p^{a-1}-jp^a}{p^{a-1}(p-1)} \right\rfloor.$$

□

The proof of the lemma gives more general

Lemma 4.2.

$$\psi^a(\pi^n [\varepsilon]^{-r}) = \sum_{j \geq 0} \pi^j \sum_{k \equiv r \pmod{p^a}} (-1)^{n-k} \binom{n}{k} \binom{(k-r)/p^a}{j}.$$

Comparing the coefficients of π^j in the lemma and the sharp estimate for ψ^a , we get

Corollary 4.3 (Sharp Congruence I). *Let $r \in \mathbb{Z}$. Then*

$$\sum_{k \equiv r \pmod{p^a}} (-1)^{n-k} \binom{n}{k} \binom{(k-r)/p^a}{j} \equiv 0 \pmod{p^{\lfloor \frac{n-p^{a-1}-jp^a}{p^{a-1}(p-1)} \rfloor}},$$

where $j \geq 0$ is a non-negative integer.

Theorem 4.4 (Sharp Estimate II). *For $n > 0$ and $a > 0$, we have*

$$\psi^a\left(\frac{1}{\pi^n} A^+\right) \subseteq \sum_{j=0}^{\lfloor \frac{(an-a+1)(p-1)}{ap-a+1} \rfloor} \frac{1}{\pi^{n-j}} p^{\lfloor \frac{(an-a+1)(p-1)-j(ap-a+1)-1}{p-1} \rfloor} A^+.$$

Proof. The theorem is true for $a = 1$. Assume now that $a > 1$ and assume that the theorem is true for $a - 1$. Then

$$\begin{aligned}
\psi^a \left(\frac{1}{\pi^n} A^+ \right) &= \psi \left(\psi^{a-1} \left(\frac{1}{\pi^n} A^+ \right) \right) \\
&\subseteq \psi \left(\sum_{j=0}^{\lfloor \frac{(a-1)n-a+2)(p-1)}{(a-1)p-a+2} \rfloor} \frac{1}{\pi^{n-j}} p^{\lfloor \frac{((a-1)n-a+2)(p-1)-j((a-1)p-a+2)-1}{p-1} \rfloor} A^+ \right) \\
&\subseteq \sum_j \sum_i \frac{1}{\pi^{n-j-i}} p^{\lfloor \frac{((a-1)n-a+2)(p-1)-j((a-1)p-a+2)-1}{p-1} \rfloor + \lfloor \frac{(n-j)(p-1)-ip-1}{p-1} \rfloor} A^+,
\end{aligned}$$

where the indices i and j satisfy

$$0 \leq j \leq \lfloor \frac{(a-1)n-a+2)(p-1)}{(a-1)p-a+2} \rfloor, \quad 0 \leq i \leq \lfloor (n-j)(p-1)/p \rfloor.$$

For fixed $i + j = k$, the exponent of p is decreasing in j and the minimum value is attained when $j = k$ and $i = 0$. It follows that

$$\begin{aligned}
\psi^a \left(\frac{1}{\pi^n} A^+ \right) &\subseteq \sum_{k \geq 0} \frac{1}{\pi^{n-k}} p^{\lfloor \frac{((a-1)n-a+2)(p-1)-k((a-1)p-a+2)-1}{p-1} \rfloor + \lfloor n-k-1 \rfloor} A^+ \\
&\subseteq \sum_{k=0}^{\lfloor \frac{(an-a+1)(p-1)}{ap-a+1} \rfloor} \frac{1}{\pi^{n-k}} p^{\lfloor \frac{(an-a+1)(p-1)-k(ap-a+1)-1}{p-1} \rfloor} A^+,
\end{aligned}$$

where we stop at $k = \lfloor \frac{(an-a+1)(p-1)}{ap-a+1} \rfloor$ in the summation as the exponent of p is zero if $k \geq \lfloor \frac{(an-a+1)(p-1)}{ap-a+1} \rfloor$. \square

Corollary 4.5 (Sharp Congruence II). *Let*

$$S_j(n, r, p^a) = \sum_{\substack{i_0 + \dots + i_{p^a-1} = n \\ i_1 + 2i_2 + \dots \equiv r \pmod{p^a}}} \binom{n}{i_0 \dots i_{p^a-1}} \binom{(i_1 + 2i_2 + \dots - r)/p^a}{j}.$$

Then for integer $j \geq 0$, we have

$$S_j(n, r, p^a) \equiv 0 \pmod{p^{\lfloor \frac{(an-a+1)(p-1)-j(ap-a+1)-1}{p-1} \rfloor}}.$$

References

- [1] L.E. Dickson, History of the Theory of Numbers, Vol. I, AMS Chelsea Publ., 1999, Chapter XI, pp. 270-275.
- [2] Z.W. Sun, Polynomial extension of Fleck's congruence, preprint, 2005, arXiv:math.NT/0507008.
- [3] Z.W. Sun and D.M. Davis, A combinatorial congruence for polynomials, arXiv:math.NT/0508087.
- [4] D. Wan, Fontaine's Rings and p -Adic L-Functions, Lecture Notes at the Morningside Center of Mathematics, 2005.
- [5] C.S. Weisman, Some congruences for binomial coefficients, Michigan Math. J., 24(1977), 141-151.